



Source details

Annals of Medicine and Surgery

Scopus coverage years: from 2012 to Present

Publisher: Elsevier

E-ISSN: 2049-0801

Subject area: Medicine: Surgery

Source type: Journal

CiteScore 2021

1.4



SJR 2021

0.373



SNIP 2021

0.965



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▼

$$1.4 = \frac{2,710 \text{ Citations 2018 - 2021}}{1,918 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$1.4 = \frac{4,237 \text{ Citations to date}}{3,082 \text{ Documents to date}}$$

Last updated on 05 October, 2022 • Updated monthly

CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine		
Surgery	#266/469	43rd

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)



Limited-Time Only To Go Coco

Give your day a Boost of Coco and go Loco today with our new Coconut Smoothies. Order now!

Boost Juice Indonesia

[Learn More](#)

Annals of Medicine and Surgery

COUNTRY

Netherlands



Universities and research institutions in Netherlands

SUBJECT AREA AND CATEGORY

Medicine
Medicine (miscellaneous)
Surgery

PUBLISHER

Elsevier BV

H-INDEX

30

PUBLICATION TYPE

Journals

ISSN

20490801

COVERAGE

2012-2021

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

annalsjournal@elsevier.com

SCOPE

Annals of Medicine and Surgery is an online-only, peer-reviewed open access journal with a global outlook and focus on those training in medicine and surgery (postgraduate and undergraduate). AMS contains a mix of original clinical and basic science research, reviews, editorials, commentary, perspectives, debate, opinion, case reports and journal club reports. The journal crosses the whole disease spectrum of medicine and surgery but does also have a special focus on the following areas: -Patient safety- Human factors, teamwork, communication and professionalism- Quality improvement science and practice- Evidence based medicine- Implementation science – clinical evidence into practice- Leadership and management- Medical education, teaching, and training- Public and global health- Healthcare policy, delivery, commissioning, and resource management- Use of technology and health informatics- Clinical ethics and medical law- Research and innovation – bedside to bench and back again – including applications like personalised medicine.






Join the conversation about this journal


Quartiles




FIND SIMILAR JOURNALS

<p>1 International Journal of Surgery NLD</p> <p>80% similarity</p>	<p>2 International Journal of Surgery Open GBR</p> <p>73% similarity</p>	<p>3 International Journal of Surgery Case Reports NLD</p> <p>67% similarity</p>	<p>4 BMC Surgery GBR</p> <p>61% similarity</p>
---	--	--	--






% International Collaboration





Citable documents

Non-citable documents



Cited documents

Uncited documents



Annals of Medicine and Surgery

Q3
Medicine (miscellaneous)
best quartile

SJR 2021
0.37

powered by scimagojr.com


Show this widget in your own website

Just copy the code below and paste within your html code:

``

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2022

K **Khalid** 10 months ago

Would you inform me what is the impact factor of this journal please?

reply



Melanie Ortiz 9 months ago

Dear Khalid, thank you very much for your comment. SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR (Check it on our website). We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team

S Sultan jarrar 2 years ago

Dear SCIMAGO team,

Could you please tell me if this journal considered to be q2 or q3 ? There is a little confusion about it .

Kind regards

reply



Melanie Ortiz 2 years ago

Dear Sultan,

thank you for contacting us. For every journal, the annual value of the SJR is integrated into the distribution of SJR values of all the subject categories to which the journal belongs. There are more than 300 subject categories. The position of each journal is different in any category and depends on the performance of the category, in general, and the journal, in particular .

Best Regards, SCImago Team

A Anton 2 years ago

Dear sir/ madam, please tell if the journal status Q2 or Q3?

reply



Melanie Ortiz 2 years ago

Dear Anton, thank you for contacting us. For every journal, the annual value of the SJR is integrated into the distribution of SJR values of all the thematic categories to which the journal belongs. There are more than 300 thematic categories. The position of each journal is different in any category and depends on the performance of the category, in general, and the journal, in particular. The next SJCR's update will be made throughout June 2020. Best Regards, SCImago Team

S **Saad** 2 years ago

Dear. Melanie
Thank you very much for your reply.
Best regards.
Saad

reply



Melanie Ortiz 2 years ago

SCImago Team

Dear Saad, thanks for your participation! Best Regards, SCImago Team

S **Saad** 3 years ago

Dear.
I want ask about annals of medicine and surgery journal. Is it Q2 or Q3 according to 2019.
Thank you.
Best regards.

reply



Melanie Ortiz 3 years ago

SCImago Team

Dear Saad,
Thank you for contacting us. Our data come from Scopus, they annually send us an update of the data. This update is send to us around April / May every year. Thus, the indicators for 2019 will be available in June 2020 and before that we can't know what will happen with this journal. Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)

I'm not a robot reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



EST MODUS IN REBUS

Horatio (Satire 1, 1, 106)

[Edit Cookie Consent](#)



ISSN: 2049-0821

ANNALS OF MEDICINE & SURGERY



EDITIO • DILIGENTIA • FIDES

www.annalsjournal.com

ScienceDirect



ScienceDirect®

Submit your article

Menu



Search in this journal

Volume 69

September 2021

[< Previous vol/issue](#)

[Next vol/issue >](#)

Receive an update when the latest issues in this journal are published

[Sign in to set up alerts](#)

Open access

Editorial Board

Article 102849

[Download PDF](#)

Original Research

FEEDBACK

[Submit your article](#)[Menu](#)[Search in this journal](#)[Download PDF](#) [Article preview](#) Research article *Open access*

Shock index as a prognosticator for emergent surgical intervention and mortality in trauma patients in Johannesburg: A retrospective cohort study

Richard Crawford, Deirdre Kruger, Maeyane Moeng

Article 102710

[Download PDF](#) [Article preview](#) Research article *Open access*

Systemic lidocaine administration influences NF-k β gene expression, NF-k β and TNF- α protein levels on BALB/c mice with musculoskeletal injury

Resiana Karnina, Syafri Kamsul Arif, Mochammad Hatta, Agussalim Bukhari, ... Cahyono Kaelan

Article 102660

[Download PDF](#) [Article preview](#) Research article *Open access*

Predictive factors of mortality related to COVID-19: A retrospective cohort study of 600 cases in the intensive care unit of the university hospital of Oujda

El Kaouini Abderrahim, Merbouh Manal, El Aidouni Ghizlane, A.A.B.D.I. Mohammed, ... Housni Brahim

Article 102711

[Download PDF](#) [Article preview](#) Research article *Open access*[FEEDBACK](#)

[Submit your article](#)[Menu](#)[Search in this journal](#)Research article *Open access*

Long- and short-term survival following laparoscopic and open pancreaticoduodenectomy for patients with periampullary tumors in Vietnam

Tran Manh Hung, Tran Que Son, Tran Hieu Hoc, Tran Thanh Tung, ... Vu Duy Kien

Article 102690

[Download PDF](#) [Article preview](#) Research article *Open access*

A clinical perspective study on the compliance of surgical safety checklist in all surgical procedures done in operation theatres, in a teaching hospital, Ethiopia, 2021: A clinical perspective study

Abraham Tarekegn Mersh, Debas Yaregal Melesse, Wubie Birlie Chekol

Article 102702

[Download PDF](#) [Article preview](#) Research article *Open access*

Ski- and snowboard related open peroneal nerve injury: A 20-year retrospective case series study

J. Metzler, E.M. Morandi, K. Schwaiger, D. Wolfram, ... C. Tasch

Article 102662

[Download PDF](#) [Article preview](#) Research article *Open access*[FEEDBACK](#)

[Submit your article](#)[Menu](#)[Search in this journal](#)Research article *Open access*

Progressive spinal cord compression technique in experimental rabbit animal model for cervical spondylotic myelopathy

Sabri Ibrahim, Wibi Riawan


Article 102603

[Download PDF](#) [Article preview](#) Research article *Open access*

Association of serum KL-6 levels on COVID-19 severity: A cross-sectional study design with purposive sampling

Titah Dhadhari Suryananda, Resti Yudhawati


Article 102673

[Download PDF](#) [Article preview](#) Research article *Open access*

Expression of AIF and Caspase-3 in New Zealand rabbit with Cervical Spondylosis Myelopathy model

Sabri Ibrahim, Abdurrahman Mousa, Wibi Riawan

Article 102604

[Download PDF](#) [Article preview](#) Research article *Open access*

Technical characteristics and quality of grafts in liver procurement from brain-dead donors: A single-center study in Vietnamese population

[FEEDBACK](#) 

Annals of Medicine and Surgery

Open access

Submit your article

Menu



Search in this journal

ISSN: 2049-0801

Copyright © 2022 IJS Publishing Group Ltd. All rights reserved



Copyright © 2022 Elsevier B.V. or its licensors or contributors.
ScienceDirect® is a registered trademark of Elsevier B.V.



FEEDBACK



Annals of Medicine and Surgery

Editorial Board

Editor-in-Chief

Riaz Agha, London, England, UK

Assistant Managing and Executive Editor

Maliha Agha, London, England, UK

Assistant Editors

Adam Frampton, London, England, UK
Jorg Kleef, Munich, Germany
Deepthi Mani, Washington, USA
David Rosin, Bridgetown, Barbados
Peter Schemmer, Heidelberg, Germany

Social Media Editor

Riaz Agha, London, UK

Managing Editor

Alexander James Fowler, London, England, UK

Statistical Editor

Lai-Chu See, Taoyuan, Taiwan

Associate Staff Editors

Sergio Acuna, Toronto, Canada
Guido Alsfasser, Rostock, Germany
Fernando Angarita, Toronto, Canada
James Clark, Plymouth, England, UK
Terry Evans, London, England, UK
Shivani Garg, Wisconsin, USA
Todd Golden, Arizona, USA
Fermin Guerrero Del Angel, Madero, Mexico
Eleni Ntouvali, Athens, Greece

Assistant Commissioning Editors

Ahmed Al-Jabir, London, UK
Zaid Alsafi, London, UK
Thomas Franchi, Sheffield, UK
Christos Iosifides, London, UK
Ahmed Kerwan, London, UK

Mehdi Khan, London, UK
Ginimol Mathew, London, UK
Maria Nicola, London, UK
Niamh O'Neill, London, UK
Catrin Sohrabi, London, UK

Junior Staff Editors

Sumita Agrawal, Jodhpur, India
Bakir Al-Dulaimy, London, UK
Mohammed Anwar, London, UK
Martinique Vella Baldacchino, Glasgow, UK
Maria Borrelli, London, UK
Kerry Anne Burke, Manchester, England, UK
Adam Cristaudo, Cairns, Australia
Buket Gundogan, London, UK
Adam Gwozdz, London, England, UK

Mike Hu, California, USA
Tobin Joseph, London, UK
Kiron Koshy, London, UK
Seon-Young Lee, Southampton, England, UK
Chris Limb, London, England, UK
Yasser Al Omran, London, England, UK
Karishma Shah, Oxford, UK
Rachel Thavayogan, London, UK

Executive Committee

Sir Graeme Catto, Aberdeen, Scotland, UK
Sir Barry Jackson, London, England, UK

David Rosin, Bridgetown, Barbados

Editorial Board

Xiao-Ping Chen, Wuhan, China
Marc Cohen, Victoria, Australia
David Green, London, England, UK
Saeed Sadiq Hamid, Karachi, Pakistan
Peter Hutchinson, Cambridge, England, UK
Peter Littlejohns, London, England, UK
Ashok Mahapatra, New Delhi, India
Judy McKimm, Swansea, Wales, UK
Dattatraya Muzumdar, Mumbai, India

Ashish Nabar, Mumbai, India
Fabio Nicoli, London, UK
John Norcini, Philadelphia, PA, USA
Christoph Paasch, Berlin, Germany
Daniel Sokol, London, England, UK
Jamsheer Talati, Karachi, Pakistan
John V. Thomas, Birmingham, AL, USA
Tim Gabe Williams, Kent, England, UK
Zhao-Fan Xia, Shanghai, China

Elsevier

Executive Publisher

Allan Ross, New York, USA

Administrative Editor

Joanne Frankland, Oxford, UK

Marketing Manager

Wendy Montalvo, New York, USA

Journal Manager

Tamsyn Hopkins, Exeter, UK

Senior Business Development

Executive (Reprints)

Emma Steel, London, UK

Advertising Sales Manager

Robert Bayliss, London, UK

Account Manager (Supplements)

Evelina Euren, The Netherlands



Cross-sectional Study

Association of serum KL-6 levels on COVID-19 severity: A cross-sectional study design with purposive sampling

Titah Dhadhari Suryananda, Resti Yudhawati*

Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia



ARTICLE INFO

Keywords:
COVID-19
KL-6
AT2

ABSTRACT

Background: The main target of SARS-CoV2 is the alveolar type II (AT2) cells of the lung. SARS-CoV2 evades the innate immune system resulting in the release of proinflammatory cytokines (IL-1 β , IL-6, TNF- α) which causes AT2 cell damage. Krebs von den Lungen (KL-6) is a specific biomarker of AT2 cell damage. KL-6 is produced in AT2 cells that are injured/regenerated.

Objective: Research that discusses the role of KL-6 in COVID-19 is still being debated and not much has been done in Indonesia.

Methods: This study was an analytical study with a prospective design on 75 COVID-19 patients who were treated. Subjects were divided into two large groups according to their degree of severity, 57 subjects with severe degrees and 18 subjects with non-severe degrees. The serum KL-6 levels were measured on days 0 and 6. Data were analyzed using paired *t*-test and independent *t*-test for data were normally distributed and Wilcoxon test and Mann Whitney test for data that were not normally distributed.

Result: In this study, the mean serum KL-6 for day 0 in the severe group was higher than the non-severe group with values of 45.70 U/mL and 44.85 U/mL. On day 6, the mean serum KL-6 in the severe group was lower than that in the non-severe group with values of 41.3 U/mL and 41.95 U/mL. Serum KL-6 in the severe group experienced an even greater decrease than the non-severe group.

Conclusion: There was no significant association between serum KL-6 values on 0 days in the severity of COVID-19.

1. Introduction

Coronavirus disease 2019 (COVID-19) is caused by the novel coronavirus severe acute respiratory syndrome coronavirus 2 (SARS-CoV2). COVID-19 was first discovered in Wuhan China at the end of December 2019, three months later the virus spread to 200 countries in the world so that the World Health Organization (WHO) declared it a global pandemic [1]. In September 2020, the worldwide incidence of COVID-19 reached 31 million. The American continent is the country with the most cases in the world, followed by Southeast Asia, Europe, the Central Mediterranean, and Africa [2]. Indonesia recorded 271,000 COVID-19 cases and 10,000 deaths, which is the highest number in Southeast Asia. East Java is the second province with the highest number of cases after DKI Jakarta, which reached 42,000 with a death rate of 3000 people [3]. SARS-CoV2 shows mild-moderate manifestations and can develop into serious cases such as severe pneumonia to

Acute Respiratory Distress Syndrome (ARDS) which is the main cause of death in COVID-19 [4].

SARS-CoV2 infects through a bond between the S protein and the host cell's angiotensin-converting enzyme 2 (ACE2) receptor. The upper respiratory tract is the first site of infection, but the main target of the virus is alveolar type II (AT2) cells. SARS-CoV2 can avoid or inhibit the innate immune system so that it easily reaches the lower respiratory tract and releases pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α). The release of proinflammatory cytokines causes damage to AT2 cells. Severe damage to AT2 cells causes cytokine storms and results in ARDS characterized by diffuse alveolar damage (DAD), hyperplasia of pneumocytes, and microvascular thrombosis [5]. Biomarker studies help understand the underlying mechanisms and pathophysiology of lung damage and thus improve therapeutic strategies and evaluation. One of the specific biomarkers of pulmonary alveolar cell damage that is currently being developed for research is von den lungen Krebs (KL-6)

* Corresponding author. Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Jl. Mayjend Prof. Dr. Moestopo No. 6-8, Airlangga, Gubeng, Surabaya, East Java 60286, Indonesia.

E-mail addresses: dhadhari1788@gmail.com (T.D. Suryananda), resti.yudhawati2021@gmail.com (R. Yudhawati).

<https://doi.org/10.1016/j.amsu.2021.102673>

Received 21 June 2021; Received in revised form 30 July 2021; Accepted 3 August 2021

Available online 12 August 2021

2049-0801/© 2021 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

[6].

KL-6 is a mucin glycoprotein which is also referred to as MUC1. MUC1 is a single glycoprotein consisting of cytoplasm, transmembrane, and extracellular domains. MUC1 has a high molecular weight with large size of 200–500 nm. KL-6 is produced on the surface of AT2 epithelial cells. KL-6 is also produced in the epithelial cells of the bronchi, basal cells, and terminal bronchioles in lower numbers than AT2. KL-6 is increased in injured AT2 cells. The production of KL-6 can increase and decrease through the activity of the TNF- α converting enzyme (TACE). The expression of KL-6 protein correlated with changes in alveolar capillary permeability, indicating the association between increased KL-6 and barrier epithelial dysfunction in ARDS. KL-6 shows a correlation between bronchoalveolar lavage (BAL) and serum so that KL-6 produced in the lungs will be visible in the bloodstream [7].

Ruiz declared that the potential use of serum KL-6 in COVID-19 is a prognosis of disease activity and fibrosis [8]. Alessandro et al. said that serum KL-6 was higher in the severe group of COVID-19 compared to the non-severe group. The cut-off value of serum KL-6 in that study was 406 U/mL with a sensitivity of 89 % and a specificity of 83 % [9]. Awano, Xue, and Deng et al. said that KL-6 serum in COVID-19 patients was higher than in healthy patients. Serum KL-6 value of the severe group of COVID-19 was higher than the non-severe group. The study also stated that serum KL-6 undergoes dynamic changes with the patient's clinical condition so that it can be used as a biomarker of severity [10–12]. Other several studies have shown less significant results. Arnold et al. declared that serum KL-6 has a sensitivity of 86 % and a specificity of 45 % [13]. Kondo et al. said that serum KL-6 in ARDS patients was only increased in BAL compared to serum [14]. Based on the description above, we are interested in analyzing the association between serum KL-6 and the degree of severity of COVID-19.

2. Methods

Participants in this study were COVID-19 patients who met the inclusion and exclusion criteria. Participant inclusion criteria included patients diagnosed with COVID-19 and aged >18 years. Participants' exclusion criteria included patients who had unsuccessful blood draws by day 6 and patients diagnosed with lung, breast, and pancreatic cancer. Participants who were willing to take part in the research first received an explanation of the rights and obligations of the participants, in which they voluntarily filled out the informed consent form.

The design of this study was observational analytic with a prospective design that used consecutive sampling. This study reported the data based on the Strengthening the Reporting of Cohort Studies in Surgery (STROCSS) 2019 guideline [15]. The number of participants in this study was 75 participants were divided into two large groups namely severe (76 %) and non-severe (24 %). Severe groups consist of severe to critical categories. Non-severe groups consist of mild to moderate categories. Data collection was carried out at Dr. Soetomo General Academic Hospital, Surabaya, Indonesia during the period May–October 2020. Data collection included participant characteristics, levels of KL-6, and the degree of severity of COVID-19.

The severity of COVID-19 in this study was assessed using WHO criteria at the time of the initial examination of the patient, which distinguished the severity of COVID-19 from being non-severe (mild-moderate category) and severe (severe-critical category). Mild: Symptomatic patient who meets the COVID-19 case definition without evidence of viral pneumonia or hypoxia. Moderate: clinical symptoms of pneumonia (fever, cough, dyspnoea, rapid breathing) but no signs of severe pneumonia, including SpO₂ \geq 90 % in room air or PaO₂ \geq 60 mmHg. (PaO₂ measurements were obtained from patient medical records). Weight: obtained clinical symptoms of pneumonia (fever, cough, shortness of breath, rapid breathing) plus one of respiratory rate >30 times/minute; severe respiratory distress or SpO₂ <90 % or PaO₂ \leq 59 mmHg. (PaO₂ measurements were obtained from patient medical records). Critically, in patients with ARDS, sepsis, and septic shock. Mild

ARDS: 200 mmHg < PaO₂/FiO₂ \leq 300 mmHg (with PEEP or CPAP \geq 5 cmH₂O). Moderate ARDS: 100 mmHg < PaO₂/FiO₂ \leq 200 mmHg (with PEEP \geq 5 cmH₂O). ARDS weight: PaO₂/FiO₂ \leq 100 mmHg (with PEEP \geq 5 cmH₂O).

KL-6 is a mucin glycoprotein that is expressed on alveolar type II (AT2) epithelial cells that are injured/regenerated. The increased height of KL-6 reflects the degree of severity of COVID-19. Serum KL-6 levels were measured using the Enzyme-linked immunosorbent assay (ELISA) method of the Bioassay technology laboratory E1980Hu brand in U/ml units. KL-6 was measured by blood serum taken on day 0, namely when the diagnosis was made and on day 6 after diagnosis.

The data are recorded in data collection sheets that have been arranged, processed, and analyzed descriptively using computer software or manually. The statistical analysis used to test the hypothesis is paired and independent *t*-test for normally distributed data, Wilcoxon test, and Mann Whitney test for data that are not normally distributed. The association between data from the results of statistical analysis is shown in tables, diagrams, and text.

3. Result

3.1. Participant characteristics

The mean age of the participants was 48.04 \pm 11.66 years with an age range of 24–73 years. Most of the male participants were 46 participants (61.3 %) and the rest were female as many as 29 participants (38.7 %). Some of the participants experienced clinical symptoms as follows: fever as many as 51 participants (68.0 %), coughing as many as 63 participants (84.0 %), dyspnoea as many as 59 participants (78.6 %), diarrhea as many as 10 participants (13.0 %), anosmia as many as 5 participants (6.0 %), 1 participant of muscle pain (2.0 %), 9 participants of nausea & vomiting (12.0 %), 9 participants of sore throat (12.0 %), 7 participants of colds (9.3 %), and 14 participants of weakness (18.6%). Most of them recovered after receiving treatment as many as 62 participants (82.7 %) and the rest died as many as 13 participants (17.3 %). Only 7 participants (9.3 %) had a history of smoking. Some of the participants also had comorbid diseases such as hypertension as many as 34 participants (45.3 %), diabetes mellitus as many as 24 participants (32.0 %), COPD as many as 1 participant (1.3 %), asthma as many as 3 participants (4.0 %), tuberculosis as many as 5 participants (6.7%), obesity as many as 8 participants (10.6 %), and heart disease as many as 8 participants (10.6 %).

The mean KL-6 level at day 0 was 56.43 \pm 31.20 U/mL with a median value of 45.1 (8.5–151.4) U/mL. The mean KL-6 level in the 6th day was 52.61 \pm 37.99 U/mL with a median value of 41.5 (4.6–163.7) U/mL. The results of the 75 participant examination showed several categories of the severity of COVID-19, as follows: 18 participants (24.0 %) moderate category, 35 participants (47.0 %) severe category, and 22 participants (29.0 %) critical categories. Details of participant characteristics can be seen in [Table 1](#).

3.2. Changes in the value of KL-6 serum on day 0 and 6

The results showed that the KL-6 value on day 0 for the non-severe group was 44.85 (11.4–151.4) U/mL and the severe group was 45.70 (8.5–131.8) U/mL. The KL-6 value on day 6 for the non-severe group was 41.95 (4.6–157.4) U/mL and the severe group was 41.30 (4.6–163.7) U/mL. Serum KL-6 in the severe group experienced an even greater decrease than the non-severe group. Based on the data obtained, there was no significant association between serum KL-6 values and the severity of COVID-19 day 0 (*p* = 0.895; [Table 2](#)).

4. Discussion

Awano et al. declared that the mean serum KL-6 value of all participants on day 0 was 229 U/mL with a minimum value of 184 U/mL and a

Table 1
Characteristics of the participant based on the severity of COVID-19.

Characteristics	Non severe n = 18	Severe n = 57	p
Age	48.5 (24.0–78.0)	52.0 (25.0–73.0)	0.084
Gender			
Male	8 (44.4)	38 (66.7)	0.158
Female	10 (55.5)	19 (33.3)	
Clinical symptoms			
Fever	12 (66.7)	39 (68.4)	1.000
Cough	14 (77.7)	49 (85.9)	0.466
Crowded	7 (38.8)	52 (91.2)	<0.001*
Diarrhea	6 (33.3)	4 (7.0)	0.001
Anosmia	2 (1.1)	3 (5.2)	0.588
Muscle ache	0 (0.0)	1 (1.7)	1.000
Nauseous vomit	2 (11.1)	7 (12.2)	1.000
Sore throat	2 (11.1)	7 (12.2)	1.000
Cold	1 (5.0)	6 (10.5)	1.000
Weak body	5 (27.7)	9 (15.7)	0.303
Outcome			
Heal	18 (100.0)	44 (77.2)	–
Died	0 (0.0)	13 (22.8)	
Smoking history			
Smoker	1 (5.5)	6 (10.5)	1.000
Comorbid disease			
Hypertension	6 (33.3)	28 (49.1)	0.367
Diabetes mellitus	4 (22.2)	20 (35.0)	0.465
COPD	0 (0.0)	1 (1.7)	1.000
Asthma	0 (0.0)	3 (5.2)	1.000
Tuberculosis	0 (0.0)	5 (8.7)	0.329
Obesity	0 (0.0)	8 (14.0)	0.186
Heart disease	1 (5.5)	7 (12.2)	0.671
Laboratory Day 0			
Leukocytes	7050 (2870–14,110)	9720 (4530–25,150)	0.006*
Lymphocytes	18.8 (6.9–48.4)	10.3 (0.7–45.6)	0.001*
Neutrophils	68.7 (35.0–91.3)	80.2 (40.8–96.8)	0.007*
Procalcitonin	0.09 (0.01–0.47)	0.23 (0.01–7.04)	0.004*
CRP	1.4 (0.01–18.9)	6.2 (0.01–39.5)	0.011*
Ferritin	323.05 (24.1–1724.7)	1180 (38.5–6498.8)	0.001*
D-Dimer	980 (310–35,200)	1380 (190–35,200)	0.368
Laboratory Day 6			
Leukocytes	7110 (5080–13,160)	11,860 (3610–29,320)	<0.001*
Lymphocytes	23.45 (5.6–39.8)	9.5 (1.1–33.8)	<0.001*
Neutrophils	64.4 (35.0–88.4)	82.6 (46.4–99.2)	<0.001*
Procalcitonin	0.07 (0.01–0.47)	0.1 (0.01–14.71)	0.049*
CRP	0.55 (0.1–8.4)	3.1 (0.1–31.9)	0.006*
Ferritin	417.8 (34.3–2116)	983 (89.4–16,321)	0.009*
D-Dimer	920 (180–21,320)	2060 (190–35,200)	0.004*
Treatment			
Lopivia	5 (27.0)	24 (42.1)	–
Oseltamivir	2 (11.0)	6 (10.5)	
Avugan	1 (5.0)	1 (1.7)	
Remdisivir	1 (5.0)	1 (1.7)	
Hyloquin	9 (50.0)	19 (33.3)	
Actemra	0 (0.0)	2 (3.5)	
Crime scene	0 (0.0)	4 (7.0)	
Vitamin C	4 (22.2)	24 (32.0)	
Vitamin D	0 (0.0)	5 (8.7)	
Steroids	1 (5.0)	34 (59.6)	
Lovenox	10 (55.5)	26 (34.6)	
Heparin	3 (16.7)	32 (56.1)	
Arixtra	0 (0.0)	1 (1.7)	
NAC	15 (8.3)	35 (61.4)	
Isoprenosine	13 (72.2)	32 (56.1)	
Zinc	1 (5.0)	4 (7.0)	
Mechanical ventilation	0 (0.0)	13 (22.8)	
Non-mechanical ventilation	0 (0.0)	13 (22.8)	

Note: non severe consisting of mild and moderate covid-19 severity; severe consisting of the severity of Covid-19, severe and critical categories; *significant if $p < 0.05$.

Table 2
The serum KL-6 value is based on the degree of severity.

KL-6	Covid-19 severity		p
	non-severe	severe	
Day 0	44.85 (11.4–151.4)	45.70 (8.5–131.8)	0.895
Day 6	41.95 (4.6–157.4)	41.30 (4.6–163.7)	–
Delta	–3.2 (–66.4–39.5)	–3.1 (–109.9–111.8)	–

Note: non severe consisting of mild and moderate covid-19 severity; severe consisting of the severity of Covid-19, severe and critical categories.

maximum of 336 U/mL. On day 6, the mean score was 283 U/mL with a minimum value of 222 U/mL and a maximum value of 540 U/mL. When compared with previous studies, the mean serum KL-6 levels in this study showed lower values. The difference can be due to differences in methods and kits where previous studies used the Nanopia KL-6 reagent kit, Sexui medical. Co., Tokyo Japan antigen agglutination technique [10]. This study used Bioassay technology laboratory E1980Hu with the ELISA method. Racial and genetic differences also affect serum KL-6 values. Horimasu et al. said that the average serum KL-6 score was higher than in Japan due to Germany. The dominant participant in Germany containing the genotype A/A and Japan G/G [16].

Mason said that as many as 80 % of patients infected with COVID-19 experience non-severe symptoms, 14 % of which require hospital treatment. As many as 20 % of patients show severe symptoms, most of whom require treatment at the hospital and 5 % of this number are experiencing a critical phase so they require treatment in an intensive care unit (ICU) [17]. This study is different from the study conducted by Mason where there was a more severe group hospitalized. The explanation for this is because Dr. Soetomo General Academic Hospital, Surabaya is a referral center hospital for eastern Indonesia. Old age is more susceptible to infection and occurs with severe manifestations due to several reasons, namely a decrease in the system called immunosenescence. Cells also experience a decrease in a function called cellular senescence. Old age also occurs with systemic inflammation called inflammaging [18].

Gender did not show a significant difference in the degree of severity even though men were infected more. Cevix said that gender is related to the immune response. Men have higher congenital cytokines and chemokines than women. Women have stronger T cell activation than men. Increasing age in men leads to decreased T cell activation. This also suggests that older men are associated with a greater degree of disease severity and risk of death [19].

Shortness of breath and diarrhea are clinical symptoms that show a significant difference in severity. Shortness of breath is caused by damage to AT2 cells by viruses. The dysregulation of the immune system causes the virus to reach target cells directly. Diarrhea was more common in the severe-critical category. Diarrhea is associated with ARDS, use of mechanical ventilators, and care in the ICU but the link between diarrhea and severity is not fully understood and requires further clarification [20]. This study found no significant difference between smoking and severity and accordance to research conducted by Rosatto et al. [21].

In this study, the laboratory value showed a significant difference in the degree of severity. The laboratory for the non-severe degree group showed normal values. In the severe group, WBC showed normal/increased values and decreased lymphocytes, while neutrophils, CRP, procalcitonin, D dimer, and ferritin increased. Biomarkers such as WBC, neutrophil procalcitonin lymphocytes, CRP, D-dimer, ferritin showed significant differences with the WHO severity [22].

Dysregulation of the immune system causes the release of proinflammatory cytokines. The release of proinflammatory cytokines causes cell damage. Damaged cells release toxins that increase the release of pro-inflammatory cytokines, causing a cytokine storm. Cytokine storms cause neutrophil lymphocytes and macrophages to be released to the site of infection. Macrophages and IL-6 release ferritin, IL-6 also releases

CRP. Stress conditions cause an increase in the hormone cortisol, as a result, lymphocytes migrate to the peripheral circulatory system, then lymphocyte destruction and apoptosis occurs. Depression of T and B cells also causes low lymphocytes. Apart from being caused by damage to lung cells, the degree of severity is also caused by damage to other organs. D-dimers are released upon activation of coagulation and fibrinolysis [23].

This study assessed serum KL-6 is based on the degree of severity. On day 0, the median KL-6 value for the non-severe group was 44.85 U/mL, while the severe group was 45.70 U/mL. On day 6, the median serum KL-6 value for the non-severe grade group was 41.95 U/mL, while the severe group value was 41.30 U/mL. The KL-6 value in the severe group was higher on that two days examination. The higher the serum KL-6 value caused by the more severe/extensive AT2 cell damage. That damage or regeneration of AT2 is the main source of serum KL-6 [6].

This study assessed changes in the KL-6 serum on days 0 and 6 based on the severity. Serum KL-6 values in both groups decreased on day 6. The main decreased serum KL-6 in the non-severe group was 2.9 U/mL and in the severe group was 4.4 U/mL. The decrease was greater in the severe group. Xue et al. said that serum KL-6 value decreased on day 4 then increased sharply on day 8 and decreased again on day 16. On days 24 and 46 the serum KL-6 value increased again but not as much as on day 16. Serum KL-6 values were consistent with the clinical condition of the participant. Changes were found in the severe group, while the non-severe group tended to be stable [11]. Deng et al. said that serum KL-6 value has increased and reaches a peak for up to 1 month, then has a gradual decrease until the 6th month [12]. Awano et al. said that serum KL-6 experienced an increase on day 6 in the severe group, while the non-severe group tended to be stable [10]. Langer said that serum KL-6 can dynamically assess disease activity and pathophysiology [24]. This study and previously conducted serially showed dynamic changes in serum KL-6.

Awano et al. Deng et al., Xue et al. and Alessandro et al. said that serum KL-6 value in the severe group was higher than that in the non-severe group. Deng et al. Xue et al. and Alessandro et al. said that serum KL-6 normal patients have the same as the non-severe group [9–12]. This study and the previous study show different results. In this study, there was no significant association between serum KL-6 value on day 0 to the degree of severity. The serum KL-6 value in the severe group was not much different from the KL-6 value for the serum in the non-severe group This difference can be caused by several reasons. First is the number of samples, this study involved more subjects (75 samples) and more severity samples. In the previous study, the sample used was less in number and more in the non-severe.

Second, the cause of the degree severity is not only due to damage to AT2 cells. Apart from AT2 cells, endothelial cells are also the main targets of SARS-CoV-2. SARS-CoV2 can infect endothelial cells directly [5]. Viral infection causes endothelial dysfunction. Endothelial dysfunction results in platelet adhesion, leukocyte aggregation, complement activation, and cytokine release leading to microvascular complications such as pulmonary embolism and deep vein thrombosis (DVT) [25]. There are also features of apoptosis, pyroptosis, and lymphocytic inflammation. These histopathologic features are associated with organ ischemia, tissue edema, and procoagulant status [5].

Limitation of the study, further research is needed on the role of serum KL-6 in COVID-19 with more average sample size. Further research is needed on the role of serum KL-6 in COVID-19 involving healthy controls. It is necessary to classify the severity of the subject on day 6.

SARS-CoV2 that enters the body destroys AT2 cells resulting in increased production of KL-6. KL-6 was mostly found in the pulmonary alveoli where the more severe the inflammation in the lungs, the higher the KL-6 level. This study is used to identify the severity of COVID-19 patients in the developing world with low resource settings to improve patient care management. In future research, it is expected that the data collection time for KL-6 can be > 6 days, the participant group

will be compared to several groups (normal, mild, and moderate severity), and the number of participants will be more and represent each age group.

5. Conclusion

Participants were mostly male (61.73 %). The mean age of the participants was 48 years. Most symptoms were cough (84 %), dyspnoea (78.6 %), and fever (68 %). As many as 9.3 % of the participants were smokers. The most common comorbidities were hypertension (45.3 %) and diabetes mellitus (32 %). The mortality rate was 17.3 % which occurred in the severe group. The mean value of KL-6 serum participants on day 0 was 56.43 ± 31.204 U/mL and day 6 of 52.61 ± 37.99 U/mL. Participants in the severe group were 76 %, while the non-severe group was 24 %. On day 6, there was a decrease in serum KL-6 values in both groups. There was no significant association between serum KL-6 on day 0 and severity.

Funding

None.

Author's contributor

All authors contributed toward data analysis, drafting, and revising the paper, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Consent

Written informed consent was obtained from the patient.

Registration of Research Studies

- Name of the registry: Health Research Ethics Committee in the Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.
- Unique Identifying number or registration ID: 1953/KEPK/IV/2020.
- Hyperlink to your specific registration (must be publicly accessible and will be checked):-.

Guarantor

Resti Yudhawati is the person in charge for the publication of our manuscript.

Ethical approval

We have conducted an ethical approval base on the Declaration of Helsinki with registration research at the Health Research Ethics Committee in Dr. Soetomo General Academic Hospital, Surabaya, Indonesia (1953/KEPK/IV/2020).

Provenance and peer review

Not commissioned, externally peer-reviewed.

Declaration of competing interest

The authors declare that they have no conflict of interest.

Acknowledgment

We would like to thank Fis Citra Ariyanto for assisting in editing our manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.amsu.2021.102673>.

References

- [1] M. Soy, G. Keser, P. Atagündüz, F. Tabak, I. Atagündüz, S. Kayhan, Cytokine storm in COVID-19: pathogenesis and overview of anti-inflammatory agents used in treatment, *Clin. Rheumatol.* 39 (7) (2020) 2085–2094, <https://doi.org/10.1007/s10067-020-05190-5>.
- [2] P. Song, W. Li, J. Xie, Y. Hou, C. You, Cytokine storm induced by SARS-CoV-2. *Clinica chimica acta, Int. J. Clin. Chem.* 509 (2020) 280–287, <https://doi.org/10.1016/j.cca.2020.06.017>.
- [3] A. Chowh, T. Milanda, C.R. Sartika, M.N. Kirana, D. Halim, A. Faried, Potency of mesenchymal stem cell and its secretome in treating COVID-19, *Regenerat. Eng. Translational Med.* (2021) 1–12, <https://doi.org/10.1007/s40883-021-00202-5>.
- [4] Q. Ye, B. Wang, J. Mao, The pathogenesis and treatment of the 'Cytokine Storm' in COVID-19, *J. Infect.* 80 (6) (2020) 607–613, <https://doi.org/10.1016/j.jinf.2020.03.037>.
- [5] S.B. Bronsahan, A.H. Jonkman, M.C. Kugler, J.S. Munger, D.A. Kaufman, COVID-19 and respiratory system disorders: current knowledge, future clinical and translational research questions, *Arterioscler. Thromb. Vasc. Biol.* 40 (11) (2020) 2586–2597, <https://doi.org/10.1161/atvbaha.120.314515>.
- [6] D.R. Janz, L.B. Ware, Biomarkers of ALI/ARDS: pathogenesis, discovery, and relevance to clinical trials, *Semin. Respir. Crit. Care Med.* 34 (4) (2013) 537–548, <https://doi.org/10.1055/s-0033-1351124>.
- [7] H. Chiba, H. Takahashi, Specific serum markers of IPF, in: H. Nakamura, K. Aoshiba (Eds.), *Idiopathic Pulmonary Fibrosis: Advances in Diagnostic Tools and Disease Management*, Springer Japan, Tokyo, 2016, pp. 61–76.
- [8] A.N. Frix, L. Schoneveld, A. Ladang, M. Henket, B. Duysinx, F. Vaillant, et al., Could KL-6 levels in COVID-19 help to predict lung disease? *Respir. Res.* 21 (1) (2020) 309, <https://doi.org/10.1186/s12931-020-01560-4>.
- [9] M. d'Alessandro, P. Cameli, R.M. Refini, L. Bergantini, V. Alonzi, N. Lanzarone, et al., Serum KL-6 concentrations as a novel biomarker of severe COVID-19, *J. Med. Virol.* 92 (10) (2020) 2216–2220, <https://doi.org/10.1002/jmv.26087>.
- [10] N. Awano, M. Inomata, N. Kuse, M. Tone, K. Takada, Y. Muto, et al., Serum KL-6 level is a useful biomarker for evaluating the severity of coronavirus disease 2019, *Respir. Invest.* 58 (6) (2020) 440–447, <https://doi.org/10.1016/j.resinv.2020.07.004>.
- [11] M. Xue, P. Zheng, X. Bian, Z. Huang, H. Huang, Y. Zeng, et al., Exploration and correlation analysis of changes in Krebs von den Lungen-6 levels in COVID-19 patients with different types in China, *Biosci. Trends* 14 (4) (2020) 290–296, <https://doi.org/10.5582/bst.2020.03197>.
- [12] K. Deng, Q. Fan, Y. Yang, X. Deng, R. He, Y. Tan, et al., Prognostic roles of KL-6 in disease severity and lung injury in COVID-19 patients: a longitudinal retrospective analysis, *J. Med. Virol.* 93 (4) (2021) 2505–2512, <https://doi.org/10.1002/jmv.26793>.
- [13] N. Kokturk, C. Babayigit, S. Kul, P. Duru Cetinkaya, S. Atis Nayci, S. Argun Baris, et al., The predictors of COVID-19 mortality in a nationwide cohort of Turkish patients, *Respir. Med.* 183 (2021) 106433, <https://doi.org/10.1016/j.rmed.2021.106433>.
- [14] T. Kondo, N. Hattori, N. Ishikawa, H. Murai, Y. Haruta, N. Hirohashi, et al., KL-6 concentration in pulmonary epithelial lining fluid is a useful prognostic indicator in patients with acute respiratory distress syndrome, *Respir. Res.* 12 (1) (2011) 32, <https://doi.org/10.1186/1465-9921-12-32>.
- [15] R. Agha, A. Abdall-Razak, E. Crossley, N. Dowlut, C. Iosifidis, G. Mathew, STROCSS 2019 Guideline: Strengthening the reporting of cohort studies in surgery, *Int. J. Surg.* 72 (2019) 156–165, <https://doi.org/10.1016/j.ijsu.2019.11.002>.
- [16] Y. Horimasu, N. Hattori, N. Ishikawa, S. Kawase, S. Tanaka, K. Yoshioka, et al., Different MUC1 gene polymorphisms in German and Japanese ethnicities affect serum KL-6 levels, *Respir. Med.* 106 (12) (2012) 1756–1764, <https://doi.org/10.1016/j.rmed.2012.09.001>.
- [17] R.J. Mason, Pathogenesis of COVID-19 from a cell biology perspective, *Eur. Respir. J.* 55 (4) (2020), <https://doi.org/10.1183/13993003.00607-2020>.
- [18] A.L. Mueller, M.S. McNamara, D.A. Sinclair, Why does COVID-19 disproportionately affect older people? *Aging* 12 (10) (2020) 9959–9981, <https://doi.org/10.18632/aging.103344>.
- [19] M. Cevik, K. Kuppalli, J. Kindrachuk, M. Peiris, Virology, transmission, and pathogenesis of SARS-CoV-2, *BMJ (Clin. Res. ed)* 371 (2020) m3862, <https://doi.org/10.1136/bmj.m3862>.
- [20] F. D'Amico, D.C. Baumgart, S. Danese, L. Peyrin-Biroulet, Diarrhea during COVID-19 infection: pathogenesis, epidemiology, prevention, and management. *Clinical gastroenterology and hepatology : the official clinical practice, J. Am. Gastroenterol. Assoc.* 18 (8) (2020) 1663–1672, <https://doi.org/10.1016/j.cgh.2020.04.001>.
- [21] M. Rossato, L. Russo, S. Mazzocut, A. Di Vincenzo, P. Fioretto, R. Vettor, Current smoking is not associated with COVID-19, *Eur. Respir. J.* 55 (6) (2020), 2001290, <https://doi.org/10.1183/13993003.01290-2020>.
- [22] S. Keddie, O. Ziff, M.K.L. Chou, R.L. Taylor, A. Heslegrave, E. Garr, et al., Laboratory biomarkers associated with COVID-19 severity and management, *Clin. Immunol.* 221 (2020), 108614, <https://doi.org/10.1016/j.clim.2020.108614>.
- [23] M. Kermali, R.K. Khalsa, K. Pillai, Z. Ismail, A. Harky, The role of biomarkers in diagnosis of COVID-19 - a systematic review, *Life Sci.* 254 (2020) 117788, <https://doi.org/10.1016/j.lfs.2020.117788>.
- [24] M. Hanaoka, Y. Katsumata, H. Kawasumi, Y. Kawaguchi, H. Yamanaka, KL-6 is a long-term disease-activity biomarker for interstitial lung disease associated with polymyositis/dermatomyositis, but is not a short-term disease-activity biomarker, *Mod. Rheumatol.* 29 (4) (2019) 625–632, <https://doi.org/10.1080/14397595.2018.1553488>.
- [25] E. Gavrilaki, P. Anyfanti, M. Gavrilaki, A. Lazaridis, S. Douma, E. Gkaliagkousi, Endothelial dysfunction in COVID-19: lessons learned from coronaviruses, *Curr. Hypertens. Rep.* 22 (9) (2020) 63, <https://doi.org/10.1007/s11906-020-01078-6>.