



Source details

Annals of Medicine and Surgery

Scopus coverage years: from 2012 to Present

Publisher: Elsevier

E-ISSN: 2049-0801

Subject area: Medicine: Surgery

Source type: Journal

CiteScore 2021

1.4



SJR 2021

0.373



SNIP 2021

0.965



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▼

$$1.4 = \frac{2,710 \text{ Citations } 2018 - 2021}{1,918 \text{ Documents } 2018 - 2021}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$1.4 = \frac{4,237 \text{ Citations to date}}{3,082 \text{ Documents to date}}$$

Last updated on 05 October, 2022 • Updated monthly

CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine		
Surgery	#266/469	43rd

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)



Limited-Time Only To Go Coco

Give your day a Boost of Coco and go Loco today with our new Coconut Smoothies. Order now!

Boost Juice Indonesia

[Learn More](#)

Annals of Medicine and Surgery

COUNTRY

Netherlands



Universities and research institutions in Netherlands

SUBJECT AREA AND CATEGORY

Medicine
Medicine (miscellaneous)
Surgery

PUBLISHER

Elsevier BV

H-INDEX

30

PUBLICATION TYPE

Journals

ISSN

20490801

COVERAGE

2012-2021

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

annalsjournal@elsevier.com

SCOPE

Annals of Medicine and Surgery is an online-only, peer-reviewed open access journal with a global outlook and focus on those training in medicine and surgery (postgraduate and undergraduate). AMS contains a mix of original clinical and basic science research, reviews, editorials, commentary, perspectives, debate, opinion, case reports and journal club reports. The journal crosses the whole disease spectrum of medicine and surgery but does also have a special focus on the following areas: -Patient safety- Human factors, teamwork, communication and professionalism- Quality improvement science and practice- Evidence based medicine- Implementation science – clinical evidence into practice- Leadership and management- Medical education, teaching, and training- Public and global health- Healthcare policy, delivery, commissioning, and resource management- Use of technology and health informatics- Clinical ethics and medical law- Research and innovation – bedside to bench and back again – including applications like personalised medicine.






Join the conversation about this journal


Quartiles




FIND SIMILAR JOURNALS

<p>1 International Journal of Surgery NLD</p> <p>80% similarity</p>	<p>2 International Journal of Surgery Open GBR</p> <p>73% similarity</p>	<p>3 International Journal of Surgery Case Reports NLD</p> <p>67% similarity</p>	<p>4 BMC Surgery GBR</p> <p>61% similarity</p>
---	--	--	--






% International Collaboration





Citable documents

Non-citable documents



Cited documents

Uncited documents



Annals of Medicine and Surgery

Q3
Medicine (miscellaneous)
best quartile

SJR 2021
0.37

powered by scimagojr.com


Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2022

K **Khalid** 10 months ago

Would you inform me what is the impact factor of this journal please?

reply



Melanie Ortiz 9 months ago

Dear Khalid, thank you very much for your comment. SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR (Check it on our website). We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team

S Sultan jarrar 2 years ago

Dear SCIMAGO team,

Could you please tell me if this journal considered to be q2 or q3 ? There is a little confusion about it .

Kind regards

reply



Melanie Ortiz 2 years ago

Dear Sultan,

thank you for contacting us. For every journal, the annual value of the SJR is integrated into the distribution of SJR values of all the subject categories to which the journal belongs. There are more than 300 subject categories. The position of each journal is different in any category and depends on the performance of the category, in general, and the journal, in particular .

Best Regards, SCImago Team

A Anton 2 years ago

Dear sir/ madam, please tell if the journal status Q2 or Q3?

reply



Melanie Ortiz 2 years ago

Dear Anton, thank you for contacting us. For every journal, the annual value of the SJR is integrated into the distribution of SJR values of all the thematic categories to which the journal belongs. There are more than 300 thematic categories. The position of each journal is different in any category and depends on the performance of the category, in general, and the journal, in particular. The next SJCR's update will be made throughout June 2020. Best Regards, SCImago Team

S **Saad** 2 years ago

Dear. Melanie
Thank you very much for your reply.
Best regards.
Saad

reply



Melanie Ortiz 2 years ago

SCImago Team

Dear Saad, thanks for your participation! Best Regards, SCImago Team

S **Saad** 3 years ago

Dear.
I want ask about annals of medicine and surgery journal. Is it Q2 or Q3 according to 2019.
Thank you.
Best regards.

reply



Melanie Ortiz 3 years ago

SCImago Team

Dear Saad,
Thank you for contacting us. Our data come from Scopus, they annually send us an update of the data. This update is send to us around April / May every year. Thus, the indicators for 2019 will be available in June 2020 and before that we can't know what will happen with this journal. Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)

I'm not a robot reCAPTCHA
[Privacy](#) - [Terms](#)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



EST MODUS IN REBUS

Horatio (Satire 1, 1, 106)

[Edit Cookie Consent](#)



ISSN: 2049-0821

ANNALS OF MEDICINE & SURGERY



EDITIO • DILIGENTIA • FIDES

www.annalsjournal.com

ScienceDirect



Annals of Medicine and Surgery

Open access


1.4


CiteScore


Articles & Issues ▾

About ▾

Publish ▾

 Search in this journal

Submit your article 

Guide for authors 

Volume 74

February 2022

[← Previous vol/issue](#)

[Next vol/issue →](#)

Actions for selected articles

Select all / Deselect all

 Download PDFs

 Export citations

Show all article previews

Contents


Original Research

Review Articles

Case Reports

Short Communications

Receive an update when the latest issues in this journal are published

 [Sign in to set up alerts](#)

[Open access](#)

Editorial Board

Article 104357

 [Download PDF](#)

Original Research

[Research article](#) [Open access](#)

Surgical patient satisfaction with a virtual hybrid care hotel model: A retrospective cohort study

Ryan M. Chadha, Margaret R. Paulson, Francisco R. Avila, Ricardo A. Torres-Guzman, ... Michael J. Maniaci

Article 105251

Actions for selected articles

[Select all](#) / [Deselect all](#) [Download PDFs](#) [Export citations](#) [Show all article previews](#)

Contents

[Original Research](#)[Review Articles](#)[Case Reports](#)[Short Communications](#)

Original Research

 Research article  Open access**Surgical patient satisfaction with a virtual hybrid care hotel model: A retrospective cohort study**Ryan M. Chadha, Margaret R. Paulson, Francisco R. Avila, Ricardo A. Torres-Guzman, ... Michael J. Maniaci
Article 103251 [Download PDF](#) [Article preview](#)  Research article  Open access**Colorectal cancer survival rates in Makassar, Eastern Indonesia: A retrospective Cohort Study**Ibrahim Labeda, Ronald Erasio Lusikooy, Mappincara, Muhammad Iwan Dani, ... Muhammad Faruk
Article 103211 [Download PDF](#) [Article preview](#)  Research article  Open access**Comparison of Doppler Ultrasound and Digital Subtraction Angiography in extracranial stenosis**Seyed Farzad Maroufi, Seyedeh Niloufar Rafiee Alavi, Mohammad Hossein Abbasi, Ali Farnouri, ... Mohammad Jafar Joghataei
Article 103202 [Download PDF](#) [Article preview](#)  Research article  Open access**The effect of protein supplements on weight loss, body composition, protein status, and micronutrients post laparoscopic sleeve gastrectomy (LSG): A Randomised Controlled Trial (RCT)**Sahar Alshamari, Mohamed Aly Elshehif, Fahad Hanna, Leyan El Akhal, ... Wahiba Elhag
Article 103220 [Download PDF](#) [Article preview](#)  Research article  Open access**Synthesis and anti-Toxoplasma activity of indole-triazole compounds on tachyzoites of RH strain**

Mohammad Saleh Bahreini, Aida Traji, Najmeh Edraki, Ali Arab Monfared, Qasem Asgari



[📄](#) Download PDF [Article preview ▼](#)

Actions for selected articles

[Select all / Deselect all](#)[📄](#) Download PDFs[📄](#) Export citations Show all article previews

Contents

[Original Research](#)[Review Articles](#)[Case Reports](#)[Short Communications](#) Research article [Open access](#)**Association of soluble receptor for advanced glycation end-products (sRAGE) serum on COVID-19 severity: A cross-sectional study**Gurtr Noor Ramadany Saputra, Resti Yudhawati, Munawaroh Fitriah
Article 103303[📄](#) Download PDF [Article preview ▼](#) Research article [Open access](#)**Survey on knowledge, attitude and practice of labor analgesia among health care providers at Debre Markos Comprehensive Specialized Hospital, Ethiopia 2021. A cross-sectional study**Amsal Shiferaw Melesse, Samuel Debas Bayable, Getamesay Demelash simogn, Yitayal Guadie Ashebir, ... Melaku Bantie Fetene
Article 103306[📄](#) Download PDF [Article preview ▼](#) Research article [Open access](#)**Postoperative poor sleep quality and its associated factors among adult patients: A multicenter cross-sectional study**Shumelis Seid Tegegne, Efrom Henta Alemnew
Article 103273[📄](#) Download PDF [Article preview ▼](#) Research article [Open access](#)**Bactericidal effect of hyperbaric oxygen therapy in burn injuries**Mendy Hatibie Oley, Maximilian Christian Oley, Louise A.J. Waworuntu Wewengkang, Billy Johnson Kepel, ... Muhammad Faruk
Article 103334[📄](#) Download PDF [Article preview ▼](#) Research article [Open access](#)**High-flow nasal oxygen therapy decrease the risk of mortality and the use of invasive mechanical ventilation in patients with severe SARS-CoV-2 pneumonia? A retrospective and comparative study of 265 cases**

Short Communications

 Short communication • Open access**Clinical characteristics and outcomes of cancer patients and their hospital course during the COVID-19 pandemic in a developing country**Muhammad Sohail Asghar, Farah Yasmin, Maryam Salma Babar, Muhammad Daim Bin Zafar, ... Muhammad Junaid Tahir
Article 103284 Download PDF Article preview  Short communication • Open access**Prasugrel versus ticagrelor for acute coronary syndrome patients undergoing percutaneous coronary intervention: A critical appraisal of randomized controlled trials**Iqbal Almas, Maryam Ehtesham, Janita Basit, Tarek Khedro, ... Jun Li
Article 103330 Download PDF



Annals of Medicine and Surgery

Editorial Board

Editor-in-Chief

Riaz Agha, London, England, UK

Assistant Managing and Executive Editor

Maliha Agha, London, England, UK

Assistant Editors

Adam Frampton, London, England, UK
Jorg Kleef, Munich, Germany
Deepthi Mani, Washington, USA
David Rosin, Bridgetown, Barbados
Peter Schemmer, Heidelberg, Germany

Social Media Editor

Riaz Agha, London, UK

Managing Editor

Alexander James Fowler, London, England, UK

Statistical Editor

Lai-Chu See, Taoyuan, Taiwan

Associate Staff Editors

Sergio Acuna, Toronto, Canada
Guido Alsfasser, Rostock, Germany
Fernando Angarita, Toronto, Canada
James Clark, Plymouth, England, UK
Terry Evans, London, England, UK
Shivani Garg, Wisconsin, USA
Todd Golden, Arizona, USA
Fermin Guerrero Del Angel, Madero, Mexico
Eleni Ntouvali, Athens, Greece

Assistant Commissioning Editors

Ahmed Al-Jabir, London, UK
Zaid Alsafi, London, UK
Thomas Franchi, Sheffield, UK
Christos Iosifides, London, UK
Ahmed Kerwan, London, UK

Mehdi Khan, London, UK
Ginimol Mathew, London, UK
Maria Nicola, London, UK
Niamh O'Neill, London, UK
Catrin Sohrabi, London, UK

Junior Staff Editors

Sumita Agrawal, Jodhpur, India
Bakir Al-Dulaimy, London, UK
Mohammed Anwar, London, UK
Martinique Vella Baldacchino, Glasgow, UK
Maria Borrelli, London, UK
Kerry Anne Burke, Manchester, England, UK
Adam Cristaudo, Cairns, Australia
Buket Gundogan, London, UK
Adam Gwozdz, London, England, UK

Mike Hu, California, USA
Tobin Joseph, London, UK
Kiron Koshy, London, UK
Seon-Young Lee, Southampton, England, UK
Chris Limb, London, England, UK
Yasser Al Omran, London, England, UK
Karishma Shah, Oxford, UK
Rachel Thavayogan, London, UK

Executive Committee

Sir Graeme Catto, Aberdeen, Scotland, UK
Sir Barry Jackson, London, England, UK

David Rosin, Bridgetown, Barbados

Editorial Board

Xiao-Ping Chen, Wuhan, China
Marc Cohen, Victoria, Australia
David Green, London, England, UK
Saeed Sadiq Hamid, Karachi, Pakistan
Peter Hutchinson, Cambridge, England, UK
Peter Littlejohns, London, England, UK
Ashok Mahapatra, New Delhi, India
Judy McKimm, Swansea, Wales, UK
Dattatraya Muzumdar, Mumbai, India

Ashish Nabar, Mumbai, India
Fabio Nicoli, London, UK
John Norcini, Philadelphia, PA, USA
Christoph Paasch, Berlin, Germany
Daniel Sokol, London, England, UK
Jamsheer Talati, Karachi, Pakistan
John V. Thomas, Birmingham, AL, USA
Tim Gabe Williams, Kent, England, UK
Zhao-Fan Xia, Shanghai, China

Elsevier

Executive Publisher

Allan Ross, New York, USA

Administrative Editor

Joanne Frankland, Oxford, UK

Marketing Manager

Wendy Montalvo, New York, USA

Journal Manager

Tamsyn Hopkins, Exeter, UK

Senior Business Development

Executive (Reprints)

Emma Steel, London, UK

Advertising Sales Manager

Robert Bayliss, London, UK

Account Manager (Supplements)

Evelina Euren, The Netherlands



Cross-sectional Study

Association of soluble receptor for advanced glycation end-products (sRAGE) serum on COVID-19 severity: A cross-sectional study

Gusti Noor Ramadany Saputra^a, Resti Yudhawati^{a,*}, Munawaroh Fitriah^b^a Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia^b Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

ARTICLE INFO

Keywords:

Serum sRAGE
 COVID-19 severity
 Infectious disease

ABSTRACT

Background: Coronavirus disease 2019 (COVID-19) is a new health problem discovered in 2019 thus requires biomarkers that can detect early tissue damage. Soluble receptor for advanced glycation end-products (sRAGE) is a biomarker that can be used to identify early lung damage.

Objective: Analyzing the association of serum sRAGE on COVID-19 severity.

Methods: This study employed a cross-sectional design with a consecutive sampling method. It was conducted from May 2020–October 2021. The number of participants in this study was 145 participants which were divided into 2 groups (non-severe = 47 and severe = 98). Association of sRAGE serum on COVID-19 severity was analyzed using the chi-square test, Fisher's exact test, independence *t*-test, Mann Withney test, and Spearman's rank test with *p*-value <0.05.

Results: The results of blood analysis showed several blood components such as leukocytes ($9896.51 \pm 4949.64/\mu\text{L}$; $z = 2.431$; $p = 0.015$), lymphocytes ($13.55 \pm 8.48\%$; $z = 2.256$; $p = 0.024$), neutrophils ($78.91 \pm 10.50\%$; $z = 2.464$; $p = 0.014$), procalcitonin ($0.92 \pm 3.22 \text{ ng/mL}$; $z = 3.323$; $p = 0.001$), CRP ($8.59 \pm 7.62 \text{ mg/L}$; $z = 2.114$; $p = 0.034$), D-dimer ($4360.29 \pm 7797.81 \text{ ng/mL}$; $z = 2.186$; $p = 0.029$), and fibrinogen ($474.58 \pm 168.90 \text{ mg/dL}$; $t = 0.383$; $p = 0.703$). There was a significant comparison in serum sRAGE values in the non-severe group ($0.78 [0.63\text{--}1.00] \text{ ng/mL}$) and severe group ($1.47 [0.97\text{--}2.25] \text{ ng/mL}$; $r = 7.154$; $p < 0.001$). There was a significant association between serum sRAGE and COVID-19 severity ($r = 0.598$; $p < 0.001$). The cut-off value for serum sRAGE between the severe and non-severe groups was 0.985 ng/mL . This study obtained sensitivity of 73.5%, specificity of 74.5% OR 8.077 and AUC 0.868 95% CI.

Conclusion: There is a significant association between serum sRAGE and COVID-19 severity and there is also a significant difference in serum sRAGE in the two groups.

1. Introduction

Coronavirus disease 2019 or better known as COVID-19 caused by SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) became a worldwide pandemic at the end of 2019 with various systemic complaints but was more dominant in respiratory disorders. The worldwide mortality rate was 2.1% by February 12, 2020 [1]. The February 2020 data by Johns Hopkins University's Center for Systems Science and Engineering (CSSE) showed a total case of more than 60,331 patients, with a total death of more than 1369 patients and an improvement of more than 6061 patients [2]. On December 27, 2020, the total number of worldwide cases was more than 79 million,

including 1,751,311 deaths. Incidents in Indonesia were 706,837 confirmed cases of COVID-19 and 20,994 cases of death [3].

The severity of COVID-19 according to WHO is divided into mild, moderate, severe, and critical [4,5]. The most frequently encountered clinical symptoms are pneumonia symptoms. Biomarkers are frequently used to determine the severity of pneumonia such as procalcitonin, C-reactive protein (CRP), copeptin, pro-ANP (atrial natriuretic peptide), adrenomedullin, cortisol, and D-dimers [6]. These biomarkers are good in determining infection in pneumonia but have not been able to detect early tissue damage, as patients often go to the hospital with a more severe condition. Recent studies in immunology have examined soluble receptors for advanced glycation end-products (sRAGE) as a biomarker

* Corresponding author. Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Jl. Mayjend Prof. Dr. Moestopo No. 6-8, Airlangga, Gubeng, Surabaya, East Java 60286, Indonesia.

E-mail address: restiyudhawati2021@gmail.com (R. Yudhawati).

<https://doi.org/10.1016/j.amsu.2022.103303>

Received 1 December 2021; Received in revised form 21 January 2022; Accepted 23 January 2022

Available online 29 January 2022

2049-0801/© 2022 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

of the severity of community pneumonia and can detect tissue damage in ARDS early [7].

Pathophysiology occurred in COVID-19 includes the inflammatory process. One of the inflammatory processes during pneumonia is characterized by an increase in receptors for advanced glycation end-products (RAGE). RAGE is one of the non-enzymatic receptors of Advanced Glycation End-Products (AGEs) which has a multi-ligand receptor, namely a V-type domain, two C-type domains, a transmembrane domain, and a cytoplasmic tail. RAGE has several ligands including AGEs, S100/calgranulins, and HMGB I which are present in different vascular cells such as endothelial cells, neuronal cells, smooth muscle cells, or inflammatory cells (monocytes). HMGB I is one of the RAGE ligands that play a role in the occurrence of sepsis which can stimulate the formation of cytokines along with TLRs in the immune system cells (B cells) [8]. The interaction between RAGE and its ligands will cause the formation of Reactive Oxygen Species (ROS) which will activate NADPH oxidation. The process will mediate the formation of inflammatory cells. Trianta et al. stated two processes of RAGE interaction with its ligands that are related to the inflammatory process, namely its interaction with leukocytes and on endothelial cells, RAGE is an adhesive receptor and directly forms inflammatory cells. The accumulation of RAGE ligands is predicted to cause chronic cell stimulation and tissue damage [9,10].

RAGE is expressed in the membrane-bound form (fl-RAGE or mRAGE) and the soluble form in the transmembrane domain. Soluble RAGE is produced by proteolytic cleavage of fl-RAGE and alternative splicing mRNA [7]. The administration of sRAGE in experimental animals can also interact with the RAGE ligand [10]. Based on these studies, the role of sRAGE becomes very important in determining COVID-19 diagnosis based on the severity quickly, so that effective and adequate treatment planning can be carried out early to reduce the morbidity and mortality of COVID-19 patients. In addition, the level of sRAGE in serum can detect early tissue damage which in turn can affect the severity of COVID-19 patients as common biomarkers have not been able to detect the process of tissue damage early. Research on sRAGE in the serum of COVID-19 patients is still limited and has never been carried out in Indonesia despite a few studies having been conducted in other countries. This biomarker is also easy to use and at a more affordable cost, so we are interested in analyzing the association of serum sRAGE on COVID-19 severity.

2. Methods

2.1. Participants

Participants in this study were COVID-19 patients diagnosed with real-time polymerase chain reaction (PCR) [5]. Participants' inclusion criteria included patients diagnosed with COVID-19 and aged >21 years. Participants' exclusion criteria included patients with a history of respiratory tract infection, myocardia infarct, cancer, and cerebral vascular attack. Participants who were willing to take part in the research first received an explanation of the rights and obligations of the participants, in which they voluntarily filled out the informed consent form.

2.2. Study design

This study used a cross-sectional design with a consecutive sampling method. It was carried out from May 2020–October 2020. This study collected participant characteristics, serum sRAGE, and COVID-19 severity. This study reported the data based on the strengthening the reporting of cohort studies in surgery (STROCSS) 2021 guideline [11]. The number of participants in this study was 145 participants that were divided into 2 groups (non-severe = 47 and severe = 98). The non-severe group consisted of participants identified as having COVID-19 in the mild and moderate category, while the severe group consisted of participants identified as having COVID-19 in the severe

and critical categories [5].

2.3. Assessment of COVID-19 severity

The severity of COVID-19 in this study was assessed using WHO criteria at the time of the initial examination of the patient, which distinguished the severity of COVID-19 from being non-severe (mild-moderate category) and severe (severe-critical categories). Mild is a symptomatic patient who meets the COVID-19 case definition without evidence of viral pneumonia or hypoxia. Moderate include clinical symptoms of pneumonia (fever, cough, dyspnoea, rapid breathing) but no signs of severe pneumonia, including SpO₂ 90% in room air or PaO₂ 60 mmHg (PaO₂ measurements were obtained from patient medical records). Severe shows clinical symptoms of pneumonia (fever, cough, shortness of breath, rapid breathing) plus one of respiratory rate >30 times/minute; severe respiratory distress or SpO₂ <90% or PaO₂ 59 mmHg (PaO₂ measurements were obtained from patient medical records). Critical when patients have ARDS, sepsis, and septic shock. Mild ARDS: 200 mmHg < PaO₂/FiO₂ 300 mmHg (with PEEP or CPAP 5 cmH₂O). Moderate ARDS: 100 mmHg < PaO₂/FiO₂ 200 mmHg (with PEEP 5 cmH₂O). ARDS weight: PaO₂/FiO₂ 100 mmHg (with PEEP 5 cmH₂O) [5].

2.4. sRAGE serum examination

The sRAGE is soluble forms in the transmembrane domain of RAGE which the serum levels of sRAGE are determined using a specific sandwich human ELISA kit BioAssay (MyBioSource Inc, San Diego, USA). The sRAGE measurement is in the range of 0.31–2.00 ng/mL. These results were obtained from taking 5 cc venous blood samples [12].

2.5. Statistical analysis

The analysis in this study used descriptive analysis and bivariate analysis. The descriptive analysis includes a descriptive presentation of the results using a distribution table, mean, median, standard deviation, maximum value, and minimum value. The analysis was conducted using IBM SPSS Statistics software version 21.0 (IBM Corp., Armonk, NY, USA). Participants' characteristic data were analyzed using the chi-square test or Fisher's exact test. Meanwhile, the data from this study were first tested for normality using the Kolmogorov-Smirnov test. Analysis of the association of sRAGE serum with COVID-19 severity using the independence *t*-test or Mann Whitney test. The comparison between the two variables is significant if *p* < 0.05. In addition, Spearman's rank test was used to analyze the association between two variables.

3. Results

3.1. Characteristics of participants

The demographic characteristics of participants included age and gender. The average age of participants was 50.54 ± 12.70 years (non-severe group = 49.11 ± 12.44 years and severe group = 51.23 ± 12.83 years). The median age of participants was 52.00 (43.00–59.00) years of which the youngest participant was 22.00 years old and the oldest participant was 80.00 years old. Most participants were in the age range of 35.00–55.00 years, consisting of 25 participants (53.2%) in the non-severe group and 51 participants (52.0%; *p* = 0.705) in the severe group. Most participants were male (90 participants; 62.1%), consisting of 25 participants (53.2%) in non-severe group and 65 participants in severe group (66.3%; OR = 0.577; *p* = 0.179; Table 1).

There were several clinical symptoms appeared, including shortness of breath in 122 participants (84.1%; 63.8% vs 93.9%; OR = 8689; *p* < 0.001), fever in 61 participants (42.1%; 46.8% vs 39.8%; OR = 0.751; *p* = 0.535), cough in 70 participants (70%; 59.6% vs. 42.9%; OR =

Table 1
Characteristics of participants.

Characteristics	COVID-19 Severity		p
	Non-severe	Severe	
Age (years)			0.705
21-35	8 (17.0)	12 (12.2)	
35-55	25 (53.2)	51 (52.0)	
55-65	8 (17.0)	24 (24.5)	
>65	6 (12.8)	11 (11.2)	
Gender			0.179
Male	25 (53.2)	65 (66.3)	
Female	22 (46.8)	33 (33.7)	
Clinical symptoms			
Shortness of breath	30 (63.8)	92 (93.9)	<0.001**
Fever	22 (46.8)	39 (39.8)	0.535
Cough	28 (59.6)	42 (42.9)	0.088
Painful swallowing	1 (2.1)	3 (3.1)	1.000
Diarrhea	3 (6.4)	4 (4.1)	0.682
Outcome			<0.001**
Recovered	41 (87.2)	47 (48.0)	
Died	6 (12.8)	51 (52.0)	
Comorbid			
Hypertension	13 (27.7)	28 (28.6)	1.000
Diabetes	20 (42.6)	46 (46.9)	0.750
Obesity	9 (19.1)	26 (26.5)	0.444

Note: *significant <0.05; **significant <0.01.

0.509; $p = 0.088$), painful swallowing in 4 participants (2.8%; 2.1% vs 3.1%; OR = 1.453; $p = 1.000$), and diarrhea in 7 participants (4.8%; 6.4% vs. 4.1%; OR = 0.624; $p = 0.682$). Based on the outcome of the COVID-19 treatment, most of non-severe participants recovered as many as 41 participants (87.2%) and most of severe participants were declared dead as many as 51 participants (52%; $p < 0.001$). Overall, 88 participants (60.7%) were recovered. Several participants were declared to have comorbidities, including hypertension as many as 41 participants (28.3%; 27.7% vs 28.6%; OR = 1.046; $p = 1.000$), diabetes as many as 66 participants (45.5%; 42.6% vs 46.9%; OR = 1.194; $p = 0.750$), and obesity as many as 35 participants (24.1%; 19.1% vs. 26.5%; OR = 1.525; $p = 0.444$; Table 1).

3.2. Association of soluble receptor for Advanced Glycation End-Products (sRAGE) serum with COVID-19 severity

The results of blood analysis showed several blood components such as leukocytes ($9896.51 \pm 4949.64/\mu\text{L}$), lymphocytes ($13.55 \pm 8.48\%$), neutrophils ($78.91 \pm 10.50\%$), procalcitonin ($0.92 \pm 3.22 \text{ ng/mL}$), CRP ($8.59 \pm 7.62 \text{ mg/L}$), D-dimer ($4360.29 \pm 7797.81 \text{ ng/mL}$), and fibrinogen ($474.58 \pm 168.90 \text{ mg/dL}$). The average value of serum sRAGE was $1.48 \pm 0.98 \text{ ng/mL}$, with a median value of 1.07 (0.85–1.84) ng/mL. The lowest and highest value of participants' serum sRAGE was 0.44 ng/mL and 5.14 ng/mL, respectively. The results of the COVID-19 severity measurement were divided into 4: mild as many as 2 participants (1.4%), moderate as many as 45 participants (31.0%), severe as many as 96 participants (66.2%), and critical as many as 2 participants (1.4%). Meanwhile, in this study, COVID-19 severity was divided into 2 groups, namely the non-severe group with 47 participants (32.88%) and the severe group with 98 participants (68.53%).

There was a significant difference in blood component in the non-severe group and the severe group as follows: leukocyte value was 8005.00 (6157.50–9687.50) vs 9840.00 (7420.00–12,830.00/ μL ; $z = 2.431$; $p = 0.015$), lymphocyte was 14.40 (8.83–21.65) vs 10.20 (6.60–16.80%; $z = 2.256$; $p = 0.024$), neutrophils was 77.40 (68.90–83.28) vs. 82.60 (76.00–87.10%; $z = 2464$; $p = 0.014$), procalcitonin was 0.11 (0.07–0.22) vs 0.27 (0.13–0.46 ng/mL; $z = 3.323$; $p = 0.001$), CRP was 4.65 (0.80–11.35) vs. 8.70 (2.30–13.60 mg/L; $z = 2.114$; $p = 0.034$), and D-dimer was 810.00 (535.00–2430.00) vs. 1460.00 (740.00–4025 ng/mL; $z = 2.186$; $p = 0.029$). Meanwhile, there was no significant difference in the levels of fibrinogen between

participants in the two groups (465.50 ± 176.04 vs. $480.06 \pm 165.92 \text{ mg/dL}$; $t = 0.383$; $p = 0.703$; Table 2).

There was a significant difference between serum sRAGE in the non-severe group and the severe group of 0.78 (0.63–1.00) vs 1.47 (0.97–2.25 ng/mL; $r = 7.154$; $p < 0.001$; Table 2). There was a significant association between serum sRAGE and COVID-19 severity ($r = 0.598$; $p < 0.001$). The cut-off value for serum sRAGE between the severe and non-severe group was 0.985 ng/mL. This study obtained sensitivity of 73.5%, specificity of 74.5%, OR of 8.077 and AUC 0.868 CI 95% (Fig. 1).

4. Discussion

This study assessed serum sRAGE based on the severity of COVID-19. The results of this study are consistent with previous studies that examined sRAGE as a biomarker for COVID-19. A study examined the association of sRAGE with severity and as an indicator of mechanical ventilation requirements, ARDS, and mortality in COVID-19 patients. The results showed an increase in serum sRAGE concentrations in COVID-19 patients based on severity [13]. These results are consistent with another study which stated a significant increase in serum sRAGE of ARDS patients admitted to non-isolated ICUs [14].

There is a significant association between serum sRAGE and COVID-19 severity. The serum sRAGE values in the severe group show a significant difference from serum sRAGE values in the non-severe group. The results are consistent with previous studies that showed an increase in serum sRAGE values in COVID-19 patients with a degree of severity. Increased sRAGE values can also help predict respiratory disorders that require mechanical ventilation and the mortality rate of COVID-19 patients [13]. Increased serum sRAGE is commonly found in ARDS patients admitted to the ICU [15]. As many as 20% of COVID-19 patients progress to the third phase called the involvement of the respiratory tract and progression to ARDS [16].

Increased serum sRAGE values can occur due to a viral infection process that will trigger an immune response, namely the innate immune system. Pattern-recognition receptors (PRR) recognize pathogen-associated molecular patterns (PAMPs) involving toll-like receptors (TLR) that detect components of infection and signaling tissue damage, one of which is HMGB1. Then it continues to the process of indirect lung tissue damage, namely damage-associated molecular patterns (DAMPs) that involve RAGE, NLR, TLR, and CLR which can exacerbate the occurrence of tissue damage that has occurred previously. The process of interaction of sRAGE with its ligand becomes more frequent due to an increase in HMGB1 that result in the increased inflammatory response in the form of IL-1 and TNF-Alpha activation [17,18].

Other tissue damage processes can also occur when SARS-CoV2 invades AT2 cells located in the periphery and subpleural so that the patient begins to feel hypoxia. SARS-CoV2 replicates in AT2 lead to cell damage and death. Dead AT2 cells release toxins and damage

Table 2
Comparison of blood component based on COVID-19 severity.

Blood Analysis	COVID-19 Severity		p
	Non-severe	Severe	
Leukocytes (n = 139)	8622.10 ± 4204.47	10,526.86 ± 5185.37	0.015*
Lymphocyte (n = 139)	15.50 ± 8.22	12.58 ± 8.49	0.024*
Neutrophile (n = 139)	76.06 ± 10.36	80.32 ± 10.34	0.014*
Procalcitonin (n = 143)	1.01 ± 4.67	0.88 ± 2.22	0.001*
CRP (n = 90)	6.52 ± 6.71	9.53 ± 7.87	0.034*
Fibrinogen (n = 85)	465.50 ± 176.04	480.06 ± 165.92	0.703
D-Dimer (n = 139)	2790.64 ± 5558.74	5162.17 ± 8641.11	0.029*
s-RAGE (n = 143)	0.82 ± 0.23	1.80 ± 1.04	<0.001**

Note: CRP = C-reactive protein; s-RAGE = soluble receptor for advanced glycation end products; *significant <0.05; **significant <0.001.




Fig. 1. Cut-off Serum sRAGE level based on severe and non-severe groups of COVID-19 patients.

surrounding cells. Infected cells send signals that are detected by the immune system which then releases cytokines such as IL-1, IL-6, and TNF- α . These cytokine release aims to kill the virus, but it also causes damage to lung cells, namely diffuse alveolar damage, formation of hyaline membranes, and multinuclear giant cells. Abnormal wound healing leads to fibrosis [16,19].

This study, however, has limitations, including the need for a future study that compares healthy individuals and pneumonia patients without COVID-19.

5. Conclusion

sRAGE is a biomarker that can be used to determine COVID-19 severity. The patients' COVID-19 severity in this study is categorized into 2, namely non-severe and severe. Based on blood component analysis, there are significant differences between the non-severe and severe groups. The differences consist of leukocytes, lymphocytes, neutrophils, procalcitonin, CRP, and D-dimer. The sRAGE values in the two groups also show a significant difference. In addition, there is a significant association between serum sRAGE and COVID-19 severity.

Ethical approval

We have conducted an ethical approval base on the Declaration of Helsinki with registration research at the Health Research Ethics Committee in Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.

Funding

Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.

Guarantor

Resti Yudhawati.

Author contributor

All authors contributed toward data analysis, drafting and revising the paper, gave final approval of the version to be published and agree to be accountable for all aspects of the work.

Provenance and peer review

Not commissioned, externally peer-reviewed.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Author contributions

All authors contributed toward data analysis, drafting and revising the paper, gave final approval of the version to be published and agree to be accountable for all aspects of the work.

Registration of research studies

1. Name of the registry: Health Research Ethics Committee in the Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.
2. Unique Identifying number or registration ID: 1954/KEPK/TV/2020.
3. Hyperlink to your specific registration (must be publicly accessible and will be checked): -.

Declaration of competing interest

The authors declare that they have no conflict of interest.

Acknowledgment

We would like to thank the COVID-19 patients and Guardian. We would also thank Dr. Soetomo General Academic Hospital as the place of our research, and our editor "Fis Citra Ariyanto".

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.amsu.2022.103303>.

References

- [1] H.A. Rothan, S.N. Byrareddy, The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak, *J. Autoimmun.* 109 (2020), 102433, <https://doi.org/10.1016/j.jaut.2020.102433>.
- [2] H. Harapan, N. Itoh, A. Yufika, W. Winardi, S. Keam, H. Te, et al., Coronavirus disease 2019 (COVID-19): a literature review, *J. of infection and public health* 13 (5) (2020) 667–673, <https://doi.org/10.1016/j.jiph.2020.03.019>.
- [3] R. Djalante, J. Lassa, D. Setiamarga, A. Sudjatma, M. Indrawan, B. Haryanto, et al., Review and analysis of current responses to COVID-19 in Indonesia: period of January to March 2020, *Progress in disaster science* 6 (2020), 100091, <https://doi.org/10.1016/j.pdisas.2020.100091>.
- [4] K.B. Son, T.J. Lee, S.S. Hwang, Disease severity classification and COVID-19 outcomes, Republic of Korea, *Bull. World Health Organ.* 99 (1) (2021) 62–66, <https://doi.org/10.2471/blt.20.257758>.
- [5] T.D. Suryananda, R. Yudhawati, Association of serum KL-6 levels on COVID-19 severity: a cross-sectional study design with purposive sampling, *Ann. Med. Surg.* 69 (2021), 102673, <https://doi.org/10.1016/j.amsu.2021.102673>, 2012.
- [6] R. Seligman, L.F. Ramos-Lima, A. Oliveira Vdo, C. Sanvicente, E.F. Pacheco, K. Dalla Rosa, Biomarkers in community-acquired pneumonia: a state-of-the-art review, *Clinics* 67 (11) (2012) 1321–1325, [https://doi.org/10.6061/clinics/2012\(11\)17](https://doi.org/10.6061/clinics/2012(11)17).
- [7] R.M. Narvaez-Rivera, A. Rendon, M.C. Salinas-Carmona, A.G. Rosas-Taraco, Soluble RAGE as a severity marker in community acquired pneumonia associated sepsis, *BMC Infect. Dis.* 12 (2012) 15, <https://doi.org/10.1186/1471-2334-12-15>.
- [8] L.J. Sparvero, D. Asafu-Adjei, R. Kang, D. Tang, N. Amin, J. Im, et al., RAGE (Receptor for Advanced Glycation Endproducts), RAGE ligands, and their role in cancer and inflammation, *J. Transl. Med.* 7 (2009) 17, <https://doi.org/10.1186/1479-5876-7-17>.
- [9] T. Chavakis, A. Bierhaus, P.P. Nawroth, RAGE (receptor for advanced glycation end products): a central player in the inflammatory response, *Microb. Infect.* 6 (13) (2004) 1219–1225, <https://doi.org/10.1016/j.micinf.2004.08.004>.
- [10] R. Clynes, B. Moser, S.F. Yan, R. Ramasamy, K. Herold, A.M. Schmidt, Receptor for AGE (RAGE): weaving tangled webs within the inflammatory response, *Curr. Mol. Med.* 7 (8) (2007) 743–751, <https://doi.org/10.2174/156652407783220714>.
- [11] G. Mathew, R. Agha, Stross 2021: strengthening the reporting of cohort, cross-sectional and case-control studies in surgery, *Int. J. Surg.* 96 (2021), 106165, <https://doi.org/10.1016/j.ijvs.2021.106165>.
- [12] A. García-Salido, G. Melen, V. Gómez-Piña, G. Oñoro-Otero, A. Serrano-González, J. Casado-Flores, et al., Circulating soluble RAGE and cell surface RAGE on peripheral blood mononuclear cells in healthy children, *J. Pediatr. Endocrinol. Metab.* : JPEM (J. Pediatr. Endocrinol. Metab.) 31 (6) (2018) 649–654, <https://doi.org/10.1515/jpem-2017-0512>.
- [13] A. Lim, A. Radujkovic, M.A. Weigand, U. Merle, Soluble receptor for advanced glycation end products (sRAGE) as a biomarker of COVID-19 disease severity and indicator of the need for mechanical ventilation, ARDS and mortality, *Ann. Intensive Care* 11 (1) (2021) 50, <https://doi.org/10.1186/s13613-021-00836-2>.
- [14] M. Jabaudon, P. Berthelin, T. Pranal, L. Roszyk, T. Godet, J.S. Faure, et al., Receptor for advanced glycation end-products and ARDS prediction: a multicentre observational study, *Sci. Rep.* 8 (1) (2018) 2603, <https://doi.org/10.1038/s41598-018-20994-x>.
- [15] M.A. van Zoelen, M. Schouten, A.F. de Vos, S. Florquin, J.C. Meijers, P.P. Nawroth, et al., The receptor for advanced glycation end products impairs host defense in pneumococcal pneumonia, *J. Immunol.* (Baltimore, Md : 1950) 182 (7) (2009) 4349–4356, <https://doi.org/10.4049/jimmunol.0801199>.
- [16] R.J. Mason, Pathogenesis of COVID-19 from a cell biology perspective, *Eur. Respir. J.* 55 (4) (2020), <https://doi.org/10.1183/13993003.00607-2020>.
- [17] M.J. Mondrinos, P.A. Kennedy, M. Lyons, C.S. Deutschman, L.E. Kilpatrick, Protein kinase C and acute respiratory distress syndrome, *Shock* 39 (6) (2013) 467–479, <https://doi.org/10.1097/SHK.0b013e318294f85a>.
- [18] H. Horiguchi, T.J. Loftus, R.B. Hawkins, S.L. Raymond, J.A. Stortz, M.K. Hollen, et al., Innate immunity in the persistent inflammation, immunosuppression, and catabolism syndrome and its implications for therapy, *Front. Immunol.* 9 (2018) 595, <https://doi.org/10.3389/fimmu.2018.00595>.
- [19] M. Carcatera, C. Caruso, Alveolar epithelial cell type II as main target of SARS-CoV-2 virus and COVID-19 development via NF-Kb pathway deregulation: a physio-pathological theory, *Med. Hypotheses* 146 (2021), 110412, <https://doi.org/10.1016/j.mehy.2020.110412>.