



# Source details

## Advances in Respiratory Medicine

Formerly known as: Pneumonologia i Alergologia Polska

Scopus coverage years: from 2015 to Present

Publisher: Via Medica

ISSN: 2451-4934 E-ISSN: 2543-6031

Subject area: Medicine: Pulmonary and Respiratory Medicine

Source type: Journal

CiteScore 2021

1.4



SJR 2021

0.251



SNIP 2021

0.455



[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### i Improved CiteScore methodology



CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▼

$$1.4 = \frac{355 \text{ Citations 2018 - 2021}}{259 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$1.6 = \frac{399 \text{ Citations to date}}{250 \text{ Documents to date}}$$


Last updated on 05 October, 2022 • Updated monthly



## CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine		
Pulmonary and Respiratory Medicine	#102/140	27th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

# Advances in Respiratory Medicine


<b>COUNTRY</b> Poland  Universities and research institutions in Poland	<b>SUBJECT AREA AND CATEGORY</b> Medicine └ Pulmonary and Respiratory Medicine	<b>PUBLISHER</b> Via Medica
<b>H-INDEX</b> <b>17</b>	<b>PUBLICATION TYPE</b> Journals	<b>ISSN</b> 25436031, 24514934
<b>COVERAGE</b> 2015-2020	<b>INFORMATION</b> <a href="#">Homepage</a> <a href="#">How to publish in this journal</a>	



**Advances in Respiratory Medicine**

**Q3** Pulmonary and Respiratory Medicine  
best quartile

**SJR 2019**  
0.29



powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```



# Advances in Respiratory Medicine

Formerly *Pneumonologia i Alergologia Polska*  
Edited since 1926

## ORIGINAL RESEARCHES

- Thoracoscopic pleural brushing — an innovative method of pleural sampling in diagnostic medical thoracoscopy
- Effect of yoga on FEV<sub>1</sub>, 6-minute walk distance (6-MWD) and quality of life in patients with COPD group B
- Diagnostic values of bronchodilator response versus 9-question questionnaire for asthma
- The role of genexpert in the diagnosis of tubercular pleural effusion in India
- Sputum culture for the diagnosis of tuberculous pleural effusion: analysis of absolute and incremental yields

## REVIEW ARTICLES

- Current treatment strategies in malignant pleural mesothelioma with a treatment algorithm

## CASE REPORTS

- Neurogenic pulmonary oedema as a rare complication of epileptic seizures
- Oligo-recurrence from anaplastic lymphoma kinase-rearranged lung adenocarcinoma

## CLINICAL VIGNETTES

- A late presentation of an uncommon congenital vascular anomaly
- An expectorated hydatid

## GUIDELINES AND RECOMMENDATIONS

- Acute subglottic laryngitis. Etiology, epidemiology, pathogenesis and clinical picture

The Journal is indexed in the following databases: Index Medicus/Medline, EMBASE, EBSCO, Emerging Sources Citation Index (ESCI), Scopus, Index Copernicus 121,1 (2018), MNiSW 2019 (20 points)

[www.journals.viamedica.pl](http://www.journals.viamedica.pl)





# Advances in Respiratory Medicine

www.journals.viamedica.pl

Formerly Pneumologia i Alergologia Polska

www.ptchp.org

## Editor-in-Chief:

Wojciech J. Piotrowski — Łódź  
wojciech.piotrowski@pneumologia.viamedica.pl

## Vice Editor-In-Chief:

Claudio Pedone — Rome  
claudio.pedone@gmail.com

## Contributing Editors:

Adam Białas — Łódź  
adam.bialas@pneumologia.viamedica.pl  
Maciej Kupczyk — Łódź  
maciej.kupczyk@pneumologia.viamedica.pl  
Sebastian Majewski — Łódź  
sebastian.majewski@pneumologia.viamedica.pl  
Sonu Sahni — New York  
sahni.sonu@gmail.com

## Statistical Editor:

Agnieszka Skoczylas — Warsaw  
Michał Poznański — Łódź  
Łukasz Mokros — Łódź

## Managing Editor:

Anna Młynarczyk

## Editorial Advisory Board

Adam Antczak — Łódź, Poland  
Ewa Augustynowicz-Kopeć — Warsaw, Poland  
Halina Batura-Gabryel — Poznań, Poland  
Andrey Belevskiy — Moscow, Russia  
Wojciech Biernacki — London, Great Britain  
Anna Bręborowicz — Poznań, Poland  
Otto Burghuber — Vienna, Austria  
Ryszarda Chazan — Warsaw, Poland  
Ivane Chkhaidze — Tbilisi, Georgia  
Joanna Chorostowska-Wynimko — Warsaw, Poland  
Elżbieta Chyczewska — Białystok, Poland  
Brendan Cooper — Birmingham, Great Britain  
Sven-Erik Dahlén — Solna, Sweden  
Wilfried De Backer — Antwerp, Belgium  
Anna Doboszyńska — Olsztyn, Poland

Antonio M. Esquinas — Murcia, Spain  
Dorota Górecka — Warsaw, Poland  
Paweł Górski — Łódź, Poland  
Sylvia Hartl — Vienna, Austria  
Raffaele Antonelli Incalzi — Rome, Italy  
Renata Jankowska — Wrocław, Poland  
Sabina Janciauskiene — Hannover, Germany  
Christer Janson — Uppsala, Sweden  
Ewa Jassem — Gdańsk, Poland  
Kozui Kida — Tokyo, Japan  
Jerzy Kozielski — Zabrze, Poland  
Piotr Kuna — Łódź, Poland  
Jan Kuś — Warsaw, Poland  
Henryk Mazurek — Rabka, Poland  
Florin Mihaltan — Bucharest, Romania  
Janusz Milanowski — Lublin, Poland  
Tadeusz Orłowski — Warsaw, Poland  
Bernard Panaszek — Wrocław, Poland  
Władysław Pierzchała — Katowice, Poland  
Tadeusz Płusa — Warsaw, Poland  
Venerino Poletti — Forlì, Italy  
Michał Poznański — Łódź, Poland  
Stephen Rennard — Omaha, United States  
Kazimierz Roszkowski — Warsaw, Poland  
Monika Szturmowicz — Warsaw, Poland  
Paweł Śliwiński — Warsaw, Poland  
Branislava Savic — Belgrade, Serbia  
Nikos Siafakas — Heraclion, Greece  
Dragan Subotic — Belgrad, Serbia  
Adam Torbicki — Otwock, Poland  
Michał Unger — Philadelphia, United States  
Arunas Valiulis — Vilnius, Lithuania  
Martina Vašáková — Prague, Czech Republic  
Jadwiga Wędzicha — London, Great Britain  
Elżbieta Wiatr — Warsaw, Poland  
Dariusz Ziara — Zabrze, Poland  
Zofia Zwolska — Warsaw, Poland

## Past Editors-in-Chief (most recent first)

Monika Szturmowicz — Warsaw  
Dorota Górecka — Warsaw  
Elżbieta Wiatr — Warsaw  
Tadeusz Płusa — Warsaw

## Opinions presented in the articles not necessarily represent the opinions of the Editors

**Advances in Respiratory Medicine** (ISSN 2451-4934) is published by VM Media sp. z o.o. VM Group sp.k., ul. Świętokrzyska 73, 80-180 Gdańsk phone: +48 58 320 94 94, fax +48 58 320 94 60, e-mail: redakcja@viamedica.pl, dim@viamedica.pl  
http://www.viamedica.pl, wap.viamedica.pl  
project manager: ewa.reda@viamedica.pl

**Editorial Address:** Biuro ZG PTChP, ul. Wronia 45, lok. 132, Warszawa, Poland

**Subscription Rates:** Paper subscription, 6 issues incl. package and postage individual 60€

Paper subscription, 6 issues incl. package and postage institutional 90€. The above prices are inclusive of regular postage costs.

Payment should be made to: Fortis Bank Polska SA, Gdańsk, Poland, Acc.: PL 15 1600 1303 0004 1007 1035 9001; SWIFT: PPABPLPK.

Single issues, subscriptions orders and requests for sample copies should be sent

to e-mail: prenumerata@viamedica.pl

Electronic orders option available at: www.um.viamedica.pl

**Advertising:** For details on media opportunities within this journal please contact the advertising sales department, ul. Świętokrzyska 73, 80-180 Gdańsk, Poland, phone: +48 58 326 78 20; e-mail: dsk@viamedica.pl

**The Editors accept no responsibility for the advertisement contents.**

All rights reserved, including translation into foreign languages. No part of this periodical, either text or illustration, may be used in any form whatsoever. It is particularly forbidden for any part of this material to be copied or translated into a mechanical or electronic language and also to be recorded in whatever form, stored in any kind of retrieval system or transmitted, whether in an electronic or mechanical form or with the aid of photocopying, microfilm, recording, scanning or in any other form, without the prior written permission of the publisher. The rights of the publisher are protected by national copyright laws and by international conventions, and their violation will be punishable by penal sanctions.

Legal note: [https://journals.viamedica.pl/pneumologia\\_i\\_alergologia\\_pol/about/legalNote](https://journals.viamedica.pl/pneumologia_i_alergologia_pol/about/legalNote)

Editorial policy and information for authors available on [www.piap.viamedica.pl](http://www.piap.viamedica.pl).

**Polish Ministry of Science and Higher Education score: 13 pts.**

**There are 5 educational pts. for the journal's subscription\***



(\* based upon the Minister of Health regulation of October 6, 2004 on a duty of professional training of physicians and dentist practitioners; Dz.U.04.231.2326 of Oct 22, 2004)



Copyright © 2019 PTChP

## Contents

### ORIGINAL RESEARCHES

**Thoracoscopic pleural brushing — an innovative method of pleural sampling in diagnostic medical thoracoscopy**

Yuvarajan Sivagnaname, Praveen Radhakrishnan, Antonious Maria Selvam..... 257

**Effect of yoga on FEV<sub>1</sub>, 6-minute walk distance (6-MWD) and quality of life in patients with COPD group B**

Resti Yudhawati, Mariani Rasjid Hs..... 261

**Diagnostic values of bronchodilator response versus 9-question questionnaire for asthma**

Mahdi Yadollahzadeh, Seyed Mohammadreza Hashemian, Arda Kiani, Fariba Ghorbani, Katayoun Najafizadeh, Fatemeh Razavi, Atefeh Abedini..... 269

**The role of genexpert in the diagnosis of tubercular pleural effusion in India**

Anushree Chakraborty, Swapna Ramaswamy, Akshata Jayachamrajpura Shivananjiah, Raghu Bokkikere Puttaswamy, Nagaraja Chikkavenkatappa ..... 276

**Sputum culture for the diagnosis of tuberculous pleural effusion: analysis of absolute and incremental yields**

Sevak Keshishyan, Viren Kaul, Anupam Gupta, Chul Ahn, Wilbert Aronow, Oleg Epelbaum ..... 281

### REVIEW ARTICLES

**Current treatment strategies in malignant pleural mesothelioma with a treatment algorithm**

Mutlay Sayan, Mehmet Fuat Eren, Apar Gupta, Nisha Ohri, Ayse Kotek, Ibrahim Babalioglu, Sedenay Oskeroğlu Kaplan, Ozge Duran, Ozlem Derinalp Or, Funda Cukurcayir, Neslihan Kurtul, Beyhan Ceylaner Bicakci, Tugce Kutuk, Sukran Senyurek, Ali Turk, Salma K Jabbour, Banu Atalar..... 289

### CASE REPORTS

**Neurogenic pulmonary oedema as a rare complication of epileptic seizures**

Milan Sova, David Franc, Filip Ctvrtlik, Petr Jakubec, Amjad Ghazal Asswad, Vitezslav Kolek ..... 298

**Oligo-recurrence from anaplastic lymphoma kinase-rearranged lung adenocarcinoma**

Yuika Sasatani, Shinichiro Okauchi, Gen Ohara, Katsunori Kagohashi, Kesato Iguchi, Koji Kawai, Hiroaki Satoh..... 301

### CLINICAL VIGNETTES

**A late presentation of an uncommon congenital vascular anomaly**

Juvva Kishan Srikanth, Abanti Das, Pranav Ish, Shibdas Chakrabarti, Nitesh Gupta ..... 305

Resti Yudhawati<sup>1</sup>, Mariani Rasjid Hs<sup>2</sup>

<sup>1</sup>Pulmonology and Respiratory Medicine, Faculty of Medicine, Airlangga University, Dr. Soetomo Hospital Surabaya, Surabaya, Indonesia

<sup>2</sup>Pulmonology and Respiratory Medicine, Faculty of Medicine, Tadulako University, Tadulako Hospital Palu, Palu, Indonesia

## Effect of yoga on FEV<sub>1</sub>, 6-minute walk distance (6-MWD) and quality of life in patients with COPD group B

### Abstract

**Introduction:** Yoga is used in the treatment of various diseases, including chronic obstructive pulmonary disease. However, no studies have assessed the effect of yoga on COPD patients in Indonesia. The difference between this study and similar studies completed in other countries lies in the type of yoga exercises completed, the method in which they were completed, and in certain, unique demographic characteristics. This study aims to analyze the effect of yoga on FEV<sub>1</sub>, 6-minute walk distance, and quality of life in patients with COPD group B in Indonesia.

**Material and methods:** This article reflects research done in the form of an experimental study using a randomized controlled trial with pre and post-test control group design. The samples were divided into 2 groups: the treatment group (yoga practice for 1 hour, 2 times a week for 12 weeks) and the control group (untreated with yoga, given lung rehabilitation brochure). Assessment of the effect of yoga exercises on lung function parameters (FEV<sub>1</sub>), 6-minute walk distance and quality of life were used using SGRQ questionnaires in COPD group B.

**Results:** 33 COPD patients fulfilled the inclusion criteria. 30 patients completed the study. Pre and post yoga results were evaluated in the treatment group versus the control group and then further assessed using statistical tests. There was a significant increase in FEV<sub>1</sub>, 6-MWD and quality of life using a SGRQ questionnaire after 12 weeks of yoga ( $p < 0.05$ ) as well as a significant change in FEV<sub>1</sub>, 6-MWD and quality of life in the treatment group ( $p < 0.05$ ) when compared with the control group ( $p > 0.05$ ).

**Conclusions:** Yoga affects FEV<sub>1</sub>, 6-MWD, and quality of life in patients with Group B COPD.

**Key words:** yoga, COPD, FEV<sub>1</sub>, 6-MWD, quality of life, SGRQ

**Adv Respir Med. 2019; 87: 261–268**

### Introduction

Chronic obstructive pulmonary disease (COPD) is one of the non-communicable diseases that is becoming an important public health issue in Indonesia. The morbidity and mortality of COPD patients is associated with periodic exacerbations or worsening of symptoms [1, 2]. As the frequency of exacerbations increases, damage to the lungs proportionately rises in severity. This is ultimately followed by a decrease in lung function [3]. Problems frequently experienced by COPD patients, especially in stages II and III, include exercise de-conditioning, muscle wasting, weight loss, depression, and isolation. Therefore, comprehensive management of COPD symptoms

in patients suffering from the disease is necessary to improve their quality of life [3].

Outside of pharmacological therapy, some COPD patients are managed non-pharmacologically (i.e. by smoking cessation and pulmonary rehabilitation) [3, 4]. The main purpose of pulmonary rehabilitation is to reduce complaints, improve physical ability to perform daily activities, improve emotions, and improve quality of life [3, 5].

Research has shown that exercises such as upper body workouts, Tai Chi, and yoga can improve COPD patients' quality of life [6]. Yoga has been included as a recommended exercise component for pulmonary rehabilitation programs. In addition to the treatment of physical therapy

**Address for correspondence:** Resti Yudhawati, Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga Jalan Mayjend Prof. Dr. Moestopo 47, Surabaya 60131, Indonesia; e-mail: restiy.apji030@gmail.com

DOI: 10.5603/ARM.2019.0047

Received: 07.05.2019

Copyright © 2019 PTChP

ISSN 2451–4934

in industrial rehabilitation programs, it has been proven to improve the coordination of mind and body. Yoga is called a “low impact” exercise that can be adapted to the needs and abilities of practitioners so that it is suitable for anyone. This includes COPD patients through activities such as asana (yoga posture) and pranayama (breathing technique). Short-term studies on yoga practice have reported an increase in pulmonary function parameters, diffusion capacity, improvement in the quality of life, and a reduction in asthma-induced stress. Yoga has also been proven to be an effective tool for coal miners with COPD [6]. This study serves to further research the impact and confirm the beneficial role of yoga in COPD based on the information that has been described above.

## Material and methods

### Study design

This article reflects research done in the form of an experimental study using a randomized controlled trial with pre and post-test control group design. The initial step was made one week before the start of the intervention by performing CAT score and mMRC score assessments to ensure that COPD group B was included in this study. Then, spirometry assessments were conducted using the Koko Legend Spirometer (606055 9.A), and 6-MWD was performed. Every result was recalculated according to GLLI-2012 and add z-score assessment to the analysis [7]. The follow-up steps were carried out one week after completion of the intervention, which lasted 12 weeks. Yoga classes were undertaken in one setting which was determined by the researcher. Classes were led by a certified yoga instructor.

### Statistical analysis

The data normality test was done using the Shapiro-Wilk test. Analysis of variance before and after treatment was carried out using a paired T-test or Wilcoxon Signed Ranks Test. Analysis of variance between yoga groups and control groups was made using the independent sample T-test or Mann-Whitney test.

### Participants

Participants were recruited from clinics and pulmonary rehabilitation centers, or by physician referral. Inclusion criteria comprised men over the age of 40, patients with stable COPD category B, patients that were willing to follow the research criteria properly and patients who agreed to informed consent. Exclusion criteria

included patients suffering from COPD which was accompanied by other diseases (i.e. neoplasm, congestive heart failure and/or asthma). Dropout criteria included subjects who refused to participate in the research, did not undergo the research protocol, and/or experienced exacerbations more than one time.

### Randomization

All participants agreed to be randomized. Stratified randomization was done using baseline measures of spirometry to ensure participant balance in disease severity. They were randomized according to time of attendance. Odd dates were included in the yoga group, and even dates were included in the control group. Participants in both groups continued to receive normal care, and those in the yoga group continued to attend the program. All participants were asked to refrain from learning or practicing anything other than what is taught during yoga interventions. Also, participants in the yoga intervention group signed a confidentiality agreement not to discuss class content with fellow participants in the control group.

### Intervention

Yoga interventions consisted of yoga classes that lasted 60 minutes twice a week, every week for 12 weeks. Yoga instructors must have had formal training. Due to the need for supplemental oxygen in order to maximize the effectiveness for persons with COPD, the class time was limited to 60 minutes. The program minimized the traditional focus on breathing during the beginning of yoga exercises to address issues related to persons with COPD. The management of non-threatening symptoms that focused on raising awareness of the dangers of dyspnea was included. *Ujjayi*, *Kaphalabati*, and *Sitkari* breathing were included. Further, a spiritual focus was introduced to augment a feeling of comfort during meditation. *Savasana* was chosen as the meditative movement exercise. This was based on the expertise of the teacher leading the intervention.

## Results

### Characteristics of research subjects

Table 1 shows the characteristics of these research subjects. The number of research subjects was 33 people, all of them being males. Subjects excluded from the research consisted of 3 people: 2 people from the treatment group, and 1 person from the control group. The num-

**Table 1. Characteristics of research subjects**

Characteristics	Groups		P-value
	Treatment (yoga)	Control	
<b>Age mean (SD)</b>	64.40 (10.453)	65.33 (8.121)	0.787
< 51 years	1 (6.7%)	0 (0.0%)	
51–60 years	4 (26.7%)	5 (33.3%)	
61–70 years	5 (33.3%)	5 (33.3%)	
71–80 years	4 (26.7%)	5 (33.3%)	
≥ 81 years	1 (6.7%)	0 (0.0%)	
<b>IMT mean (SD)</b>	23.59 (5.366)	20.42 (3.770)	0.072
< 18.5 (less)	1 (6.7%)	4 (26.7%)	
18.5–24.9 (normal)	3 (20.0%)	10 (66.7%)	
25–29.9 (excess)	9 (60.0%)	1 (6.7%)	
≥ 30 (obesity)	2 (13.3%)	0 (0.0%)	
<b>Comorbid</b>			
Non comorbid	11 (73.3%)	11 (73.3%)	0.557
Hypertension	1 (6.7%)	1 (6.7%)	
DM	3 (20.0%)	1 (6.7%)	
Coronary heart disease	0 (0.0%)	2 (13.3%)	
<b>Smoking duration mean (SD)</b>	33.21 (8.781)	25.62 (12.984)	0.085
< 20 years	0 (0.0%)	2 (15.4%)	
20–39 years	10 (71.4%)	8 (61.5%)	
≥ 40 years	4 (28.6%)	3 (23.1%)	
Number of cigarettes per day (median/range)	24 (12 until 48)	12 (12 until 30)	0.144
< 20	5 (35.7%)	8 (61.5%)	
20–29	6 (42.9%)	4 (30.8%)	
> 30	3 (21.4%)	1 (7.7%)	
<b>Brinkman Index</b>			
0–199 (light)	0 (0.0%)	2 (15.4%)	0.104
200–599 (medium)	6 (42.9%)	8 (61.51%)	
≥ 600 (high)	8 (57.1%)	3 (23.1%)	
<b>Obstruction degree</b>			
GOLD 1	1 (6.7%)	1 (6.7%)	0.895
GOLD 2	5 (33.3%)	3 (20.0%)	
GOLD 3	4 (26.7%)	5 (33.3%)	
GOLD 4	5 (33.3%)	6 (40.0%)	

ber of subjects that took part in the study until its completion was 30 people. Based on the degree of COPD, in the treatment group there were 5 people in category GOLD 2 and 5 people in category GOLD 4 (33.3%). Meanwhile, in the control group, there were 6 people (40%) in the GOLD 4 obstruction degree.

The results of the normal distribution test using the Shapiro-Wilk test showed that the data

on differences in 6-MWD in the yoga and control groups and the impact data of the yoga group were not normally distributed ( $p < 0.05$ ). Therefore, the paired T-test was used to find out differences in observations before and after treatment was given in each group, except for the difference in 6-MWD using the Wilcoxon Signed Rank Test. Meanwhile, the T-test of 2 free samples was used to observe differences in findings between yoga



**Table 2. Results of the T-test of FEV<sub>1</sub> (L) and FEV<sub>1</sub> (%) before and after treatment given in each group and T-tests among groups**

		Mean (standard deviation)		P-value
		Yoga group	Control group	T-test of 2 free samples
		n = 15	n = 15	
FEV <sub>1</sub> (L)	Before	1.025 (0.507)	0.941 (0.488)	0.647
	After	1.402 (0.629)	1.017 (0.534)	0.082
	Difference	0.377 (0.364)	0.077 (0.273)	0.016
	P-value	0.001	0.295	
	Paired T-test			
FEV <sub>1</sub> (%)	Before	43.53 (20.625)	40.87 (22.309)	0.736
	After	58.93 (22.799)	44.60 (22.344)	0.093
	Difference	15.40 (15.371)	3.733 (10.964)	0.024
	P-value	0.002	0.208	
	Paired T-test			
FEV <sub>1</sub> , Z-score	Before	-3.558 ± 1.089	-3.409 ± 0.986	0.697
	After	-2.613 ± 1.302	-3.162 ± 1.191	0.239
	Different	0.945 ± 0.786	0.247 ± 0.617	0.011
	P-value	0.001	0.144	
	Paired T-test			

and control groups except for the difference in the impact after treatment, which was done using the Mann-Whitney test.

**FEV<sub>1</sub> change**

FEV<sub>1</sub> assessment was carried out in the two groups. Mean FEV<sub>1</sub> in the treatment group before yoga was 1.025 L (0.507), while in the control group was 0.941 L (0.488). FEV<sub>1</sub> re-assessment was carried out after 12 weeks. In the treatment group there was an increase to 1.402 L (0.629) (p = 0.001) while in the control group there was an increase to 1.017 L (0.534) (p = 0.295).

Table 2 shows the results of the paired T-test of FEV<sub>1</sub> (L) and FEV<sub>1</sub> (%) before and after treatment was given in each group and before T-test among these groups. There were significant differences in FEV<sub>1</sub> (L) before and after treatment in the yoga group (p < 0.05), whereas no significant differences were noted in the control group (p > 0.05). The results of the T-test of 2 free samples between the yoga and control groups indicated that there were no significant differences in FEV<sub>1</sub> (L) both before and after treatment (p > 0.05), but there were significant differences in the value of FEV<sub>1</sub> (L) (p < 0.05).

As can be seen in Table 2, the mean FEV<sub>1</sub> (% prediction) in the treatment group before yoga

was 43.53% (20.625). In the control group, it was 40.87% (22.309). FEV<sub>1</sub> re-assessment was carried out after 12 weeks. In the treatment group there was an increase of 58.93% (22.799) (p = 0.002), and in the control group there was an increase of 44.60% (22.344) (p = 0.208). The results of the paired T-test showed significant differences in FEV<sub>1</sub> (%) before and after treatment in the yoga group (p < 0.05), whereas there were no significant differences in the control group (p > 0.05). The results of the T-test of 2 free samples between the yoga and control groups indicated that there were no significant differences in FEV<sub>1</sub> (%) both before and after treatment (p > 0.05), but there were significant differences in the value of FEV<sub>1</sub> (%) (p < 0.05).

**6-MWD change**

As can be seen in Table 3, there was an increase in mean 6-MWD after yoga in the treatment group. The Wilcoxon Signed Rank Test results in the treatment group showed no significant difference in 6-MWD value before and after yoga with a p-value of 0.001. It can be concluded that patients with COPD treated with yoga have a significantly increased 6-MWD value, whereas in the control group there was a decrease in the mean of 6-MWD after 12 weeks. The Wilcoxon Signed

**Table 3. Results of T-test of 6-MWD (m) before and after treatment in each group as well as T-test among groups**

6-MWD (m)	Median (IQR)		P-value Mann- Whitney test
	Yoga group n = 15	Control group n = 15	
Difference	19 (28)	0 (85)	0.016
Before	289 (75)	258 (147)	
After	334 (66)	258 (69)	
P-value Wilcoxon Signed Rank Test	0.001	0.328	

6-MWD — 6-minute walk distance

Rank Test results in the control group showed no significant difference in 6-MWD value before and after yoga with a p-value of 0.328 (NS). It can be concluded that COPD patients who did not get yoga treatment have an insignificantly declining 6-MWD value. The 6-MWD change in the yoga group was greater than in the control group. Test results using Mann-Whitney test showed a significant difference in 6-MWD value change between the two groups with a p-value of 0.016. It can be concluded that patients with COPD treated with yoga have a significantly greater 6-MWD value change when compared with patients suffering from COPD not treated with yoga.

The results of the Wilcoxon test showed significant differences in 6-MWD (m) before and after treatment in the yoga group ( $p < 0.05$ ), whereas there was no significant difference ( $p > 0.05$ ) in the control group. The results of the Mann Whitney test on the difference in the 6-MWD between the yoga and control groups showed significant differences ( $p < 0.05$ ).

### Change in quality of life

Table 4 shows the results of symptoms, activity, impact, and SGRQ total before and after treatment given in each group. In the treatment group after yoga, there was a decrease in the mean value of the SGRQ questionnaire. The paired T-test results in the treatment group showed a significant difference in the SGRQ questionnaire value between before and after yoga with a p-value of  $< 0.001$ . It can be concluded that patients with COPD treated with yoga have a significantly decreased SGRQ questionnaire value, while in the control group there was an insignificantly decreased mean value of the SGRQ questionnaire after 12 weeks. The paired T-test result in the control

group showed no significant difference in the SGRQ questionnaire value in terms of symptoms before and after yoga with a p-value of  $> 0.05$ . It can be then concluded that patients with COPD not treated with yoga have an insignificantly decreased SGRQ questionnaire value in terms of symptoms.

### Discussion

The number of subjects screened was 74 people who are patients with stable outpatient COPD at Asthma-COPD Polyclinics of Dr. Soetomo Hospital Surabaya and Airlangga University Hospital Surabaya. 24 people were excluded, and 17 people refused to participate in the research. 33 people of male sex were randomly divided into two groups. 17 people were placed into the treatment group and 16 people were placed into the control group. Patients who were excluded from this research consisted of 2 people from the treatment group who were unable to regularly exercise according to the schedule, and 1 person from the control group who refused to continue the taking part in the research study. Patients who completed taking part in the research study until it concluded amounted to 30 people.

Our results suggested that  $FEV_1$  change (L) in the treatment group is greater than in the control group. It can be concluded that patients with COPD treated with yoga therapy have a significantly greater change in  $FEV_1$  value (L) compared to COPD patients not treated with yoga therapy.  $FEV_1$  change (% prediction) in the treatment group is greater than in the control group. It can be concluded that patients with COPD treated with yoga therapy have a significantly greater change in  $FEV_1$  value (% prediction) compared with COPD patients who did not get yoga treatment.

This research is in line with meta-analysis [8] literature by Katiyar [9] and Donesky [10] which found that there was a significant increase in  $FEV_1$  in patients with COPD treated with yoga therapy for 12 weeks. Madanmohan [11] stated that yoga therapy for a 12 week period would increase the maximum expiratory pressure, maximum inspiratory pressure, length of time holding breath during inspiration, length of time holding breath during expiration, and grip strength. This result is also in line with the research by Joshi *et al.* in 1992 which found that there was an increase in ventilation function in the form of respiratory rate reduction, increase in FVC,  $FEV_1$ , MVV and ability to hold breath. during 6 weeks of *Pranayama* exercises.

**Table 4. Result of symptoms, activity, impact and SGRQ total before and after treatment given in each group**

		Mean (standard deviation)		P-value
		Yoga group	Control group	T-test of 2 free samples
		n = 15	n = 15	
Symptoms	Before	54.26 (16.257)	50.63 (21.031)	0.601
	After	23.26 (14.941)	48.91 (26.145)	0.003
	Difference	-30.99 (21.398)	-1.73 (17.155)	< 0.0001
	P-value	< 0.0001	0.703	
	Paired T-test			
Activity	Before	56.08 (21.860)	56.27 (23.602)	0.981
	After	28.62 (24.037)	54.48 (21.843)	0.005
	Difference	-27.46 (21.481)	-1.80 (12.999)	0.001
	P-value	< 0.0001	0.601	
	Paired T-test			
Impact	Before	37.20 (18.784)	48.29 (23.259)	0.162
	After	13.21 (12.787)	43.85 (22.605)	
	Difference	-23.99 (14.533)	-4.44 (14.851)	0.001
	P-value	< 0.0001	0.266	
	Paired T-test			
Total score	Before	45.76 (16.318)	51.10 (21.135)	0.445
	After	19.55 (14.305)	47.93 (20.405)	< 0.0001
	Difference	-26.20 (14.514)	-3.17 (12.884)	< 0.0001
	P-value	< 0.0001	0.357	
	Paired T-test			
<b>Median (IQR)</b>				<b>Mann-Whitney test</b>
Impact	After	8.14 (17.48)	45.63 (38.07)	0.001

The effect of yoga that was demonstrated in this research study relates to the deep breathing technique (*Pranayama*) and meditation, leading to a respiratory rate reduction. This has many beneficial effects: it enables the modulation of airway reactivity, improves breathing sensation through regulating breathing pattern, reduces oxygen consumption, reduces the incidence of hypoxia and hypercapnia, improves blood oxygenation without increasing ventilation, improves respiratory resistance, improves muscle strength and modulates autonomic function by decreasing heart rate at rest and sympathetic activity [3, 10].

The result of this research is in line with the results obtained by Katiyar [9] and Donesky [10] which showed that there was a significant increase in 6-MWD in a COPD group which took part in yoga exercises. Ranjita [12] also obtained a similar result showing a significant increase in

COPD patients symptoms after completing yoga exercises over a 12 week period. The increase in 6-MWD is due to the effect of yoga on the musculoskeletal and cardiorespiratory systems, improving cardiovascular efficiency and homeostatic control of the body. The effect on muscle during continuous yoga stretch posture helps to increase oxidative capacity and skeletal muscle strength, flexibility, endurance, coordination, power, static and dynamic balance. It also helps to reduce glycogen utilization which improves physical performance, walking speed, and step length. Relaxation techniques have been found to increase cardiopulmonary resistance through body and breathing control, which manifests clinically as lung capacity increase, oxygen delivery increase and breathing rate reduction. Therefore, overall, it can increase exercise capacity at 6-MWD [12].

Quality of life is assessed by the SGRQ questionnaire which consists of symptoms, activity, impact, and total score of each group. The assessment of the quality of life increases when the SGRQ value is decreased. The British Thoracic Society (BTS) recommends using SGRQ as it is more sensitive in assessing clinical change. The SGRQ questionnaire can also be used to detect the response to medical treatment or non-medical treatment, as is the case in a lung rehabilitation program. Clinical change is considered to be significant if the SGRQ value decreases by 4% [4].

The SGRQ change in the yoga group is greater than in the control group in each component of the SGRQ questionnaire. The test result showed a significant difference in the change of SGRQ questionnaire value (symptom, activity, impact, total score) with a p-value of  $< 0.05$ . It can be concluded that patients with COPD treated with yoga have a significantly bigger decrease in SGRQ questionnaire values in some components (symptom, activity, impact and total score) compared to patients with COPD untreated with yoga.

This research is in line with previous completed studies. In the investigation completed by Fulambarker [13] and Katiyar [9], there was a significant change in the quality of life through the assessment of the SGRQ questionnaire in the group given yoga treatment for 12 weeks (24 meetings). There was a decrease in the mean value of SGRQ in the treatment group for 12 weeks in terms of symptoms, activity, impact, and total score. It means that there is an improvement in the quality of life in the treatment group. Lacasse *et al.*, as part of their research via a meta-analysis, concluded that pulmonary rehabilitation will reduce shortness of breath symptoms and improve COPD patient's ability to be active, which will improve functional capacity and quality of life [14]. Berry *et al.* explained that pulmonary rehabilitation would increase the maximum oxygen consumption and maximum working capacity, thereby increasing the functional capacity and quality of life [15].

This study also consistent with Kulpati *et al.* Patients who received yoga training were seen to have the best maintenance of function, reduction in the respiratory rate ( $p < 0.001$ ) and heart rate ( $p < 0.01$ ) compared with a group who received conventional therapy, while a group who received breathing exercise that only intermediate in their response, probably indicated the efficacy of yogasanas [16].

Yoga exercise will improve the functioning of bodily systems, namely in the nervous system

and in all organs of the body. It also affects the psychological and spiritual aspects. Yoga exercise is a process of integrating aspects of oneself, including physical, psychological, and spiritual aspects. The optimal integration process enables a person to achieve the quality of life that they expect. A healthy body condition is indicated by the enhancement of psychological well-being (positive psychological conditions such as mood and happiness improve, whilst at the same time decreasing negative psychological symptoms including stress, anxiety, and depression). Practicing yoga also affects spiritual aspects. It allows for greater self-respect, respect of others and the surrounding environment, allows one to be closer to God and to have a meaningful life [17].

This study is consistent with Villien F's research [18], which concluded that changes in breathing patterns could be used as a measure of participating in meditational interventions. However, it is possible that emotions can mediate the relationship between respiratory rate and level of mindfulness. Respiratory patterns have been demonstrated to be stable over time and fluctuate in a consistent manner during changes in respiratory demand, such as exercise. Despite this stability, respiration patterns have been shown to change at rest in response to meditational practices in persons without pulmonary disease. After attending a meditational intervention which focused on *Ujjayi* breathing, it was found that healthy persons demonstrated a decrease in respiratory rate and an increase in expiratory time [18].

## Conclusions

Yoga has beneficial effects in COPD Group B. FEV<sub>1</sub>, 6-MWD, and the quality of life increase after yoga exercise. Therefore, yoga can be used as an option for pulmonary rehabilitation in patients with COPD category B. The pulmonary rehabilitation program should be given continuously in patients with COPD. However, pulmonary rehabilitation needs to be performed together with the instructor within a set exercise schedule in order to achieve the proper movements that correspond to the set procedures.

## Ethical clearance

This study follows the principles of the Declaration of Helsinki. This study has received ethical clearance from Dr. Soetomo General Hospital before the study began (Ethical Clearance Number

300/Panke.KKE/IV/2017). All subjects gave their informed consent prior to their inclusion in the study. Before signing the informed consent, information for informed consent was given. Details that might disclose the identity of the subjects who took part in the study were omitted.

### Authors' contributions

MA and RY designed the study. MA and RY collected samples. MA and RY gathered data. MA and RY analyzed the data. MA and RY made Tables and Figures. MA and RY wrote the manuscript, and all authors contributed to review and revision and have been approved the final version.

### Acknowledgment

We are truly thankful to the patients who participated in the study and the authorities and staff of Dr. Soetomo Hospital, Surabaya, Indonesia, who helped and supported us during the study.

### Conflict of interest

All authors declare no conflict of interest.

### References:

- World Health Organization. Chronic respiratory diseases. Burden of COPD. Available at: <https://www.who.int/respiratory/copd/burden/en> [access: 14.09.2019].
- Sethi S, Evans N, Grant BJB, et al. New strains of bacteria and exacerbations of chronic obstructive pulmonary disease. *N Engl J Med.* 2002; 347(7): 465–471, doi: [10.1056/NEJMoa012561](https://doi.org/10.1056/NEJMoa012561), indexed in Pubmed: [12181400](https://pubmed.ncbi.nlm.nih.gov/12181400/).
- Global Initiative for Chronic Obstructive Lung Disease. GOLD 2017 Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease. Available at: <https://goldcopd.org>. [access: 14.09.2019].
- Ikalius Y, Yunus F, Suradi F, et al. Changes in the quality of life and functional capacity of patients with chronic obstructive pulmonary disease after pulmonary rehabilitation. *Maj Kedokt Indones.* 2007; 57(12): 447.
- Duerden M. The management of chronic obstructive pulmonary disease. *MeRec Bull.* 2006; 16(5): 17–20.
- Chandra F. Respiratory practices in yoga. *Behavioral and Psychological Approaches to Breathing Disorders.* 1994: 221–232, doi: [10.1007/978-1-4757-9383-3\\_16](https://doi.org/10.1007/978-1-4757-9383-3_16).
- Quanjer PH, Stanojevic S, Cole TJ, Stocks J. GLL-2012 Desktop Software for Individual Calculations. Version 3.3.1 build 5. 2014. Available at: <https://www.ers-education.org/guidelines/global-lung-function-initiative/spirometry-tools/desktop-individual-calculator.aspx> [access: 10.10.2019].
- Liu XC, Pan L, Hu Q, et al. Effects of yoga training in patients with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *J Thorac Dis.* 2014; 6(6): 795–802, doi: [10.3978/j.issn.2072-1439.2014.06.05](https://doi.org/10.3978/j.issn.2072-1439.2014.06.05), indexed in Pubmed: [24977005](https://pubmed.ncbi.nlm.nih.gov/24977005/).
- Katiyar SK, Bihari S. Role of pranayama in rehabilitation of COPD patients - a randomized controlled study. *Indian J Allergy Asthma Immunol.* 2006; 20(2): 98–104.
- Donesky-Cuenco D, Nguyen HQ, Paul S, et al. Yoga therapy decreases dyspnea-related distress and improves functional performance in people with chronic obstructive pulmonary disease: a pilot study. *J Altern Complement Med.* 2009; 15(3): 225–234, doi: [10.1089/acm.2008.0389](https://doi.org/10.1089/acm.2008.0389), indexed in Pubmed: [19249998](https://pubmed.ncbi.nlm.nih.gov/19249998/).
- Thombre DP, Balakumar B, Nambinarayanan TK, et al. Effect of yoga training on reaction time, respiratory endurance and muscle strength. *Indian J Physiol Pharmacol.* 1992; 36(4): 229–233, indexed in Pubmed: [1291472](https://pubmed.ncbi.nlm.nih.gov/1291472/).
- Ranjita R, Hankey A, Nagendra HR, et al. Yoga-based pulmonary rehabilitation for the management of dyspnea in coal miners with chronic obstructive pulmonary disease: A randomized controlled trial. *J Ayurveda Integr Med.* 2016; 7(3): 158–166, doi: [10.1016/j.jaim.2015.12.001](https://doi.org/10.1016/j.jaim.2015.12.001), indexed in Pubmed: [27545747](https://pubmed.ncbi.nlm.nih.gov/27545747/).
- Fulambarker A, Farooki B, Kheir F, et al. Effect of yoga in chronic obstructive pulmonary disease. *Am J Ther.* 2012; 19(2): 96–100, doi: [10.1097/MJT.0b013e3181f2ab86](https://doi.org/10.1097/MJT.0b013e3181f2ab86), indexed in Pubmed: [21048431](https://pubmed.ncbi.nlm.nih.gov/21048431/).
- Lacasse Y, Wong E, Guyatt GH, et al. Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. *Lancet.* 1996; 348(9035): 1115–1119, doi: [10.1016/S0140-6736\(96\)04201-8](https://doi.org/10.1016/S0140-6736(96)04201-8), indexed in Pubmed: [8888163](https://pubmed.ncbi.nlm.nih.gov/8888163/).
- Berry MJ, Rejeski WJ, Adair NE, et al. Exercise rehabilitation and chronic obstructive pulmonary disease stage. *Am J Respir Crit Care Med.* 1999; 160(4): 1248–1253, doi: [10.1164/ajrcm.160.4.9901014](https://doi.org/10.1164/ajrcm.160.4.9901014), indexed in Pubmed: [10508815](https://pubmed.ncbi.nlm.nih.gov/10508815/).
- Kulpati DD, Kamath RK, Chauhan MR. The influence of physical conditioning by yogasanas and breathing exercises in patients of chronic obstructive lung disease. *J Assoc Physicians India.* 1982; 30(12): 865–868, indexed in Pubmed: [7184909](https://pubmed.ncbi.nlm.nih.gov/7184909/).
- Kinasih A. The influence of yoga practice on improving the quality of life. *Bull Psychol.* 2002; 18(1): 1–12.
- Villien F, Yu M, Barthélémy P, et al. Training to yoga respiration selectively increases respiratory sensation in healthy man. *Respir Physiol Neurobiol.* 2005; 146(1): 85–96, doi: [10.1016/j.resp.2004.11.010](https://doi.org/10.1016/j.resp.2004.11.010), indexed in Pubmed: [15733782](https://pubmed.ncbi.nlm.nih.gov/15733782/).



**KOMITE ETIK PENELITIAN KESEHATAN  
RSUD Dr. SOETOMO SURABAYA**

**KETERANGAN KELAIKAN ETIK  
(" ETHICAL CLEARANCE ")**

**300 / Panke.KKE / IV / 2017**

**KOMITE ETIK RSUD Dr. SOETOMO SURABAYA TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA PENELITIAN DENGAN JUDUL :**

**" Pengaruh Yoga terhadap FEV1, 6-Minute Walk Distance (6-MWD) dan Kualitas Hidup pada Pasien Penyakit Paru Obstruktif Kronik Kelompok B "**

**PENELITI UTAMA : Resti Yudhawati, dr., Sp.P (K)**

**PENELITI LAIN : Mariani Rasjid HS, dr**

**UNIT / LEMBAGA / TEMPAT PENELITIAN : RSUD Dr. Soetomo Surabaya**

**DINYATAKAN LAIK ETIK**

**SURABAYA, 19 APR 2017**



**(Dr. Elizeus Hanindito, dr., Sp.An, KIC, KAP)**  
**NIP. 19511007 197903 1 002**