

In Vivo Study of Sealing Capability of Raw Propolis Extract and Calcium Hydroxide on Dentin Surface

by Muhammad Roelianto

Submission date: 22-Jan-2022 08:46PM (UTC+0800)

Submission ID: 1745937105

File name: Raw_Propolis_Extract_and_Calcium_Hydroxide_on_Dentin_Surface.pdf (4.65M)

Word count: 3325

Character count: 17793

13
In vivo Study of Sealing Capability of Raw Propolis Extract and Calcium Hydroxide on Dentin Surface

14

This article was published in the following Dove Press journal:
Clinical, Cosmetic and Investigational Dentistry

Muhamad Roelianto
Dian Agustin Wahjuningrum 
Yusuf Bagus Pamungkas
Nanik Zubaidah
Ryza Indah Permatasari
Eska Dianti

Department of Conservative Dentistry,
Faculty of Dental Medicine, Universitas
Airlangga, Surabaya, Jawa Timur 60132,
Indonesia

Aim: This research aimed to investigate the differences in the sealing capability of propolis extracts and Ca(OH)₂ on dentin surfaces.

Materials and Methods: Eighteen mandible incisors of Wistar rats were prepared at a depth of 0.5 mm after which they were randomly divided into 3 groups (n=6) including control (C), pulp-capped with Ca(OH)₂ group (P1), and indirect pulp-capped with propolis extract group (P2). All the cavities were sealed with glass ionomer restorative cement. Moreover, the sections of the teeth were obtained after six Wistar Rats from each group were sacrificed on the second day after treatment. The rats' incisors were cleaved transversally to the area where Ca(OH)₂ and propolis extract bonded with dentin for the SEM (scanning electron microscope) analysis and examined using microphotographs test with 5000x magnification. All the samples were measured and examined with spectrophotometry test to determine the bonding distance between Ca(OH)₂ and dentin, as well as propolis extract and dentin. The result from the two tests was analyzed with SPSS using an independent *t*-test at *p* < 0.05.

Results: There were significant differences between the calcium hydroxide and propolis extract groups (*p* < 0.05) based on the results obtained in the form of spectrums chemical functional groups of spectrophotometry examination. No new chemical bonding or compound was also observed to have been formed between propolis extract with dentin as well as calcium hydroxide with dentin.

Conclusion: The adaptation or sealing capability of propolis extract was found to be better than calcium hydroxide to the dentin surface.

Keywords: pulp capping, sealing ability, calcium hydroxide, propolis, scanning electron microscope, spectrophotometry

Introduction

Dental caries is an infectious disease affecting dental hard tissues and the process usually starts with the release of chemical product produced by bacteria to damage the inorganic and organic components. This chemical product is organic acids produced as a by-product of carbohydrate metabolism¹ and also has the ability to pass through dentinal tubules thereby affecting the pulp vitality which functions to initiate several defense mechanisms to protect the body from bacterial invasion. The infected dentin in contact with pulp tissue is likely to encourage inflammation and discourage reparative dentin formation. Pulp capping is one of the treatments aimed to sustain pulp tissue vitality and avoid more extensive approaches dictated by extraction or endodontic treatment.^{2,3} The process can be applied as a direct pulp cap with the medicament placed directly over the exposed pulp, or as an indirect

Correspondence: Dian Agustin Wahjuningrum
Department of Conservative Dentistry,
Faculty of Dental Medicine, Universitas
Airlangga, Mayjen Prof. Dr. Moestopo 47,
Surabaya, Jawa Timur 60132, Indonesia
Tel +62 315030252
Fax +62 315022472
Email dian-agustin-w@fkg.unair.ac.id


11

submit your manuscript | www.dovepress.com

    
<http://dx.doi.org/10.2147/CCIDE.S243714>

Clinical, Cosmetic and Investigational Dentistry 2020: 12 335–341

335

 © 2020 Roelianto et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at <https://www.dovepress.com/terms.php> and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0) License (<http://creativecommons.org/licenses/by-nc/3.0/>). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (<https://www.dovepress.com/terms.php>).

pulp cap, in which a cavity liner or sealer is placed over the affected dentin. This treatment is applied to maintain pulp vitality⁴ and the possibility of failure in the process is mainly caused by bacterial contamination due to the microleakage under the capping materials.

Microleakage is a phenomenon where there is no bond formation or release of adhesions between the restorative material and the tooth structure, enamel, and dentin. It is, however, possible to have good adaptations or sealing capability with dentin surface and prevent the micro-leakage through the use of a material with power of attachment or adhesion to create an optimal adaptation between the two substances due to the inherent tension force existing between them.⁵ This is achievable either mechanically or chemically, with the chemical aspect observed to be attached to the diffusion and absorption or ion exchange mechanism initiated by the material's contact with the dental tissues. Meanwhile, the mechanical attachment mechanism involves the material being mechanically interlocked into the micro undercut on the tooth surface.⁶ This, therefore, means the sealing capability of pulp capping material on the dentin surface plays an important role in preventing bacterial contamination by forming a bridge.

Calcium hydroxide ($\text{Ca}(\text{OH})_2$) is widely used as a pulp-cap material but it has been discovered not to have a good adaptation to the dentin surface nor does it consistently stimulate the differentiation of odontoblasts and has also proven to be cytotoxic to cell cultures.⁷ Moreover, it is not stable under restorations after long-term usage, thereby, leaving the cavity unprotected and the restoration unsupported. This happens when $\text{Ca}(\text{OH})_2$ has a contact with dentinal fluid which dissolves the material and affect its effectiveness.⁸ The solubility of calcium hydroxide is very low which is 0.16 grams in 100 mL of water at 30° C and this decomposes to produce insoluble particles including hydroxyl ions (OH^-) and calcium ions (Ca^{2+}).⁹ The hydroxyl ions make the environment alkaline with the pH increased to 12.5 at 37° C and are antimicrobial.^{10,11} This further affects the adaptation of the compound in the dentin.

Propolis is a resinous yellow-brown to dark brown compound collected by bees (*Apis mellifera*) from tree buds, sap flows, shrubs, or other vegetable sources. It is a potent antimicrobial, antioxidant, and anti-inflammatory agent,¹² and being a natural substance it is believed to have fairly good adaptability to the dentin surface. Propolis also contains flavonoids which allegedly has the ability to produce crystals required to seal the dentinal tubules and reduce fluid movement in the dentin and, consequently, its sensibility.¹³

Propolis extract has been reported to have low cytotoxicity in gingival fibroblasts and this was considered toxic at 92.70 $\mu\text{g}/\text{mL}$ concentration.¹⁴

There are limited studies on propolis as an indirect pulp capping material; therefore, this study was conducted to determine the material with better sealing capability between propolis and calcium hydroxide. An in vivo approach was used with the results obtained through the simulation of the biological conditions found in a live subject because an in vitro study has been previously conducted on the antibacterial attributes of propolis and the finding produced a successful indirect pulp capping.¹⁵

Materials and Methods

Preparation of Propolis Extract

Propolis extract was obtained by maceration from raw Propolis collected from *Apis Mellifera* sp bee farms in Malang-Indonesia. This involved the maceration of 500 grams of raw propolis with 500 mL 96% ethanol in a closed container and the mixture shaken using a shaker at a speed of 80 rpm. The process was stopped after 7 days and the content was filtered and this was followed by 24-hour treatment repeated for another 7 days with the ethanol color observed not to have changed. This, therefore, makes the total maceration time to be 14 days. Moreover, the ethanol solution was evaporated using a vacuum evaporator at 50–60° C to produce a thick brownish liquid from the propolis extract. This was further diluted with distilled water to have 1 mg/mL concentration for the indirect pulp capping material which was later placed as a thin layer on the dentin surface.

Preparation of the Samples

The procedures used in this study were in accordance with the ethical standards instituted by the Responsible Committee on animal care and use experimentation in the Faculty of Dental Medicine No: 141/KKEPK.FKG/IX/2014. The research involved the use of 18 male Wistar rats weighing 200–230 g and aged 10–12 weeks as samples and they were further divided into 3 groups (n=6) including Control (P0) which was without any treatment, cavity pulp-capped with $\text{Ca}(\text{OH})_2$ (Pulpdent, Watertown, USA) known as (P1), and cavity pulp-capped with propolis extract known as (P2). The anesthetic injections used were ketamine HCl 50 mg/kg BB and xylazine HCl 10 mg/kg which were injected intramuscularly in the femoral caudal extremity. Furthermore, the cavity class V

was drilled up to 0.5 mm in depth without exposing the pulp in the mandible incisor using a slow-speed handpiece with fissure diamond bur (Coltene, Altstätten, Switzerland) at 1500 rpm speed before it was irrigated with sterile saline and dried with a cotton pellet. The pulp cap material was applied in a thin layer on the surface of the dentin incisor mandible after which the cavity was sealed using a Dental GC Fuji II Glass Ionomer Restorative Cement. Each group consisted of six samples and observed 2 days after treatment and the mandible teeth were decapitated. All the groups of the experimental animals were provided uniform standard feed during adaptation up to the end of the treatment. On the second day after the treatment, they were sacrificed to retrieve the mandible with the control and treated group of rats anesthetized using ketamine HCl at a dose of 50 mg/kg BB and xylazine HCl 10 mg/kg injected intramuscularly in the femoral caudal extremity. They were laid on the dissecting tray after they showed no painful response to the anesthetization with each leg fixed using a needle. The incisor teeth including the mandible were cut using a blade and fixed in a pot filled with formalin.

SEM Examination

The samples were prepared for SEM examination by cutting the mandible incisor teeth transversally with the area limited to the places covered with the pulp capping materials and dentin. They were fixed and coated before being inserted into the SEM machine. Three points of each sample and the distance between the pulp capping material with dentin were measured and calculated based on the P1 image taken at 5000x magnification. The adhesion of the Ca(OH)₂ and Propolis extract materials to the dentin surface was determined using the Scanning Electron Microscope (Hitachi, Krefeld, Germany) while spectrophotometry (Hitachi, Tokyo, Japan) test was applied to evaluate the chemical bonding.

Spectrophotometry Test

The Wistar rat incisor tooth samples prepared and grounded to a powder were placed into a test machine for examination while KBr powder (potassium bromine) was applied as a conductor.

Statistical Analysis

The SEM and spectrophotometry results were analyzed with SPSS version 21 (IBM, New York, USA) using an independent *t*-test at $p < 0.05$.

Results

The interpretation of Propolis and Ca(OH)₂ pulp capping material SEM image. The microleakage of the group treated with propolis was found to be significantly smaller with 0.64 μm mean value than those treated with Ca(OH)₂ ($P=0.002$) with 1.71 μm mean value.

Discussion

The SEM analysis showed there is a significant difference in the distance between the propolis extract and calcium hydroxide on the surface of the Wistar rats tooth dentin at $p = 0.02$ and mean value of $p < 0.05$. This means there are meaningful differences in the kinds of materials to be used for pulp capping.

The 5000x magnification field of view used in the SEM to obtain a clearer picture for the group treated with Calcium Hydroxide did not affect the results due to the use of a scale to measure the visible distance. The group had a greater distance with 1.71462 μm compared to the group treated with propolis extract which was 0.64581 μm . Moreover, the interpretation of SEM image of calcium hydroxide-treated group shown in Figure 1 describes did not provide a good adaptation to the dentin and was also unstable under restoration leaving unprotected cavity and unsupported restoration.

The solubility of calcium hydroxide in water was also found to be low with 0.16 grams in 100 mL of water at 30°C and this produced a lot of insoluble particles including calcium ions and hydroxyl ions. Its application makes the dentin surface to be wet thereby reducing its sealing ability.^{8,9} Meanwhile, the interpretation of SEM image of propolis extracts treated group shown in Figure 2 showed




Figure 1 SEM image of group treated with Ca(OH)₂ at 5000x magnification.




12
Figure 2 SEM image of group treated with propolis extract at 2000x magnification.

better adaptation to the dentin. Propolis has hydrophilic properties and this makes them have better sealing ability. This observation is in line with Sforcin and Bankova's⁶ findings that the compound has good solubility in water.

The quite good adaptability of the propolis extract with the dentine surface was associated with the bee wax and flavonoids contained in it which was allegedly reported to

have the ability to produce crystals after reacting with dentin which later binds and close the open tubules.¹⁶

The spectrophotometry analysis shown in Figures 3–7 describes the spectrum of functional groups contained in each sample. The treatment group with the propolis extract was observed to have produced a new compound with a mixture of weak chemical bonds including Van Der Waal and hydrogen bonds which are noncovalent and differ in geometry, strength, and specificity.¹⁷ The treatment group with calcium hydroxide did not produce any new compound but also had the same weak chemical bonds¹⁸ with the Van Der Waals obtained from the Ca^{2+} ionic charge ions on calcium hydroxide and the O^- ions on dentin while the hydrogen bond was based on calcium ions from calcium hydroxide and OH^- from dentin.^{18,19} Moreover, the weak chemical and alleged presence of ionic bond between the propolis extract and dentin were associated with the bees' wax with the formula $\text{C}_{13}\text{H}_{27}\text{CO}_2\text{C}_{26}\text{H}_{53}$ and 12 types of flavonoids with different chemical formula but generally having C, H, and O elements found in the propolis extract which reacted with the hydroxyapatite ($\text{Ca}_{10}(\text{PO}_4)_6\text{OH}_2$) contained in the dentine.²⁰



8
Figure 3 Spectrum image of dentin.




Figure 4 Spectrum image of propolis extract.




Figure 5 Spectrum image of dentin treated with propolis extract.




Figure 6 Spectrum image of calcium hydroxide.




Figure 7 Spectrum image of dentin treated with calcium hydroxide.

Conclusion

The sealing capability of propolis as an indirect pulp capping material is better than calcium hydroxide.

1

Disclosure

The authors report no conflicts of interest for this work.

References

1. Featherstone J. Dental caries: a dynamic disease process. *Aust Dent J*. 2008;53:286–291. doi:10.1111/j.1834-7819.2008.00064.x
2. Bansal P, Kapur S, Ajwani P. Effect of mineral trioxide aggregate as a direct pulp capping agent in cariously exposed permanent teeth. *Saudi Endod J*. 2014;4:135–40. doi:10.4103/1658-5984.138146
3. Ho C, Argáez C. Endodontic therapy interventions for root canal failure in permanent dentition: a review of clinical effectiveness, cost-effectiveness, and guidelines; 2017.
4. Hilton TJ. Keys to clinical success with pulp capping: a review of the literature. *Oper Dent*. 2009;34:615–625. doi:10.2341/09-132-0
5. Von Fraunhofer JA. Adhesion and cohesion. *Int J Dent*. 2012;8.
6. Sforzin JM, Bankova V. Propolis: is there a potential for the development of new drugs? *J Ethnopharmacol*. 2011;133:253–260. doi:10.1016/j.jep.2010.10.032
7. Bogen G, Kim JS, Bakland LK. Direct pulp capping with mineral trioxide aggregate. *J Am Dent Assoc*. 2008;139:305–315. doi:10.14219/jada.archive.2008.0160
8. Francisconi LF, Freitas APD, Scaffa PMC, Mondelli RFL, Francisconi PAS. Water sorption and solubility of different self-adhesive cements. *J Appl Oral Sci*. 2009;17:472. doi:10.1590/S1678-77572009000500014
9. Safavi K, Nakayama TA. Influence of mixing vehicle on dissociation of calcium hydroxide in solution. *J Endod*. 2000;26:649–651. doi:10.1097/00004770-200011000-00004
10. Solak H, Öztan MD. The pH changes of four different calcium hydroxide mixtures used for intracanal medication. *J Oral Rehabil*. 2003;30:436–439. doi:10.1046/j.1365-2842.2003.01062.x
11. Komabayashi T, D'souza RN, Dechow PC, Safavi KE, Spångberg LSW. Particle size and shape of calcium hydroxide. *NIH Public Access*. 2009;35:284–287.
12. Garg P, Tyagi SP, Sinha DJ, Singh UP, Malik V, Maccune ER. Comparison of antimicrobial efficacy of propolis, *Morinda citrifolia*, *Azadirachta indica*, triphala, green tea polyphenols and 5.25% sodium hypochlorite against *Enterococcus faecalis* biofilm. *Saudi Endod J*. 2014;4:122–7. doi:10.4103/1658-5984.138141
13. Hongal S, Torwane NA, Goel P, Chandrashekar B. The effect of 30% ethanolic extract of Indian propolis on replica of human dentin compared against commercially available desensitizing agent: a methodological SEM study in vitro. *Pharmacognosy Res*. 2014;6:113. doi:10.4103/0974-8490.129026
14. Mooduto L, Fredline C, Sampoemo G, Goenharto S, Puteri FH, Wahjuningrum DA. Cytotoxicity of sodium hypochlorite, chlorhexidine and propolis on human periodontal ligament fibroblast cell. *J Int Dent Med Res*. 2019;12:476–480.
15. Wahjuningrum DA, Subijanto A. Influence of extract propolis on the adherence of enterococcus faecalis as a candidate root canal irrigation solution. *Int J Sci Clin Lab*. 2015;8:1–11.
16. Sales-peres SHDC, Carvalho FND, Marsicano JA, et al. Effect of propolis gel on the in vitro reduction of dentin permeability. *J Appl Oral Sci*. 2010;318–323.
17. Hubbard RE. Hydrogen bonds in proteins: role and strength. *eLS*. 2001;1–7.
18. Gribova V, Crouzier T, Picart C. A material's point of view on recent developments of polymeric biomaterials: control of mechanical and biochemical properties. *J Mater Chem*. 2011;21:14354–14366. doi:10.1039/c1jm11372k
19. Estrela C, Holland R. Calcium hydroxide: study based on scientific evidences. *J Appl Oral Sci*. 2003;11:269–282. doi:10.1590/S1678-77572003000400002
20. Zheng Y-Z, Zhou Y, Liang Q, Chen D-F, Guo R, Lai R-C. Hydrogen-bonding interactions between apigenin and ethanol/water: a theoretical study. *Sci Rep*. 2016;6:34647. doi:10.1038/srep34647

Clinical, Cosmetic and Investigational Dentistry

Dovepress

Publish your work in this journal

Clinical, Cosmetic and Investigational Dentistry is an international, peer-reviewed, open access, online journal focusing on the latest clinical and experimental research in dentistry with specific emphasis on cosmetic interventions. Innovative developments in dental materials, techniques and devices that improve outcomes and patient

satisfaction and preference will be highlighted. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-cosmetic-and-investigational-dentistry-journal>

In Vivo Study of Sealing Capability of Raw Propolis Extract and Calcium Hydroxide on Dentin Surface

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 ro.uow.edu.au 3%
Internet Source
- 2 Christopher Okunseri. "Patient characteristics and trends in nontraumatic dental condition visits to emergency departments in the United States", *Clinical Cosmetic and Investigational Dentistry*, 01/2012 3%
Publication
- 3 Murlidhar Rajagopalan, Amar Jeet Chitkara, Samir Dalwai, Abhishek De et al. "Burden of Disease, Unmet Needs in the Diagnosis and Management of Atopic Dermatitis: An Indian Expert Consensus", *Clinical, Cosmetic and Investigational Dermatology*, 2021 1%
Publication
- 4 Y Sakata. "Temocapril prevents transition to diastolic heart failure in rats even if initiated after appearance of LV hypertrophy and diastolic dysfunction", *Cardiovascular Research*, 03/01/2003 1%
Publication

5	saudiendodj.com Internet Source	1 %
6	"Modern Operative Dentistry", Springer Science and Business Media LLC, 2020 Publication	1 %
7	Robert Karlinsey. "Remineralization of eroded enamel by a NaF rinse containing a novel calcium phosphate agent in an in situ model: a pilot study", Clinical Cosmetic and Investigational Dentistry, 08/2010 Publication	1 %
8	Hongbin Lei, Botao Wang. "Real-Time Recognition Algorithm of Vehicle External Specific Sounds Based on Spectrum Characteristics", Journal of Physics: Conference Series, 2020 Publication	1 %
9	Shiny Mounica Penumudi, Ramesh Babu Mandava, Disha Saraswathi D, Vajja Santhi, Swetha B, Bhavana Gandhi. "Antimicrobial Efficacy of Herbs in Endodontics", Journal of Advanced Oral Research, 2018 Publication	1 %
10	Karima, Fahd AA, Asma Sultana, Tazdik G Chowdhury, Md Samsul Alam, and Mohammed Wahiduzzaman. "Ledermix Pulp	1 %

treatment in a permanent molar: A case report", Update Dental College Journal, 2014.

Publication

11

Yashika Jain. "A comparison of the efficacy of powered and manual toothbrushes in controlling plaque and gingivitis: a clinical study", Clinical, Cosmetic and Investigational Dentistry, 2013

Publication

<1 %

12

Perihan Oyar, Farzin Asghari Sana, Rukiye Durkan. "Comparison of mechanical properties of heat-polymerized acrylic resin with silver nanoparticles added at different concentrations and sizes", Journal of Applied Polymer Science, 2018

Publication

<1 %

13

www.dovepress.com

Internet Source

<1 %

14

doi.org

Internet Source

<1 %

15

www.thieme-connect.com

Internet Source

<1 %

16

Miloni S Vora, Nidhi K Nihal, Jaya Agali Ramachandra. "Root Canal Irrigants in Primary Teeth", World Journal of Dentistry, 2015

Publication

<1 %

17 Qi-long Sun, Xuguang Zhang, Quan-tai Xing, Peng Ding, Jinbo Feng, Zhanmin Wang, Xiaopeng Wu. "A study of mifepristone/IFN- γ -induced apoptosis of human cholangiocarcinoma cell line FRH-0201 in vitro", *OncoTargets and Therapy*, 2012
Publication

18 Taciana Marco Ferraz Caneppele, Letícia Grilo de Souza, Manuela da Silva Spinola, Felipe Eduardo de Oliveira et al. "Bacterial levels and amount of endotoxins in carious dentin within reversible pulpitis scenarios", *Clinical Oral Investigations*, 2020
Publication

19 Amy LOW, Hamidah MOHD YUSOF, Fazal REZA, Asma ABDULLAH NURUL et al. "Gypsum-based biomaterials: Evaluation of physical and mechanical properties, cellular effects and its potential as a pulp liner", *Dental Materials Journal*, 2015
Publication

20 bmcpregnancychildbirth.biomedcentral.com
Internet Source

21 tessera.spandidos-publications.com
Internet Source

22 Sardana, Divesh, KR InduShekar, Sheetal Manchanda, Bhavna Gupta Saraf, and Neha

Sheoran. "Role of propolis in dentistry: review of the literature", Focus on Alternative and Complementary Therapies, 2013.

Publication

23

Vessela P. Stefanova, Georgi T. Tomov, Snezhana Ts. Tsanova. "Morphological Study Of Border Area Of Pulp-Capping Materials And Er:YAG Laser Prepared Hard Dental Surface", Folia Medica, 2015

Publication

<1 %

24

www.cdejournal.com

Internet Source

<1 %

25

www.elynnmed.com

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

In Vivo Study of Sealing Capability of Raw Propolis Extract and Calcium Hydroxide on Dentin Surface

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
