



# BERKALA ILMU KESEHATAN KULIT DAN KELAMIN (PERIODICAL OF DERMATOLOGY AND VENEROLOGY)

FAKULTAS KEDOKTERAN UNIVERSITAS AIRLANGGA

P-ISSN : 25494082 <> E-ISSN : 25494082 Subject Area : Health, Science

0 Impact Factor

518 Google Citations


Sinta 2 Current Accreditation

- Google Scholar
- Garuda
- Website
- Editor URL

### History Accreditation



Citation Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2017
Citation	518	466
h-index	9	8
i10-index	7	6

e-journal-en

# BERKALA ILMU KESEHATAN KULIT DAN KELAMIN



Nowadays, scientific journals plays a big role in displaying the growth in scientific fields. It functions as a media between researchers and readers, to deliver the outcome of a research.

nu Kesehatan Kulit dan Kelamin (BIKKK) was first published in August 9th, 1989 by Staf Medik Fungsional Dermatovenereology of the Faculty of Medicine, Airlangga University – Dr. Soetomo Academic General Hospital Surabaya.

For more than 30 years, BIKKK has been consistently releasing issues every 4 months, with articles encompassing a variety of topics in Dermatovenereology. At first, BIKKK serve as a platform for the staff and residents of Department/Staf Medik Fungsional Dermatovenereology of the Faculty of Medicine, Airlangga University – Dr. Soetomo Academic General Hospital Surabaya to publish their internal reports and cases. As time went by, it has transformed and broadened its scope, receiving articles from the external parts, but still focusing on the field of Dermatovenereology.

Since 2014, BIKKK has changed it's title to "BIKKK, Berkala Ilmu Kesehatan Kulit dan Kelamin Periodical of Dermatology and Venereology". This abbreviation is intended for the publication to be more internationally acknowledged, and providing an easier way for researchers to cite the journal.

Having been nationally accredited in 2016 (Nomor: 36a/E/KPT/2016, ISSN 1978-4279) and as Sinta 2 (Nomor: 30/E/KPT/2018) in 2018 and then 2020 (Nomor : 200/M/KPT/2020), by the Indonesian Ministry of Research, Technology & Higher Education, BIKKK has begun to receive article submissions via its website, <https://e-journal.unair.ac.id/BIKK>. And starting in 2020, BIKKK has been receiving articles exclusively in English.

BIKKK's editorial board consists of colleagues from Indonesian and International board of dermatovenereology. This is done to ensure that we emphasize the quality of the journal and that we are open to suggestions, in order to improve the journal performance.

We welcome your articles – researchers, pre-graduate students, residents, and lecturers can submit their articles (Clinical research, Review articles, Case report) to our website.

For submission, please kindly open [HERE](#).

## Other Journals




PREVIOUS

MOZAIK HUMANIORA

NEXT

GLOBAL STRATEGIS






**Vol. 33 No. 2 (2021): AUGUST**

**Current Issue**

p-ISSN 1978 - 4279  
e-ISSN 2549 - 4082  
Vol. 33 / No.2 / August 2021

# Berkala Ilmu Kesehatan Kulit dan Kelamin



*Periodical of Dermatology and Venereology*

Vol. 33 No. 2 (2021): AUGUST


Published: 2021-07-31




## Research / Retro

### A Retrospective Study: Risk Factor Analysis of Secondary Bacterial Infection in Pediatric Atopic Dermatitis Patients

 DOI : 10.20473/bikk.V33.2.2021.83-87

 Desiana Widityaning Sari , Sawitri Sawitri , Muhammad Yulianto Listiawan , Dwi Murtiastutik , Linda Astari , Alpha Fardah Athiyyah , Afif Nurul Hidayati

 83-87


 Abstract : 1149


 PDF : 529




### Contact Dermatitis in Tertiary Hospital: A 2-year Retrospective Study

 DOI : 10.20473/bikk.V33.2.2021.88-92

 Efenina Ginting , Damayanti Damayanti , Deasy Fetarayani , Afif Nurul Hidayati

 88-92


 Abstract : 1243


 PDF : 635



### Contact Dermatitis Knowledge Level in Batik Workers of Desa Batik, Tanjung Bumi, Bangkalan, Madura

 DOI : 10.20473/bikk.V33.2.2021.93-95

 Cita Rosita Sigit Prakoeswa , Rahmadewi Rahmadewi , Trisniartami Setyaningrum , Damayanti Damayanti , Hasnikmah Mappamasing , Sylvia Anggraeni , Menul Ayu Umborowati

 93-95

 Abstract : 880


 PDF : 292



### Cutaneous Adverse Drug Reactions in Hospitalized HIV/AIDS Patients

 DOI : 10.20473/bikk.V33.2.2021.96-102

 Yusuf Wibisono , Indropo Agusni , Afif Nurul Hidayati , Rahmadewi Rahmadewi , Maylita Sari , Astindari Astindari , Septiana Septiana , Dwi Murtiastutik

 96-102

 Abstract : 741

 PDF : 313




### In Vitro Comparison of Antifungal Activity between Epigallocatechin Gallate (EGCG) and Nystatin on Candida Sp. Stored Isolates in HIV/AIDS Patients with Oral Candidiasis

 DOI : 10.20473/bikk.V33.2.2021.103-109

 Yusuf Wibisono , Afif Nurul Hidayati , Sawitri Sawitri , Cita Rosita Sigit Prakoeswa , Iskandar Zulkarnain , Evy Ervia , Rahmadewi Rahmadewi , Esti Hendradi , Pepy Dwi Endraswari , Dwi Murtiastutik

 103-109

 Abstract : 1314

 PDF : 722

 PDF

## The Effect of Lifestyle on Skin Aging

 DOI : 10.20473/bikk.V33.2.2021.110-115

 Yasmin Adzra Nabila , Damayanti Damayanti , Samsriyaningsih Handayani , Trisniartami Setyaningrum

 110-115


 Abstract : 2367


 PDF : 802

 PDF

## The Profiles of Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN) Patients in Tertiary Hospital

 DOI : 10.20473/bikk.V33.2.2021.116-122

 William Andrew Isaac , Damayanti Damayanti , Nurmawati Fatimah , Afif Nurul Hidayati

 116-122

 Abstract : 947

 PDF : 510

 PDF

## The Role of Connexin in Cutaneous Adverse Drug Reactions (CADRs) in Patients with Increasing Aspartate Aminotransferase (AST) and Alanine Aminotransferase (ALT)

 DOI : 10.20473/bikk.V33.2.2021.123-128

 Dyah Ayu Mira Oktarina , Gilang Baswara , Hardyanto Soebono

 123-128


 Abstract : 531

 PDF : 219

 PDF

## Risk Factor Profile and Quality of Life of Psoriasis Vulgaris

 DOI : 10.20473/bikk.V33.2.2021.129-134

 As'ad Naufal , Damayanti Damayanti , Ety Hary Kusumastuti , Afif Nurul Hidayati

 129-134

 Abstract : 1141

 PDF : 464


 PDF

## Literature Review

### Clinical Manifestation and Management of Terrestrial Animal Bites

 DOI : 10.20473/bikk.V33.2.2021.135-140

 Emma Hidayati Sasmito , Sawitri Sawitri

 135-140

 Abstract : 793

 PDF : 307


 PDF

## Case Report

### Erythrodermic Manifestation due to Hyperinfestation of Scabies

 DOI : 10.20473/bikk.V33.2.2021.141-144

 Sandra Widaty , Martinus Martinus , Yenny Rachmawati

 141-144

 Abstract : 625


 PDF : 338

 PDF

### Secondary Syphilis During Pregnancy: The Importance of Screening and Clinical Management

 DOI : 10.20473/bikk.V33.2.2021.145-149

 Indah Purnamasari , Jusuf Barakbah , Sunarko Martodiharjo , Dwi Murtiastutik , Maylita Sari , Astindari Astindari , Septiana Septiana , Afif Nurul Hidayati

 145-149

 Abstract : 676

 PDF : 403

 PDF

## Journal Policy

Focus and Scope

Publication Ethics

Peer Review Process

Peer Review

Article Processing Charge

Editorial Team

Open Access Policy


Archiving

Plagiarism

Copyright

Contact

Old Website



### Editorial Team



**Prof. Dr. Cita Rosita Sigit Prakoeswa, dr., Sp.KK(K),  
FINS DV, FAADV**

**Editor In Chief**

Faculty of Medicine, Airlangga University Surabaya, Indonesia, Indonesia

200000-0003-3232-095X  
pL7tjnEAAAAJ&hl=en  
57189894608



**dr. Damayanti Damayanti, Sp.KK(K), FINS DV**

**Editorial Manager**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

-



**dr. Irmadita Citrashanty, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

57204810428



**dr. Maylita Sari, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

-



**dr. Septiana Widyantari, Sp.KK, FINS DV**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

57210805038







**dr. Medhi Denisa Alinda, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

57202009260



-



**dr. Hasnikmah Mappamasing, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

57210793193



-



**dr. Sylvia Anggraeni, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

ugPVNPQAAAAJ&hl

57210634125



-



**Dr. Afif Nurul Hidayati, dr., Sp.KK(K), FINS DV, FAADV**

**Editorial Board**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

ctqbEVQAAAAJ

57196050143



-



**dr. Evy Ervianti, Sp.KK(K), FINS DV, FAADV**

**Editorial Board**

Faculty of Medicine, Airlangga University Surabaya, Indonesia, Indonesia, Indonesia

-

Qsz4JHgAAAAJ

57201074072



-



**dr. Linda Astari, Sp.KK(K), FINS DV, FAADV**

**Editorial Board**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

kLaGucOAAAAJ&hl

57196052464



-





**dr. Astindari Astindari, Sp.KK**

**Editorial Board**

Faculty of Medicine, Airlangga University Surabaya, Indonesia



**Prof. Dr. med. Isaak Effendy**

**Editorial Board**

Academic Teaching Hospital Bielefeld Klinikum Bielefeld, Germany  
Dermatology and Allergology, Germany



7005243490



**Prof. dr. Taruna Ikrar, M.Pharm, MD., Ph.D**

**Editorial Board**

Biomedical Science, The National Health University, California, United States



xtqyDXsAAAAJ&hl  
14621460200



**dr. Sri Manovita Pateda, S.Ked., M.Kes**

**Editorial Board**

Ehime University, Japan, Japan



**dr. Jimmy Rusdian Masjkur, M.Sc., Ph.D**

**Editorial Board**

University Hospital Carl Gustav Carus Dresden, Germany, Germany



36621238000



**Robert Tungadi, S.Si., M.Si., Apt**

**Editorial Board**

Aachen University, Germany, Germany



710v6wAAAA18:hl  
56609263400



**dr. David Sudarto Oeiria, Sp.KK, FINSDV, FAADV**

**Editorial Board**

Division of Dermatologic Surgery, Dept. of Dermatology, Huashan Hospital -  
Fudan University, Shanghai, China



**dr. Yuri Widia, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia



vgSrq4oAAAAJ&hl  
57210799099



**dr. Menul Ayu Umborowati, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia



**dr. Bagus Haryo Kusumaputra, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia



57202017415



**dr. Putri Hendria Wardhani, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia



57210789350



**dr. Regitta Indira Agusni, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia





ID -

JWEo4rIAAAAJ&hl



## Journal Policy

Focus and Scope	Publication Ethics
Peer Review Process	Peer Review
Article Processing Charge	Editorial Team
Open Access Policy	Archiving
Plagiarism	Copyright
Contact	Old Website

## Meet Our Editorial Team



**Prof. Dr. Cita Rosita Sigit Prakoeswa, dr., Sp.KK(K), FINS DV, FAADV**  
Editor In Chief  
Airlangga University Surabaya, Indonesia, Indonesia  
57189894608



**dr. Damayanti Damayanti, Sp.KK(K), FINS DV**  
Editorial Manager  
Airlangga University Surabaya, Indonesia  
-



**dr. Irmadita Citrashanty, Sp.KK**  
EDITORIAL HANDLING  
Airlangga University Surabaya, Indonesia  
57204810428



## Instruction for Author

---

[Author Guidelines](#)

[Online Submission](#)

## Certificate

---



## Author Guidelines

---



## Template

---



## Author Declaration Form

---



## Tools

---



## Visitors

---



## A Retrospective Study: Risk Factor Analysis of Secondary Bacterial Infection in Pediatric Atopic Dermatitis Patients

Desiana Widityaning Sari<sup>1</sup>, Sawitri<sup>1</sup>, M. Yulianto Listiawan<sup>1</sup>, Dwi Murtiastutik<sup>1</sup>, Linda Astari<sup>1</sup>, Alpha Fardah Athiyyah<sup>2</sup>, Afif Nurul Hidayati<sup>1</sup>

<sup>1</sup>Department of Dermatology and Venereology, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Universitas Airlangga, Surabaya

<sup>2</sup>Department of Pediatrics, Faculty of Medicine, Universitas Airlangga, Surabaya

### ABSTRACT

**Background:** Atopic Dermatitis (AD) is a complex multifactorial disease that includes defects in skin architecture, immune dysregulation, and changes of skin flora, and it predominantly occurs in infancy and childhood. The defects in skin barrier structures are mentioned as one of the factors that facilitates bacterial colonization. Bacterial infection in AD can worsen the inflammation. It requires treatment with antibiotics, which takes longer therapy time, higher costs, and ultimately affects the patient's quality of life and his/her family members. **Purpose:** To find out the epidemiology, diagnosis, management of AD, and analyze the risk factors of secondary bacterial infection in new AD patients at the Pediatric Dermatology Division, Dermatovenereology Outpatient Clinic, Dr. Soetomo General Academic Hospital. **Methods:** This was a retrospective study with a total sampling method. The research data were medical records of new AD patients at the Pediatric Dermatology Division, Dermatovenereology Outpatient Clinic, Dr. Soetomo General Academic Hospital Surabaya from January 2012 to December 2018. **Result:** There were 404 new patients with AD at the Pediatric Dermatology Division from January 2012 to December 2018, and 210 of them were accompanied by secondary bacterial infection. There was a correlation between a history of dry skin ( $p=0.000$ ) with the incidence of secondary bacterial infection in AD patients. **Conclusion:** In this study, AD patients with a history of dry skin had a risk of complication such as bacterial secondary infection.

**Keywords:** atopic dermatitis, risk factor, infectious disease.

Correspondence address: Afif Nurul Hidayati, Department of Dermatology and Venereology Faculty of Medicine, Universitas Airlangga / Dr. Soetomo General Academic Hospital / Universitas Airlangga, Surabaya, Jl. Mayjen Prof. Dr. Moestopo No. 6-8 Surabaya 60131, Indonesia. Phone: (031) 5501609, e-mail: affif\_nurulhidayati@fk.unair.ac.id.

### BACKGROUND

Atopic Dermatitis (AD) is a complex multifactorial disease that includes defects in skin barrier structures, immune dysregulation, and changes in skin flora which mostly occur in infancy and childhood. Based on the clinical picture, AD can be divided into 3 forms, namely AD in infants (2 months–2 years), children (2–12 years), and adolescents (over 12 years). In AD, defects in the skin barrier structure, decreased functional integrity, and reduced regenerability affect immune responses and specific inflammatory reactions.<sup>1</sup>

A phase III study by ISAAC (International Study of Asthma and Allergies in Childhood) involving 1.2 million school children from 98 countries reported that the prevalence of AD in children aged 6–7 years was 7.9% and 7.3% in children over 12 years old. This shows an increase in the prevalence of AD, especially in Asia in the last few decades, which is influenced by several factors such as urbanization, changes in

lifestyle and dietary habits, as well as an increase in the standard of living and education. Skin barrier defect is mentioned as one of the factors that facilitates bacterial colonization. Infection in AD skin lesions can worsen the inflammation, and it requires treatment with topical or systemic antibiotics. This would require longer therapy times, higher costs, the risk of antibiotic resistance, and ultimately affecting the quality of life for patients and their families.<sup>2,3,4</sup>

In 2014, a cross-sectional study in China reported that 52.3% of AD patients of all ages also had secondary infections. AD patients over 11 years old have a higher incidence of infection caused by *Staphylococcus aureus* compared to children under age groups. In this study, the ratio of AD patients with secondary bacterial infection is half of the total number of AD patients. Therefore, a secondary bacterial infection in AD is a problem that needs to be addressed because this complication can lead to recurrence in AD. If left untreated, it can lead to several systemic

infections such as osteomyelitis, septicemia, and endocarditis.<sup>5,6,7</sup>

The risk factors for secondary bacterial infection in AD have not been determined. Research in China reported the risk factors associated with the incidence of secondary bacterial infection in AD patients: history of allergies, history of AD in infancy, and involvement of the flexor area. This study analyzes the risk factors associated with the incidence of secondary infection in AD patients, which are individual history of atopy, familial history of atopy, dry skin, and involvement of the flexor area.<sup>8</sup>

**METHODS**

This was an analytic retrospective study aimed to find out the epidemiology, diagnosis, treatment, and analyze the risk factors of secondary bacterial infection in AD patients. The data were obtained from the electronic medical records of AD patients. The data contained identity, history taking, examination, diagnosis, and treatment with a cross-sectional approach. The sample was patients admitted at the Pediatric Dermatology Division, Dermatovenereology

Outpatient Clinic Dr. Soetomo General Academic Hospital from January 2012 to December 2018. The ethical clearance has been obtained from the Ethical Committee of Dr. Soetomo General Academic Hospital Surabaya Number 1758/KEPK/I/2020.

**RESULT**

There were 404 new AD patients at the Pediatric Dermatology Division, Dermatovenereology Outpatient Clinic Dr. Soetomo General Academic Hospital from 2012 to 2018, contributing 13.2% of the patients of the Pediatric Dermatology Division and 1.8% of the total patients admitted in the Dermatovenereology Outpatient Clinic Dr. Soetomo General Academic Hospital. Of the 404 new AD patients, 210 of them were presented with secondary bacterial infection, contributing 6.9% of the patients of the Pediatric Dermatology Division and 0.9% of the total patients admitted to the Dermatovenereology Outpatient Clinic. Thus, the ratio of AD patients and those with secondary bacterial infection was 1: 0.5, and the ratio of AD patients and those presented with other diseases was 1: 0.2.

**Table 1.** The distribution of new atopic dermatitis patients and new atopic dermatitis patients with secondary bacterial infection at the Pediatric Dermatology Division, Dermatovenereology Outpatient Clinic, Dr. Soetomo General Academic Hospital

New Patients	Year							Number (%)
	2012 (%)	2013 (%)	2014 (%)	2015 (%)	2016 (%)	2017 (%)	2018 (%)	
Atopic Dermatitis	84 (20.7)	63 (15.5)	46 (11.3)	60 (14.8)	51 (12.6)	53 (14.1)	47 (11.6)	404 (1.8)
Atopic Dermatitis & Secondary Bacterial Infection	35 (16.6)	24 (11.4)	27 (12.8)	28 (13.3)	23 (10.9)	35 (16.6)	38 (18.0)	210 (0.9)
Atopic Dermatitis & other diseases	17 (18.8)	10 (11.1)	12 (13.3)	10 (11.1)	10 (11.1)	12 (13.3)	19 (21.1)	90 (0.4)
Pediatric Dermatology Division	597 (19.6)	536 (17.6)	489 (16.0)	383 (12.5)	332 (10.9)	340 (11.1)	365 (11.9)	3,042 (14.2)
Dermato Venereology Outpatient Clinic	4211 (19.7)	3840 (17.9)	3358 (15.7)	2673 (12.5)	2545 (11.9)	2556 (12)	2179 (10.2)	21,362 (100)

There were 216 female patients (53.5%) and 188 male patients (46.5%) admitted at the Pediatric Dermatology Division, Dermatovenereology Outpatient Clinic, Dr. Soetomo General Academic Hospital during 2012–2018. Most AD patients with

secondary bacterial infection were males, 112 patients (53.3%) than women, 98 patients (46.7%). Itching and painful vesicles were the chief complaints in AD patients with secondary bacterial infection, as observed in 89 patients (42.4%). Also, 133 patients (63.3%)

reported that most complaints appeared in less than 1 month.

Most AD patients with secondary bacterial infection, as many as 90 patients (42.9%), reported no history of atopy. However, most patients, 69 (32.9%), had individual history of atopy. This means that the patient has a history of atopy, but none of his/her family has. The atopy type in AD patients with secondary bacterial infection was dominated by allergic rhinitis in 53 patients (65.4%). Meanwhile, familial history of atopy in AD patients was dominated by bronchial asthma and atopic dermatitis, as many as 34 patients each (66.6%).

In AD patients with secondary bacterial infection, most lesions appeared on the hands; 196 patients (93.3%), followed by legs; 175 patients (83.3%), and feet; 149 patients (70.9%). Also, in AD patients with secondary bacterial infection, the lesions were dominated by erythema, as many as 176 patients (83.8%), followed by crusting as many as 164 patients (78.1%), and xerosis, as many as 163 patients (77.6%).

Most AD patients with bacterial secondary, 134 patients (63.8%), received treatments with oral antibiotics, NaCl compresses, moisturizers, and antihistamines. Erythromycin was the most opted antibiotic therapy, as observed in 102 patients (58.6%), followed by amoxicillin, as observed in 46 patients (26.4%), and cloxacillin, as observed in 26 patients (14.9%).

This study also analyzed the relationship between individual history of atopy, familial history of atopy, dry skin, and involvement of flexor area as the risk factors of secondary bacterial infection in AD patients. In this study, dry skin had a p-value < 0.05 (Table 2). Therefore, there was a positive correlation between dry skin and the incidence of secondary bacterial infection. The univariate analysis was then continued with the multivariate analysis. The confidence interval for dry skin was 3.803–8.774 (Table 3), reaffirming that dry skin is a risk factor for secondary bacterial infection in AD patients.

**Table 2.** Univariate analysis of variables as the risk factors in atopic dermatitis patients with secondary bacterial infection at the Pediatric Dermatology Division, Dermatovenereology Outpatient Clinic, Dr. Soetomo General Academic Hospital

Variables	Secondary Bacterial Infection		X <sup>2</sup>	p
	Yes	No		
History of atopy in patients or their families			0.275	0.600
Yes	122	245		
No	88	159		
Dry Skin			95.87	0.000
Yes	162	142		
No	48	262		
Involvement of the flexor area			3	0.083
Yes	181	324		
No	29	80		

**Table 3.** Multivariate analysis of variables as the risk factors in atopic dermatitis patients with secondary bacterial infection at the Pediatric Dermatology Division, Dermatovenereology Outpatient Clinic, Dr. Soetomo General Academic Hospital

Variables	Estimate	p	OR	CI
Dry Skin				
Yes	1.754	0.000	5.776	3.803–8.774
No				
Involvement of the flexor area				
Yes	0.309	0.263	1.363	0.793–2.343
No				

OR= Odds Ratio; CI = Confidence Interval.



## DISCUSSION

The risk factors for secondary bacterial infection in AD are not fully determined. In 2014, a cross-sectional study in China reported that 52.3% of AD patients of all ages had secondary bacterial infection.<sup>6</sup> This study analyzed the relationship between risk factors and the incidence of secondary bacterial infections in AD. The risk factors examined were individual/familial history of atopy, dry skin, and involvement of the flexor area.

The skin is continuously in contact with the surrounding environment, exposing it to several organisms.<sup>8</sup> In addition, the surface of the skin contains many products of skin keratinization, such as fat, nitrogen-containing ingredients, and minerals. This is a good environment for the growth of microorganisms. *S. epidermidis* is a dominant non-pathogenic organism on the skin, and *S. aureus* is a pathogenic organism that often causes infection. *S. aureus* is more susceptible to attack skin with chronic epithelial integrity disorders, such as AD. *S. aureus* causes disease both by producing toxins and inducing pyogenic inflammation. The exotoxin produced by *S. aureus* can cause secondary infection or exacerbation of AD. Superantigens produced by *S. aureus* can penetrate the epidermis and dermis, where they interact with immune cells such as T and APC cells to cause inflammation.<sup>9</sup>

We found that secondary bacterial infection is more common in AD patients with a familial history of atopy (122 patients) than those without a familial history of atopy (88 patients). However, the univariate analysis showed a p-value of  $> 0.05$ . It means no correlation between individual and familial history of atopy. In several studies, individual and familial history of atopy has been associated with the incidence of AD itself. In 2017, research in Padang stated that most AD patients have a familial history of atopy.<sup>10</sup> Research at the Dr. Mohammad Hoesin Palembang from 2011 to 2013 stated that familial history of AD includes AD (25.8%), allergic rhinitis (6.5%), asthma (12.9%), and allergic conjunctivitis (3.2%).<sup>11</sup> However, no research has discussed the correlation between a history of atopy and the incidence of secondary infection in AD patients.

This study also analyzed dry skin as a risk factor for secondary bacterial infection in AD patients. It is showed that 162 patients experienced dry skin, and the univariate analysis resulted in a p-value  $< 0.05$ . It means that there was a correlation between dry skin and secondary bacterial infection in AD patients. Furthermore, a multicenter study in China in 2016 presented that dry skin is one of the risk factors for secondary bacterial infection in AD.<sup>12</sup> Dry skin in AD is associated with the mutation of filaggrin (FLG).

There are two roles of FLG in the skin: a barrier and a growth modulator for *S. aureus*. FLG is converted into hygroscopic amino acids in the skin, including urocanic acid (UCA) and pyrrolidone carboxylic acid (PCA). UCA and PCA are components of natural moisturizing factors (NMF) that hydrate and maintain the pH of the stratum corneum acid.<sup>7</sup> Reduced NMF results in abnormalities in the stratum corneum such as dry skin, scaly skin, wrinkled skin, and fissures.

This study found that most lesions involved the flexor area, as observed in 181 patients. However, the univariate analysis resulted in a p-value  $> 0.05$ , which means there is no correlation between the involvement of the flexor area and the incidence of secondary bacterial infection. There is still no study that further discusses the incidence of secondary bacterial infection and the location of the lesions. The generally known theory is the predilection location of AD according to the patient's age. At 0–2 years of age, the predilection location is on the face (forehead, cheeks). At 2–12 years of age, the location of predilection is in the flexors and folds such as the folds of the elbows, knees, neck, and wrists. At over 12 years of age, the distribution of the lesions is less characteristic but more frequent on the trunk and extensors.

The management of AD consists of improving the skin barrier, reducing inflammation, reducing itching, avoiding the triggering factors of AD, providing information, education, and psychological counseling or intervention and consultation. Education for parents and caregivers is an important part of managing AD, especially what is expected to control the flare rather than to cure. Furthermore, the main goal for atopic dermatitis is to control the symptom, reduce the risk of complications, and prevent relapse.<sup>13,14,15,16</sup> In this study, 189 patients (89.5%) received emollient therapy in the form of urea (69.5%), and the rest was in the form of bio cream. Emollients effectively repair the skin barrier as they reduce transepidermal water loss (TEWL) and increase the skin's ability to retain water. A total of 23 patients (11%) received oral steroid therapy, and 42 patients (20%) received topical steroid therapy. AD patients with secondary bacterial infection also received additional therapies, such as oral antibiotics, as observed in 186 patients (88.6%), and erythromycin was the most opted antibiotic, as observed in 149 patients (71%). This was in accordance with the Clinical Practice Guideline of Dr. Soetomo General Academic Hospital. Additional therapies include NaCl dressings, as observed in 169 patients (80.5%), and topical antibiotic therapy, as observed in 60 patients (28.7%).

From 2012 to 2018, 210 AD patients with secondary bacterial infection were admitted to the

Pediatric Dermatology Division, Dermatovenereology Outpatient Clinic, Dr. Soetomo General Academic Hospital. This contributed to 51.9% of all AD patients, 6.9% of total patients admitted to the Pediatric Dermatology Division, and 0.9% of the total patients admitted in the Dermatovenereology Outpatient Clinic. To conclude, dry skin was the risk factor for secondary bacterial infection in AD patients (OR 5.776, 95% CI 3.803–8.774,  $P < .05$ ). Accurate and proper medical records of AD cases are important for better history taking, diagnosis establishment, monitoring, and follow-up treatment. The records should cover the identity of the patient, risk factors, and SCORAD.

## REFERENCESS

- Kennedy K, Heimall J, Spergel JM. Advances in atopic dermatitis in 2017. *J Allergy Clin Immunol* 2017;142(6):1740–7.
- Lubis M. Association between bacterial infection and severity of atopic dermatitis. *Indian J Med Res* 2018;7(5):18–20.
- Fishbein AB, Mueller K, Kruse L. Sleep disturbance in children with moderate/severe atopic dermatitis: A case-control study. *J Am Acad Dermatol* 2018;78(2):336–41.
- Nowicka D, Grywalska I. The role of immune defects and colonization of *Staphylococcus aureus* in the pathogenesis of atopic dermatitis. *Anal Cell Pathol* 2018, Article ID 1956403.
- Pyun BY. Natural history and risk factor of atopic dermatitis in children. *Allergy Asthma Immunol Res* 2015;7(2):101–5.
- Wang X, Shi XD, Li LF, Zhou P, Shen YW. Classification and possible bacterial infection in outpatient with eczema and dermatitis in China. *Medicine* 2017;96:35(e7955).
- Ong PY, Leung DYM. Bacterial and viral infections in atopic dermatitis: a comprehensive review. *Clin Rev Allergy Immunol* 2016;51(3):329–37.
- Lyons JJ, Milner JD, Stone KD. Atopic dermatitis in children: clinical features, pathophysiology, and treatment. *Immunol Allergy Clin North Am* 2015;35(1):161–8.
- Breuer K, Haussler S, Kapp A, Werfel T. *Staphylococcus aureus*: colonizing features and influence of an antibacterial treatment in adults with atopic dermatitis. *Br J Dermatol* 2002;147(1):55–61.
- Lubbe, J. Secondary infections in patients with atopic dermatitis. *Am J Clin Dermatol* 2013;4(9):641–54.
- Eliska N, Thaha MA, Anwar C. Faktor risiko pada dermatitis atopik. *J Ked Kes* 2015; 2(1): 143–9.
- Mei X, Li L. Risk factor for secondary infection in foot eczema: a multicenter study. *Int J Clin Exp Med* 2016;9(6):11974–81.
- Rubel, D, Thirumoorthy T, Soebaryo RW, Weng SCK, Gabriel TM, Villafuerte LL, *et al*. Consensus guidelines for the management of atopic dermatitis: An Asia-Pacific perspective. *J Dermatolog*, 2013; 40(3): 160–71.
- Xu Y, Zhi-Qiang S, Wei L, Yun-Sheng L, Yan Z, Hua C, *et al*. Guidelines for Diagnosis and Treatment of Atopic Dermatitis in China. *Int. J. Dermatol. Venereol.* 2021;4(1):1–9.
- Sathishkumar D, Gupta A, Saini K. Atopic dermatitis in children: An update for pediatricians. *Curr Med Issues* 2020;18(3):17–25.
- Mirrakhimova MK. Improving methods of Treatment of Atopic Pathology in Children. *JCR* 2020; 7(12):190–2.