



**KOMISI ETIK PENELITIAN
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA
*Animal Care and Use Committee (ACUC)***

**KETERANGAN KELAIKAN ETIK
“ ETHICAL CLEARANCE ”**

No : 1.KE.107.06.2019

**KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE)
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA,
TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG
DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :**

- PENELITIAN BERJUDUL** : Potensi Ekstrak Rumput Kebar (*Biophytum
petersianum*) Dalam Mencegah Penurunan Fungsi
Otak Anak Mencit (*Mus musculus*) Masa Laktasi yang
Terpapar Karbofuran
- PENELITI UTAMA** : Epy Muhammad Luqman
- UNIT/LEMBAGA/TEMPAT
PENELITIAN** : Departemen Anatomi Veteriner
Fakultas Kedokteran Hewan Universitas Airlangga
- DINYATAKAN** : LAIK ETIK

Surabaya, 27 Juni 2019

Mengetahui,
Dekan FKH-Unair



Prof. Dr. Pudji Sianto, M.Kes.,Drh.
NIP. 195601051986011001

Ketua,

Dr. Nusdianto Triakoso, M.P.,Drh.
NIP. 196805051997021001

ISSN: 0974-201X (Print)
ISSN: 0974-2028 (Online)

Research Journal of Pharmacy and Technology

RJPT

An International Peer-reviewed
Journal of Pharmaceutical Sciences

Indexed / Abstracted in:

- ISI: Indian Science Abstracts
- CAS: Chemical Abstracts Service (CAS)
- CAB: Abstract
- Google Scholar
- Scopus

 RJPT

Editor In Chief

Dr. Mrs. Monika S. Daharwal

Editor In Chief

A & V Publications, RJPT House, Lokmanya GrihNirman Society, Rohanipuram, In-front of Sector- 1, Pt. Deendayal Upadhyay Nagar, Raipur 492 010. (CG) India

Email: editor.rjpt@gmail.com

Associate Editor

Marwan Mahmood Saleh

Associate Editor

Anbar-Ramadi- Habbaniya- 4-4-17

Email: bio.marwan92@gmail.com

Dr.rer.nat Arli Aditya Parikes

Associate Editor

Department of Bioinformatics School of Life Sciences Indonesia International Institute for Life Sciences Jl. Pulomas Barat Kav.88 Jakarta 13210

Email: arli.parikesit@i3l.ac.id

Dr G Kumaraswamy

Associate Editor

Dr.Kumara Swamy. Gandla Prof.& HeadDept.of Pharmaceutical AnalysisCare College of Pharmacy, Warangal, Telangana.Mobile: +91-9000973789

Email: kumaraswamy.gandla@gmail.com

Dr. G. Manikandan

Associate Editor

Dr. G.Manikandan Assistant Professor Department of Botany Sri Kaliswari College (Autonomous) Sivakasi - 626130 Tamil Nadu India

Email: rgmani.19@gmail.com

Dr.S.Mohanasundaram

Associate Editor

Department of Biochemistry, Sri Sankara Arts and Science College (Autonomous), Kanchipuram - 631561, Tamilnadu, India

Email: sbmohan2007@gmail.com

Dr Shaeesta K. Bhavikatti

Associate Editor

College of Dentistry, King Khalid University, Abha, Saudi Arabia

Email: drshaeesta@gmail.com

Dr Karteek Eswara

Associate Editor

T2,staff quarters, ksr Institutions, ksr kalvi nagar, Tiruchengode-637215, Tamilnadu

Email: karteekeswara@gmail.com

Dr. Chukwuebuka Emmanuel Umeyo

Associate Editor

Department of Pharmaceutics and Pharmaceutical Technology, Faculty of Pharmaceutical Sciences, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria

Email: ec.umeyor@unizik.edu.ng

Academic Editor

Dr. Bharti Ahirwar

Associate Professor, SLT Institute of Pharm. Sciences, Guru Ghasidas University, Bilaspur CG India

Email: ah_bharti@yahoo.com

Dr. Ashok A. Hajare

Professor and Head, Department of Pharm. Tech., Bharati Vidyapeeth College of Pharmacy, Kolhapur (M. S.), India Pin - 416013

Email: ashok.hajare@bharativedyapeeth.edu

Dr. Ajay Kumar Meena

Captain Srinivasa Murthy Regional Ayurveda Drug Development Institute, Arumbakkam, Chennai – 600 106

Email: ajaysheera@gmail.com

Editors

Dr Avinash B Darekar

KVNNSPS, Institute of Pharmaceutical Education and Research, Canada Corner, Nashik, Maharashtra-422002.

Email: avibdarekar@gmail.com

Dr Sudip Kumar Mandal

Dr. B. C. Roy College of Pharmacy & Allied Health Sciences, Durgapur, West Bengal, India

Email: gotosudip79@gmail.com

Dr K Manikandan

Dr.K.Manikandan, M.Pharm., Ph.D., Associate Professor, SRM College of Pharmacy, SRM University, Kattankulathur, Kancheepuram - 603203 Mobile Number - 0 9444708710

Email: manikank2@srmist.edu.in

Dr Shaik Harun Rasheed

professor & Head Srikrupa Institute of Pharmaceutical Sciences Velikatta, siddipet -502277
Telangana.

Email: shaikharunrasheed@gmail.com

Dr. Ashish Kumar

Professor and HOD, Department of Chemistry, Lovely Professional University, Jalandhar road Phagwara

Email: drashishchemlpu@gmail.com

Dr. Shakta Mani Satyam

Department of Pharmacology, Melaka Manipal Medical College (Manipal Campus), Manipal Academy of Higher Education, Manipal- 576104, District- Udupi, State- Karnataka (India)

Email: smsatyam21@gmail.com

Articles in Volume - 15, Issue - 5

Online Since: Monday, May 30, 2022

QbD Stressed Development and Validation of Stability-Indicating RP-HPLC Method for the Simultaneous Estimation of Linagliptin and Metformin HCl in Pharmaceutical Dosage Form

Author(s): Khushbu Patel, Ujashkumar A. Shah, Hirak V. Joshi, Jayvadan K. Patel, Chhaganbhai N. Patel

DOI: [10.52711/0974-360X.2022.00319](https://doi.org/10.52711/0974-360X.2022.00319)

Cite: Khushbu Patel, Ujashkumar A. Shah, Hirak V. Joshi, Jayvadan K. Patel, Chhaganbhai N. Patel. QbD Stressed Development and Validation of Stability-Indicating RP- HPLC Method for the Simultaneous Estimation of Linagliptin and Metformin HCl in Pharmaceutical Dosage Form. Research Journal of Pharmacy and Technology. 2022; 15(5):1917-3. doi: [10.52711/0974-360X.2022.00319](https://doi.org/10.52711/0974-360X.2022.00319)

Strategic archetype of Herbal Medicine Product (HMP) in Regulated and Emerging Market

Author(s): Swagat Tripathy, PN Murthy, BP Patra, Vikram, Sumit Sanduria, Harish Dureja

DOI: [10.52711/0974-360X.2022.00328](https://doi.org/10.52711/0974-360X.2022.00328)

Cite: Swagat Tripathy, PN Murthy, BP Patra, Vikram, Sumit Sanduria, Harish Dureja. Strategic archetype of Herbal Medicine Product (HMP) in Regulated and Emerging Market. Research Journal of Pharmacy and Technology. 2022; 15(5):1973-0. doi: [10.52711/0974-360X.2022.00328](https://doi.org/10.52711/0974-360X.2022.00328)

New stability indicating RP-UFLC method for the quantification of Efavirenz in pharmaceutical dosage forms

Author(s): Sai Ganeswari Aluri, Mukthinuthalapati Mathrusri Annapurna

DOI: [10.52711/0974-360X.2022.00329](https://doi.org/10.52711/0974-360X.2022.00329)

Cite: Sai Ganeswari Aluri, Mukthinuthalapati Mathrusri Annapurna. New stability indicating RP-UFLC method for the quantification of Efavirenz in pharmaceutical dosage forms. Research Journal of Pharmacy and Technology. 2022; 15(5):1981-8. doi: [10.52711/0974-360X.2022.00329](https://doi.org/10.52711/0974-360X.2022.00329)

Antibiotic Sensitivity Pattern of Clinically Isolated Klebsiella Species in C. U. Shah Hospital, Surendranagar

Author(s): Vaibhavi Patel, Priya Mehta, Kunjan Kikani, Girish K. Goswami

DOI: [10.52711/0974-360X.2022.00330](https://doi.org/10.52711/0974-360X.2022.00330)

Views: 0 (pdf), 34 (html)

Cite: Vaibhavi Patel, Priya Mehta, Kunjan Kikani, Girish K. Goswami. Antibiotic Sensitivity Pattern of Clinically Isolated Klebsiella Species in C. U. Shah Hospital, Surendranagar. Research Journal of Pharmacy and Technology. 2022; 15(5):1989-4. doi: [10.52711/0974-360X.2022.00330](https://doi.org/10.52711/0974-360X.2022.00330)

Production and Characterization of Micro-Collagen from Carp Scales Waste (Cyprinus carpio)

Author(s): Rasmi Zakiah Oktarlina, Syaiful Bahri, Andi Nafisah Tendri Adjeng

DOI: [10.52711/0974-360X.2022.00331](https://doi.org/10.52711/0974-360X.2022.00331)

Cite: Rasmi Zakiah Oktarlina, Syaiful Bahri, Andi Nafisah Tendri Adjeng. Production and Characterization of Micro-Collagen from Carp Scales Waste (*Cyprinus carpio*). Research Journal of Pharmacy and Technology. 2022; 15(5):1995-2. doi: [10.52711/0974-360X.2022.00331](https://doi.org/10.52711/0974-360X.2022.00331)

[Read More](#)

Effect of combination of some Polymers with Carbopol 940 on Pregabalin Release Rate from Emulgels

Author(s): Nawar Haddad, Jameela Hasian

DOI: [10.52711/0974-360X.2022.00332](https://doi.org/10.52711/0974-360X.2022.00332)

Cite: Nawar Haddad, Jameela Hasian. Effect of combination of some Polymers with Carbopol 940 on Pregabalin Release Rate from Emulgels. Research Journal of Pharmacy and Technology. 2022; 15(5):2003-9. doi: [10.52711/0974-360X.2022.00332](https://doi.org/10.52711/0974-360X.2022.00332)

The effect of Adrenaline on the contractility of the Right Ventricular Myocardium in rat pups with altered Serotonin concentration in Ontogenesis

Author(s): Akhmetova M., Nigmatullina R., Mindubayeva F., Tykezhanova G.

DOI: [10.52711/0974-360X.2022.00333](https://doi.org/10.52711/0974-360X.2022.00333)

Cite: Akhmetova M., Nigmatullina R., Mindubayeva F., Tykezhanova G. The effect of Adrenaline on the contractility of the Right Ventricular Myocardium in rat pups with altered Serotonin concentration in Ontogenesis. Research Journal of Pharmacy and Technology. 2022; 15(5):2010-6. doi: [10.52711/0974-360X.2022.00333](https://doi.org/10.52711/0974-360X.2022.00333)

Preparation of Apamarg Kshar Tablet and Assessment of In Vitro Potential against Urolithiasis (Mutrakrichra)

Author(s): Dileep Singh Baghel, Saurabh Singh, Narendra Kumar Pandey, Bimlesh Kumar, Amit Mittal, Anand Kumar Chaudhary, Anu Mittal, Amrik Singh

DOI: [10.52711/0974-360X.2022.00334](https://doi.org/10.52711/0974-360X.2022.00334)

Cite: Dileep Singh Baghel, Saurabh Singh, Narendra Kumar Pandey, Bimlesh Kumar, Amit Mittal, Anand Kumar Chaudhary, Anu Mittal, Amrik Singh. Preparation of Apamarg Kshar Tablet and Assessment of In Vitro Potential against Urolithiasis (Mutrakrichra). Research Journal of Pharmacy and Technology. 2022; 15(5):2017-7. doi: [10.52711/0974-360X.2022.00334](https://doi.org/10.52711/0974-360X.2022.00334)

The Effect of Kebar Grass Extract (*Biophytum petersianum* Klotzch) on Total Growth of Ovarian Follicles on Mice (*Mus Musculus*) during Lactation period were Exposed by Carbofuran

Author(s): Athaya K. Mulyadi, Tri W. Suprayogi, Maslichah Mafruchati, Imam Mustofa, Eka P. Hestianah, Juliano M. Ntoruru, Epy M. Luqman

DOI: [10.52711/0974-360X.2022.00335](https://doi.org/10.52711/0974-360X.2022.00335)

Cite: Athaya K. Mulyadi, Tri W. Suprayogi, Maslichah Mafruchati, Imam Mustofa, Eka P. Hestianah, Juliano M. Ntoruru, Epy M. Luqman. The Effect of Kebar Grass Extract (*Biophytum petersianum* Klotzch) on Total Growth of Ovarian Follicles on Mice (*Mus Musculus*) during Lactation period were Exposed by Carbofuran. Research Journal of Pharmacy and Technology. 2022; 15(5):2028-4. doi: [10.52711/0974-360X.2022.00335](https://doi.org/10.52711/0974-360X.2022.00335)

Evaluation of early Menopause symptoms in Post-Hysterectomy and Premature Ovarian insufficiency in women of reproductive age group

Author(s): Vandana Rani, Jaspreet Kaur, Mamta Devi, Parminder Nain, Shaveta Garg, Prerna Sarup

DOI: [10.52711/0974-360X.2022.00336](https://doi.org/10.52711/0974-360X.2022.00336)

Cite: Vandana Rani, Jaspreet Kaur, Mamta Devi, Parminder Nain, Shaveta Garg, Prerna Sarup. Evaluation of early Menopause symptoms in Post-Hysterectomy and Premature Ovarian insufficiency in women of reproductive age group. Research Journal of Pharmacy and Technology. 2022; 15(5):2035-0. doi: [10.52711/0974-360X.2022.00336](https://doi.org/10.52711/0974-360X.2022.00336)

Diabetic Wound Healing in IL-1 β expression by Nano Herbal of Zanthoxylum acanthopodium and Rhodomyrtus tomentosa

Author(s): Risma D. Manurung, Syafruddin Ilyas, Salomo Hutahaean, Rosidah rosidah, Putri C. Situmorang

DOI: [10.52711/0974-360X.2022.00337](https://doi.org/10.52711/0974-360X.2022.00337)

Cite: Risma D. Manurung, Syafruddin Ilyas, Salomo Hutahaean, Rosidah rosidah, Putri C. Situmorang. Diabetic Wound Healing in IL-1 β expression by Nano Herbal of Zanthoxylum acanthopodium and Rhodomyrtus tomentosa. Research Journal of Pharmacy and Technology. 2022; 15(5):2041-6. doi: [10.52711/0974-360X.2022.00337](https://doi.org/10.52711/0974-360X.2022.00337)

Attributes for Discrete Choice Experiment on Pharmacy-based Alcohol Brief Intervention Service in Thailand

Author(s): Sukunta Muadthong, Nusaraporn Kessomboon

DOI: [10.52711/0974-360X.2022.00320](https://doi.org/10.52711/0974-360X.2022.00320)

Cite: Sukunta Muadthong, Nusaraporn Kessomboon. Attributes for Discrete Choice Experiment on Pharmacy-based Alcohol Brief Intervention Service in Thailand. Research Journal of Pharmacy and Technology. 2022; 15(5):1924-2. doi: [10.52711/0974-360X.2022.00320](https://doi.org/10.52711/0974-360X.2022.00320)

A Simple and Rapid RP-HPLC Method for simultaneous estimation of Paracetamol and Pamabrom in Combinational Dosage Form

Author(s): Karthika Paul, Jaswanth Gowda B.H., S.J. Shankar

DOI: [10.52711/0974-360X.2022.00338](https://doi.org/10.52711/0974-360X.2022.00338)

Cite: Karthika Paul, Jaswanth Gowda B.H., S.J. Shankar. A Simple and Rapid RP-HPLC Method for simultaneous estimation of Paracetamol and Pamabrom in Combinational Dosage Form. Research Journal of Pharmacy and Technology. 2022; 15(5):2047-2. doi: [10.52711/0974-360X.2022.00338](https://doi.org/10.52711/0974-360X.2022.00338)

RESEARCH ARTICLE

The Effect of Kebar Grass Extract (*Biophytum petersianum* Klotzch) on Total Growth of Ovarian Follicles on Mice (*Mus Musculus*) during Lactation period were Exposed by Carbofuran

Athaya K. Mulyadi¹, Tri W. Suprayogi¹, Maslichah Mafruchati¹, Imam Mustofa¹, Eka P. Hestianah¹, Juliano M. Ntoruru², Epy M. Luqman^{1*}

¹Department of Veterinary Science, Faculty of Veterinary Medicine, Universitas Airlangga Jl Muyorejo Surabaya, 60115 Indonesia.

²Research Assistant, Meru University of Science and Technology, Kenya.

*Corresponding Author E-mail: epy-m-1@fkh.unair.ac.id

ABSTRACT:

The study aimed to know the effect of Kebar grass extraction amounts of mice (*Mus musculus*) follicles ovary exposed by carbofuran. This study was an experimental study with a completely randomized design. Forty two lactation mice used in this study were divided into seven groups, that were: C was control (Aquadest), P1 was injected with carbofuran 1/4 LD₅₀ (0.0125mg/day), P2 was injected with carbofuran 1/8 LD₅₀ (0.00625mg/day), P3 was injected with carbofuran 1/4 LD₅₀ (0.0125mg/day) and Kebar grass extract (0.135mg/day), P4 was injected with carbofuran 1/8 LD₅₀ (0.00625mg/day) and Kebar grass extract (0.135mg/day), P5 was injected with carbofuran 1/4 LD₅₀ (0.0125mg/day) and vitamin C (0.2ml/day) and P6 was injected with carbofuran 1/8 LD₅₀ (0.00625mg/day) and vitamin C (0.2ml/day) with a treatment period of 14 days. Then on the 15 day of treatment, mice were sacrificed, and the ovary was taken. Then, histology preparation with HE (Hematoxylin-eosin) staining was made, and follicles ovary were counted. Data were analyzed by One Way ANOVA followed by the Duncan test. This study showed that exposed carbofuran was a significant decrease in the growth of ovarian follicles ($p < 0.05$). The administration of the Kebar grass extraction exposure to carbofuran is proven to increase primary and secondary follicles. The Kebar grass extraction administration gave a better effect than vitamin C ($p < 0.05$).

KEYWORDS: Kebar grass extract, Follicles, Ovary, Carbofuran, Lactation, Pesticide stress.

INTRODUCTION:

The insecticide has an essential role in agriculture activities to protect plants from pests. By the rules, the application should not cause environmental pollution, health problem, poisoned, and residues in food products¹. According to World Health Organization (WHO), every year occur cases of insecticide poisoned by farmers cause dead. 80% of poisoned reported happening in a developing country. Insecticide groups of carbamates have many uses in Indonesian to eradicate the pest. Group of carbamates, which is often used carbofuran, carbaryl, and aldicarb.

The risk of carbofuran can causes residues on rice, rice field water, and soil field in Java islands in 1997. The resulting residues on rice are 0.5-1.3 ppb with maximum residue limit according to Codex Alimentarius commission is 0.1 ppm². Carbofuran residue risk can cause teratogenic, adrenal damage, and reproductive system disorders on *Passer domesticus*^{3,4}.

Damage due to exposure carbofuran affects Reactive Oxygen Species (ROS) activities that cause dead brain cells during the mother lactation period⁵. ROS or free radicals is a molecule that is not pairs and stable that molecule will take an electron from another molecule and very reactive^{6,7}. Free radicals induce reproduction toxicity and infertility because insecticide exposure causes oxidative stress who induce lipid peroxidation and DNA damage on fish, rats, and pregnant women⁸. Exposure carbofuran shows an increase in the total

atretic follicle and decreases in a total of healthy follicles⁹. A decrease in healthy follicles' total resulted from granulosa cells in the ovary have apoptosis and necrosis¹⁰ causes atretic follicles. Insecticide also affects vacuolar degeneration on cells follicle, so that looks increase empty follicles in the ovary¹¹.

The lactation period is when mother secretion milk from the mammary gland after giving birth to their child¹². In this period, mothers need lots of nutrition compared when pregnant^{13,14}. Environmental pollutions like insecticide and heavy metal cause a change in the immune system. Toxic substances cause the suppression of immunomodulatory components, which indirectly affect organs as nerve, reproduction, respiratory, and endocrine¹⁵. Insecticide causes damage to the mother and their child before, during, and after birth¹⁶. Measurement of follicles during the lactation period is thought to have the function of identifying mothers that are potentially relatively infertile after weaning¹⁷.

The body has natural defenses to neutralize free radicals by increase the production of antioxidants endogenous. However, if there are more free radical concentrations in the body, it is necessary to balance with antioxidants exogenous^{18,19}. Flavonoids are antioxidants derived from phenolic compounds derived from plants^{20,21}. These compounds are useful in counteracting free radicals and are also proven to fight diseases caused by free radicals²². Kebar grass (*Biophytum petersianum Klotzch*) is a plant that contains flavonoids. Other antioxidants found in Kebar grass are vitamin E and A. Vitamin E is the primary defense against free radicals and can also attack lipid peroxide, which is thought to be a degenerative cause of cells²³. In contrast, vitamin A plays an essential role in stabilizing free radicals²².

Research on the Kebar grass extraction on the reproduction of mother during lactation has not been widely studied, especially the effect on the number of follicles in the ovaries exposed to carbofuran. This study was conducted to show that giving Kebar grass extraction can increase the number of healthy ovarian follicles and decrease the number of atretic follicles exposed by carbofuran.

MATERIAL AND METHODS:

The research tools that are used is experimental animal cages, feed and drinking containers, sonde needles, syringe 3ml, surgical equipment (surgical scissors, scalpel, tweezers), object glass, cover glass, ointment pots, tissue processor automatic, water bath, hot plate, microtome, blade and Olympus® CX-23 microscopes. The research materials that are used is mice, carbofuran (2, 3-Dihydro-2,2-dimethyl-7-benzofuranol N-methyl-carbamate 98%) from Aldrich Chemistry dengan

Bellstain Registry number 1428746, Product of USA, dry Kebar grass in packaging from Papua, vitamin C used as a reference substance, pellets, drinking water, aquadest, CMC Na, 70% etanol, husks for cage tools, ether solution, 10% buffer formalin and alcohol.

Preparation of Experimental Animals:

Pregnant mice strain Balb/C were divided into seven groups is K, P1, P2, P3, P4, P5, and P6. There are six repetitions for each group.

Method:

The research procedure was conducted by testing the code of ethics of experimental animals with the number 1.KE.107.06.2019 at the Faculty of Veterinary Medicine, Airlangga University. Mice were divided into seven groups, that were: K was control (aquadest), P1 was injected with carbofuran 1/4 LD₅₀ (0.0125mg/day), P2 was injected with carbofuran 1/8 LD₅₀ (0.00625 mg/day), P3 was injected with carbofuran 1/4 LD₅₀ (0.0125mg/day) and Kebar grass extract (0.135mg/day), P4 was injected with carbofuran 1/8 LD₅₀ (0.00625 mg/day) and Kebar grass extract (0,135mg/day)^{24, 25}, P5 was injected with carbofuran 1/4 LD₅₀ (0.0125mg/day) and vitamin C (0.2ml/day) and P6 was injected with carbofuran 1/8 LD₅₀ (0.00625mg/day) and vitamin C (0.2ml/day) with a treatment period of 14 days. Then on the 15day of treatment, mice were sacrificed and the ovary was taken. Then, histology preparation with HE (Hematoxylin-eosin).

Histopathological preparations:

Counting the number of ovarian follicles was each treatment group in five different fields of view on each histopathological preparation. Microscopic observation of the ovary in histopathology preparations using an Olympus® CX-23 microscope with a magnification of 100X. Observations were made by counting the number of the primary, secondary, tertiary, Graafian, and atretic follicles. The calculation of each field of view in one preparation is added up, then the average is calculated, then statistical analysis.

Data Analysis:

The data were processed and analyzed statistically using the ANOVA (Analysis of Variance) test if normally distributed data were obtained, followed by the Duncan test. If it is found that the data is not normally distributed, the data is analyzed using the Kruskal Wallis test.

RESULTS:

Primary Follicle:

The analysis results with the ANOVA test showed that the primary follicle was significantly different ($p < 0.05$), so it was continued using the Duncan test. Table 1 shows

that group C has a significant difference in the number of primary follicles in groups P1, P2, and P4. Group P1 did not significantly differ with P2, but there was a significant difference with P5, P6, K, P3, and P4. Group P4 did not have a significant difference with P3, but there was a significant difference with P1, P2, P5, P6, and C. Group P6 did not have a significant difference with P5, C, and P3. However, there was a significant difference between P1, P2, and P4.

Table 1: Duncan test results for the average number of the primary, secondary, tertiary, and atretic follicles.

Group	Primary Follicle	Secondary Follicle	Tertiary Follicle	Atretic Follicle
C	5.13 ^c ±0.70	4.86 ^c ±0.64	1.13 ^{a,b} ±0.41	1.93 ^d ±0.41
P1	2.93 ^a ±0.61	2.26 ^a ±0.41	0.40 ^a ±0.34	3.93 ^f ±0.30
P2	3.67 ^{a,b} ±0.90	2.73 ^a ±0.30	0.46 ^a ±0.41	3.46 ^f ±0.50
P3	5.67 ^{c,d} ±0.23	5.20 ^c ±0.40	1.20 ^{a,b} ±0.20	2.26 ^{d,e} ±0.30
P4	6.40 ^d ±0.34	5.93 ^d ±0.30	1.60 ^b ±0.52	2.00 ^{d,e} ±0.20
P5	4.60 ^{b,c} ±0.20	4.06 ^b ±0.50	0.67 ^a ±0.57	2.60 ^e ±0.20
P6	4.86 ^c ±0.64	4.46 ^{b,c} ±0.11	0.80 ^{a,b} ±0.40	2.46 ^{d,e} ±0.30

Note: Different superscripts show significant differences (p<0.05). C (Control), P1 (Carbofuran 1/4 LD₅₀), P2 (Carbofuran 1/8 LD₅₀), P3 (Kebard grass extraction + Carbofuran 1/4 LD₅₀), P4 (Kebard grass extraction + Carbofuran 1/8 LD₅₀), P5 (Vitamin C + Carbofuran 1/4 LD₅₀), P6 (Vitamin C + Carbofuran 1/8 LD₅₀).




Fig 1 Diagram of the average number of ovarian follicles in each treatment.

Secondary Follicle:

The analysis results with the ANOVA test showed significantly different (p <0.05) and then continued using the Duncan test. Based on Table 1 shows that in group C there was a significant difference in the number of secondary follicles in groups P1, P2, P5, and P4. Group P1 did not have a significant difference with P2, but there was a significant difference with groups P5, P6, C, P3, and P4. Group P6 did not significantly differ with C, P3, and P5, but there was a significant difference between P1, P2, and P4.

Tertiary Follicle:

The ANOVA test, it was found that were significantly different (p <0.05) then continued by using the Duncan test. Table 1 shows that there was no significant difference in the number of tertiary follicles in the P1, P2, P3, P4, P5, and P6 groups in the C group. The P4 group had a significant difference with P1, P2, and P5.

Atretic Follicle:

The analysis results with the ANOVA test showed that they were significantly different (p <0.05), so it was continued using the Duncan test. Based on Table 1 show that in group C there were differences in the number of atretic follicles that were significant with P5, P2, and P1. Group P5 did not significantly differ with P4, P3, and P6, but there was a significant difference between C, P2, and P1. Groups P1 and P2 had significant differences with C, P4, P3, P6, and P5.




Fig 2 The histopathology of ovaries with Hematoxylin Eosin staining using an Olympus CX-23 microscope with 100x magnification in groups K, P1, P2, P3, P4, P5, and P6.

DISCUSSION:

Primary Follicle:

Primary follicles are the beginning of folliculogenesis. They are characterized by granulosa cells that are cuboidal and oocyte in the middle. The process of development is influenced by extra ovarian and intrafollicular signals. The intrafollicular signals from growth factors, while the extra ovarian signals from gonadotropin hormones²⁶. The receptors receive these signals on the granulosa cells of the ovarian follicles. Primary follicles express FSH receptors are on granulosa cells^{27,28}.

The results showed that giving carbofuran caused a decrease in the number of primary follicles in the P1 and P2 groups allegedly because the toxic effects caused an increase in ROS that formed free radicals. P1 experienced a lower decrease than P2; this could be due to the higher carbofuran dose at P1 (0.0125mg) compared to P2 (0.00625mg). Carbofuran entered the body and metabolized cytochrome P₄₅₀ to produce 3-hydroxycarbofuran, causing increased toxic activity because it is systemic²⁹. Insecticides can indirectly affect gonadal function by influencing the action of the hypothalamus-pituitary-gonads or directly by oxidative stress mechanisms³⁰. Oxidative stress results from an increase in free radicals that can damage cell membranes

through lipid peroxidation²⁰. This cell damage causes the primary follicle to decline to not develop into the next stage, namely the secondary follicle.

The increase in the number of primary follicles in the P4 group was due to the antioxidants contained in Kebar grass, namely flavonoids, vitamins E and A. Flavonoids have a role in reducing the effects of ROS by trapping free radicals such as superoxide and hydroxyl anions³¹. Vitamins E and A are non-enzymatic antioxidants that protect membranes cell and improve reproductive function. Vitamin E changes free radicals into hydroperoxides and neutralizes ROS, likes superoxide radicals, hydrogen peroxide, and hydroxyl radicals. Vitamin A is a fat-soluble antioxidant that scavenges free radicals from plasma lipoproteins and cell membranes⁷ and can protect cells from superoxide radicals³². The P3 group showed inhibition in decreasing the number of primary follicles. The difference in giving the Kebar grass extract with the same dose (3.375mg) could be due to differences in the dose of exposure to carbofuran in each group. This study showed that the Kebar grass extract could ward off free radicals caused by carbofuran, but the P3 group could not increase the number of primary follicles.

The administration of vitamin C showed inhibition in reducing the number of primary follicles, both in the P5 and P6 groups. Vitamin C is a hydrophilic antioxidant that can trap free radicals to protect cell membranes from oxidative stress³³. Administration of vitamin C in aging mice showed an increase in the number of the primary, secondary, and antral follicles³⁴. This study indicates that the antioxidants from Kebar grass extract provide a more significant effect than giving vitamin C because the compounds contained in Kebar grass are flavonoids. Flavonoids are phenolic compounds with more than one hydroxyl group to reduce free radicals such as superoxide anion radicals, hydroxyl radicals, peroxy radicals, and alkoxyl radicals³¹. The antioxidant ability of flavonoids is more effective than vitamin C and vitamin E^{22,35}.

Secondary Follicle:

Secondary follicles are the next developmental stage of primary follicles, consisting of granulosa cells, internal theca cells, zona pellucid, and oocytes. Like primary follicles, secondary follicles in their development are influenced by intrafollicular signals and extra ovarian signals. Secondary follicles express FSH receptors on granulosa cells and express LH receptors on theca cells²⁷. Granulosa cells and theca cells in ovarian follicles have an essential role in synthesizing the hormone estrogen. Besides, granulosa cells play a role in the growth and maturation process of follicles³⁶. If there is damage to these cells, it can cause impaired growth and maturation of ovarian follicles.

The results of this study showed a decrease in the number of secondary follicles in the P1 and P2 groups. This decrease is thought to occur in the same way as primary follicles due to exposure to carbofuran, causing free radicals that increase ROS production in the body, resulting in oxidative stress. Reactive oxygen species compounds such as O₂·, H₂O₂, and OH· have an essential role in oxidative damage to cells⁷. Carbofuran is lipid-soluble that will form lipid peroxidation after binding to lipids from the cell membrane³⁷. The final result of lipid peroxidation is malondialdehyde (MDA). Increased MDA level is one factor that indicates a decrease in the number of ovarian follicles leading to infertility³⁸. It is known that giving carbofuran to mother mice during lactation resulted in an increase of MDA levels, necrosis of Purkinje cells, and a decrease in ChE⁵.

The increase in the number of secondary follicles in the P4 group is thought to contain in content Kebar grass is flavonoids, vitamins E and A; this is following a study that states that the flavonoids found in plants can increase the number of antral follicles exposed to the insecticide cypermethrin³⁹. Administration of insecticides with vitamin E antioxidant therapy can reduce MDA levels and increase rat ovaries' GSH levels. Vitamin E can defend cells from oxidative stress⁴⁰ by acting as a protector against lipid peroxidation through its chain-breaking action⁸. Vitamin A in Kebar grass has been shown to protect cells from free radical attack and maintain the epithelial layer of the reproductive tissue⁴¹. As with primary follicles, the P3 group study results showed that it could maintain the number of secondary follicles; this could be because the dose of exposure to carbofuran in the P3 group was lower than the P4 group.

Administration of vitamin C showed a decrease in the number of secondary follicles in the P5 group, while the P6 group maintained the secondary follicle count compared to the control group. The P5 group had the highest exposure dose in this study. Based on antioxidants' actions, it is better to give antioxidants in combination compared with a single form⁷. In the P5 group, there was a decrease in the number of secondary follicles. Antioxidants in vitamin C have not been able to provide a good effect in the P5 group. However, the P6 group showed that vitamin C could maintain the number of secondary follicles; this is the same as the research results from the number of primary follicles.

Tertiary Follicle:

Tertiary follicles are the next development stage of secondary follicles, which consist of granulosa cells, internal and external theca cells, zona pellucid, oocytes, and small spaces between granulosa cells called call Exner bodies. Its development of these follicles is affected by intrafollicular signals and extraovarian signals²⁷.

The results of this study showed a decrease in the number of tertiary follicles in the P1 and P2 groups, but there was no significant difference with the control group. The decrease in the number of tertiary follicles is the same as the decrease in the number experienced by primary and secondary follicles, which is thought to occur because carbofuran's toxic effect causes an increase in ROS in oxidative stress. ROS binds to plasma lipoproteins and cell membrane unsaturated fats, resulting in lipid peroxidation formation⁷. There was no significant difference with the control group, presumably because the mice used in this study were in the lactation period. During lactation, the follicles in the ovaries experience a decrease in the release of gonadotropin hormones from breastfeeding children⁴².

The administration of Kebar grass extract showed that it maintained the number of tertiary follicles in the P3 and P4 groups compared to the control group because the antioxidants present in Kebar grass are flavonoids that can inhibit lipid peroxidation by reducing peroxyl radicals and overcoming superoxide anions³¹. Vitamin E, a lipid antioxidant, can overcome free radicals and functions to stabilize cell membranes⁴³ and vitamin A which can stabilize free radicals⁷. The provision of vitamin C showed the same results as the P1 and P2 groups. There was no significant difference with the control group; this can be affected because vitamin C's antioxidant action has not worked maximum. After all, it is given in a single form⁷.

Graafian follicle:

Graafian follicles are mature follicles and ready to be ovulated. The extra ovarian signal (gonadotropin hormone) plays an essential role in reaching the mature follicle. Granulosa cells have FSH receptors and theca cells have LH receptors²⁷. A large space is formed at this stage called follicular antrum, which contains follicular fluid rich in protein and estrogen. Ovulation occurs when the LH level increases, causing the oocyte to leave the follicle ready for fertilization⁴². Based on Table 1, the results showed no Graafian follicle development in each treatment group; this is thought to be due to cell damage in primary, secondary, and tertiary follicles so that the follicles do not reach the maturation stage. A study showed that giving carbofuran decreased growing follicles, Graafian follicles, and increased atretic follicles⁹. Another study conducted on mice exposed to carbofuran showed the results of fewer developing follicles, more atretic follicles, and hemorrhage⁴⁴.

The control group did not show Graafian follicles' development; this can be caused because the mice used in this study were mother mice during the lactation period. During lactation, the release of gonadotropin hormones decreases and inhibits estrus during labor

because the anterior pituitary secretes the hormone prolactin for breastfeeding⁴². After post-partum estrus, the rats do not re-estrus until the weaning period is complete, called postweaning estrus⁴⁵. The weaning period in mice ranged from 18-21 days⁴⁶. The lactation period will suppress fertility for some time after birth; this causes limited follicular growth because, in this period, there is recovery from the pituitary gonadotropins due to the suppressive effect of increasing steroid hormones during pregnancy. During breastfeeding activity, LH hormone levels' secretion is erratic and lower than that of the standard follicular phase^{47,48}. This statement is presumably the reason why this study does not show any development of the Graafian follicle.

Atretic Follicle:

The analysis results with the ANOVA test showed that they were significantly different ($p < 0.05$), so it was continued using the Duncan test. Based on Table 1 show that in group C there were differences in the number of atretic follicles that were significant with P5, P2, and P1. Group P5 did not significantly differ with P4, P3, and P6, but there was a significant difference between C, P2, and P1. Groups P1 and P2 had significant differences with C, P4, P3, P6, and P5.

An atretic follicle is a degenerative process that causes oocyte, not ovulation. It is suspected that the ovarian oocyte will disappear at a particular stage of development. Atretic follicles occur affected by nutrition, impaired ovarian blood supply, pregnancy, and breastfeeding⁴⁹. Atretic follicles occur accompanied by damage to cell membranes, reduced granulosa cells, and pycnosis⁵⁰.

The results of this study indicated an increase in the number of atretic follicles in the P1 and P2 groups. This increase occurs because exposure to carbofuran results in apoptosis of granulosa cells caused by oxidative stress, thereby increasing free radicals' formation⁸. Oxidative stress in the ovaries causes folliculogenesis disorders, follicles do not develop and atretic⁵¹. The decreasing number of ovarian follicles is thought to be due to an increase in atretic follicles so that only a few follicles develop⁵². In addition to an increase in atretic follicles caused by oxidative stress, this increase can result from disorders of the hypothalamus-pituitary-ovarian system with a hormonal imbalance so that the developing follicles were decreased and atretic follicles increased in mice exposed by carbofuran⁹. This is consistent with the statement that ovarian toxicity caused by insecticides can be affected by endocrine disruption and oxidative stress³⁰. Other insecticides such as malathion show decreased antioxidant enzyme activity in healthy antral follicles. They exerted an inhibitory effect on SOD and

CAT activity with increased lipid peroxidation leading to atretic follicles⁵³.

The inhibition of increasing the number of atretic follicles in the P3 and P4 groups was thought to be due to the antioxidant effect of the Kebar grass, which can prevent cell damage in ovarian follicles due to oxidative stress. The combination of the antioxidants present in Kebar grass has a good effect on the number of ovarian follicles. Antioxidant administration is preferably given in combination based on its antioxidant action⁷. The combination of primary antioxidants that act in scavenging/reducing free radicals with phenolic antioxidants that reduce free radicals and as a chelating agent gives maximum results in oxidative stress^{22,54}. In the Kebar grass, the primary antioxidants are vitamin A and vitamin E, while those that act as phenolic antioxidants are flavonoids. A study conducted using local rabbits showed that the extract of Kebar grass could stimulate follicles' formation in the ovaries^{55,56}.

Administration of vitamin C showed inhibition of increasing the number of atretic follicles in the P6 group. In contrast, the P5 group showed an increase in atretic follicles, presumably due to the higher carbofuran exposure dose in this group. Oxidative stress causes follicles that do not develop into atretic follicles⁵¹. In its single form, the administration of vitamin C causes this antioxidant not to work optimally²². Vitamin C in the P6 group showed inhibition in increasing the number of atretic follicles. This dose showed that the antioxidant vitamin C gave a better effect than the P5 group.

This study indicated that giving carbofuran was shown to reduce follicular development significantly ($p < 0.05$). The administration of Kebar grass extraction to mice exposed to carbofuran increased the number of primary and secondary follicles at a carbofuran dose of 0.00625 mg. The application of Kebar grass extract gave a better effect than vitamin C. The extract of Kebar grass with a dose of 3.375mg had a more significant effect on exposure to carbofuran with a dose of 0.00625mg compared to a dose of 0.0125mg.

ACKNOWLEDGEMENT:

The authors express sincere thanks to the Ministry of Research, Technology and Higher Education of the Republic of Indonesia for funding the research, and the Dean of Faculty of Veterinary Medicine Universitas Airlangga for providing all necessary facilities and funds in conducting this research work.

CONFLICT OF INTEREST:

The authors declare no conflict of interest.

REFERENCES:

1. Indrianingsih. Effect of Carbamate Insecticide on Animal Health and its Products. *Wartazoa Journal*. 2008; 18(2): 101-114
2. Ardiwinata N, Jatmiko SY, Harsanti ES. Insecticide residue monitoring, West Java. Minutes of Seminar on Results of Research on Greenhouse Gas Emissions and Increasing Rice Productivity in Paddy Fields. Bogor. 1999; p. 91 – 105.
3. Wadhwa V, Bakre PP, Bakre VP. Avian Adrenal Respons to Furadan Sp 50. *Journal of Environmental Biology*. 1991; 12(1):1-7.
4. Luqman EM, Widjiati, Mafruchati M, Rahardjo BPS, Hestianah EP. Crosstalk between Necrosis and Apoptosis of Embryonal Cerebral Cortex Neuron Mice (*Mus musculus*) Caused by Carbofuran Exposure. *Research Journal of Pharmacy and Technology*. 2019; 12(11):5492-5498.
5. Luqman EM, Widjiati, Yustinasari LR. Brain Cells Death on Infant Mice (*Mus Musculus*) Caused by Carbofuran Exposure During The Lactation Period. *Kafkas Universitesi Veteriner Fakultesi Dergisi*. 2018; 24(6): 845-852.
6. Arimbi, Azmiah A, Darsono R, Plumeriastuti H, Widyatmo TV, Legowo D. *Veterinary General Pathology Textbook Ed 2nd*. Airlangga University Press. Surabaya. 2015; p. 19
7. Widayati E. Biology Oxidation, Free Radicals, and Antioxidant. *Scientific Magazine Sultan Agung*. 2012; 50(128): 26-32.
8. Bhardwaj JK, Mittal MP, Saraf MP, Kumari P. Pesticide Induced Oxidative Stress and Female Infertility: A Review. *Toxin Reviews*. 2018; 1-13.
9. Baligar PN, Kaliwal BP. Reproductive Toxicity of Carbofuran to the Female Mice: Effects on Estrous Cycle and Follicles. *Industrial Health*. 2002; 40(4): 345-352.
10. Sharma D, Gurinder KS, Kuldeep SK. Triazopho Induced Oxidative Stress and Histomorphological Changes in Ovary of Female Wistar Rats. *Pesticide Biochemistry and Physiology*. 2015; 117: 9-18.
11. Suffiriyanto, Indradji M. Evaluation ff The Effect of Anti-Steroidal Diazinon Contamination on The Ovaries and Adrenal Cortex in Animal Models. *Animal Production*. 2002; 4(2): 101 – 111
12. Kalaimathy A, Vidhya K. Surrogacy and Breastfeeding: Inducing Lactation. *Asian Journal of Nursing Education and Research*. 2011; 1(3): 94-95.
13. Moran VH, Lowe N, Crossland N, Berti C, Cetin I, Hermoso M, Koletzko B, Dykes F. Nutritional Requirements During Lactation. Towards European Alignment of Reference Values: the EURRECA network. 2010; 6(2) 39-54.
14. Danasu R. Effectiveness of Reflex Zone Stimulation on initiation and maintenance of Lactation among Lactation Failure Mothers at SMVMCH, Kalitheerthalkuppam, Puducherry. *Asian Journal of Nursing Education and Research*. 2015; 5(4): 505-512.
15. Pajewska-Szmyt M, Sinkiewicz-Darol E, Gadzala-Kopcluch. The Impact of Environmental Pollution on The Quality of Mother's. *Environmental Science and Pollution Research*. 2019; 26: 7405-7427.
16. Alio A. Toxic Stress and Maternal and Infant Health: a Brief Overviews and Tips for Community Health Worker. New York. 2017;6.
17. Lopes TP, Padilla L, Bolarin A, Martinez HR, Roca J. Ovarian Follicle Growth during Lactation Determines the Reproductive Performance of Weaned Sows. *Animals*. 2020; 10: 1-12.
18. Werdhasari A. The Role of Antioxidants for Health. *Indonesian Journal of Biotech Medisiana*. 2014; 3(2): 59-68
19. Zambare KK, Thalkari AB, Tour NS, Reddy KV. Herbal Remedies for treatment of Polycystic Ovary Syndrome. *Asian Journal of Pharmacy and Technology*. 2019; 9(2):112-114.
20. Khaira K. Counteract free radicals with antioxidants. *Journal of Science and Technology*. 2010; 2(2): 183-187.
21. Babu A, Veerasamy R, Sivadasan S. Metformin- A Drug of Plant Origin. *Research Journal of Pharmacy and Technology*. 2018; 11(6): 2701-2708.
22. Sayuti K, Rina, Y. Natural and Synthetic Antioxidants. *Andalas University Press*. Padang. 2015; p. 32, 38-39, 52-53, 70.
23. Lamid A. Vitamin E as antioxidant. *Media Research and*

- Development. 1995; 5(1): 14-16.
24. Luqman EM, Hestianah EP, Tehupuring BC, Yustinasari LR. 2020. Interaction Between Autophagy, Apoptosis and Necrosis of Infant Mice (Mus Musculus) Brain Cells From Its Carbofuran Exposed Mothers During Lactation Periods. *Acta Veterinaria Brno* 89 (2), 195-200.
 25. Labib MF, Widjiati, Hernawati T, Luqman EM, Kurnijasanti R, Suprayogi TW. Kebar grass (*Biophytum petersianum* K.) The effect in maintaining mice (*Mus musculus*) sperm quality exposed to dioxin. *International Journal of Research in Pharmaceutical Sciences*. 2020, 11(3), 4977-4981.
 26. AbbasHadi AH, Dohan KH. Effect of Potassium Dichromate on Ovary and Uterus in Female Mice. *Research Journal of Pharmacy and Technology*. 2018; 11(2):429-433.
 27. Figueiredo JR, Lima LF, Silva JRV, Santos R. Control of Growth and Development of Preantral Follicle: insight from in vitro culture. *Animal Reproduction*. 2018; 15(1): 648-659.
 28. Revathi R, Julius A. Effect of LH/FSH Ratio and its Correlation with Insulin Resistance in PCOS Obese woman of Reproductive age Group. *Research Journal of Pharmacy and Technology*. 2018; 11(6): 2217-2219.
 29. Gupta RC. Carbofuran toxicity. *Journal of Toxicology*. 1994; 43: 383-418.
 30. Sharma RK, Singh P, Setia A, Sharma AK. Insecticides and Ovarian Functions: Review. *Environmental and Molecular Mutagenesis*. 2020; 61: 369-392.
 31. Widowati W, Safitri R, Rumumpuk R, Siahaan M. Screening of Superoxide Dismutase Activity in Various Plants. *Journal of Public Health*. 2005; 5(1): 33-38
 32. Luddi A, Capaldo A, Focarelli R, Gori M, Morgante G, Piomboni P, Leo VD. Antioxidants reduce Oxidative Stress in Follicular Fluid of Aged Women Undergoing IVF. *Journal of Reproductive Biology and Endocrinology*. 2016; 14(1): 1-7.
 33. Mustaf SJ, Salih TA, Yasseen HA, Marouf BH, Mohammed AI. Effect of Monosodium Glutamate on Mice Ovaries and The Possible Protective Role of Vitamin C. *Annals of Applied Biology*. 2015; 2(4): 100-105.
 34. Abdollahifar MA, Azad N, Sajadi E, Mofarahe ZS, Zare F, Moradi A, Rezaee F, Gholamin M, Abdi S. Vitamin C Restores Ovarian Follicular in a Mouse Model of Aging. *Anatomy & cell biology*. 2019; 52: 196-203.
 35. Prior RL, Cao G. Antioxidant Phytochemicals in Fruits and Vegetables: Diet and Health Implications. *HortScience*. 2000; 35(4): 588-592.
 36. Puriastuti AC, Maramis MM, Annas JJ. Chronic stress reduces the thickness of the granulosa cell layer of the Antral Follicle *Rattus norvegicus*. *Journal of biomedical practitioners*. 2017; 19(3): 268-275.
 37. Purnomo E, Luqman EM, Ratnani H, Plumeriastuti H, Mafruchati M, Dhamayanti Y. The Increasing Damage in Tubular Cell of Infant Mice's Kidneys from Carbofuran Exposure during its Mothers' Lactation Period. *International Journal of Pharmaceutical Research*. 2020; 12(4): 17-24.
 38. Rohmawati H, Sujuti H, Made PD, Setyo HY. The Effect of Extra Virgin Oil on Folliculogenesis and Ovarian Malondialdehyde Levels in Rats Exposed to Rhodamine B. *Indonesian Journal of Health Information*. 2018; 4(2): 120-127.
 39. Sari MP, Winarsih S, Raras TYM, Mintaroem K. Ethanol Extract of Green Tea (*Camellia sinensis*) Increases Levels of Follicle Stimulating Hormone (FSH) and Amounts of Antral Follicles in Female Rats Exposed to Spermethrin. *Qanun Medika*. 2018; 2(2): 39-47.
 40. Sargazi Z, Nikravesh MR, Jalali M, Sadeghnia HR, Anbarkeh FR, Mohammadzadeh L. Diazinon-induced ovarian toxicity and protection by vitamin E. *Iranian Journal of Toxicology*. 2014; 8(26): 1130-1135.
 41. Sembiring B, Darwati I. Identification of the chemical components of Kebar grass (*Biophytum petersianum*) accession from Papua and Java. *The Litro Bulletin*. 2014; 25(1): 37-44
 42. Ismudiono, Srianto P, Anwar H, Madyawati SP, Samik A, Safitri E. Reproductive Physiology in Animal Textbook. Airlangga University Press. Surabaya. 2010; p. 109-114.
 43. Lung JKS, Destiani DP. Antioxidant Activity Test of Vitamins A, C, E using the DPPH method. *Farmaka*. 2017; 15(1): 53-59.
 44. Bhalodiya SJ, Patel BJ, Joshi DV, Raval SH, Patel JG, Zapadia BB, Jagoda MD. Toxicopathology of Induced Carbofuran Toxicity in Wistar Rats (*Rattus norvegicus*). *AGRES*. 2014; 3(3):217-230.
 45. Hanson AF, Berdoy M. Rats. In: Tynes VV (ed.) Behavior of Exotic Pets. Oxford: Wiley-Blackwell. 2010; p. 104 - 116.
 46. Kusumawati D. Friendly with Experimental Animals. Gajah Mada University Press. Yogyakarta. 2016; p. 8-10, 49-50.
 47. McNeilly AS. Neuroendocrine Changes and Fertility in Breast-Feeding Women. *Prog Brain Res*. 2001; 133(14): 207-214.
 48. Kaushik S, Satapathy T, Roy A, Gupta PP, Purabiya P. Endocannabinoid activation and polycystic ovary syndrome: A systematic review. *Research J. Pharm. and Tech*. 2020; 13(1):448-452. doi: 10.5958/0974-360X.2020.00087.6
 49. Maheshwari H, Satyaningtjas AS, Harlina E, Cahyaningsih U, Effendi M, Mustofa MA, Bekalani YK. The Role of Fennel Infusion on Estrous Cycle and Follicles Development of White Rats. *Indonesian Journal of Pharmaceutical Sciences*. 2016; 14(1): 19-25.
 50. Nurjanah, Widiyaningrum P. Development of the ovaries of mice exposed to x-ray radiation. *Journal of Natural Science Mathematics*. 2016; 39(2): 85-91.
 51. Agarwal A, Mellado AA, Premkumar BJ, Shaman A, Gupta S. The Effects of Oxidative Stress on Female Reproduction: a Review. *Journal of Reproductive Biology and Endocrinology*. 2012; 10: 1-31.
 52. Ramadhani SA., Supriatna I, Karja NWK., Winarto A. Ovarian Folliculogenesis Control by Giving Cotton Seed Extract. *Journal of Veterinary Science*. 2017; 35(1): 71-80.
 53. Revathi R, Julius A. A Biological Effect of Sex Hormone Binding Globulin and Testosterone in Polycystic Ovary Syndrome (PCOS) Obese Women. *Research Journal of Pharmacy and Technology*. 2017; 10(7): 2143-2145.
 54. Bhardwaj, J.K. and P. Saraf. Granulosa Cell Apoptosis by Impairing Antioxidant Defense System and Cellular Integrity in Caprine Antral Follicles Post Malathion Exposure. *Environ Toxicol*. 2016; 31(12): 1944-1954.
 55. Sajid H, Guzar, Enaas S, Jawad, Mohammed A, Altahan, Nabaa N, Hameed. The Effects of Trace Element levels on Polycystic Ovary Syndrome in human female Sat Thi-Qar governorate/Iraq. *Research Journal of Pharmacy and Technology*. 2019; 12(9):4447-4453.
 56. Riyadi, Humaidah N, Kalsum U. Effect of Different Doses of Kebar Grass Flour (*Biophytum petersianum*) on Reproduction Performance of Local Female Rabbits. *Journal of Animal Husbandry Renewal*. 2019; 2(1): 160-163.