

Calcium (Ca²⁺) expression and intensity in cumulus-oocyte complex (COCs) in Kacang goat after vitrification

by Epy M. Luqman

Submission date: 20-Dec-2022 10:59AM (UTC+0800)

Submission ID: 1984819278

File name: Icium_Ca2_Expression_and_Intensity_on_Cumulus-Oocyte_Complex.pdf (299.75K)

Word count: 5084

Character count: 27714

Calcium (Ca^{2+}) expression and intensity in cumulus-oocyte complex (COCs) in Kacang goat after vitrification

W. Widjiati¹, Z. Faizah², N. Darsini², V.F. Hendrawan³, H.N. Karima⁴,
C. Chotimah⁴, S.B. Sumitro⁵, L.R. Yustinasari⁶, A.A.M.N. Kasman⁷,
J.M. Ntoruru⁸, E.M. Luqman⁶

1

¹ Post Graduate School of Universitas Airlangga Surabaya, Indonesia² Department of Biomedical Science, Faculty of Medicine Universitas Airlangga Surabaya, Indonesia³ Department of Reproduction, Faculty of Veterinary Medicine, Universitas Brawijaya Malang, Indonesia⁴ Bio-Science Central Laboratory, Universitas Brawijaya Malang, Indonesia⁵ Department of Biology, Faculty of Science, Universitas Brawijaya Malang, Indonesia⁶ Department of Veterinary Science, Faculty of Veterinary Medicine,

Universitas Airlangga Surabaya, Indonesia

⁷ Faculty of Health Science, Universitas Muhammadiyah Mataram, Mataram⁸ Research Assistant, Meru University, Kenya

Abstract

The process of vitrification of the cumulus-oocyte complex (COCs) often results in cold shock. When warming, heat shock occurs which can disrupt the balance of intracellular calcium (Ca^{2+}) intensity. Drastic changes in temperature cause Reactive Oxygen Species (ROS), affecting changes on Ca^{2+} in COCs. The role of calcium is needed for oocyte activation in the fertilization process. The purpose of this study was to examine the expression of Ca^{2+} and the intensity of Ca^{2+} in COCs after vitrification. The study was divided into 2 groups, the control group (C) of fresh COCs, and the treatment group (T) of COCs after vitrification. After vitrification for 24 hours, then thawing, the expression of Ca^{2+} was examined using the Immunocytochemistry (ICC) method and the intensity of calcium (Ca^{2+}) with a Confocal Laser Scanning Microscope (CLSM). The research data obtained were analyzed statistically by T-Test. The results showed that the expression of Ca^{2+} in the control group (12.00 ± 0.00) was different from the treatment group (0.35 ± 0.79). The intensity of Ca^{2+} in the control group (1059.43 ± 489.59) was different from the treatment group (568.21 ± 84.31). The conclusion of this study is that cryopreservation affects calcium in COCs; there were differences in the expression and the intensity of Ca^{2+} between fresh COCs and COCs after vitrification. Ca^{2+} intensity of COCs after vitrification was concentrated in the nucleus, while in fresh COCs it was concentrated in the cytoplasm.

Key words: Calcium (Ca^{2+}), confocal laser scanning microscope, cumulus-oocyte complex (COCs), food production, Kacang goat

13

Correspondence to: W. Widjiati, e-mail: widjiati@fkh.unair.ac.id, tel.: +62315992785

Introduction

The Kacang goat is one of Indonesia's germ-plasm, where its body shape is small and may not be profitable economically compared to foreign goats, but it has several advantages, including being easy to adapt because it is native to Indonesia and maintenance costs are not high. The breakthrough technology to produce goat embryos quickly can be done *in vitro* using the In vitro fertilization (IVF) method, either conventionally by adding mature eggs with spermatozoa outside the body or using the intra-cytoplasmic sperm injection (ICSI) method. The main obstacles to goat embryo production through conventional IVF are low oocyte quality and limited oocyte sources for *in vitro* goat embryo production (Nasar et al. 2007).

Oocytes as a source of female gamete cells can be stored before maturation as an oocyte bank. The oocyte bank is a breakthrough for providing gamete cells for *in vitro* fertilization. Storage of oocytes for a long time for oocyte bank purposes can be done by cryopreservation. The best oocyte cryopreservation method is the vitrification method (Rienzi et al. 2017). The vitrification method is often used because there is no crystal ice formation and the processing time is faster than the slow freezing method (Munck and Vajta 2017). According to Tavukcuoglu et al. (2012), the drawback of the vitrification method is that osmotic stress and toxic content in cryoprotectants can reduce the level of oocyte quality.

Chen and Yang (2009) explained that both the slow-freezing method with increased sucrose concentration and new vitrification techniques significantly improve the results of cryopreservation of human oocytes. The survival of cryopreserved oocytes ranged from 74% to 90% using the slow-freezing method, and from 84% to 99% using the vitrification method. Overall, the survival rate of oocytes from vitrification (95%, 899/948) appeared higher than that of the slow-freezing method (75%, 1,275/1,683). For both protocols, the cells must first be treated with a combination of cell-permeating and non-permeating agents to minimize ice formation which can severely damage the cell. The stark difference between the success of oocyte and embryo cryopreservation has yet to be strictly explained, but differences in cell size and membrane permeability have been proposed (Stachecki and Cohen 2004, Leibo 2008). Previously, high concentrations of cryoprotectant were required for vitrification, but novel containers and submicroliter volumes result in super-fast cooling and subsequent warming rates, so that concentrations closer to those used in slow-freezing can now be used (Mukaida et al. 2002, Kasai and Mukaida 2004).

During the vitrification process, cryoprotectants are needed to protect oocytes both extracellularly and intracellularly from drastic temperature changes. This temperature change will cause an increase in Reactive Oxygen Species (ROS). Likewise, when warming up, a heat shock will occur. The heat shock occurs when there is a change in temperature from cold to warm. Increased ROS during thawing will cause damage to the mitochondrial function and changes in Ca^{2+} ion influx (Favetta et al. 2017).

Changes in temperature due to vitrification and during thawing will highly affect the intensity of Ca^{2+} ions. The role of Ca^{2+} ions in oocytes is very important and is related to the oocyte maturation process. In the maturation process, there will be maturation of the nucleus and cytoplasm. In the nucleus maturation process, there will be an increase in Ca^{2+} influx into the oocyte nucleus, so that protein regulators will run for the nuclear maturation process to occur. Calcium (Ca^{2+}) is necessary for the fertilization process and embryo development. The purpose of this study was to measure the expression of Ca^{2+} and the intensity of Ca^{2+} in cumulus-oocyte complex.

The immunocytochemistry (ICC) method indicates an increase in Ca^{2+} expression. The extracellular Ca^{2+} expression can only be observed using the ICC method, while the intracellular Ca^{2+} intensity requires the use of the Confocal Laser Scanning Microscope (CLSM) method. The combination of the immunocytochemistry method and the (CLSM) can accurately measure the content and position of Ca^{2+} in oocytes, due to the freezing process using the vitrification method.

36

Materials and Methods

Ethical eligibility

The ethical eligibility of this research was obtained from the Faculty of Veterinary Medicine, Universitas Airlangga, ethics number 1.KE.061.04.2019.

Sample

The sample for this study was an ovary obtained from the waste of a slaughterhouse in Surabaya. Ovarian samples, after being cleaned of tissue and blood, were then washed with PBS. The clean ovary was placed in a bottle containing PBS solution and brought to the laboratory in a warm container at 37°C. The sample unit is cumulus-oocyte complex (COCs) obtained by follicular aspiration with a follicular surface diameter of 8-12 mm. The aspirated COCs are further classified based on the number of layers of cumulus cells surrounding the COCs. Only COCs coated with two or

more layers of cumulus cells were used for this study. From the results of the COCs classification obtained, it is used as a research sample unit. The sample unit was used to test the calcium intensity qualitatively by immunocytochemistry and quantitatively for the calcium intensity by CLSM. In the ICC process, much COCs was lost due to the staining process immediately, so that at the end of the ICC process, the total COCs obtained as a research sample unit was 17 COCs.

Research site

This study examined the expression of calcium (Ca^{2+}) between fresh COCs and COCs after vitrification using an immunocytochemistry method, and also examined the intensity of calcium (Ca^{2+}) between fresh COCs and COCs after vitrification using the confocal laser scanning microscope (CLSM) ^{for 20 h}. The immunocytochemistry examination was carried out at the Division of Veterinary Pathology, Faculty of Veterinary Medicine, Universitas Airlangga. The CLSM examination was carried out at the Central Laboratory of Biological Sciences, Universitas Brawijaya.

Experimental design

This study was replicated 5 times, consisting of 3 replicates to the immunocytochemistry examination sample and 2 replicates to the CLSM examination sample. This research was carried out in stages after all the immunocytochemistry examinations were completed, and the research then continued for CLSM examination. In ovarian sampling, it was taken 4-6 ovaries to obtain a number of COCs samples. After aspiration, 40 to 45 COCs were obtained but not all of these COCs qualified as sample units. From the COCs obtained, they were then selected according to the criteria for the sample unit. On average, each collection obtained 24-25 COCs that met the requirements. COCs that met the criteria for the sample unit were then divided into 2 for immunocytochemistry examination of fresh COC samples without being vitrified, and some for vitrification of them being vitrified. 24-25 COCs that met the requirements, the 12 vitrified COCs were divided into 3 straws and 12-13 oocytes for immunocytochemistry examination of fresh non-vitrified COCs. Vitrification was carried out for 24 hours, and the COCs were then thawed and examined using immunocytochemistry. Post thawing and after immunocytochemistry examination, a total of 17 sample units were obtained for each control and treatment group. With immunocytochemistry examination there are always COCs that are lost because the staining and suction process is too strong. At the end of staining only half of the successful oocytes could be observed. The CLSM examination

was repeated twice, and the obtained COCs were divided into 2; 12 were vitrified COCs and 12-13 fresh COCs were not vitrified, and CLSM examinations were immediately performed. For the vitrified COCs, 24 hours later it was thawed and CLSM examination was performed. The total number of eligible ovaries was 20, and COCs as sample units that met the requirements were 104. After immunocytochemistry examination, the total sample units obtained for each group was 17.

Cumulus-oocyte complex collection

The ovaries of goat were cleaned of ligaments and fatty tissue and washed with 0.9% physiological NaCl, which was added to 50 l of gentamicin sulfate. They were then put in a sterile tube and heated in a water bath at a temperature of 30-35°C. COCs are collected by follicular aspiration with a follicular surface size of 8-12 mm in diameter. Collected using a 10 ml disposable syringe with an 18 G needle previously filled with MEM media. The COC that had been collected was washed with MEM media (Widjiati et al. 2012). From 20 eligible ovaries with good ovarian condition, 34 COCs met the criteria as sample unit (the immunocytochemistry process obtained 17 COCs and CLSM as many as 17 COCs).

Cumulus-oocyte complex vitrification

The COC samples were immersed in the equilibration medium for 18-20 minutes then immersed in the vitrification medium 1 for 30 seconds and the vitrification medium 2 for 30 seconds. 30 seconds in vitrification medium 1 and 2 is the optimum time for the entry of intra and extra cellular cryoprotectants into the COC, and the COC form has returned to its initial form and does not shrink. The soaked COC was placed at the end of the hemistraw as a place for COCs during vitrification. Furthermore, the hemistraw which already contained COC was evaporated over liquid N_2 , then put into liquid N_2 , then put in a canester and stored in a liquid N_2 container for 24 hours. After 24 hours, they were thawed for immunocytochemistry and CLSM examination. Post thawing, if there is a degenerated COC, it is not used as a research sample unit.

Cumulus-oocyte complex warming

Warming of COCs was carried out at room temperature. COCs removed from the hemistraw were immersed in preheated thawing medium. The warming medium contained sucrose solution with a gradual concentration of 0.5 Mol and 1 Mol each for 1-3 minutes until the COCs returned to their original shape, then

Table 1. Assessment of immunoc[26]hemistry examination scoring according to Novack et al (200[2] semi-quantitatively based on the IRS value (AxB), which is the result of multiplying the percentage score of positive cells (A) with the color reaction intensity score (B).

A	B
Score 0: no positive cell	Score 0: no color reaction
Score 1: Positive cells less than 10%	Score 1: low color intensity
Score 2: Positive cells between 11% - 50%	Score 2: medium color intensity
Score3: Positive cells between 51% - 80%	Score 3: strong color intensity
Score 4: Positive cells more than 80%	

Table 2. Qualitative examination of calcium (Ca^{2+}) intensity by immunocytochemistry between fresh cumulus-oocyte complexes (COC) without vitrification and COC after vitrification.

Group	Calcium Expression					
	Total COCs	Mean	Standard Deviation	Minimum	Maximum	P
Control	17	0.351	0.79	0	2	0.000*
Treatment	17	12	00.00	12	12	

* significant at $p<0.05$; C: fresh COCs without vitrification; T: COCs after vitrification.

Description: C: Fresh COCs without vitrification, T: COCs after vitrification.

transferred to the culture medium and ready for the stage of immunocytochemistry and CLSM examination.

Examination of Ca^{2+} expression by immunocytochemistry

The fresh COCs was fixed on a glass slide, then rehydrated with graded alcohol, then washed with PBS, then soaked in 3% hydrogen peroxide H_2O_2 (in DI water) for 20 minutes, 1% BSA in PBS for 30 [12]utes at room temperature, primary antibody Ca^{2+} 1:1000 overnight, cold temperature 4°C, Biotin-labeled secondary antibody (Ar[2] Rat IgG Biotin Labelled) and Ca^{2+} primary antibody, 1 hour at room temperature, SA-HRP (Streptavidin-Horseradish Peroxidase), 60 minutes, room temperature, Chromogen DAB (3,3-diaminobenzidine tetrahydrochloride), 20 minutes, room temperature, Counterstain (Aceto orcein), 3 minutes, room temperature then examined under a microscope (Widjiati et al. 2017). Semi-quantitative [6]ervation of the expression of Ca^{2+} was carried out according to the modified Remmele method (Novak et al. 2007). The Remmele scale index (IRS) is the result of multiplying the percentage score of immunoreactive cells with the color intensity score produced in the cells (Table 1).

Calcium (Ca^{2+}) intensity examination with confocal laser scanning microscope (CLSM)

The oocytes were added to 10 l Fluo-3 40 M and incubated for 30 minutes, immersed in 100 l PBS (Ca-Mg) for 30 minutes, then coverslipped and observed with a confocal microscope.

Data Analysis

[28] Expression and intensity of calcium (Ca^{2+}) data were analyzed non-parametrically using the T-Test.

Results

Examination of calcium²⁺ cumulus-oocyte complex expression with immunocytochemistry

Calcium²⁺ intensity data were qualitatively examined by immunocytochemistry staining to investigate the effect of cold shock and heat shock on indications of changes in calcium ions. The total [27]mple of calcium intensity, qualitatively, can be seen in Table 2.

The data obtained in Table 2 showed the average qualitative examination of calcium (Ca^{2+}) intensity of Fresh COCs without vitrification and COCs after vitrification. [10]

Based on the results of the different test, the T-Test showed that there was a significant difference between the COCs of the Fresh COCs group without vitrification and the COCs after vitrification group ($p<0.05$).

Examination of calcium intensity (Ca^{2+}) in cumulus-oocytes complex with the confocal laser scanning microscope (CLSM) method

To visualize the intensity of calcium ions, the COCs examination was performed using the confocal laser scanning microscope method.




Fig. 1. Qualitative examination of calcium (Ca^{2+}) intensity indicated by a very strong brown color (arrow) in COCs (arrow). On slide (A) the control group of fresh COCs without vitrification, showed a number of immunoreactive cells >80% (score 4) with very strong colocalization (score 17). In (B) the group of COCs after vitrification showed a low expression with Score 1 (immunocytochemistry staining; 400x magnification; Olympus BX-50, Pentax optio 230; 2.0 megapixel Digital Camera).




Fig. 2. Calcium intensity (Ca^{2+}) in COC after vitrification was concentrated in the nucleolus (B), in fresh COC without vitrification it was concentrated in the cytoplasm (A); White arrows: COCs are in good condition, and arrows show fluorescent luminescence in the nucleus and cytoplasm (Olympus Microscope, Japan Type FV1000, Olympus Software FLUoview version 4.2a; x100).

Table 3. Quantitative examination of calcium (Ca^{2+}) intensity by confocal laser scanning microscope between fresh COC without vitrification and COCs after vitrification.

Group	Total COCs	Mean	Std. Deviation	Sig.
Control	17	1059.43 ^a	489.59	0.006
Treatment	17	568.21 ^b	484.31	

Description: Different superscripts in the same column show significant differences ($p<0.05$).

C: Fresh COCs without vitrification, T: COCs after vitrification.

Discussion

The results obtained qualitatively indicated that the intensity of calcium in fresh COCs without vitrification was higher than in vitrified COCs. Likewise, the quantitative examination of calcium intensity in fresh COCs without vitrification was higher than that of vitrified COCs. The results of qualitative examination of calcium intensity showed brown color expression in COCs in the group of fresh oocytes without vitrification (a number of immunoreactive cells with very strong color intensity) and COCs after vitrification exhibited a low expression shown in Fig. 1 above. Fig. 2 shows that calcium intensity (Ca^{2+}) in fresh COC without vitrification was concentrated in the cytoplasm (A) and in COC after vitrification was concentrated in the nucleolus [5].

The results of statistical analysis showed that the average calcium intensity in fresh oocytes without vitrification was significantly different from vitrified COCs ($p < 0.05$). Expression of calcium (Ca^{2+}) was higher in the fresh oocyte group compared to COCs after vitrification (Table 3). This is in accordance with the research of Bonte et al. (2020), who stated that rat oocytes after vitrification-thawing showed decreased Ca^{2+} potential during oocyte activation induced by cryoprotectant exposure.

From the research data that has been obtained on vitrified COCs there is a qualitative decrease in calcium intensity. The results of this study are consistent with Sanaei et al. (2018), who state that calcium is needed to maintain oocyte quality. Likewise, according to the results of Marques et al. (2018), the use of vitrified solution with calcium chelator can increase the ability of COCs after vitrification to maintain mitochondrial distribution and function as well as chromosomal segregation. The release of Ca^{2+} ion concentration in vitrified media is the most effective protocol for oocyte cryopreservation (Marques et al. 2018).

The decrease in calcium intensity qualitatively by immunocytochemistry examination in this study was influenced by many factors, among others, heat shock when the COCs were thawing. According to Jang et al. (2017) and Orieif et al. (2005), post-thawing greatly affects the quality of COCs. Many factors affect the

quality of post-thawing COCs; including the length of exposure to the cryoprotectant, the type of cryoprotectant used, temperature, and viscosity. At the time of thawing there will be a change in temperature from frozen to melting, and this condition will affect the release of free radicals. The release of free radicals will affect changes in calcium oscillations and cause disturbances in mitochondrial function (Roth 2018).

The success rate of cryopreservation is influenced by lipid content in the cytoplasm. For example, centrifugation to remove lipids increases the survivability of post-freezing embryos. The specific mechanism of lipid accumulation in [11] is still not fully understood, but the most probable mechanism is lipid peroxidation, which is amplified in the cryopreservation process due to excessive production of free radicals, and the inhibition of ROS production in the cytoplasm increases viability (Barceló-Fimbrisand 2007).

According to Barceló-Fimbrisand (2007), unsaturated fatty acids such as albumin-bound linoleic are a useful additive during culture, although they do not affect embryonic development. Unsaturated fatty acids increase post-vitrification viability. Unsaturated fatty acids can increase the fluidity of the membrane.

In this study, COCs were not matured first. Vitrification of COCs was at the germinal vesicle stage. Germ-stage vesicle COCs were more tolerant of the effects of stress during vitrification. The genetic material is still encased in the nuclear membrane (Daddangadi et al. 2020).

The results obtained from quantitative examination of calcium intensity with CLSM showed that the intensity in the fresh COCs group without vitrification was higher than in the vitrified COCs group ($p < 0.05$).

In this study, using immunocytochemistry examination, Ca^{2+} expression was seen only in COCs, but not in granulosa cells. Calcium (Ca^{2+}) present in COCs regulates COCs membrane balance and plays a role in the COC maturation process. The results of this study are in accordance with the theory that intracellular Ca^{2+} plays a role in ATP regulation and the regulation of cytokines present in the oocyte nucleus is related to the maturation process in the oocyte nucleus and cytoplasm.

The immunocytochemistry method has not been

able to fully describe the intensity of Ca^{2+} levels in oocytes. Calcium (Ca^{2+}) expression is only expressed descriptively on the surface of oocytes, while oocytes are 3-dimensional cells. For this reason, it is necessary to measure the intensity of Ca^{2+} using the confocal laser scanning microscope (CLSM) method, so that the intensity of Ca^{2+} can be measured in the intracellular part of the oocyte.

Intracellular Ca^{2+} intensity in fresh COCs without vitrification and vitrified COCs can be detected using a confocal laser scanning microscope (CLSM), using a 40 M Fluo 3 calcium probe in Phosphate Buffer Saline (-Ca, -Mg) medium. Calcium (Ca^{2+}) was detected using a 488 nm argon laser. The intensity value is determined by determining the cut-off, by adjusting the wavelength of the instrument used. The cut-off value is the intensity value of the reflected light. Calcium (Ca^{2+}) oscillation in oocytes is an important factor because it determines the success of fertilization and subsequent embryonic development. In the process of fertilization, spermatozoa will release Phospholipase C-Zeta (PLC ζ) to induce the release of Ca^{2+} in the cytoplasm before the occurrence of fertilization (Ramdan et al. 2012, Chithiwala et al. 2015). The increase in intracellular Ca^{2+} is a signal that indicates the occurrence of the meiotic stage, and an increase in Ca^{2+} allows interactions between oocytes and spermatozoa in the fertilization process (Borgers et al. 2009, Kang et al. 2015, Karabulut et al. 2018). The combination of the immunocytochemistry method to visualize the expression of Ca^{2+} and the CLSM method to measure the intensity of Ca^{2+} is an appropriate method for accurately determining the increase in Ca^{2+} levels in oocytes.

In thawing COCs after vitrification, there will be an increase in HSP 70 to protect the oocyte from changes in cold temperatures to warm temperatures, so that the cells do not experience heat shock. If HSP 70 is not able to protect cells from temperature changes, there will be a decrease in Ca^{2+} which will cause a change in the ratio of Ca^{2+} and Sodium (Na). Changes in Ca^{2+} will affect the quality of the COCs.

Calcium (Ca^{2+}) is an important factor in the process of oocyte development. Changes in Ca^{2+} will affect ATP regulation and impaired Ca^{2+} stabilization intracellularly (Feng et al. 2018). The decrease in the intensity of intracellular Ca^{2+} in post-warming oocytes will cause changes in proteins in the oocyte nucleus and will activate the pro-apoptotic cascade.

The biochemical and structural changes that occur during the COCs maturation process affect the oocyte's ability to increase the calcium concentration in the cytoplasm. The structure of the Endoplasmic Reticulum will change during the freezing and warming process and will affect its ability to release calcium. Increasing

the concentration of calcium is important in maintaining oocyte quality (Fraser 1982, Wang and Machaty 2013).

Calcium (Ca^{2+}) is important in the regulation of cytokine signaling in oocytes for the COCs maturation process. Calcium (Ca^{2+}) oscillation is important in the preparation of the COCs maturation process, which will activate MPF and MAPK. When the COCs are mature and have released polar body II, the concentration of Ca^{2+} is high which indicates good COCs quality, whereas, in poor COCs quality, the level and intensity of Ca^{2+} will decrease. High concentrations of Ca^{2+} in the intracellular will support successful fertilization and will indirectly improve the quality of embryo development (Fernandez et al. 2012, Cheon et al. 2013).

Conclusion

There was a qualitative and quantitative difference in Ca^{2+} intensity between fresh non-vitrified COC and vitrified COC. The intensity of calcium (Ca^{2+}) was qualitatively higher in fresh COCs without vitrification. The intensity of calcium (Ca^{2+}) in COCs vitrification was concentrated in the nucleus and fresh COCs without vitrification were concentrated in the cytoplasm.

Acknowledgements

³² This study was supported by a Mandat Grant (SK 1408/UN3/2019) from Universitas Airlangga (Surabaya, Indonesia).

References

- Barceló-Fimbre M, Seidel GE Jr (2007) Effects of fetal calf serum, phenazine ethosulfate and either glucose or fructose during in vitro culture of bovine embryos on embryonic development after cryopreservation. Mol Reprod Dev 74: 1395-1405.
- Bonte D, Thys V, de Sutter P, Boel A, Leybaert L, Heindryckx B (2020) Vitrification negatively affects the Ca^{2+} releasing and activation potential of mouse oocytes, but vitrified oocytes are potentially useful for diagnostic purposes. Reprod Biomed Online 40: 13-25.
- Borges E Jr, Braga DP, de Sousa Bonetti TC, Iaconelli A Jr, Franco JG Jr (2009) Artificial oocyte activation using calcium ionophore in ICSI cycles with spermatozoa from different sources. Reprod BioMed Online 18: 45-52.
- Chen SU, Yang YS (2009) Slow freezing or vitrification of oocytes: their effects on survival and meiotic spindles, and the time schedule for clinical practice. Taiwan J Obstet Gynecol 48: 15-22.
- Cheon B, Lee HC, Wakai T, Fissore RA (2013) Ca^{2+} influx and the store-operated Ca^{2+} entry pathway undergo regu-

- lation during mouse oocyte maturation. *Mol Biol Cell* 24: 1396-1410.
- Chithiwala ZH, Lee HC, Hill DL, Jellerette-Nolan T, Fissore R, Grow D, Dumescic DA (2015) Phospholipase C-zeta deficiency as a cause for repetitive oocyte fertilization failure during ovarian stimulation for in vitro fertilization with ICSI: a case report. *J Assist Reprod Genet* 32: 1415-1419.
- Daddangadi A, Uppangala S, Kalthur G, Talevi R, Adiga SK (2020) Germinal stage vitrification is superior to MII stage vitrification in prepubertal mouse oocytes. *Cryobiology* 93: 49-55.
- De Munck N, Vajta G (2017) Safety and efficiency of oocyte vitrification. *Cryobiology* 78: 119-127.
- Favetta V, Colombo RC, Júnior JFM, de Faria RT (2017) Light sources and culture media in the in vitro growth of the Brazilian orchid *Microelaia lundii*. *Semin Cienc Agrar* 38: 1775-1783.
- Fraser LR (1982) Ca^{2+} is required for mouse sperm capacitation and fertilization in vitro. *J Androl* 3: 412-419.
- Gómez-Fernández C, López-Guerrero AM, Pozo-Guisado E, Álvarez IS, Martín-Romero FJ (2012) Calcium signaling in mouse oocyte maturation: the roles of STIM1, ORAI1 and SOCE. *Mol Hum Reprod* 18: 194-203.
- Jang TH, Park SC, Yang JH, Kim JY, Seok JH, Park US, Choi CW, Lee SR, Han J (2017) Cryopreservation and Its clinical applications. *Integr Med Res* 6: 12-18.
- Kang HJ, Lee SH, Park YS, Lim CK, Ko DS, Yang KM, Park DW (2015) Artificial oocyte activation in intracytoplasmic sperm injection cycles using testicular sperm in human in vitro fertilization. *Clin Exp Reprod Med* 42: 45-50.
- Karabulut S, Aksünger Ö, Ata C, Sağıroglu Y, Keskin I (2018) Artificial oocyte activation with calcium ionophore for frozen sperm cycles. *Syst Biol Reprod Med* 64: 381-388.
- Kasai M, Mukaida T (2004) Cryopreservation of animal and human embryos by vitrification. *Reprod Biomed Online* 9: 164-170.
- Leibo SP (2008) Cryopreservation of oocytes and embryos: optimization by theoretical versus empirical analysis. *Theriogenology* 69: 37-47.
- Marques CC, Santos-Silva C, Rodrigues C, Matos JE, Moura T, Baptista MC, Horta AEM, Bessa RJB, Alves SP, Soveral G, Pereira RMLN (2018) Bovine oocyte membrane permeability and cryosurvival: Effects of different cryoprotectants and calcium in the vitrification media. *Cryobiology* 81: 4-11.
- Mukaida T, Takahashi K, Kasai M (2002) Blastocyst cryopreservation: ultrarapid vitrification using cryoloop technique. *Reprod BioMed Online* 6: 221-225.
- Nowak M, Madej JA, Dziegel P (2007) Intensity of COX2 expression in cells of soft tissue fibrosarcomas in dogs as related to grade of tumour malignancy. *Bull Vet Inst Pulawy* 51: 275-279.
- Orief Y, Schultze-Mosgau A, Dafopoulos K, Al-Hasani S (2005) Vitrification: Will it replace the conventional gamete cryopreservation techniques? *Middle East Fertil Soc* 10: 171-184.
- Rahman ANA, Abdullah R, Embong WK (2007) Goat embryo development from in vitro matured oocytes of heterogeneous quality through intracytoplasmic sperm injection technique. *Biotechnol* 6: 373-382.
- Ramadan WM, Kashir J, Jones C, Coward K (2012) Oocyte activation and phospholipase C zeta (PLC ζ): diagnostic and therapeutic implications for assisted reproductive technology. *Cell Commun Signal* 10: 1-20.
- Rienzi L, Gracia C, Maggiulli R, LaBarbera AR, Kaser DJ, Ubaldi FM, Vanderpoel S, Racowsky C (2017) Oocyte, embryo and blastocyst cryopreservation in ART: systematic review and meta-analysis comparing slow-freezing versus vitrification to produce evidence for the development of global guidance. *Hum Reprod Update* 23: 139-155.
- Roth GA, Abate D, Abate KH, Abay SM, Abbafati C, Abbas N (2018) Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 392: 1736-1788.
- Sanaei B, Movaghari B, Valojerdi MR, Ebrahimi B, Bazrgar M, Jafarpour F, Nasr-Esfahani MH (2018) An improved method for vitrification of in vitro matured ovine oocytes; beneficial effects of ethylene glycol tetraacetic acid, an intracellular calcium chelator. *Cryobiology* 84: 82-90.
- Stachecki JJ, Cohen J (2004) An overview of oocyte cryopreservation. *Reprod Biomed Online* 9: 152-163.
- Wang C, Machaty Z (2013) Calcium influx in mammalian eggs. *Reproduction* 145: R97-R105.
- Wang F, Yuan RY, Li L, Meng TG, Fan LH, Jing Y, Zhang RR, Li YY, Liang QX, Dong F, Hou Y, Schatten H, Sun QY, Ou XH (2018) Mitochondrial regulation of $[\text{Ca}^{2+}]_{\text{i}}$ oscillations during cell cycle resumption of the second meiosis of oocyte. *Cell Cycle* 17: 1471-1486.
- Widjiati W, Aulanni'am A, Hendrawan VF (2017) The effect of vitrification of oocytes cumulus complex apoptosis of mice (*Mus musculus*) to apoptosis, rate of fertilization and embryo quality. *Int J Pharm Clin Res* 9: 179-182.
- Widjiati W, Boediono A, Sumitro SB, Hinting A, Aulani'am, Susilowati T (2012) Isolation and identification of transforming growth factor β from in vitro matured cumulus oocyte complexes. *Hayati J Biosci* 19: 6-10.

Calcium (Ca²⁺) expression and intensityin cumulus-oocyte complex (COCs) in Kacang goat after vitrification

ORIGINALITY REPORT



PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | nexusacademicpublishers.com
Internet Source | 2% |
| 2 | Submitted to iGroup
Student Paper | 2% |
| 3 | mafiadoc.com
Internet Source | 2% |
| 4 | intl.reproduction-online.org
Internet Source | 1% |
| 5 | repository.pertanian.go.id
Internet Source | 1% |
| 6 | crjim.ub.ac.id
Internet Source | 1% |
| 7 | www.ommegaonline.org
Internet Source | 1% |
| 8 | M. G Larman. "Calcium-free vitrification reduces cryoprotectant-induced zona pellucida hardening and increases fertilization | 1% |

rates in mouse oocytes", Reproduction,

01/01/2006

Publication

9 www.reproduction-online.org

Internet Source

1 %

10 M M S Engel, I Kusumastuty, K W Anita, D Handayani. "The Effect of High Fat High Fructose Diet (Modification of AIN-93M) on Nuclear Factor Kappa Beta Expression in the Liver Tissue of Male Sprague Dawley Rats", Journal of Physics: Conference Series, 2019

Publication

<1 %

11 M. Barceló-Fimenes. "Effects of fetal calf serum, phenazine ethosulfate and either glucose or fructose during in vitro culture of bovine embryos on embryonic development after cryopreservation", Molecular Reproduction and Development, 11/2007

Publication

<1 %

12 repository.unair.ac.id

Internet Source

<1 %

13 www.veterinaryworld.org

Internet Source

<1 %

14 Liqiong Hu, Hao Chen, Xingliang Zhang, Zhencheng Feng, Haifeng Zhang, Qingqi Meng. "Rosiglitazone ameliorates radiation-induced intestinal inflammation in rats by

<1 %

inhibiting NLRP3 inflammasome and TNF- α production", Journal of Radiation Research, 2020

Publication

- | | | |
|----|--|------|
| 15 | jukuri.luke.fi
Internet Source | <1 % |
| 16 | www.eajournals.org
Internet Source | <1 % |
| 17 | www.rjpbc.com
Internet Source | <1 % |
| 18 | "OS01: Lung Cancer 1 : Oral Sessions",
Respirology, 2013.
Publication | <1 % |
| 19 | Abu Nasar Md. Aminoor Rahman ., Ramlie Bin
Abdullah ., Wan Khadijah Wan Embong ..
"Goat Embryo Development from in vitro
Matured Oocytes of Heterogeneous Quality
Through Intracytoplasmic Sperm Injection
Technique", Biotechnology(Faisalabad), 2007
Publication | <1 % |
| 20 | phcogj.com
Internet Source | <1 % |
| 21 | scholarworks.waldenu.edu
Internet Source | <1 % |
| 22 | www.jgpt.co.in
Internet Source | <1 % |

- 23 Hui Yang, Qiuyu Sun, Feifei Chong, Xiaoru Jiang, Yanli Wang, Kedi Xu, Yuanlin Zou, Linping Xu, Chunhua Song. "Polymorphisms in lncRNA MIR2052HG and susceptibility to breast cancer in Chinese population", Aging, 2021 <1 %
- Publication
-
- 24 Somayeh Keshavarzi, Azadeh Dokht Eftekhari, Hajar Vahabzadeh, Marzieh Mehrafza et al. "Post-warming survival rates and clinical outcomes of human cleavage stage embryos vitrified/warmed using CryoTouch and Cryotop methods", Middle East Fertility Society Journal, 2021 <1 %
- Publication
-
- 25 prr.hec.gov.pk <1 %
- Internet Source
-
- 26 Dwikora N. Utomo, Ferdiansyah Mahyudin, Kukuh D. Hernugrahanto, Heri Suroto, Muhammad Zaim Chilmi, Fedik Abdul Rantam. "Implantation of platelet rich fibrin and allogenic mesenchymal stem cells facilitate the healing of muscle injury: An experimental study on animal", International Journal of Surgery Open, 2018 <1 %
- Publication
-
- 27 Shalini Srivastava, S.B. Agrawal, M.K. Mondal. "Biosorption isotherms and kinetics on <1 %

removal of Cr(VI) using native and chemically modified Lagerstroemia speciosa bark",
Ecological Engineering, 2015

Publication

-
- 28 eprints.eudl.eu <1 %
Internet Source
-
- 29 f1000research.com <1 %
Internet Source
-
- 30 harvest.usask.ca <1 %
Internet Source
-
- 31 "Poster Presentation Abstracts", International Journal of Rheumatic Diseases, 2016 <1 %
Publication
-
- 32 "Reproductive and Developmental Strategies", Springer Nature, 2018 <1 %
Publication
-
- 33 D M Astutik, L Sulmartiwi, E Saputra, D Y Pujiastuti. "The effect addition of kappa carrageenan flour to the level of gel strength and acceptability of dumpling from threadfin bream fish (*Nemipterus nematophorus*) surimi", IOP Conference Series: Earth and Environmental Science, 2020 <1 %
Publication
-
- 34 Elisabetta Tosti, Yves Ménézo, Aitken, Check et al. "Gamete activation: basic knowledge" <1 %

and clinical applications", 'Oxford University Press (OUP)'

Internet Source

- 35 "In Vitro Fertilization", Springer Science and Business Media LLC, 2019 <1 %
Publication
-
- 36 A. A. Muhammad Nur Kasman, Budi Santoso, Widjiati Widjiati. "The effect of vitrification after warming on the expressions of p38, CDK1, and cyclin B in immature goat oocytes followed by in vitro maturation", Veterinary World, 2020 <1 %
Publication
-

Exclude quotes Off

Exclude bibliography On

Exclude matches Off

Calcium (Ca^{2+}) expression and intensity in cumulus-oocyte complex (COCs) in Kacang goat after vitrification

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
