



**KOMISI ETIK PENELITIAN
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA
*Animal Care and Use Committee (ACUC)***

**KETERANGAN KELAIKAN ETIK
“ ETHICAL CLEARANCE ”**

No : 1.KE.061.04.2019

**KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE)
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA,
TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG
DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :**

PENELITIAN BERJUDUL : Penyelamatan Genetik Kambing Lokal Melalui Metode
Intracytoplasmik Sperm Injection dan *Preimplantation
Genetik Diagnosis* Sebagai Upaya Konservasi dan
Ketahanan Pangan

PENELITI UTAMA : Widjati

UNIT/LEMBAGA/TEMPAT PENELITIAN : Penelitian Terapan Unggulan Perguruan Tinggi
Fakultas Kedokteran Hewan Universitas Airlangga

DINYATAKAN : LAIK ETIK

Surabaya, 18 April 2019

Ketua,

Dr. Nusdianto Triakoso, M.P.,Drh.
NIP. 196805051997021001

Mengetahui,
Dekan FKH-Unair,


Prof. Dr. Rudi Srianto, M.Kes.,Drh.
NIP. 195601051986011001

Journal of the British Ecological Society

ECOLOGY

ENVIRONMENT & CONSERVATION

ISSN 0014-1801



AN INTERNATIONAL JOURNAL

Ecology, Environment and Conservation Editorial Advisory Board

Chief Editor

Prof.(Dr.) R.K.Trivedy, Pune, India

Email id- rktrivedy@gmail.com

Tel: 91-20-46745119

Mobile No.-9975703363

EDITORIAL ADVISORY BOARD

1. Dr. Prof. Moses Inbaraj, Professor, Department of Botany (Retired), Madras Christian College, Chennai, India
2. Dr. Prof. D.J. Lee, Professor, Department of Chemical Engineering, National Taiwan University, Taiwan
3. Dr. T. Bahorun, Professor, Department of Biosciences, Mauritius University, Mauritius
4. Dr. Prof. A.R. Ghosh, Professor, Department of Environmental Sciences, University of Burdwan, Burdwan, India
5. Dr. Prof. M. Zaman, Professor Department of Botany, Dhaka University, Dhaka, Bangladesh
6. Dr. Marcantonio Bragdin, Professor Department of Chemistry, University of Venice, Venice, Italy
7. Dr. Z. Fuat Topark, Professor Dicle University, Diyarbakir, Turkey
8. Dr. Wilson S. Tisera, Associate Professor Christian University, Kupang, Indonesia
9. Dr. Prof. Philip C. Reid, Scientist Marine Biological Association, U.K
10. Dr. Prof. Mohd. Yusuf, Professor Deptt. of Environmental Engineering, University Sans Malaysia, Malaysia
11. Dr. Prof. Azni Idris, Deputy Vice-Chancellor, Research and Innov., Univ. of Putra Malaysia, Malaysia
12. Dr. Vikas Sharma, Associate Professor, RARS of SKUAST, Jammu, J&K, India
13. Dr. A.C. Pandey, Senior Scientist, Birsa Agricultural University, Ranchi, India
14. Dr. Prof. Agoes Soegiarto, Professor, Dept. Biology, Airlangga Univ. Kampus C, Surabaya, Indonesia
15. Dr. Ahmed L. Mahmoudi, Professor, Water Studies Centre, King Faisal University, Saudi Arabia
16. Dr. Siroop Chaudhuri, Associate Professor Center for Environment, Sustainability and Human Development (CESH) O.P. Jindal Global University, Sonipat, India
17. Dr. Amin L. Setyo, Professor, Graduate School of International Crop, Hiroshima University, Hiroshima, Japan
18. Dr. Francis Gbogbo, Associate Professor, Department of Animal Biology, University of Ghana, Ghana
19. Dr. Prof. Deenbandhu Sahoo, Professor Department of Botany, university of Delhi, Delhi, India
20. Dr. Prof. J.K. Ladha, Professor Emeritus University of California, Davis, California, U.S.A.

21. Dr. Prof. M.Z.M. Nomani, Professor Faculty of Law, Aligarh Muslim University, Aligarh, India
22. Dr. G.R. Pathade, Principal (Retired) H.V. Desai College, Pune, India
23. Dr. Alfred Maroyi, Associate Professor Department of Botany, University of Fort Hare, Alice 5700, South Africa
24. Dr. Prof. A.V.L.N.S.H. Hariharan, Professor Dept. of Chemistry, College of Science, GITAM University, Visakhapatnam, India
25. Dr. Zeinab El-Sayed Zayed, Associate Professor Department of Biotechnology, The Central Lab of Date Palm Researches and Development Agriculture Research Center, Cairo, Egypt
26. Dr. Raj Singh, Professor Department of Botany, MMD University, Mullana, Ambala, Haryana, India
27. Dr. Prof. Mohammed Latif Khan, Professor, Department of Botany Dr. Hari Singh Gour Vishwavidyalaya Sagar, India
28. Dr. R. Dinesh Principal Scientist, ICAR-Indian Institute of Spices Research, Marikunnu, P.O. Kozhikode, India
29. Dr. Suresha Gududappa, Senior Environmental Scientist, Saudi Arabia
30. Dr. Azma Hanim Ismail, School of Biological Sciences, Universiti Sains Malaysia, 11800 USM, Pulau Pinang, Malaysia.

ECOLOGY, ENVIRONMENT AND CONSERVATION

VOL. 28 (2) : 2022

CONTENTS

- 589–594 Disaster Risk Reduction for Students through Development of Educational Games Lusiana
—*Karunia Puji Hastuti, Akhmad Munaya Rahman and M. Feindhi Ramadhan*
- 595–601 Antioxidant Activity of Ethanolic Extract of *Bauhinia scandens*
—*Lianah Lianah, Rita Ariyana Nur Khasanah, Flori R. Sari, Krisantini, Neneng Windayani and Mashuri Masri*
- 602–605 *Acanthaster planci* a Potential Source of Protein and Amino Acids for Fish Feed Ingredients
—*Muhammad Safir, Novalina Serdiati and Kasim Mansyur*
- 606–609 Assessment of the treatment efficiency of some polycyclic aromatic hydrocarbon (PAHs) in the water by doped TiO₂
—*To Xuan Quynh and Vu Duc Toan*
- 610–614 Knowledge, Attitudes and Practices towards the Risk of Open Defecation among the Inhabitants of Rural area
—*Mohamed Osman Elamin, Aida Zaid Ali, Hatim Rahimtallah and Hatim Y. Natto*
- 615–620 Feasibility and synergistic effect of glucoamylase hydrolysis and microbial fuel cell on electricity generation from potato-waste
—*Md. Nurul Islam Siddique, B.K. Zaied and Mohd. Zamri Bin Ibrahim*
- 621–625 Habitat Distribution of Fungal Species Collected at Lung NGOC Hoang Nature Reserve
—*M.T. Duong, V. L. Loi and D. H. Truong*
- 626–630 First report on the morphological and molecular identification of *Lactocollybia variicystis* from Salahadin Governorate, North Central Iraq
—*Talib O. Al-Khesraji, Milad A. Mezher and Ahmed H. M. Shugran*
- 631–634 The Use Values of Macro Fungi in Lung Ngoc Hoang Nature Reserve, Vietnam
—*T.M. Duong, D.H. Truong and V.L. Loi*
- 635–641 Calcium Ionophore Supplementation in *In vitro* Fertilization Medium towards Fertilization Rates of Kacang Goats (*Capra hircus*)
—*Azizah Yasita Najiati, Tjuk Imam Restiadi, Sri Pantja Madyawati, Imam Mustofa, Maslichah Mafruchati, Widjiati, Viski Fitri Hendrawan and Epy Muhammad Luqman*
- 642–648 Interpretation Method for Cultural and Climatic Expression Performance Acceptance for Envelope Building in Phinisi Tower, Makassar, Indonesia
—*Yuri Hermawan Prasetyo, Antariksa and Bachtiar Fauzy*
- 649–663 Mobility and Risk Interaction: Vulnerability of Accessibility of the Algiers Metropolis
—*Nait Amara Farid and Baouni Tahar*
- 664–671 Analysis of Growth and Water Quality Dynamics in vannamei white Shrimp (*Litopenaeus vannamei*) Cultivation Using the Millennial Shrimp Farming System in Indonesia
—*Zulkisam Pramudia, Abd. Rahem Faqih and Andi Kurniawan*
- 672–676 Adsorption of Nutrient Ions into Biofilm Matrices of Natural Microbial Consortium
—*Yogita Ayu Dwi Susanti, Muhammad Musa and Andi Kurniawan*
- 677–684 Assessment of *Aspergillus niger* and *Aspergillus flavus* in Inoculant Induced Composting to Remove Antibiotics in Poultry Manure
—*Roy Searca Jose P. Dela Cruz, Florencio C. Ballesteros, Jerwin R. Undan, Lani Lou Mar A. Lopez, Ma. Eloisa Faye C. Cortez and Fe L. Porciuncula*
- 685–691 Assessment of butterfly diversity status and conservation issues in Agartala urban localities of Tripura, North East, India
—*Dipak Das*

- 692–696 A Comparative Change Detection Analysis Using Remote Sensing and Geographical Information System for Coimbatore City, Tamilnadu
—Varatharajaperumal Thangavel, E. Sarojini, R. Vignesh, C.B. Anagha and G. Subashini
- 697–701 A Survey on Household Solid Waste Management in Trivandrum City, India
—Nimmi Babychan
- 702–706 Impact of frontline demonstrations of Broad Bed Furrow (BBF) on yield and economics of soybean (*Glycine max* (L.) Merrill) under Rainfed conditions of Northern Telangana Zone, India
—Raghuveer M., G. Shiva Charan, Y. Praveen Kumar, M. Sunil Kumar, A. Ramadevi, A. Poshadri and K. Mamatha
- 707–718 Climate Change Impact Assessment on Stream flow of Ganjal River, India
—Adarsh Singh and Geeta Singh
- 719–726 Efficient use of agricultural wastes as Biosorbent in the removal of Fuchsin basic dye from aqueous solutions
—S.D. Patil, S.T. Morbale and N.T. Patel
- 727–735 CO₂ emission, economic growth and energy consumption in India: A VAR analysis
—Bishnu Charan Behera
- 736–740 Remote Sensing based Noise Level Assessment: A Case Study of Coimbatore City, India
—E. Sarojini, S.D. Anitha Selvasofia, S. Kanchana and B. Kamal
- 741–746 Impact of Urbanization on Nest Site Selection and Fecundity Rate of House Sparrow, *Passer domesticus* Population
—Balaji Sundarmahalingam and Martina Jeevamani
- 747–751 Moisture Dependent Physical Properties Assessment of Deenanath Grass Seeds
—Sanjay Kumar Singh, Sheshrao Kautkar, P.K. Pathak and Amit Kumar Patil
- 752–757 Phycoremediation of Dyes: An Overview on Mechanism, Challenges, and Prospects
—M. Chugh and N. Bharadvoaja
- 758–763 Experimental Investigation of Increasing the Strength of Pervious Concrete by Partially using Flyash as A Replacement Material of Cement
—Varatharajaperumal Thangavel, Karthickraja, G., Ajith, N. and Mohamed Shameem. K
- 764–767 Poverty Eradication of Jhumias through Rubber Plantation in Tripura: an Case Study
—Suman Kalyan Chaudhury and Sukanta Sarkar
- 768–773 Native Plants Usage in Indian Ornamental Landscaping Scenario: A Review
—Rupkatha Roy Chowdhury and Shagufi Narula
- 774–778 Characterization of Soil Quality Parameters under Plain and Hilly Region of East Siang District, Arunachal Pradesh
—Tuli Taga, Ramawatar Meena, Varadaraju Viveka and Kamini Kumari
- 779–784 Effect of elevated temperature and moisture stress on gas exchange parameters of *Trianthema portulacastrum* L.
—Govindaraj T. and P. Murali Arthanari
- 785–789 Assessment of Soil Fertility Status of Cuddapah Block in YSR District, Andhra Pradesh, India
—Varadaraju Viveka, R. Meena, P. Vennela, Thaiseen, R. Karmakar and Tuli Taga
- 790–801 Influence of Urbanization on Particulate Matter Pollution in Coimbatore City, India
—M. Arunkumar, A. Manisekar and S. Dhanakumar
- 802–805 Influence of Integrated Nutrient Management in Rice (*Oryza sativa*) on Performance of Wheat (*Triticum aestivum*) under Long Term Rice-Wheat Cropping System
—Shivani Ranjan, Sumit Sow, Sanjay Kumar and Swaraj Kumar Dutta

- 806–808 Effect of Integrated Nutrient Management on Yield attributes and Yield of Broccoli (*Brassica oleracea* var. *italica*) under Mid-Hills condition of Uttarakhand
—Kuriyal H. and Goswami G.
- 809–814 Delineation of Coastal Regulation Zone of an Ecologically Sensitive Coastal Region in the Southern Indian Peninsula: A GIS Approach
—Gurugnanam Balasubramanian, Bairavi Swaminathan, Glitson Pereira, Santonu Goswami and Saroj Choudhury
- 815–823 Adsorption of Ni (II) ions from wastewater on adsorbents derived from natural materials: Kinetic and Thermodynamic study
—S.D. Patil, S.T. Morbale and N.T. Patel
- 824–827 Effect of Natural Biopesticides Neem and Commercial Chemical Pesticide Fipronil on Microbial Diversity of Earthworm Gut
—Snehalata Ankaram
- 828–830 First Record of Partial Xanthism in Bronze Featherback *Notopterus synurus* (Osteoglossiformes: Notopteridae)
—Priyankar Chakraborty and Kranti Yardi
- 831–852 Influence of predator incited fear and interference competition in the dynamics of prey-predator system where the prey species are protected in a reserved area
—Narayan Mondal, Hussam Alrabaiiah, Dipesh Barman, Jyotirmoy Roy and Shariful Alama
- 853–862 Which Indian Metropolitan City Poses Highest Human Health Risk from Ambient PM_{2.5}?
—Dikshant and Lovleen Gupta
- 863–868 Split application of GA₃: Effective to improve growth, yield and quality of capegooseberry (*Physalis peruviana* L.)
—Madhvendra Bahadur Singh, Pawan Kumar, Manoj Kundu, Sanjay Sahay and Vivek Kumar
- 869–873 Study on Hydro Priming of Pomegranate Seeds for Germination and Seedling Vigour
—Tanumita Mondal and Ankan Das
- 874–878 Thermal Characteristics of Vernacular Architecture in Warm Humid Region
—Mihir N. Vakharia and Mahendra Joshi
- 879–885 Silicon Fertilizer – An Imperative Source for Enhancing Yield and Phytolith Content of Maize Hybrid in Desilicated Soil (*Typic rhodustalf*)
—A.M. Prabha, P. Christy Nirmala Mary, P. Saravana Pandian, T. Sivakumar and M. Shanthi
- 886–890 Extraction of Waterbodies using Support Vector Machine and Spectral Indices
—Shiva Chandra Vaddiraju, K. Bhavana, Apsana Shaik, R. Akhila and K. Sreeja
- 891–898 Responses of Physico-chemical Variables to Hypolimnetic Aeration in Lake Naukuchiyatal, Central Himalaya, India
—Mamta Pant, Pragya Singh and P.K. Gupta
- 899–904 Benthic Macro Invertebrate Diversity of Tropical River Kolar, from Nagpur, (M.S.), India
—Rawlekar K.N. and Sawane A.P.
- 905–906 Evaluation of Pigeon Pea (*Cajanus cajan* L. Millsp) Genotypes to bud Weevil Resistance
—B.D. Gite, P.V. Patil and R.M. Ambenagre
- 907–911 Evaluation of improved varieties of rice (*Oryza sativa* L.) under environmental condition of Namsai, Arunachal Pradesh, India
—Hibu Sunya, Avinash Sharma, Chowlani Manpoong, V.S. Devadas, Himanshu Pandey and Devendra Singh
- 912–914 Effect of Plant Growth Regulators (PGR) on flowering and fruiting of Pomegranate (*Punica granatum* L.) cv. Bhagwa under Prayagraj agro - climatic condition
—Saket Mishra, Jampani Nihari and Ravi Kumar Singh

- 915–921 Experimental Investigation and Modeling on Impacts of Leachate from Bio-manures and Fertilizers Used in the Agricultural Fields Along Noyyal River
—*E. Sarojini, Varatharajaperumal Thangavel, S. Kanchana, S.D. Anitha Selvasofia and S.T. Sudharsan*
- 922–927 An Urban Built Form and its Microclimate on Urban Heat Island - A Review
—*Nikhil P. Mehare and Mahendra Joshi*
- 928–934 Influence of Nitrogen and Boron Levels as Well as Macro and Micro Nutrient Content in Leaf on Production Potential of Guava
—*Bibha Kumari, Suraj Prakash and U.S. Jaiswal*
- 935–941 An Assessment of Genetic variability and association studies for yield and its components of soybean MAGIC population lines (*Glycine max* (L.) Merrill.)
—*Rajendar Reddy M., Sreedhar Chauhan, Uma Reddy R. and Sumalini K.*
- 942–945 Establishment of Pomegranate (*Punica granatum* L.) and observing the effect of organic and inorganic fertilizers on growth and development of Ganesh variety
—*Saket Mishra, Maddirala Bhargavi and Ravi Kumar Singh*
- 946–956 Boron in Soil Plant System and Its Significance in Indian Agriculture
—*MPS Khurana, Jatinder Singh, Vandna Chhabra, Talwinder Singh and Ankush Tarkha*
- 957–963 Global Challenges of Small Mammalian Carnivores: A Review
—*Samujjal Saharia, Malabika Kakati Saikia and Prasanta Kr. Saikia*
- 964–974 Morphometric Analysis of Baghlan Province for Amu River Basin in Afghanistan using Remote Sensing and GIS
—*Safiullah Sherzad and Tharavathy N.C.*
- 975–978 Environmental and Health Impact of Agriculture Chemicals: Paragquat, Chlorpyrifos and Glyphosate in Thailand
—*N. Vattanaprateep*
- 979–983 Noise Mapping During Deepawali Festival in Raipur City of Chhattisgarh, India
—*Vishal Kumar, A.V. Ahirwar and A.D. Prasad*
- 984–988 Soil physical properties as influenced by different agroforestry systems in Haryana
—*Garima Dahiya, K.K. Bhardwaj, Isha Ahlawat, R.S. Dhillon, Charan Singh and Sonia Devi*
- 989–999 Zooplankton Assemblages in a Heavily Polluted River Ecosystem (Pinang River, Malaysia): Impacts of Anthropogenic Stressors on the Communities
—*Azma Hanim Ismail, Ng Yang Qing and Wan Nor Afiqah Zulkarnain*
- 1000–1004 Nutritional value of six wild edible plants traditionally used as vegetables in Morocco
—*Ridwane Ghanimi, Ahmed Ouhammou, Hakim Baazizi, Khaoula Houguig and Mohamed Cherkaoui*
- 1005–1012 Potential use of Algerian natural diatomite as pozzolanic and insulator material in cement mortar
—*Bouksani Omar, Abba Fatiha, Belkacemi Marwa, Kharchi Fattoum, Benhadji Maissan and Belhamel Farid*
- 1013–1018 Agricultural Suitability and Static Modelling: A GIS Approach
—*Arushi Rana and Rashmi Sharma*
- 1019–1026 Exploring indigenous bacterial isolate from dye effluent sites for decolorization of Blue HERD dye
—*Shete U.R., Bhuktar J. J. and Jadhav A.G.*
- 1027–1030 The Effect of Salt Percentage Difference on the Nutritional Quality of Fermented Climbing Perch (*Anabas Testudineus* Bloch) Wadi from Central Kalimantan
—*Petrus Senas*

-
- 1031–1036 **Breeding success of Black kite (*Milvus migrans*) on Mobile Towers of Malappuram and Calicut Districts of Kerala**
—*Binu Chullakkattil and K. Seedikkoya*
- 1037–1041 **Impacts of Insurgency and Sensationalism on Eco-tourist Decision Making (TDM)**
—*Hafizullah Dar and Parwaiz Ahmad Najar*
- 1042–1047 **Spatial Variation in Elemental and Isotopic Compositions of Carbon and Nitrogen in a Coastal Lagoon Reveals Natural and Anthropogenic Influences**
—*Dang Thi Nhu Y.*
- 1048–1050 **Phenol colour reaction and its biochemical basis in cultivated wheat genotypes**
—*Payal Mathur and S. K. Chakrabarty*

Calcium Ionophore Supplementation in *In vitro* Fertilization Medium towards Fertilization Rates of Kacang Goats (*Capra hircus*)

Azizah Yasita Najati¹, Tjuk Imam Restiadi¹, Sri Pantja Madyawati¹, Imam Mustofa¹, Maslichah Mafruchati¹, Widjiati¹, Viski Fitri Hendrawan² and Epy Muhammad Luqman^{1*}

¹Department of Veterinary Science Faculty of Veterinary Medicine, Universitas of Airlangga, Kampus C Unair Jl. Mulyorejo Surabaya, 60115, Indonesia

²Department of Animal Reproduction Faculty of Veterinary Medicine, Universitas Brawijaya, Malang 65151, Indonesia

(Received 19 September, 2021; Accepted 20 October, 2021)

ABSTRACT

In vitro fertilization is assisted reproduction technology that can overcome infertility problems. This method can produce embryos with high quality in large quantities. In its application, the success rate of IVF is still relatively low. This is due to the poor quality of oocytes and spermatozoa, which can cause IVF failure. The absence of oocyte activation during fertilization also results in failure of fertilization. Calcium ionophore is known to be able to increase the oocyte activation process. This study aims to prove that Calcium ionophore can increase the rate of fertilization in goat oocytes *in vitro*. This study was divided into two groups: the control group was fertilized oocytes using a medium without Calcium ionophore supplementation, the treatment group was fertilized oocytes using a medium supplemented with Calcium ionophore at a dose of 5.2 µl/ml. Fertilization rates are observed based on the formation of zygote or two pronuclear (2PN) in the cytoplasm of oocytes and observed under an inverted microscope. The data obtained were analyzed using the T test. The results of the study after being analyzed statistically showed that there were significant differences between the control group and the treatment group ($P < 0.05$). Fertilization rates in the treatment group (93.12 ± 13.7) showed higher results than the control group (65.96 ± 28.36). Based on these results it can be concluded that calcium ionophore supplementation at a dose of 5.2 µl/ml in the fertilization medium can increase oocyte activation thereby increasing the rate of oocyte fertilization of the goat nuts.

Key words : Food production, Infertility, Oocyte activation, Medium, Supplementation.

Introduction

In the field of reproduction, infertility is a problem that is still commonly found in both humans and animals. According to the World Health Organization (WHO) in 2012, there were around 50-80 million couples experiencing infertility in the world. Efforts that can be made in tackling infertility are assisted reproductive technologies that are now in-

creasingly developing. One of the assisted reproductive technologies that are now popular for tackling the problem of infertility is *In vitro* Fertilization (IVF). IVF is a very profitable method, because besides being able to overcome infertility problems, this method can produce embryos with high quality in large numbers (Amin, 2000). Kacang goat is a local Indonesian goat whose existence began to be marginalized because of its small shape different

Corresponding author's email: epy-m-l@fkh.unair.ac.id

from outside goats like Boer goats. Although Kacang goat is an Indonesian germplasm, people are less interested because of the small body shape of the goat so that the number of carcasses is less. The success of IVF methods in conventional goats is still relatively low. With IVF technology, it is expected to be able to increase the population of Indonesian Kacang goats.

The success rate of IVF methods in conventional goats is still relatively low. Research conducted by Boediono *et al.* (2000) states that the success rate of *in vitro* fertilization carried out in goats only reached 40.91%. In addition, oocyte failure after fertilization to reach the blastocyst phase of *in vitro* fertilization is high, approximately until 60% (Gilchrist and Thompson, 2007). One of the factors that mostly support the success of IVF is the quality of spermatozoa and oocytes used. The quality of oocytes can be seen by the presence or absence of cumulus cells surrounding the oocytes, which act as chemo attractant spermatozoa to reach the oocytes so the fertilization occurs.

Oocytes with intact cumulus cells provide essential factors during the process of maturation, play a role in maintaining oocytes, during the stages of meiotic division and have the potential to be fertilized. Besides those spermatozoa also have an important role in the success of *in vitro* fertilization. During the fertilization process, spermatozoa make a very important contribution to the process of division and development of the embryo by providing oocyte activation factors, centrosome components, and paternal chromosomes (Dogan *et al.*, 2015). Oocytes activation is one of the factors that influence the success of fertilization. Oocyte activation can occur with complex interactions triggered by the entry of spermatozoa into the oocyte. An early indicator of oocyte activation is characterized by repeated increase of intracellular calcium concentrations (Karabulut *et al.*, 2018).

Calcium ionophore (CaI) A23187 is a compound commonly used to increase intracellular calcium (Ca^{2+}). CaI A23187 is a chemical compound that acts as a divalent cation ionophore, which allows these ions to pass through cell membranes that are generally inaccessible (Nalley and Hine, 2015). This allows the administration of CaI A23187 to increase intracellular Ca^{2+} levels in oocytes. An increase in intracellular calcium is an initiator mechanism of oocyte activation that can be observed after a complex interaction which triggered by the entry of

sperm cells into the oocyte during the fertilization process (Karabulut *et al.*, 2018). During fertilization, the endoplasmic reticulum in the oocyte releases Ca^{2+} ions which are known as important triggers for the development of oocytes into embryos (Hardy 2002). Increased Ca^{2+} levels in the cytoplasm of the oocyte will initiate pronucleus formation, which is a sign that the oocyte has been fertilized.

An increase in intracellular calcium levels can be increased by administering CaI A23187. Therefore, it is necessary to conduct research on supplementation of CaI A23187 in *in vitro* fertilization medium so that intracellular Ca^{2+} levels in oocytes can be increased.

Materials and Methods

This research went through several stages, i.e. oocyte collection, oocyte maturation, spermatozoa preparation, *in vitro* fertilization and observation of fertilization rates. This research has obtained ethical eligibility with the number: 1.KE.061.04.2019 from the Faculty of Veterinary Medicine, Universitas Airlangga.

Oocyte Collection

Kacang goat ovaries as a source of oocytes were obtained from Surabaya Abattoir as many as 54 ovaries. The ovaries are cleaned from the mesovary, then washed with 0.9% physiological NaCl to clean the remaining fat and blood attached to the ovaries and stored in 0.9% physiological NaCl which has been added with 50 $\mu\text{g}/\text{ml}$ gentamycin sulfate, at a temperature of 30-35 °C. The ovaries were brought to the laboratory to be washed again thoroughly using 0.9% physiological NaCl which had been added with 50 $\mu\text{g}/\text{ml}$ gentamycin sulfate. Furthermore, the oocyte is taken out from follicles with a follicular surface diameter of 3-5 mm, by aspirating ovarian follicular fluid using a 10 ml disposable syringe that has been filled with aspiration media namely Dulbeccos Modified Eagle Medium (DMEM-Sigma-Aldrich, Inc, St. Louis, Missouri United States). The aspiration follicular fluid is accommodated in a sterile tube, waited 5-10 minutes until sediment of oocyte show up. After 10 minutes the supernatant is removed and the sediment is poured in a petri dish and added with DMEM medium. The oocyte collection is then carried under inverted microscope. Oocytes from the collection are washed with DMEM medium until clean. The clean oocytes are selected and only oocytes in grade A, where it

coated with more than 3 layers of cumulus cells used for *in vitro* maturation were chosen.

Maturation of Oocytes

The maturation medium used was DMEM added with Pregnant Mare Serum Gonadotrophin (Folligon, PT Intervet Indonesia, Jakarta Indonesia) 50 µg / ml medium. The maturation medium is prepared 20 hours before being used in a warm way that is incubated in a 5% CO₂ incubator at 38.5 °C. Medium maturation in the form of medium drops with a size of 50 µl and covered with mineral oil (Sigma-Aldrich, Inc, St. Louis, Missouri United States). Grade A oocytes that have been selected after oocyte collection, are then washed using the maturation medium before being inserted into the drops of the maturation medium. In 1 drop, 50 µl maturation medium is filled with 5-6 oocytes. Then drop the maturation medium which has been filled with oocytes incubated in incubator of 5% CO₂ at 38.5 °C for 22 hours.

Preparation and Capacitation of Spermatozoa

Rosset medium in the form of large pits connected by bridges to small pits is prepared 20 hours before being used for spermatozoa preparation. The Rosset medium was incubated in a 5% CO₂ incubator at 38.5 °C to warm. The spermatozoa used for *in vitro* fertilization are derived from frozen semen of the goat nuts obtained from the Insemination Center for Animal Husbandry Education at the Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya. Spermatozoa preparation and spermatozoa capacitation were performed two hours before *In vitro* Fertilization. The frozen semen was thawed at 35°C, removed from the straw and put in a centrifuge tube. The centrifuge tube that has been filled with cement plus DMEM medium with a ratio of 1: 1, then centrifuged at a speed of 2500 g for 10 minutes. The supernatant is removed using a carefully sterilized Pasteur pipette. After that, spermatozoa pellets were dropped 150 µl into a large well on a prepared rosset medium. Medium form of rosset that already contains spermatozoa is left for 30 minutes to give the spermatozoa an opportunity to move towards a small well. After 30 minutes the motile spermatozoa will move towards small wells.

In vitro fertilization

Fertilization medium is prepared 20 hours before fertilization *in vitro*. In this study, prepared 2 Petri

dish *in vitro* fertilization medium consisted of 1 petri dish *in vitro* fertilization medium for the control group (P0): DMEM medium without supplementation, and 1 more petri dish for the treatment group (P): DMEM medium that was given CaI A23187 at a dose of 5.2 µl/ml. before use incubated in incubator of 5%CO₂ temperature 38.5°C. Oocytes from *in vitro* maturation were transferred to the fertilization medium which had previously been added with spermatozoa at a dose of 500,000-1,000,000 spermatozoa. After the spermatozoa and oocytes are mixed in the Fertilization medium then reinserted in a 5% of CO₂ incubator at 38.5 °C for 24 hours to give the spermatozoa time to fertilize the oocytes.

Observation of Fertilization Rates

Observation of fertilization rates were done 24 hours after the *in vitro* fertilization process and observe based on the formation of two pronuclear (2 PN) or zygote (Widjiati *et al.*, 2012). Oocytes that are successfully fertilized are characterized by the formation of two pronuclear (2 PN) in the cytoplasm of the oocyte which can be observed 24 hours after fertilization. Observation of the number of zygotes formed using inverted microscope Nikon Diaphot 300 with magnification of 40 times.

Statistical analysis

Data obtained from the number of fertilized and non-fertilized oocytes between the control group and the treatment group were analyzed using the statistical T Test. They were considered to be significantly different if P < 0.05.

Results and Discussion

The study compared fertilization rates performed on two fertilization mediums, namely the control group (P0) using the medium without supplementation and the treatment group using the medium with calcium ionophore supplementation at a dose of 5.2 µl / ml with each group performed 9 replications. The number of oocytes used as a source of gametes in this study is limited, because the number of goat's kacang ovaries obtained by the abattoir is little related to the goat kacang population which is starting to decline. Observation of oocyte number of goat nuts can be observed 24 hours after *in vitro* fertilization culture. Fertilization rates can be determined by observing the formation of 2PN in the oocyte cytoplasm, which can be observed under an

inverted Nikon Diaphot 300 type microscope with a magnification of 40 times. Based on data obtained using the Mann Whitney U test on SPSS, the results of the research from the two research groups showed significantly different results ($P < 0.05$). The results are shown in Table 1 and Fig. 1.

In the treatment group, the oocytes were fertilized using CaI A23187 supplementation medium showed a higher percentage of fertilization rates than the oocyte group which was fertilized with the medium without supplementation (Table 1). Based on the table above shows that the control group was significantly different from the treatment group ($P < 0.05$), this shows that CaA A23187 supplementation on the fertilization medium was proven to increase the rate of fertilization by the method of *in vitro* fertilization. In the treatment group, ie oocytes were fertilized using CaI A23187 supplementation medium showed a higher percentage of fertilization rates than the oocyte group which was fertilized with medium without supplementation (Table 1). Based on the table above shows that the control group was significantly different from the treatment group ($P < 0.05$), this showed that supplementation of CaI A23187 in the fertilization medium was proven to be able to increase the rate of fertilization by *in vitro* fertilization methods.

In vitro fertilization culture involves the fusion of mature oocytes and spermatozoa that are cultured in a medium. Mature oocytes can be obtained by maturation *in vitro* with medium maturation. After the *in vitro* maturation process, the mature oocytes are characterized by the expansion of the accumulation cells, which can be observed under an inverted microscope (Fig.1A). The mature oocytes are transferred in a fertilization medium which is then reunited with the spermatozoa which are capacitated. Capacitated spermatozoa move towards the oocyte

and will bind to the oocyte pellucida zone. After binding to the oocyte pellucida zone, spermatozoa will penetrate the pellucida zone which will then occur fusion between spermatozoa and oocytes. The fusion process between spermatozoa and oocytes is called the fertilization process (Fig. 1B). The process of fertilization *in vitro* can be declared successful if in the cytoplasmic the oocyte forms 2 pronucleus which can be observed 24 hours after the culture of fertilization under microscopic inverted (Fig. 1C).

In vitro Fertilization (IVF) is a reproductive biotechnology that is commonly used to overcome infertility problems that have been widely applied both to humans and animals. IVF is a method of fertilization that is carried out outside the parent body in a medium. This method is a beneficial method because it can improve the genetic quality of the embryos produced and overcome problems related to infertility. The occurrence of fertilization has a close relationship with oocyte activation. Oocyte activation is initiated by repeated oscillations or intracellular Ca^{2+} levels. This oscillation is initiated by the release of Ca^{2+} from inositol 1,4,5-trisphosphate (InsP3) which plays a role in increasing Ca^{2+} concentration by activating the InsP3 Receptor (InsP3R) which causes rapid calcium release and results in increased Ca^{2+} levels in the cytoplasm (Dutta, 2000). Increased intracellular Ca^{2+} levels contribute to the resumption of meiotic division and the formation of pronuclei in cytoplasmic oocytes, which is a sign that the oocyte has been fertilized (Nomikos *et al.*, 2017).

Oocyte activation is the transition from oocytes to zygotes where a series of intracellular Ca^{2+} oscillations will occur initiated by the fusion between spermatozoa and oocytes. Calcium oscillations play an important role in fertilization, and calcium ions act as secondary messengers in different metabolic pathways and cellular processes during oocyte acti-

Table 1. Mean, standard deviation and percentage of effect of post fertilization CaI A23187 supplementation on *in vitro* fertilization rates.

Observation	Group of Treatment		P
	P0 (Without CaI Supplementation)	P1 (With CaI Supplementation)	
Fertilizes Oocytes	65,9 ^b ±28,36	93,1 ^a ± 13,70	0.028*
Infertilized Oocytes	34,04 ^a ±26,74	6,88 ^a ± 13,70	0,244

* p < 0.05 indicates there is a significant difference

Information:

P0 = Control group without CaI supplementation 23187

P1 = Treatment group (With CaI supplementation 23187)

vation. Upon oocyte activation, calcium ions are released from intracellular storage in the endoplasmic reticulum; free in the cytosol, which functions as an intracellular messenger and acts as a process modulator in the early stages of fertilization and embryo development (Hojnik and Kovacic, 2019). Under normal circumstances, the presence of sperm in the cytoplasm of the oocyte will evoke characteristic patterns of intracellular calcium oscillation, where extrusion of polar bodies will occur, exocytosis of cortical granules, cytoskeletal rearrangement, recruitment of maternal mRNAs, and formation of pronucleus (Yeste *et al.*, 2017).

In the development of embryos in the IVF method, oocytes are placed in a medium that is made to resemble the conditions in the female reproductive tract and contain nutrients needed for each development. The composition of nutrients in the culture medium is made close to the composition of nutrients, electrolytes, and macromolecules that are in the female reproductive tract (Widjiati *et al.* 2012). In the process of fertilization, spermatozoa must pass through cumulus cells, the pelucida zone, and

the vitellin membrane. Entry of spermatozoa into the egg cell involves attaching the head of spermatozoa cells to the surface of the vitellin membrane is a very important period because this is when oocyte activation occurs. Oocyte activation is initiated by an increase in intracellular Ca^{2+} levels which is a signal that contributes to the resumption of meiosis and early embryonic development (Karabulut *et al.* 2018).

In a culture medium in IVF, conditions of the female reproductive tract cannot be fully replicated *in vitro*, but the ability of the medium used in the *in vitro* fertilization method can be improved by adding various substances and compounds to the culture medium. Modification of *in vitro* culture conditions is one technique that can be used to increase the rate of fertilization and viability of blastocysts (Widjiati *et al.*, 2017). The addition of supplements to the culture medium aims to increase the success rate of IVF depending on the function of substances and compounds used.

Calcium ionophore (CaI) A23187 is a compound commonly used in research to increase intracellular




Fig. 1. (A) Mature oocytes characterized by expansion of cumulus cells (1), (B) Spermatozoa that penetrate the oocyte pelucida zone walls (2), (C) and fertilized shell oocytes which are characterized by the presence and 2 pronucleus (3) observed under an inverted microscope at 40x magnification

calcium. This substance has been shown to be able to activate cells by inducing Ca^{2+} signaling, intracellular Ca^{2+} mobilization, and activating Ca^{2+} entry (Dedkova *et al.*, 2000). In the field of *in vitro*, CaI A23187 is often used in the method of intra cytoplasmic sperm injection (ICSI) which is an IVF method by injecting spermatozoa on the cytoplasm of oocytes using a micromanipulator. In the ICSI method, CaI A23187 plays a role in activation after spermatozoa injection (Eftekhar *et al.*, 2012). The study compared the use of fertilization medium without supplementation (control group) and medium supplemented with calcium ionophore (treatment group). Supplementation was carried out using Dulbeccos Modified Eagle's Medium (DMEM) as a base medium supplemented with CaI A23187 at a dose of 5.2 $\mu\text{l}/\text{ml}$. In the treatment group, the mature oocytes were cultured on supplementation medium and then incubated in CO_2 incubators for 24 hours. DMEM media was a basal medium consisting of vitamins, amino acids, salts, glucose, and pH indicators. This media does not contain protein or growth agents. This media requires complete supplementation such as Fetal Bovine Serum (FBS) (Rashid and Coombs, 2019).

In this study, IVF culture with supplementation medium was administered using doses with low concentrations and was the optimum dose according to the research conducted by Maitra (2018) which was 5.2 $\mu\text{l} / \text{ml}$. Giving calcium ionophore with excessive doses can cause oocyte damage because prolonged exposure to CaI A23187 can cause toxins (Swann, 2018). In addition, increased levels of Ca^{2+} with high concentrations can affect embryonic development. This is consistent with research conducted by Nalley and Hine (2015) which states that post-exposed CaI A23187 embryos with high concentrations indicate lower cleavage rates.

The results showed that oocytes cultured by IVF using a supplemented medium CaI A23187 showed a higher fertilization rate (93.12 ± 13.7) compared to oocytes cultured using a medium without supplementation that is (65.96 ± 28.36). This result is higher than research conducted by Taylor *et al.* (2008) where the number of oocyte fertilization reached (53.2 ± 19.8) in humans. Research conducted by Eftekhar *et al.* (2012) using CaI A23187 at a dose of 5 μL given after ICSI showed fertilization rates reaching 72.5%. The success of fertilization with the method of *in vitro* fertilization shows varied results, which depend on several influential factors. These

influential factors are the quality of the oocytes and spermatozoa used, and also the factors in the culture medium which include temperature, pH, osmotic pressure, and CO_2 levels.

This shows that oocytes fertilized using a medium supplemented with calcium ionophore have been shown to increase fertilization rates. Ionophore calcium can increase intracellular calcium levels in oocytes both internally and externally, which internally CaI A23187 works in internal storage through channels in the endoplasmic reticulum and externally depends on extracellular Ca^{2+} entry (Ebneretal, 2014). Repeated increases in intracellular Ca^{2+} levels are signals that initiate oocyte activation and pronucleus formation. In mammals, repeated increases of Ca^{2+} in oocytes that occur during fertilization are signals that play a role in early embryonic development, and without the presence of Ca^{2+} signals during fertilization can cause failure in embryo development (Ducibella *et al.*, 2006). The conclusions from this study shows that these supplementation of Calcium Ionophore (CaI) A23187 in the vitro fertilization medium at a dose of 5.2 $\mu\text{l}/\text{ml}$ can increase oocyte activation so as to increase the oocyte fertilization rate of goat nuts

Conflict of Interest

Authors declare that they have no conflict of interest.

Acknowledgments

The authors express sincere thanks to the Ministry of Research, Technology and Higher Education of the Republic of Indonesia for funding research and Dean Faculty of Veterinary Medicine for providing all necessary facilities and fund for conducting research work.

Authors' Contribution

AYN designed the study, interpreted the data, and drafted the manuscript. TIR, SPM, IM, MM and VFH were involved in collection data and also contributed in manuscript preparation. AYN, EML and WW took part in preparing and critical checking of this manuscript.

References

Amin, M. 2000. Molecular mechanism of fertilization process in animals. *Hayati*. 7 : 117-120.

- Boediono, A., Rusiantono, Y., Mohamad, K., Djuwita, I. and Herliatien, 2000. Goat Oocyte Development After Maturation, Fertilization, and *In vitro* Culture. *Media Vet.* 7 : 1-17.
- Dedkova, E. N., Sigova, A. A. and Zinchenko, V. P. 2000. Mechanism of action of calcium ionophores on intact cells: Ionophore-resistant cells. *Membr. Biol.* 13: 357-368.
- Dogan, S.P., Vargovic, R., Oliveira, L. E., Belser, A., Kaya, A., Moura, A., Sutovsky, P., Parrish, J., Topper, E. and Memili, E. 2015. Sperm protamine-status correlates to the fertility of breeding bulls. *Biol Reprod.* 92: 1-9.
- Ducibella, T., Schultz, R. M. and Ozil, J. P. 2006. Role of calcium signals in early development. *Semin Cell Dev Biol.* 17 : 324-332.
- Dutta D 2000. Mechanism of store-operated calcium entry. *J. Biosci.* 25 : 397-404.
- Ebner, T., Mayer, R. B., Oppelt, P. and Gruber, I. 2014. Treatment with Ca²⁺ ionophore improves embryo development and outcome in cases with previous developmental problems: A prospective multicenter study. *Human Reprod.* 30 : 97-102.
- Eftekhari, M. F., Yousefnejad, F., Khani, P. and Aflatoonian, A. 2012. Effect of calcium ionophore on unfertilized oocytes after ICSI cycles. *Iran J Reprod Med.* 10 : 83-86.
- Gilchrist, R. and Thompson, J. 2007. Oocyte maturation: Emerging concepts and technologies to improve developmental potential *in vitro*. *Theriogenology.* 64: 6-15.
- Hardy DM 2002: Fertilization. California: Academic Press.
- Karabulut, S., Aksünger, O., Ata, C., Sađýroglu, Y. and Keskin, I. 2018. Artificial oocyte activation with calcium ionophore for frozen sperm cycles. *Syst Biol Reprod Med.* 64 : 381-388.
- Ksiazkiewicz, L. K., Opiela, J. and Rynska, B. 2007. Effects of oocyte quality, semen donor and embryo co-culture system on the efficiency of blastocyst production in goats. *Theriogenology.* 68 : 736-744.
- Maitra, D. W. 2018. Post Denudation Procedures: Basic ICSI Overview. PIFE 1st Meeting: Fertility 360°.
- Nalley, W. M. and Hine, T. M. 2015. Mice Parthenogenetic Embryo Activation and Development Rate After Exposure to Calcimycin and Ionomycin. *Jurnal Veteriner.* 16 : 576-584.
- Nomikos, M., Kashir, J. and Lai, F. A. 2017. The role and mechanism of action of sperm PLC-zeta in mammalian fertilisation. *Biochem J.* 474 : 3659-3673.
- Ondho, Y. S. and Anwar, S. 2012. Handling and Storage of Ovaries from Slaughterhouses as a Source of Oocytes for *In vitro* Embryo Production. *Proceeding of National Seminar on Zootechniques for Indigenous Resources Development.*
- Rashid, M. and Coombs, K. M. 2019. Serum-reduced media impacts on cell viability and protein expression in human lung epithelial cells. *J Cell Physiol.* 234(6): 7718-7724.
- Swann, K. 2018. The role of Ca²⁺ in oocyte activation during *in vitro* fertilization: insight into potential therapies for rescuing failed fertilization. *Mol Cell Res.* 1865 : 1830-1837.
- Taylor, T. H., Wright, G., Jones-Colon, S., Mitchell-Leef, D., Kort, H. I. and Nagi, Z. P. 2008. Comparison of ICSI and conventional IVF in patients with increased oocyte immaturity. *Reprod Bio Mede Ol.* 17 : 46-52.
- Widjiati, Pusporini, S. E. and Arifin, M. Z. 2012. Perbandingan Angka Fertilitas and Hambatan Perkembangan Embrio Mencit yang Dikultur Dalam Medium M16 and Human Tubal. *Fluid. J. Vet.* 13 : 227-234.
- Widjiati, Luqman, E. M. and Tehupuring, B. C. 2017. Effectivity of Insulin Transferrin Selenium and Bovine Serum Albumin Addition on *In vitro* Culture Medium on Fertilization and Blastocyst Rate of Mice (*Mus musculus*). *J. Int. Dent. Medical Res.* 10 : 1080-1083.
- Yeste, M., Jones, C., Amdani, S. N. and Coward, C. 2017. Oocyte Activation and Fertilization: Crucial Contributors from the Sperm and Oocyte. In: Signaling-Mediated Control of Cell Division. Springer Link. 59: 213-239.
-