

QUARTERLY INTERNATIONAL JOURNAL


# ECOLOGY

ENVIRONMENT & CONSERVATION

ISSN 0971-785 X



(Supplement Issue)  
February, 2022



IEM INTERNATIONAL

# Quarterly International Journal Ecology, Environment and Conservation

ISSN 0972-3005

Editor: Dr. R.K.Trivedy, Pune, India

## EDITORIAL ADVISORY BOARD

1. Dr. Teresa Ferreira, Portugal
2. Dr. Rakesh Kumar, AICTE, New Delhi, India
3. Mr. Sandeep Singhal, Chennai, India
4. Dr. M.M. Amin, Rajshai, Bangladesh
5. Dr. D.J. Lee, Taiwan
6. Dr. J.K. Dutta, Burdwan, India
7. Dr. K. Venkateshwarlu, Anantpur, India
8. Dr. Linda Blackwell, Australia
9. Dr. G. Zellner, Netherlands
10. Dr. I.C. Onyema, Nigeria
11. Dr. M.F. Hamoda, Kuwait
12. Dr. H.A. Abrahamse, South Africa
13. Dr. Rowshan Mumtaz, Bangladesh
14. Dr. Hassan Poorbahaei, Iran
15. Dr. C.K. Yap, Malaysia
16. Dr. Hassan Ibrahim Ali, Sudan
17. Dr. M.T. Udo, Nigeria
18. Dr. Mohd. Nural Anwar, Bangladesh
19. Dr. A.K. Kumaraguru, Madurai, India
20. Dr. A.U. Pagarkar, Ratnagiri, India
21. Dr. Ganesh Iyer, Mumbai, India
22. Dr. A.R. Ghosh, Burdwan, India
23. Prof. M. Zaman, Bangladesh
24. Dr. Marcantonio Bragdin, Venice, Italy
25. Dr. Ramesh Thatikunta, Hyderabad, India
26. Dr. Z. Li, Bonn, Germany
27. Dr. Anju Singh, Mumbai, India
28. Dr. Ing. Agr. Mario Ridardo Sabbatini, Argentina
29. Dr. Philip C. Reid, U.K.
30. Dr. Amarjeet Kaur, New Delhi, India
31. Dr. B.N. Prasad, Noida, India
32. Dr. Oswaldo A. Fernandez, Argentina
33. Dr. A. Olawale, Nigeria
34. Dr. Ms. Mirela Tulik, Warsaw, Poland
35. Dr. Piroz Azizi, Iran
36. Dr. Ashwini Wangenao, Bhopal, India
37. Dr. G. R. Pathade, Pune, India
38. Dr. P.S. Panesar, Longowal, India
39. Dr. Meenakshi C. Mahajan, Pune, India
40. Dr. Mohd. Yusuf, Malaysia
41. Dr. A.K. Pnigrabi, Berhampur, India
42. Ms. Aditi Trivedi, Pune
43. Dr. Azni Idris, UPM, Malaysia
44. Dr. T. Bahorus, Mauritius
45. Dr. B.K. Sharma, Shilong, India
46. Dr. E. Amiri, Iran
47. Dr. T.A. Yakeen, Nigeria
48. Dr. L.L. Chukwu, Nigeria
49. Dr. Michael Ukwuru, Nigeria
50. Dr. Sabrina Naz., Bangladesh

### Ecology, Environment and Conservation is Abstracted in -

- > Chemical Abstracts, U.S.A.
- > Ecological Abstracts, U.S.A.
- > Cambridge Science Abstracts, U.S.A.
- > Ecodisc-CD-Rom, Netherlands
- > Geological Abstract, Netherlands
- > Pollution Abstract, U.S.A.
- > International Dev. Abstracts, U.S.A.
- > Paryavaram Abstracts, India
- > Current Awareness in Biological Sciences, Elsevier
- > World Textiles Abstracts, Netherlands
- > Fluid Abstracts, Netherlands
- > Environmental Abstracts, India
- > Indian Science Abstracts, India
- > Cambridge Science Abstracts

### Impact Rating 3.5 (NAAS 2011)

**COPYRIGHT:** This journal is registered at the COPYRIGHT CLEARANCE CENTRE INC. (CCC), 222, Rosewood Drive, Suite 910, Danver, MA01923, U.S.A. The copyright owner consents that in the U.S.A. copies of the articles may be made for personal or internal use, of specific clients, on payment of fee. All requests for permission to photocopy should be addressed to the copyright owner ([www.copyright.com](http://www.copyright.com))

The journal figures in Ulrich International Periodicals Directory, U.K., Gale Directory, U.K. & SAARC Directory of periodicals.

### SUBSCRIPTION RATES

	INDIA	OVERSEAS
Individuals	Rs : 1000.00	US\$ : 150.00
Institutional	Rs : 2000.00	US\$ : 600.00

Note : All payments in favour of EM INTERNATIONAL (Postage included in Subscription)

Publishers : EM INTERNATIONAL, Prakriti, Balewadi, Baner 411 045, M.S., India

☎( 91-2164- 220369; 20- 46745119; E-mail : [rktm@pn3.vsnl.net.in](mailto:rktm@pn3.vsnl.net.in)

# ECOLOGY, ENVIRONMENT AND CONSERVATION

## VOL. 28 (February Suppl. Issue) : 2022

### CONTENTS

- S1–S5 Parental investment of adult Barn swallow *Hirundo rustica rustica* at nestling feeding in the urban locality of El Bouni (Northeast, Algeria)  
—*Dadci Walid, Sakraoui Rym, Boukheroufa Mehdi, Abdallah Khadidja Wissal, Sakraoui Ferial and Tahar Ali*
- S6–S13 Land-Use/ Land-Cover Changing Trends and their Relation to Landslides: A Case Study of Hali-Elain Badulla District, Sri Lanka  
—*A.M.R.N.K. Alahakoon, K.L. Thisara Sathsara, Udara S.P.R. Arachchige and Shyamantha Subasinghea*
- S14–S20 Regional and Seasonal Variations of Polychlorinated Biphenyls (PCBs) in Water of Shatt AL-Arab River, Iraq  
—*Rafid A.T. Al-Zabad, Ayad H.D. Al-Khafaji and Hamid T. AL-Saad*
- S21–S27 Synthesis and Characterization of Nanocellulose from *Cladophora* sp.  
—*S.W. Suciayati, P. Manurung, Junaidi, S. Sembiring and R. Situmeang*
- S28–S34 A Teacher's Perception of Green Building Technology: Implications to Sustainable Construction and Environmental Education in Anambra State, Nigeria  
—*Daniel Uchenna Chukwu, Hyginus Osita Omeje, Godwin Keres Okereke, Chinwe Patricia Eze and Ikechukwu Chidiebere Odogwu*
- S35–S40 Antitumor and antioxidant activity of *Aloe Vera* leaf gel extract  
—*Worood Kamil Shalash*
- S41–S45 Effect of Polyphenol Fraction Exposure of *Aloe vera* on the Change in Cortisol of Koi Fish (*Cyprinus carpio*) as Stress Response  
—*Sri Andayani, M Sulaiman Dadiono, Widya Tri Elwira, Husna Nabila, Rodhyansyah, Nelvan Subayu and Ashari Fahrurrozi*
- S46–S51 Regulatory Framework for the Disposal and Management of Personal Protective equipment (PPE) During Covid-19 in Bangladesh  
—*Farhana Helal Mehtab, Md. Safiullah and Md. Arif Mahmud*
- S52–S58 Solar Cell as Energy Chlorinator for Disinfection of Flowing Water  
—*Beny Suyanto, Denok Indraswati, Tuhu Pinardi and Sujangi*
- S59–S61 Production Performance of Broiler Chicken Supplemented with *Lactobacillus plantarum* and *Lactobacillus casei* Incubated in Different Medium Infusion  
—*Nabila Hanina Hibatul Haqqi, Fatihah Istafaro Maulidiya, Galuh Hesti Dwi Indrawati<sup>1</sup>, Wenny Nur Azizah, Herinda Pertiwi and Loh Teck Chwen*
- S62–S73 Motivators and Inhibitors of Green Building Technology Integration and Advancement: Nigerian Lecturers' perspective  
—*D. U. Chukwu, H. O. Omeje, G. K. Okereke, B. A. Omeje, A. A. Okekpa and J. A. Okereke*
- S74–S78 The Impact of Electromagnetic Pollution on Human Health and Environment: Recommendation for an Effective Regulatory Framework in Bangladesh  
—*Arif Mahmud*
- S79–S84 Differences effect of red and big white ginger extract as anti-inflammatory agents by *In vitro*  
—*E.B. Aksono, T.D.K. Wungu, C.H. Wijaya, R.R.F. Sarassina, N. Hidayatik, H. Pertiwi and N. Puspitasari*
- S85–S92 Visualization of Coral Reef Cover with Photogrammetry Method at Coastal Waters of Lemon Island, Manokwari, Indonesia  
—*Dimas Algutomo, Ricardo F. Tapilatu, and Aradea Bujana Kusuma*

- S93–S98 Restoring River Water Pollution from the Impact of Water Hyacinth (*Eichornia crassipes*) through Utilization as Raw Material for Fertilizer  
—*Hery Koesmantoro, Karno, Sunaryo and Suparji*
- S99–S102 Effect of Slope on Landslide Potential at Kuranji Watershed  
—*Andriani, Hendri Gusti Putra, Masril Syukur and Ulung Sadewo*
- S103–S110 Isolation of Fungi from Oil-contaminated Soil from Maysan Province, Southern Iraq  
—*Ahmed R. Mossa and Ali A. Kasim*
- S111–S119 Ecological Diagnosis of the Algerian West Coast Marine Environment by Using Benthic Macroalgae  
—*Imene Bouri, Omar Rouane-Hacene and Zitouni Boutiba*
- S120–S124 Effectiveness of Adsorbents Chicken Eggshells and Cassava Skin to Reduce Pollutants in the Electroplating Industry  
—*Muhammad Al Kholif, Pungut, Sugito and Djoko Adi Walujo*
- S125–S127 Impact of Lockdown on Dengue in Kolkata: A Socio-Ecological Study  
—*Abhishek Mondal and Saheli Guha Neogi Ghatak*
- S128–S130 Potable Water Quality of Thenhlum Village in Lunglei District, Mizoram, India  
—*Malsawmtluanga*
- S131–S134 Heavy Metal Concentration in Surface Water around the Industrial Area M.I.D.C. Mahad, Maharashtra, India  
—*Srikant Kekane and R.P. Chavan*
- S135–S142 Study of Microbial Diversity and their Role in the Deterioration of Tarkeshwar Temple, Jaipur, India  
—*Mansvi Yadav and Varsha Gupta*
- S143–S146 Diversity of Mangrove Ecosystem from Vaitarna Estuary (Palghar), Maharashtra, India  
—*Harshad S., Vanmali and Rahul N. Jadhav*
- S147–S152 Assessment of Physico-chemical Characteristics of Vairtarna Estuary of Palghar, (M.S.) India  
—*Harshad Vanmali and Rahul Jadhav*
- S153–S162 Linkages of Sustainable Development Goals with Integrated Coastal Zone Management Project in India: An Approach to Enhance Efficacy  
—*Satyendra Kumar and C. R. Suthar*
- S163–S167 The Influence of Environmental Shock Initiation on Vegetative and Spore Production of *Bacillus megaterium*  
—*I Made Dedi Mahariawan, Wahyu Endra Kusuma, Ating Yuniarti and Anik Martinah Hariati*
- S168–S174 Impacts of the Changing Housing Patterns on the Populations of House Sparrow *Passer domesticus* (Passeriformes: Passeridae) in the Rural Hills of Uttarakhand: A Case Study  
—*Ritika Waldia and Jay Prakash Bhatt*
- S175–S179 An Overview on Facile Green Synthesis Approaches and Optical Sensing Applications of Silver Nanoparticles  
—*Sugandha Sangar and Kulvinder Singh*
- S180–S182 Detection of Valuable Material-protein, in Faecal Sludge  
—*Hansa K., M.C. Sampath Kumar and Priya V.*
- S183–S187 Effect of Value Added Admixture Zeolite on Setting Time and Compressive Strength of Magnesia Cement  
—*R.N. Yadav and Ramkishan*
- S188–S193 Genetic Variability Studies for Yield and Yield Attributing Traits of Wheat (*Triticum aestivum* L.)  
—*Mayur Roshan Sahoo, Aarti Sharma and Hemant Kumar Jaiswal*

- S194–S205 **Life Without Antibiotics? - A Review on Increasing Threat of Antibiotic Resistance with A Broader Look on Agricultural aspect**  
—*Debanjana Debnath, Chinmayee Mohapatra, Snehasish Routray and Suraj Goldar*
- S206–S209 **Assessing Land -use, Misuse and Abuse in Nilgiris District of Tamil Nadu Using Remote Sensing Data**  
—*M. Suguna Devakumari, R. Alex Immanuel Jeyasingh, S. Praveena Katharine and S. Carolin Jeeva*
- S210–S212 **Anthropogenic Drives of Emerging Infectious Diseases and Possible Mitigation Approaches**  
—*Pinak Dutta and Mita Dutta*
- S213–S216 **Eco-sensitive Zones and Ecologically Fragile Lands in Kerala: Known and Unknown**  
—*Liji Samuel*
- S217–S226 **Traditional Medicinal Values of *Rubus ellipticus* with Biological Activities Observed from its Crude Extract: A Review**  
—*Vinit Prakash, Anjana Kumari, Harpreet Kaur, Manoj Kumar, Sumeet Gupta, Ritu Bala, Dinesh Gupta*
- S227–S232 **Study on Development of Consortia for NPK Bacteria and its Effect on Growth of Selected Crops**  
—*Nileema S. Gore, Vaishali Hiwale, Vaibhav Jagdale and Sanjay Harke*
- S233–S241 **Effect of Organic Fertilizer on Growth, Yield and Quality of *Pisum sativum* L.: A Review**  
—*Sarbeswar Behera, B. Jyotirmayee, Ushashee Mandal, Ananya Mishra, Pratyasha Mohanty and Gyanranjan Mahalik*
- S242–S250 **Kriging Interpolation Based Error Analysis of Epic Model with Respect to Standard K Factor Method for Gurushikhar, Rajasthan, India**  
—*Narendra Kumar Maurya, Prakash Singh Tanwar and Yusuf A.S.*
- S251–S257 **A Review of techniques to eliminate fluoride from water**  
—*Amit Punia, M.S. Narwal and A.K. Berwal*
- S258–S264 **Enzyme Analysis and Histological Studies of Earthworms in the Treatment of Different Organic Wastes**  
—*G. Priya, Vimala Bangarusamy, Nasreen Najeeb and P. Mahalakshmi*
- S265–S272 **Study of soil moisture distribution under different mulching methods**  
—*Nitin M Changade and Raj Kumar*
- S273–S274 **Augmenting the Geotechnical Characteristics of Soil by Dry Bagasse and Fly Ash**  
—*Tanu Kumar and Ashok Kumar Gupta*
- S275–S278 **Green Energy Bricks: A Step Towards Smart and Sustainable Structures**  
—*Pratibha Singh, Vivek Sharma, Sandeep Gupta, Savitha H.S., Asha Upadhyay and Nupoor Srivastava*
- S279–S283 **A Questionnaire-based Survey on Groundwater Characterization**  
—*N. Sridhar, K. Ezhisaivallabi and S. Poongothai*
- S284–S290 **Assessment of Rooftop Rainwater Harvesting Potential in Karnal Smart city, (Haryana) using Geospatial Technology**  
—*Meenakshi, K.E. Mothi Kumar and Nisha Kumari*
- S291–S293 **Correlation of Chemical Attributes of Soil with Yield of Wheat Crop**  
—*Amit Kumar Pandey, Ashutosh Singh and Umesh Singh*
- S294–S299 ***Eria meghasaniensis* (S. Misra) S. Misra, an endemic and endangered orchid of Similipal Biosphere Reserve, Odisha, India**  
—*D. Behera, S.S. Jena, B.T. Panda and H.N. Subudhi and S.P. Panda*

- S300–S306 Evaluation of Heavy Metals Contamination in Suspended Sediments of Dikrong River, Assam, India  
—*Satya Ranjan Goswami, B. J. Saikia, A. Goswami, A. J. Dev Sarma, Md. Y. Hassan and M. Buragohain*
- S307–S312 Temperature trend and variation analysis in the eastern plain zone of Uttar Pradesh  
—*Devegowda S.R., Saket Kushwaha, P.S. Badal, Avdhesh Sharma and Nagaveni M.*
- S313–S321 Optimal Annual Allocation Model of Large Storage Based Irrigation Project by Multi Level Approach  
—*A. Srinivasa Prasad, G.K. Viswanadh, P.V.S. Maruthi Krishna and K. Leela Krishna*
- S322–S327 Hydrobiological Study of Lotic Water of Yamuna River at Yamuna Nagar, Haryana, India during 2018-2019  
—*Babita and Sushil Kumar Upadhyay*
- S328–S339 Sanitation Condition and its Association with Menstrual Hygiene of Non-Tribal and Tribal Communities in Selected Districts of West Bengal  
—*Lopamudra Ganguly and Lakshminarayan Satpati*
- S340–S343 Effect of Air Pollution and Oxidative Stress in Development of Cataract- A Study in Rural Regions of West Bengal  
—*Somnath Ghosh, Sayani Pal Choudhury, Gopeswar Mukherjee and Animesh Dey*
- S344–S350 Impact of Jhum Cultivation on Forest Ecosystem and Environment Management Policies in Meghalaya, India  
—*Trishna Moni Tamuli and Ashok K. Bora*
- S351–S362 Plant disease diagnosis using image processing techniques – A review on machine and deep learning approaches  
—*Kalpesh Patel and Atul Patel*
- S363–S366 Performance of contingent crops under deferred sowing in transitional plain of luni basin in Western Rajasthan  
—*Dharmendra Meena, Santosh Kumari Meena, Saresh, N.V., Asu Singh Godara, Nema Ram, Arjun Lal Bijarnia and Dileep Kumar*
- S367–S371 A Study on Air Quality Index for Tiruchirappalli City Corporation  
—*Vignesh PM, J. Fowmithabanu, P. Kalaipriya, K. Sridhar and M. Govindaraju*
- S372–S381 Ameliorative effect of *Nelumbo nucifera* seeds and wheat grass formulated feed against tannery effluent toxicity and stress in freshwater fish *Oreochromis mossambicus*  
—*M. Sankar, T. Kumaran and R. Saravanan*
- S382–S386 Multinutrient Extraction – Need of the Hour – A Review  
—*Praveena Katharine Stephen, Suguna Devakumari, M. and Susan Poonguzhali, R.*
- S387–S398 Comprehensive Study on Raniganj Coalfield Area, India: A Review  
—*Dibyendu Saha, Jai Prakash Keshri and Nimai Chandra Saha*
- S399–S405 Extraction of Bioactive Compounds in *Ganoderma lucidium* from Himachal Pradesh  
—*K. Theresa, Vimalin Hena<sup>2</sup> and Jasmin Dora*
- S406–S411 Macronutrient availability and microbial population in fly ash amended alluvial soil  
—*K. Theresa and J. Vimalin Hena*
- S412–S424 Potential of Sodium Hydroxide Pre treatment of *Prosopis juliflora* for Fermentable Sugar Production  
—*Vijayakumar Palled, M. Anantachar, M. Veerangouda, K.V. Prakash, C.T. Ramachandra, Nagaraj M. Naik and R.V. Beladadhi*
- S425–S435 Population dynamics of *Maydelliathelphusa masoniana* Henderson 1893 from freshwater bodies of Jammu region (Jammu & Kashmir, India)  
—*Harmeen Kour, Seema Langer, and Nipoon Sharma*

- S436–S443 A Multi Criterion Decision Approach for Identification of Potential Flood Risk Zones in Hussainsagar Watershed using Geomatics – A Case Study  
—*Padmaja G. and Giridhar M.V.S.S.*
- S444–S451 Diversity and abundance of Macrozoobenthic communities in a Gangetic flood plain wetland of West Bengal, India  
—*D. Sarkar (Paria)*
- S452–S458 Analyzing River Bank Erosion Vulnerability of Sadiya Region, Assam, India  
—*Ratul Das and Barnali Gogoi*
- S459–S469 Forecasting the Emission of Carbon-di-oxide Equivalent in Key Sectors of India Using Arima Model  
—*Deepa S. and Vijayasree G.*
- S470–S478 Seed Source Determines Germination and Survivability in *Saussurea obvallata* (Dc.) Edgew an Endemic Herb of Indian Himalayan Region, India  
—*Vijay Kant Purohit, Pradeep Dobhal, Bhagat Singh Mengwal, Jaidev Chauhan, Mohan Chandra Nautiyal, Anant Ram Nautiyal and Pratti Prasad*
- S479–S487 The fragile Environment, ever growing tourism and the quest for sustainability: A case study from Duars, Alipurduar, West Bengal  
—*Gobinda Chandra Roy, Manjil Gupta and Siddharthasankar Banerjee*
- S488–S491 Physico-chemical Investigation of Neera River Water Near Sarola Bridge Shirval, District Pune (Maharashtra), India  
—*S. D. Jadhav and M.S. Jadhav*
- S492–S498 Environmental impact of paper consumption in the educational sector and digital learning - examination system post Covid-19  
—*Ashok Verghese, A. K. Parvathy, S. Sathyalakshmi, Beena James and M. Prabakaran*
- S499–S505 Eco-friendly approach toward preparation of plantable paper like material from organic sugarcane bagasse waste  
—*Minakshi Bhattacharjee, Susanta Mog and Nabin Soren*
- S506–S511 Study of water management and scarcity of drinking water in Alwar City (Rajasthan)  
—*R.N. Yadav and Ajit Singh*
- S512–S519 Distribution of Avian Diversity in Different Habitats from Gangetic Plains of West Bengal, India  
—*Kausik Ghosh*
- S520–S527 Hydrothermal Carbonization of Paper Mill ETP Sludge and Assessing its Potential Application in Agriculture  
—*K. Blessy Monica, E. Parameswari, V. Davamani, K. Sivakumar and P. Kalaiselvi*
- S528–S533 Contribution to the Study of the Soil's Diversity of the Algerian North-east, Case of the Transect «Souk-ahras -tebessa»  
—*Samai Ibtissem, Amri Naziha, Chouba Ibtissem, Zentar Amina and Remita Ferial*
- S534–S538 A Study on the Economic Impact of Covid 19 among Different Sections of People  
—*Kanchan Devi*
- S539–S543 Biochemical and molecular identification of *Listeria monocytogenes* bacteria isolated from Raw vegetables in Madhya Pradesh, India  
—*Shivendra Verma, Abhishek Gupta and Shalini Singh*
- S544–S547 Effect of plant extracts against stripe rust of barley caused by *Puccinia striiformis* f. sp. *hordei*  
—*Harshraj Kanwar, Pradeep Singh Shekhawat and Brajnandan Singh Chandrawat*

# Differences effect of red and big white ginger extract as anti-inflammatory agents by *In vitro*

E.B. Aksono<sup>1,7\*</sup>, T.D.K. Wungu<sup>2</sup>, C.H. Wijaya<sup>3</sup>, R.R.F. Sarassina<sup>4</sup>, N. Hidayatik<sup>1</sup>,  
H. Pertiwi<sup>5</sup> and N. Puspitasari<sup>6</sup>

<sup>1</sup>Department of Basic Veterinary Science Faculty of Veterinary Medicine, Universitas Airlangga Indonesia

<sup>2</sup>Department of Physics, Nuclear Physics and Biophysics Research Group Faculty of Mathematics and Natural Sciences, Institute Teknologi Bandung Indonesia

<sup>3</sup>Department of Food Science and Technology Faculty of Agriculture Engineering, Institute Pertanian Bogor Indonesia

<sup>4</sup>Department of Economics and Business Vocational School, Universitas Gadjah Mada Indonesia

<sup>5</sup>Department of Health Studies Faculty of Vocational Studies, Universitas Airlangga Indonesia

<sup>6</sup>Department of Statistic and Population Faculty of Public Health, Universitas Airlangga Indonesia

<sup>7</sup>Natural Sciences and Engineering Institute, Universitas Airlangga Indonesia

(Received 7 July, 2021; Accepted 30 July, 2021)

## ABSTRACT

This research was conducted to evaluate the possibility of different anti-inflammatory effects from the extract of red ginger and big white ginger by measuring their protein (bovine serum albumin) denaturation inhibitory capability. This study is a quasi experimental, using a model reaction between 0.2 % BSA were heated to form protein denaturation by five groups: red ginger water extract group, red ginger ethanol extract group, big white ginger water extract group, and big white ginger ethanol extract group as the test group (200 ppm, 100 ppm, 10 ppm, and 1 ppm) and diclofenac sodium as the standard group (40 ppm, 20 ppm, 10 ppm, 5 ppm, 2.5 ppm, and 1.3 ppm). Potential inhibition of protein denaturation is known to determine the value of IC<sub>50</sub>. The results of this study indicate that the red ginger water extract has the IC<sub>50</sub> value of 107.4513 ppm (r=0.967), the red ginger ethanol extract has the IC<sub>50</sub> value of 193.4338 ppm (r=0.959), the big white ginger water extract has the IC<sub>50</sub> value 160 ppm (r=0.969), the big white ginger ethanol extract has the IC<sub>50</sub> value 160.9536 ppm (r=0.994), of whereas for diclofenac sodium by 27.1133 ppm (r=0.874). These results indicate that the red ginger dan big white ginger as a potential inhibitor of protein denaturation for inflammatory processes *in vitro*.

**Key words:** Anti-inflammatory, Ginger, Bovine serum albumin, Denaturation, Protein

## Introduction

Indonesia is a country rich in natural resources including resources for traditional medicine as out of 30,000 plant species from a total of 40,000 plant species in the world, 940 of them are medicinal plants

(Rahayu and Andini, 2019). Red ginger and big white ginger of Zingiberaceae are widely known as medicinal plants in Indonesia. In Asia, varieties of ginger are spread across wet tropical regions. Ginger is abundant in North Sumatera, Bengkulu, West Java, Central Java and East Java. Ginger is divided

Corresponding author's email: eduardus-b-a-h@fkh.unair.ac.id



broadly into three kinds, red ginger, big white ginger and small white ginger. The difference between the three is based on their essential oil quantity. The red ginger contains the most essential oil and small white ginger contains the least essential oil.

Red ginger (*Zingiber officinale var. rubrum*) has 42-43 mm diameter, 52-104 mm in height and 123-126 mm in length while the big white ginger (*Zingiber Officinale var. officinarum*) has 48-85 mm, 62-113 mm in height and 158-327 mm in length. The red ginger is very spicy with a strong aroma which contains 2.6-3.9% essential oil. Big white ginger on the other hand has less spicy taste and less strong compared to the red ginger and small white ginger. It also contains 0.82-2.8 % essential oil so both of them are used as medicinal plants. Fresh ginger is categorized into two kinds, volatile and non-volatile. The volatile ginger contains sesquiterpene hydrocarbons and monoterpenoids which give a distinctive aroma and taste. In contrast, the compounds which are not easily evaporated include gingerol, shogaol, paradol, and zingerone and they have anti-inflammatory, anti-oxidant, anti-bacterial and anti-thrombocytopenic properties (Jolad *et al.*, 2004; Williams and Lamprecht, 2008; Mashadi *et al.*, 2013; Chandra *et al.*, 2012).

Inflammation is a body response from an injury or infection marked by fever, reddened skin, pain, swell and physiological function disorder. It is a normal protection response against tissue injury caused by physical trauma, chemical substance or dangerous microbes (Perumal *et al.*, 2008). In inflammation, denatured protein has a role as autoantigen. It is commonly found in some illnesses such as rheumatism. Various therapies and treatments have been developed to treat the inflammation process, such as NSAIDs (Non Steroidal Anti Inflammatory Drugs) with diclofenac drug group. Diclofenac is an anti-inflammatory drug which works by inhibiting COX, the COX-2 inhibitor (Perumal *et al.*, 2008; Williams *et al.*, 2008). Other than NSAIDs, medicinal plants can also be utilized. The natural materials are expected to contain a certain compound with significant anti-inflammatory activity. The advantages from the natural material (herbal medicine) are for its accessibility, inexpensiveness and lesser side-effects. One of the natural resources with bioactive compounds and potential for development is red or big white ginger (Perumal *et al.*, 2008).

This research is conducted to evaluate the possibility of different effects between extract of red gin-

ger and big white gingers to protein denaturation in their function as anti-inflammatory agents.

## Materials and Methods

### Media Preparation

The samples used in this study were red ginger (*Zingiber officinale var. Rubrum*) and big white ginger (*Zingiber officinale var. Officinarum*) from traditional markets in Surabaya-East Java. To create tris buffered saline solution or TBS, mix 1.21 g tris base and 8.7 g sodium chloride (NaCl) and dissolve into sterile aquadest until it weighs 900 ml. Adjust the solution pH level to 6.2-6.5 with glacial acetic acid and then add more sterile aquadest until the solution volume at 1000 ml (Erianti *et al.*, 2015). 0.2% Bovine Serum Albumin (BSA) is made by mixing 0.2 gram BSA crystal with TBS solution until the volume reaches 100 ml and then vortex the solution to completely dissolve the BSA crystal (Aditya *et al.*, 2015).

### Treatment group

The negative control used for this research is 1 ml 0.2% BSA solution in a microtube and labeled. The positive control used for this research is diclofenac sodium solution made by mixing 4 mg diclofenac sodium with TBS solution until the volume reaches 1 ml and dissolving it with a vortex for several minutes. The process is done with a microtube to produce a stock solution of positive control with 4,000 ppm. From the stock solution, some other solutions are created with different concentrations; 2,000 ppm, 1,000 ppm, 500 ppm, 250 ppm and 130 ppm. Each solution is made in microtubes (Aditya *et al.*, 2015). The testing solution in this research is made by mixing 20 mg dry extract with TBS solution until the volume is at 1 ml in the microtube then dissolving it with the vortex machine. From this procedure, the stock testing solution with 20,000 ppm is acquired. From this stock solution, some serial solutions are made with different concentrations; 10,000 ppm, 1,000 ppm, and 100 ppm (Erianti *et al.*, 2015).

### Protein Anti Denaturation Effect Analysis

After all series of treatment groups and positive control solutions are prepared, the samples from all concentrated solutions are taken about 10 µl each. Mix each sample with 0.2% BSA solution until the volume reaches 1 ml and then shake it slowly to make sure that it dissolves well. From this procedure, se-

ries of concentrated solutions are made for testing solutions (200 ppm, 100 ppm, 10 ppm, and 1 ppm) and for the diclofenac sodium (positive control) solutions (40 ppm, 20 ppm, 10 ppm, 5 ppm, 2.5 ppm, and 1.3 ppm). With the negative control prepared beforehand, incubate all the samples of solutions mixed with BSA for 30 minutes at 25 °C temperature. After the incubation procedure, heat all the samples of testing solutions, positive control solutions and negative control using the heatblock for 5 minutes at 72 °C temperature. After the heating procedure, incubate them again at 23 °C for 25 minutes. After the second incubation, let them cool down and then use the vortex machine to make the mixture in the microtubes homogenous. After that, they are analyzed with a microplate spectrophotometer. Move all the samples of solutions to well microplates. Each well is filled with 250 µl solution from the series of testing solutions, positive control solutions and negative control solutions. There are three readings conducted (1 microtube = 3 wells). The readings by this spectrophotometer are conducted with 655 nanometer wavelengths (Erianti *et al*, 2015). The percentage of protein denaturation inhibition is calculated with a formula as follows:

$$\text{Inhibition (\%)} = \frac{\text{Negative Control Absorbance} - \text{Sample Absorbance}}{\text{Negative Control Absorbance}} \times 100\%$$

In vitro inflammatory activity is expressed by determining the amount of IC50 (inhibition Concentration 50) by making a linear graph  $y=a+bx$  with  $y$ =absorbance and  $x$ =concentration of red ginger (*Zingiber officinale var. Rubrum*) and big white ginger (*Zingiber officinale var. Officinarum*) (Erianti *et al*, 2015). A compound which is able to inhibit protein denaturation more than 20%, it is predicted to have strong anti-denaturation effect and the compound can be developed as an alternative anti-inflammatory drug. (Aditya *et al.*, 2015)

## Results

### Anti-Inflammatory Effect of Red Ginger Water Extract

The effect of red ginger water extract as anti inflammatory agent can be seen from the Figure 1 and 5. At 1-200 ppm concentration, the extract has shown 23.7%-65.48% inhibition percentage. It is higher than the diclofenac sodium as positive control which only showed 16.33-57.34% from 1-40 ppm concentration.




Fig. 1. Differences percentage inhibition of bovine serum albumin in vitro due to diclofenac sodium (positive control) and red ginger water extract

From the calculation of IC50 red ginger water extract and diclofenac sodium are 107.4513 ppm and 27.1133 ppm.

### Effect of Red Ginger Ethanol Extract

The effect of red ginger ethanol extract as anti-inflammatory agent can be seen from the Figure 2 and 5. At 1-200 ppm concentration, the extract has shown 21.31%-52.40% inhibition percentage while the diclofenac sodium as positive control only showed 16.33-57.34% from 1-40 ppm concentration.




Fig. 2. Differences percentage inhibition of bovine serum albumin in vitro due to diclofenac sodium (positive control) and red ginger ethanol extract

From the IC50 calculation, red ginger ethanol extract and diclofenac sodium are 19.4338 ppm and 27.1133 ppm.

### Anti-Inflammatory Effect of Big White Ginger Water Extract

The effect of big white ginger water extract as anti




Fig. 3. Differences percentage inhibition of bovine serum albumin in vitro due to diclofenac sodium (positive control) and big white ginger water extract

inflammatory agent can be seen from the Figure 3 and 5. At 1-200 ppm concentration, the extract has shown 19.58%-57.78% inhibition percentage while the diclofenac sodium as positive control only showed 16.33-57.34% from 1-40 ppm concentration. From the IC50 calculation, big white ginger water extract and diclofenac sodium are 160 ppm and 27.1133 ppm.

**Anti-Inflammatory Effect of Big White Ginger Ethanol Extract**

The effect of big white ginger ethanol extract as anti inflammatory agent can be seen from the figure 4 and 5. At 1-200 ppm concentration, the extract has shown 18.58%-58.89% inhibition percentage while the diclofenac sodium as positive control only showed 16.33-57.34% from 1-40 ppm concentration. From the IC50 calculation, big white ginger ethanol extract and diclofenac sodium are 160.9536 ppm and 27.1133 ppm

**Discussion**

To produce herbal medicine, preclinical tests to ani-

mals are needed. However, there are some problems on using animals for testing, such as ethical problem, the lack of reasons using it when there are still other methods, and the cross-species extrapolation is difficult to perform due to the lack of isomorphisms between the animals and humans, especially at cellular and molecular level where the illness started (Erianti *et al.*, 2015). Previous researches have shown that some natural products are able to stabilize bovine serum albumin at pathological pH. This finding was then used to detect anti-inflammatory activity by *in vitro* method. Williams *et al.* (2008) have studied the properties of some natural resources to stabilize bovine serum albumin at pathological pH (pH level 6.5-6.8). This finding was then used to detect anti-inflammatory activity by *in vitro* method.

Higher absorbance found from the samples compared to the control solution showed the protein denaturation inhibition which was induced by heat (albumin) of ginger extract and positive control of diclofenac sodium. In this study red ginger and big white ginger extracts are more active than diclofenac sodium, which is an active lower concentration. This




Fig. 4. Differences percentage inhibition of bovine serum albumin in vitro due to diclofenac sodium (positive control) and big white ginger ethanol extract




Fig. 5. Differences percentage inhibition of bovine serum albumin in vitro due to diclofenac sodium (positive control), red ginger water extract, red ginger ethanol extract, big white ginger water extract, and big white ginger ethanol extract

result showed that there was a change in turbidity due to viscosity change. This result was supported by (Chandra *et al.*, 2012) where the viscosity changes because protein solution viscosity is increasing at denaturation.

Protein denaturation is the protein's secondary, tertiary and quaternary structure change without the breaking of covalent bonds and is one of the causes of inflammation. The factors which cause protein denaturation are stress, temperature, pH, agitation, electricity, chemical substance, alcohol and reducing agents. Heat is one of the causes of protein denaturation because heat can disrupt the hydrogen bond so its capability of bonding the water decreased (Aditya *et al.*, 2015). It happens because high temperature increases the kinetic energy and causes the molecules to vibrate so rapidly and violently that the bonds are disrupted. Protein denaturation occurs steadily and unchanged. Denatured protein loses its solubility so it is easy to settle (Nijveldt *et al.*, 2001). When the protein denatures, some of its biological function might be lost. In addition, Autoantigen produced by certain illnesses related to inflammation may be caused by denaturation of protein. Protein denaturation is the loss of a structure and protein function caused by stress, chemical substance or heat. There is also loss of biological function when the protein denatures (Aditya *et al.*, 2015).

Red ginger and big white ginger extracts are capable of anti-inflammatory activity with protein denaturation inhibition percentage higher than 20%. Besides, as seen from the potential IC<sub>50</sub>, positive r values indicate the existence of a positive relation between concentration and inhibiting protein denaturation potential. A compound with anti-inflammatory is flavonoid. The anti-inflammatory activity mechanism can be done through some ways, Cyclooxygenase (COX) and lipoxygenase (LOX) enzyme inhibition, leukocyte accumulation inhibition, neutrophil degranulation inhibition, histamine release inhibition (Riansyah *et al.*, 2015). Anti-inflammatory activity from the flavonoid with Cyclooxygenase (COX) and lipoxygenase (LOX) enzyme inhibition can cause leukotriene and prostaglandin synthesis inhibition which then inhibit mucus secretion which protect the stomach wall. Leukocyte accumulation inhibition during the inflammation process will lower the body response against inflammation. This inhibition happens because Cyclooxygenase (COX) is inhibited so thromboxane

causing the leukocyte modulation is inhibited. Histamine release inhibition happens because flavonoid can inhibit mast cell release (Hidayati *et al.*, 2008).

Beside flavonoid compound, tannin compound also reported to have anti-inflammatory property (Verma *et al.*, 2011; Erianti *et al.*, 2015) and saponin compound has the potential to be anti-inflammatory agent by protein denaturation inhibition (Erianti *et al.*, 2015; Ahmad, 2009). Proteins of a vulnerable body can experience denaturation caused by free radical formation which causes an inflammation mechanism through inflammatory mediator release. Denaturation of protein is a process where protein loses its secondary or tertiary structure due to external compounds such as strong acid or base, inorganic salt, organic solvent and heat (Verma *et al.*, 2011). The possibility of interaction or bond between molecules in red ginger and big white ginger extracts can inhibit denaturation of protein. At high temperature, protein in BSA solution tends to form big molecules so it cannot go through the membrane and this phenomenon is known as denaturation of protein. If the BSA solution is heated, it will turn murky as the solubility of protein is decreased and protein molecules form aggregate so the molecule protein becomes bigger from the previous size (Ahmad, 2009).

One of the potentials of some non-steroid anti-inflammatory drugs is their ability to stabilize (prevent denaturation) albumin which has been given heat treatment at physiological pH (pH: 6.2-6.5) (Erianti *et al.*, 2015; Williams *et al.*, 2008). Therefore, with this initial research, we can conclude that ginger has an anti-inflammatory effect in vitro marked by protein denaturation. Further research is needed to ensure the mechanism and constituent of its anti-inflammatory function

## Conclusion

It could be concluded that that red ginger extract has higher anti-inflammatory activity compared to the big white ginger in protein denaturation inhibition by in vitro with the highest inhibition percentage from the red ginger water extract which is at 65.48% and also IC<sub>50</sub> is 107.4513 ppm.

## Conflict of Interest

The authors declares that there are no conflicts of interest regarding the publication of this article

## Acknowledgement

This work was supported by grant from Indonesia Collaborative Research Program Kemenristekdikti and Universitas Airlangga 2018

## References

- Aditya, M. R. T., Marisa, D. and Suhartono, 2015. Anti-inflammatory potential of mangosteen fruit juice (*Garcinia mangostana*) against in vitro protein denaturation. *Jurnal Berkala Kedokteran*. 11 : 149-156. doi: 10.20527/jbk.v11i2.138.
- Ahmad, S. 2009. Membrane Ultra filtration performance test with bovine serum albumin as bait solution. Bandung. *Jurnal Sains Materi Indonesia*. 11 : 6-9. doi: 10.17146/jsmi.2009.11.1.4541.
- Chandra, S., Chatterjee, P., Dey, P., Battacharya, S. 2012. Evaluation of *in vitro* anti inflammatory activity of coffee againts the denaturation of protein. *Asia Pasific Journal of Tropical Biomedicine*. S178-S180. doi: 10.1016/S2221-1691(12)60154-3.
- Erianti, F., Marisa, D., Eko Suhartono. 2015. Potential anti-inflammatory juice of starfruit (*Averrhoa Carambola* L.) towards the denaturation of protein *in vitro*. *Jurnal Berkala Kedokteran*. 11, 33-39. doi: 10.20527/jbk.v11i1.183.
- Hidayati, N.A., Listyawati, S., Setyawan, A.D. 2008. Chemical content and anti-inflammatory test from the ethanol extract of *Lantana camara* L. On Male Albino Rats (*Rattus novergicus* L.). *Bioteknologi*. 5(1): 10-14.
- Jolad, S.D., Lantz R.C., Solyom, A.M., Chen, G.J., Bates, R.B. and Timmermann, B.N. 2004. Fresh organically grown ginger (*Zingiber officinale*): Composition and effects on LPS-induced PGE2 production. *Phytochemistry*. 65 : 1937-1954. doi: 10.1016/j.phytochem.2004. 06.008.
- Mashadi, N.S., Ghiasvand, R., Askari, G., Hariri, M., Darvishi, L. and Mofid, M.R. 2013. Anti-oxidative and anti-inflammatory effects of ginger in health and physical activity: Review of current evidence. *Int. J. Prev. Med*. 4 : S36-S42. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3665023/>.
- Nijveldt, R.J., Nood, E.V., Hoorn, D. EC.V., Boelens, P.G., Norren, K.V. and Leeuwen, P.A.M.V. 2001. Flavanoid a review of probables mechanisms of action and potential applications. *American Journal of Clinical and Nutrition*. 74. 418-425. doi: 10.1093/ajcn/74.4.418.
- Perumal, R., Dey, A. and Manavalan, R. 2008. Inhibition of albumin denaturation and anti inflammatory activity of furfuryl substituted pyrimidino imidazolinones. *Int. J. Chem. Sci*. 6(4) : 2016-2022
- Rahayu, S. M. and Andini, A. S. 2019. Ethnobotanical Study on Medicinal Plants in Sesaot Forest, Narmada, West Lombok, Indonesia. *Biosaintifika: Journal of Biology & Biology Education*. 11(2) : 234-242.
- Riansyah, Y., Mulqie, L. and Choerina, R. 2015. The anti-inflammatory activity test of leaf ethanol extract purple sweet potato (*Ipomoea batatas* L. Lamk) on male Wistar rats. *Research Proceeding from SPeSIA*, Unisba.
- Sembiring, S. and Sismudjito, 2015. The knowledge and utilization of treatment methods in Sukanalu village, Barus Jahe subdistrict. *Jurnal Perspektif Sosiologi*. 3 : 104-117.
- Verma, A.M. Kumar, A.P., Kavitha. D. and Anurag, K.B. 2011. Anti denaturation and antioxidant activities of *Annona cherimola in-vitro*. *International Journal of Pharma and Bio Sciences*. 2(2).
- Williams, C.A. and Lamprecht, ED. 2008. Some commonly fed herbs and other functional foods in equine nutrition : A Review. *The Veterinary Journal*. 178(1) : 21-31. doi: 10.1016/j.tvjl.2007.06.004.
- Williams, L.A.D., Connar, A.O., Latore, L., Denis, O., Ringer, S., Whittaker, J.A., Conrad, J., Vogler, B., Rosner, H. and Kraus, W. 2008. The *in vitro* anti-denaturation effects induced by natural product and non-steroidal compounds in heat treated (immunogenic) bovine serum albumin is proposed as a screening assay for the detection of anti-inflammatory compounds, without the use of animals, in the early stages of the drug discovery process. *West Indian Medical Journal*. 57 (4) : 327-331. doi: 10.1215/9780822388630-010.