


ECOLOGY

ENVIRONMENT & CONSERVATION

ISSN 0971-765 X

Vol. 21
No. (2)
2015



EM INTERNATIONAL

QUARTERLY INTERNATIONAL JOURNAL

20

ECOLOGY

ENVIRONMENT & CONSERVATION

ISSN 0971-765 X

Vol. 21
No. (2)
2015



EM INTERNATIONAL

198

Quarterly International Journal Ecology, Environment and Conservation

ISSN 0972-3005

Editor: Dr. R.K.Trivedy, Pune, India

EDITORIAL ADVISORY BOARD

1. Dr. Teresa Ferreira, Portugal
2. Dr. Rakesh Kumar, AICTE, New Delhi, India
3. Mr. Sandeep Singhal, Chennai, India
4. Dr. M.M. Amin, Rajshai, Bangladesh
5. Dr. D.J. Lee, Taiwan
6. Dr. J.K. Dutta, Burdwan, India
7. Dr. K. Venkateshwarlu, Anantpur, India
8. Dr. Linda Blackwell, Australia
9. Dr. G. Zellner, Netherlands
10. Dr. I.C. Onyema, Nigeria
11. Dr. M.F. Hamoda, Kuwait
12. Dr. H.A. Abrahamse, South Africa
13. Dr. Rowshan Mumtaz, Bangladesh
14. Dr. Hassan Poorbabaei, Iran
15. Dr. C.K. Yap, Malaysia
16. Dr. Hassan Ibrahim Ali, Sudan
17. Dr. M.T. Udo, Nigeria
18. Dr. Mohd. Nural Anwar, Bangladesh
19. Dr. A.K. Kumaraguru, Madurai, India
20. Dr. A.U. Pagarkar, Ratnagiri, India
21. Dr. Ganesh Iyer, Mumbai, India
22. Dr. A.R. Ghosh, Burdwan, India
23. Prof. M. Zaman, Bangladesh
24. Dr. Marcantonio Bragdin, Venice, Italy
25. Dr. Ramesh Thatikunta, Hyderabad, India
26. Dr. Z. Li. Bonn, Germany
27. Dr. Anju Singh, Mumbai, India
28. Dr. Ing. Agr. Mario Ridardo Sabbatini, Argentina
29. Dr. Philip C. Reid, U.K.
30. Dr. Amarjeet Kaur, New Delhi, India
31. Dr. B.N. Prasad, Noida, India
32. Dr. Oswaldo A. Fernandez, Argentina
33. Dr. A. Olawale, Nigeria
34. Dr. Ms. Mirela Tulik, Warsaw, Poland
35. Dr. Piroz Azizi, Iran
36. Dr. Ashwini Wangenao, Bhopal, India
37. Dr. G. R. Pathade, Pune, India
38. Dr. P.S. Panesar, Longowal, India
39. Dr. Meenakshi C. Mahajan, Pune, India
40. Dr. Mohd. Yusuf, Malaysia
41. Dr. A.K. Panigrahi, Berhampur, India
42. Ms. Aditi Trivedi, Pune
43. Dr. Azni Idris, UPM, Malaysia
44. Dr. T. Bahorun, Mauritius
45. Dr. B.K. Sharma, Shilong, India
46. Dr. E. Amiri, Iran
47. Dr. T.A. Yakeen, Nigeria
48. Dr. L.L. Chukwu, Nigeria
49. Dr. Michael Ukwuru, Nigeria
50. Dr. Sabrina Naz., Bangladesh

Ecology, Environment and Conservation is Abstracted in -

- > Chemical Abstracts, U.S.A.
- > Ecological Abstracts, U.S.A.
- > Cambridge Science Abstracts, U.S.A.
- > Ecodisc-CD-Rom, Netherlands
- > Geological Abstract, Netherlands
- > Pollution Abstract, U.S.A.
- > International Dev. Abstracts, U.S.A.
- > Paryavaram Abstracts, India
- > Current Awareness in Biological Sciences, Elsevier
- > World Textiles Abstracts, Netherlands
- > Fluid Abstracts, Netherlands
- > Environmental Abstracts, India
- > Indian Science Abstracts, India
- > Cambridge Science Abstracts

Impact Rating 3.5 (NAAS 2011)

COPYRIGHT: This journal is registered at the COPYRIGHT CLEARANCE CENTRE INC. (CCC), 222, Rosewood Drive, Suite 910, Danver, MA 01923, U.S.A. The copyright owner consents that in the U.S.A. copies of the articles may be made for personal or internal use, of specific clients, on payment of fee. All requests for permission to photocopy should be addressed to the copyright owner (www.copyright.com)

The journal figures in Ulrich International Periodicals Directory, U.K., Gale Directory, U.K. & SAARC Directory of periodicals.

SUBSCRIPTION RATES

	INDIA	OVERSEAS
Individuals	Rs : 1000.00	US\$: 150.00
Institutional	Rs : 2000.00	US\$: 600.00

Note : All payments in favour of EM INTERNATIONAL (Postage included in Subscription)

Publishers : EM INTERNATIONAL, Prakriti, Balewadi, Baner 411 045, M.S., India

☎ (91-2164- 220369; 20- 46745119; E-mail : rktem@pn3.vsnl.net.in)

ECOLOGY, ENVIRONMENT AND CONSERVATION

VOL. 21 (2) : 2015

CONTENTS

- 591-602 Driving mechanism of environmental variables for biomass accumulation in Tibetan large-scale grassland
—*Hong Sheng Liu, Chad Larson and Zewei Miao*
- 603-613 Composting leachate treatment by an anaerobic migrating blanket reactor
—*Asghar Ebrahimi, Hassan Hashemi, Mohammadmehdi Amin, Bijan Bina, Afshin Ebrahimi, Hamidreza Pourzamani and Yaghoob Hajizadeh*
- 615-626 Biological systems as tools to mitigate climatic change in the tropics
—*Muhamad Hanif Md Nor, Chin Hong Neoh, Nurul Farhana Ahmad Rapani, Noor Aina Baijan, Norahim Ibrahim, Adibah Yahya, Shafinaz Shahir, Madihah Salleh, Chun Siong Chong, Wan Rosmiza Zana Wan Dagang, Mohd Firdaus Abdul Wahab, Huszalina Hussin, Haryati Jamaluddin, Fazilah Abdul Manan, Shaza Eva Mohamad, Zaharah Ibrahim*
- 627-632 Evaluation of A Hydrus-3d model for phosphorus transport in soil column
—*Yaser Hoseini and Alireza Pilpayeh*
- 633-639 An investigation of genetic diversity in endangered populations of birch in Iran based upon sequencing of chloroplast matK gene
—*Mohammad Esmaeilpour, Kambiz Taheri Abkenar, Ali Aalami, Amireslam Bonyad and Mostafa Ghaderi-Zefrehei*
- 641-644 Predictors of applying soil conservation practices by farmers at Kurdistan Township, Iran
—*Shabnam Feizi Khazaei*
- 545-651 The study of modifications in land usage in the City of Shandiz towards ecological city realization
—*Mohamad Rahim. Rahnma and Foruzan. Taheri*
- 653-656 The impact of globalization on social and economic inequalities between Urban and rural communities in less developed countries : Case study: Shirvan County (North Khorasan Province), Iran
—*Yahya. Jebraili and Mahdi. Jahani*
- 657-660 The project of Nomadic Tourism hamlet establishing as a suitable strategy for Tribal culture and traces preservation (Case Study: Northern Khorassan)
—*Reza. Rowshanzamir, Seyed Rahim and Moshiri, Ali. Dadar*
- 661-664 The role of villagers' participation and organizations' and administrations' cooperation in the performance of local management organizations (case study: rural mayors of Koohsorkh district of Kashmar)
—*Reza Saberi Tavalaei and Hamid Jafari*
- 665-668 Empowerment: an approach to improve the performance of rural managers (Case study: rural mayors of Kouhsorkh district of Kashmar)
—*Reza Saberi Tavalaei and Hamid Jafari*
- 669-671 Effective factors on rural emigration in golmakan rural District, Chenaran County
—*Mahdi. Jahani and Abolfazl. Vatandoust*
- 673-677 An assessment of socio-economic conditions in citrus genus using GIS techniques in Ramsar, Iran
—*A. Ahmad, H. Zabihi, M. Golmohammadi and B. Golein*
- 679-684 Investigation of the effective factors in destruction of water exploitation wells in Salmas plain
—*Esfandiyar Abbas Novinpoor and Ehsan Alizadeh*

- 685-689 Conservation impact on molecular genetic changes on Java green peacock (*Pavo muticus*) through mitochondrial D-loop marker
—E. Bimo Aksono H. and Herry Agoes Hermadi
- 691-698 The relationships among carbon sequestration, erosion and some soil properties in different land-use types
—Sayed Fakhreddin Afzali and Maryam Soleymani
- 699-706 Evaluation of climate fluctuations using meteorological drought index and its vulnerability risk mapping (Case study : South of Fars Province, Iran)
—Mohammad Reza Honar and Karim Hejazi Jahromi
- 707-714 Reliability survey of metropolitan areas of Iran for for converting them into creative Cities
—Simin Abafat, Mahdi Abafat and Meysam Abafat
- 715-721 Investigation of effect of roughness on riprap stability around the bridge pier in 180 degree river
—Alireza Masjedi, Amir Taeedi, Bita Jafari and Mohsen Mojtahed
- 723-730 A study on effects of ship ballast water discharges on the zooplanktons in calabar coastal waters
—S.U. Jatau, M.C. Emere, M.S. Abdulsalam, Y. Umar and N.A. Etangetuk
- 731-737 Nematicidal activities of *Quercus brantii* L., *Myrtus communis* L. and *Citrullus colocynthis* (L.) SCHRAD
—Abbas Salahi Ardakani
- 739-746 Experimental study on scour hole around submerged vanes at a 180 degree flume bend
—Alireza Masjedi, Hamid Abyar, Seyed Vahab Belyadian and Bita Jafari
- 747-750 Disinfection of drinking water using UV-LED
—P. Niroopa and S. Kanmani
- 751-757 Effects of drought on the agriculture sector from the perspective of agricultural extension Experts in Kermanshah Township
—Afsaneh Salmani, Roya Eshraghi and Alireza Poursaeed
- 759-761 Assessment of losses due to pulse beetle *Callosobruchus chinensis* L. in chickpea
—V. Venkatesham, R.S. Meena and M.A. Laichattiwar
- 763-767 Genetic diversity analysis of castor (*Ricinus communis* L.) using microsatellite (SSR) Markers
—M. Sudharani, S. Nagalakshmi, M. Madhavi and N.V. Naidu
- 769-771 Evaluation of newer insecticides against red rust thrips (*Cheatanophothrips signipennis* Bagnell)
—N.M. Patil and N.B. Shaikh
- 773-776 Study of genetic diversity in chickpea (*Cicer arietinum* L.) using microsatellite markers
—M. Sudharani, S. Nagalakshmi, M. Madhavi and N.V. Naidu
- 777-781 Reductive dehalogenation of aldrin by cast Iron
—Yangdup Lama, Alok Sinha and Gurdeep Singh
- 783-788 Effectiveness of biorational insecticides for the management of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee and on yield
—Laishram Laishana Devi, Tushar M. Ghule, M.L Chatterjee and A.K. Senapati
- 789-794 Sustainability of basmati-wheat sequence under intensive management of nutrients and its effect on yield and uptake of Zn, Cu, Fe and Mn
—J.K. Dhaliwal, G.S. Saroa and S.S. Dhaliwal

- 795–800 **Effect of letrozole loaded PLGA nanoparticles on sex reversal of *Cortes swordtail Xiphophorus cortezi* (Cyprinodontiformes, Poeciliidae)**
—Harshavardhan D. Joshi, Milind B. Katare, J.K. Pathan, Tanveer Hussain, M.P. Bhendarkar and Karan Kumar Ramteke
- 801–805 **Seasonal variations in chlorophyll-*a* concentration in coastal waters off veraval, Gujarat coast**
—Gunjan Motwani, Rahul Rajan, Mini Raman and Hitesh Soalnki
- 807–811 **Uv irradiation effects on seed quality and storability of chickpea genotypes**
—S.N. Vasudevan, Rajarajeshwari, Vijayakumar, A. Amaregouda and S.R. Doddagoudar
- 813–817 **Quality characterization of market available Indian butter catfish (*Ompok bimaculatus*) in Tripura, India**
—Jag Pal and A.U. Muzaddadi
- 819–821 **Integrated nutrient management studies in Pumpkin**
—A. N. Alekar, A.P. Wagh, R. P. Kharde and P. S. Hirave
- 823–826 **Effect of plant growth regulators on rooting in cuttings of fig (*Ficus carica*) cv. Dinkar**
—L.S. Khapare, M.H. Dahale, L.S. Deshmukh and R.B. Bhusari
- 827–829 **Bacteriological investigation of the air in a private college**
—Ritika Charaya and Kavita Naruka
- 831–837 **Variation in pulp content and physical characters of *Elaeocarpus serratus* L. fruits in different landuse patterns of Western Ghats, India**
—Vikas Kumar, Ajeesh R., Vidyasagaran K. and Sreejith Babu
- 839–849 **Eco-labeling for environmental friendly production in agriculture sector: An International and Indian perspective**
—A.V. Manjunatha, Y. Nagaraju and Sudarshan Sampath
- 851–860 **Interbreeding status of *Helicoverpa armigera* hubner (Lepidoptera: Noctuidae) collected from different hosts and agroclimatic zones**
—Arshdeep K. Gill and Ramesh Arora
- 861–863 **Effect of planting dates on growth and yield of Garlic (*Allium sativum*)**
—Swati Barche and Kamal S. Kirad
- 865–872 **A review of mulching on vegetable crops production**
—Swati Barche, Reena Nair and P.K. Jain
- 873–876 **Performance of gladiolus (*Gladiolus grandiflorus* Hort.) genotypes for flowering, flower quality and corm yield under hill zone of Karnataka**
—H.P. Sudeep, B.S. Reddy, S. Latha, B. Divya and Kallappa Narode
- 877–879 **Breeding pole type french bean (*Phaseolus vulgaris* L.) for rust disease (*Uromyces phaseoli* Reben wint) resistance using SCAR markers**
—B. Divya, T. S. Aghora, N. Mohan, C. Aswath and H.P. Sudeep
- 881–883 ***In vitro* effects of plant extracts, on Mycelial growth or percent inhibition of blossom blight fungi**
—S. V. Kolase, Tejesh Kamble and Sunil Jogdand
- 885–890 **Growth regulators influencing yield and quality of pineapple [*Ananas comosus* (L) Merr]**
—R. Pal, S. K. Mahato, Binoy Chhetri and C.P. Suresh
- 891–896 **Protected cultivation technology (PCT) in Karnataka – an economic analysis of Gerbera**
—P. B. Gamanagatti and B.L. Patil
- 897–899 **Bioefficacy of fungicides for management of blossom blight of Mango (*Mangifera indica*)**
—S.V. Kolase, S.M. Jogdand, Tejas Kambale and S.S. Kulkarni

- 901-904 **Agroforestry systems for fodder production under rainfed conditions**
—N. Kaushik, R.P.S. Deswal and Suman Malik
- 905-909 **Growth performance of *Pangasius sutchi* in HDPE lined pond during drought like situation in Bihar**
—Pankaj Patial, A.P. Mishra and M. A. Hassan
- 911-914 **Fluoride in groundwater of satlasana Taluka, Mehsana District, Gujrat, India**
—Parikh Punita and Vyas Payal
- 915-919 **Character association and path analysis for yield contributing traits in quality protein maize (*Zea mays* L.)**
—V. Ulaganathan, S.M. Ibrahim and A. Gurusamy
- 921-923 ***Per se* performance of gladiolus (*Gladiolus* spp. L) varieties for yield and quality under Shevaroy's conditions**
—T. Vetrivel, V. Lakshmanan and M. Jawaharlal
- 925-928 **Effect of different sowing dates on growth and yield of wheat (*Triticum aestivum*) varieties under humid sub tropical zone of Uttarakhand, India**
—Jai Prakash Meena and R.S. Verma
- 929-932 **Establishment of correlation factor between total dissolved solids and electrical conductivity for groundwater quality analysis**
—Mayur C. Shah, Sarita K. Parmar and Prateek G. Shilpkar
- 933-938 **Efficacy of sulfonylurea herbicides for broad spectrum weed control in transplanted rice (*Oryza sativa* L.)**
—G. Uma, M. Venkata Ramana, T. Ram Prakash and Sharana basava
- 939-941 **Effect of GA₃ and 6-Benzyl adenine in production of scions for grafting in papaya cv. 9-1(D)**
—S. Senthilkumar, N. Kumar, K. Soorianathasundaram and P. Jeyakumar
- 943-946 **Influence of different postharvest treatments on physico-chemical characteristics of grapefruit at ambient conditions**
—Harmanpreet Kaur, Pushpinder Singh Aulakh, S.K. Jawandha and H. Singh
- 947-953 **Land use characterization of a mid-altitudinal village landscape in Garhwal Himalaya, India**
—J.S. Chandrashekar
- 955-962 **Impact of influential and outlier observations on ordinary least square model for Indian agricultural workers**
—Banti Kumar, Manish Sharma, S. E. H. Rizvi and S.P. Singh
- 963-973 **An interpretative account of voyages to address climate change under united Nations framework convention on climate change: gradual development of policies-prospects and protests**
—Kapil Joshi and Vinay Sharma
- 975-979 **Effect of iron and sulphur on Mustard crop**
—Raman Kumar Pachouri, Anil Kumar Pal, Manoj Pandey and S.K. Trivedi
- 981-984 **Meat recovery, biochemical composition of raw and processed Shrimp (*Litopenaeus vannamei*) cultured in saline waters of Punjab**
—Ajeet Singh, Ravneet Singh, Meera D. Ansal and S.N. Datta
- 985-989 **Investigation on processing quality of commercial Mango varieties and their blends**
—S.P. Salvi, M.M. Kulkarni and P.P. Relekar
- 991-999 **Price spread, marketing efficiency and constraints in Guava (*Psidium guajava* L.) marketing in Uttar Pradesh, India**
—S.H. Malik, Hakim Shabir Ahmad, S.A. Saraf and Wilson Kispotta

- 1001–1006 Wetland soil of kasaragod district of Kerala – Its prospects and aspects for human health
—M. Mini and P.M. Beebi Razeena
- 1007–1012 Changes in physico-chemical composition and sensory qualities of cashew (*Anacardium occidentale* L.) bar during storage
—S. M. Sarkate, K. H. Pujari and P. P. Relekar
- 1013–1019 Economic evaluation of protected cultivation technology for Capsicum
—P. B. Gamanagatti and B. L. Patil
- 1021–1025 Path coefficient and correlation analyses of some yield associated traits in finger millet (*Eleusine coracana* (L.) Gaertn.) germplasm
—V. Ulaganathan and A. Nirmalakumari
- 1027–1032 Status of an extreme dry fish market: A study of dry fish market, on the coastal region of Saurashtra, Gujarat
—R.A. Khileri, S.R. Lende, Vikas, S.A. Muley and G.P. Deshmukh
- 1033–1037 Study of genetic diversity in rice genotypes (*Oryza sativa* L.) using microsatellite (SSR) markers
—M. Sudharani, S. Nagalakshmi, M. Madhavi, P.S. Rao and N.V. Naidu
- 1039–1047 Isolation of phosphate solubilizing bacteria from Rhizospheric soil and its Physiological studies
—Lochan Singh and Sanjay Jha
- 1049–1052 Variation of fluoride concentration in aquifers of different depths: possible geologic reasons
—Anurag Tewari and Ashutosh Dubey
- 1053–1057 Management of hazardous waste generated in dental radiology- A review
—Virender Gombra and Mandeep Kaur
- 1059–1066 An indicator based vulnerability assessment of wilayats to development
—B.S. Choudri, Mahad Baawain, A. Al-Sidairi and H. Al-Nadabi
- 1067–1071 Identification and suggestion of sustainable income generation activities for sthree shakthi group members in rural areas for quality livelihood
—J.M. Saraswathi
- 1073–1076 Strengths, Weaknesses, Opportunities and Threats (SWOT) of sthree shakthi groups of Dharwad and Tumkur Districts
—J.M. Saraswathi and P.R. Sumangala
- 1077–1081 Characteristics and soil fertility appraisal of forest nursery soils of Bilaspur circle in Himachal Pradesh
—Neetan Khajuria, D. Tripathi, Raj Kumar, Vipin Sharma and Shashi Suman Pathania
- 1083–1087 Studies on effect of osmotic dehydration on quality and storage behaviour of ripe jackfruit blubs (*Artocarpus heterophyllus* L.)
—Wanve B.V., K. H. Pujari and P. P. Relekar
- 1089–1094 Assumptions and premises of heating systems hydraulic calculation methods
—Vsevolod Afanas'evich, Seminenko Artem Sergeevich, Alifanova Alla Ivanovna, Elistratova Julija Vasil'evna and Tkach Ljudmila Vladimirovna
- 1095–1100 Evaluation of kriging and co-kriging methods in estimating soil saturated hydraulic conductivity by using soil texture (Case Study: Fath Ali Irrigation and Drainage Network, Ardabil, Iran)
—Yaser Hoseini
- 1101–1106 Therapeutic effect of wood apple on hypertension and diabetes
—S. Anitha, Umadevi S. Hiremath, Savita Hulimani, Geetha Dandin and K. Srinivas

-
- 1107-1111 Variability, heritability genetic advance, correlation and path analysis between yield and yield components in Potato (*Solanum tuberosum* L.)
—*Patram Singh, P. K. Sharma, N.C. Banjara, Naresh P. Sahu and Ruchi Sharma*
- 1113-1116 Studies on genetic divergence in potato (*Solanum tuberosum* L.)
—*Patram Singh, P.K. Sharma, P.K. Joshi, S.K. Nair and Amit Dixit*
- 1117-1120 Interaction effects of nitrogen and potassium on soil available nutrient status, yield and economics of Okra (*Abelmoschus esculentus* L.)
—*B. Naveen Kumar, G. Padmaja and P. Chandrasekhar Rao*
- 1121-1127 Virulence analysis and genetic diversity of *Xanthomonas oryzae* pv. *oryzae* strains in Tamilnadu state, Southern India
—*S. Bharathkumar and S.S. Gnanamanickam*
- 1129-1132 The effect of different olive pulp aqueous extracts on seedling growth of three wheat cultivars, Chamran, Karkhe and Triticale
—*Azadeh Niroomand and Mozhan Vafaei*

Conservation impact on molecular genetic changes on Java green peacock (*Pavo muticus*) through mitochondrial D-loop marker

E. Bimo Aksono H¹* and Herry Agoes Hermadi*

¹Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia

¹Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia

(Received 9 December, 2014; accepted 15 February, 2015)

ABSTRACT

The study of genetic diversity of species Java Green Peacock (*Pavo muticus*) captured from the wild and the conservation-bred in East Java is aimed at investigating the impact of conservation on the molecular genetic changes of Java Green Peacock (*Pavo muticus*). This study utilized samples from 10 (ten) wild Java Green Peacocks (*Pavo muticus*) from the forest and 6 (six) Java Green Peacocks (*Pavo muticus*) bred in the conservation in East Java. The materials examined were their feathers (calamus). The technique of *Polymerase Chain Reaction* (PCR) with D-loop mitochondrial based-designed primer was used to investigate the genetic diversity of species among Java Green Peacock (*Pavo muticus*). PCR products were analyzed using electrophoresis gel 1% . Then they were purified and labelled for data sequencing. The sequencing data obtained were in a form of base order that enabled the analysis of genetic diversity among species (Phylogenetic Analysis) by using a software program from Genetix Mac Ver 10.0. Green Peacock (*Afropavo congensis*) was used as the reference species with the code DQ 834507. The result showed that the 10 (ten) samples of wild Java Green Peacock (*Pavo muticus*) feathers were genetically different from the Green Peacock reported to the GenBank with code DQ 834507 as well as from the 6 (six) conservation-bred species Java Green Peacock (*Pavo muticus*). Compared to wild Java Green Peacock (*Pavo muticus*) from the forest, the conservation-bred Java Green Peacock (*Pavo muticus*) experienced mutations since the result of phylogenetic analysis indicated that the conservation-bred Java Green Peacock species and the wild ones were not in the same cluster. Thus, peacocks kept in conservation, regardless the sophisticated and modern facilities, may lost their inherited genetic characteristics which in return would eliminate Indonesian superior germs for Java Green Peacock (*Pavo muticus*) in Indonesia

Key words : Java Green Peacock, PCR, Mitochondrial D-Loop, Conservation, Indonesia

Introduction

Genetic conservation is mainly used to support scientists to define, describe and prioritize scarce units for conservation (Carty *et al.*, 2009). Globally, the number of endangered species increased significantly in recent years (IUCN, 2008). According to Carty *et al* (2009) the percentage of hindrances will

reach at least 18% by 2050 due to climate change. Adequate conservation measures must be taken to maximize the ability of species to adapt to rapid environmental change.

There are two types of peacocks in the world; the blue peacocks (*Pavo cristatus*) that spread over India and Sri Lanka, and the green peacocks (*Pavo muticus*) that spread in Burma, Thailand, Indochina,

*Corresponding author's email: baksono@yahoo.com

Malaysia and Java island. Green Peacock consists of three subspecies which are green peacocks of Burma (*Pavo muticus spicifer*) spread in Burma (extinct), Indochina Green Peacocks (*Pavo muticus imperator*) spread in Indochina, and Java Green Peacock (*Pavo muticus muticus*) spread in Malaysia (already extinct) and Java (Hernowo, 2011). Java Green Peacocks (*Pavo muticus muticus*) Linnaeus 1758, currently only live in Java. This bird spreads in a relatively small number of population in various habitat types. Java Green Peacocks can be found in some protected areas (Natural Reserves, Wildlife and National Parks) and unprotected areas (productive forests, plantations) (Hernowo, 2011).

On the other hand according to Wan *et al.* (2004), quoting from several sources, they stated that the conservation process led to some differences and similarities with its ancestors. This study is aimed at examining whether the peacocks bred in conservation experience genetic variations. It is very important to consider the genetic changes in order to keep the wealth of Indonesia superior germ plasma as well as to avoid genetic quality degradation.

Therefore, in a bid to maintain Java Green peacocks (*Pavo muticus*), the management factor is needed to keep the existence of the population through genetic enrichment programs, in which the basic information can be identified through phylogenetic reconstruction (Moritz *et al.*, 1996). High genetic variability enable the species to cope with the pressure from environmental changes (Avice, 1994; Hartl, 2000). In addition, the high heterozygosity of a population is a good basic for further developing the population. Appropriate molecular markers to reconstruct population phylogenetic are mitochondrial DNA sequence variations. The characteristics of mitochondrial DNA are: (1) the relatively high speed of evolution (5-10 times) compared to nuclear DNA; (2) Transmittable through the maternal line between generations without experiencing recombination, so that the whole molecules can be considered as a single genetic unit with many alleles (Sudoyo, 1995); and (3) the relatively small size that is easy to observe (Li and Graur, 1991; Taberlet, 1996). In observing the closely-related types, the use of nucleotide sequences of mtDNA control region (D-Loop) can provide good resolution, since the mtDNA D-loop sequences contain a variety of sequences which mutation rate are of 4-5 times faster compared to other parts of mtDNA (Horai *et al.*, 1993).

Materials and Methods

The samples were taken from the feather (Calamus) of 6 (six) Java Green Peacocks (*Pavo muticus*) from conservation areas and 10 (ten) Java Green Peacocks from the forest in Probolinggo, East Java,

DNA extraction was performed on 25 mg (3-4 Calamus), from the apex of the feathers by first adding 180 μ l of lysis buffer (100 ml of 1 M Tris, 200 ml of 0.5 M EDTA, 2 ml of 5 M NaCl, 100 ml of 10 % SDS), 25 μ l of 100 mg/ml DTT (dithiothreitol) and 20 μ l of 10 mg/ml proteinase K. The resulting mixture was further incubated at 50 ° C, to dissolve all parts of the feathers (calamus) in approximately (3-5 hours). Then 400 μ l phenol (Tris-HCl pH 8.0) was added and the mixture was re-incubated for another 30 minutes. Next, it was centrifuged at 13,000 rpm speed for 3-5 minutes, and the top layer was removed to another vial which already contained 400 μ l of chloroform Isoamyl alcohol (24: 1) which was then incubated for 10 minutes. After that, it was centrifuged at 13,000 rpm for 3-5 minutes and the layer formed at the bottom of vial was removed. Next, it was mixed with absolute ethanol and 40 μ l of sodium acetate and incubated for 45 minutes at -20°C. Then it was centrifuged at 13,000 rpm for 30 minutes at 4°C and then cleansed with 10% ethanol and re-centrifuged at the speed of 13,000 rpm for 10 minutes at 4°C. DNA obtained was re-dissolved with 100 μ l TBE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA) to determine the level of purity and spectrophotometer with a wavelength of 260 and 280 nm (Sefc *et al.*, 2003; Leeton *et al.*, 1993; Malago *et al.*, 2002; Kimball *et al.*, 1999).

PCR reaction was using the primer design already prepared which was GPDF (5'-GGGGGTATACTATGCATAATC GTG -3') and GPDR (5'- AAAGAATG GGCCTGAAGCTAGT -3'). To amplify, the materials used were: 25 μ l total volume which consists of 1.5 μ l of 10 \times PCR buffer (Sigma), 1.5 μ l 25 mM MgCl₂, 100 ng for each primer, 200 μ M of each dNTP, 0.5 U Taq DNA polymerase, and 250 ng sample DNA. PCR amplification conditions consist of stage pre-denaturation 94°C for 1 minute 30 seconds, 35 cycles that consist of 94°C for 30 seconds, 57°C for 30 1 minute, and 72°C for 31 minutes, and the final stage elongation 72°C for 10 minutes. The resulting PCR product was 330 bp (Wiwegwean and Meckvichai, 2011).

PCR result 330 bp was visualized by agarose gel 1% that contains ethidium bromide (Sefc *et al.*, 2003;

a separate cluster (MJM-1; MJMJ-1; MJA -1; MJMA-1). Similar condition also observed in the population of Java Green peacock (*Pavo muticus*) from the wild. Yet, despite unclear data identity, there is an assumption F1 and F2 are different, and that several clusters present which are first cluster (M), second cluster (M-2), third cluster (M-3, M-4 and M-5) and fourth cluster (M-6, M-7, M-8, M-9 and M-10) (Fig. 3).

Discussion

The theory of neutral molecular evolution (Wan *et al.*, 2004) suggested that most of the changes in nucleic acid and protein sequences occur selectively. Although still controversial, this theory still highlights the need to consider the act of natural selection at the molecular level.

Along with the decline of peacocks population, especially the Java green peacocks, several attempts have been conducted including breeding in conservation efforts. According to Wiwegwean and Meckvichai (2011) many research have been carried out, especially on *Pavo muticus* in Thailand in relation to genetic variation caused by breeding in conservation. In this case, the genetic diversity of *Pavo muticus* breeding in conservation will vary among populations depending on the number and geographic range of the mothers. Differences in mitochondrial DNA (mtDNA) haplotypes in a population will illustrate some of the origins of the lineage of the female parent. Thus, the study of mtDNA variations may provide insight on the origin of the female parent of the peacocks bred in conservation and the genetic structure of natural or wild populations. In addition, the study can provide some evidences about the origin of the hybridization through mtDNA introgression between *Pavo muticus* and related species.

According to Tarwiningsih (2009), quoting from several sources, she suggested that to detect the molecular basis of genetic diversity may utilize mitochondrial DNA (mtDNA). This method is widely used to examine animal genetic diversity and systematic relationships at different levels of hierarchy (Lamb and Osentoski, 1995) due to the maternal nature of mtDNA, which is purely descended from the female parent. Mitochondrial genome also has a relatively small size, approximately $\pm 16,500$ bp and has a rapid rate of evolution, especially in the control area (D-loop), giving rise to a high diversity in

mtDNA sequences intra-species (Avice, 1994).

The existence of genetic variations that appear in an organism is a result of an evolutionary process in which genetic variation can occur due to a change in gene frequency. According to Tarwiningsih (2009), there are four basic factors that cause changes in gene frequency, i.e. natural selection, mutation, migration and genetic deviation. Natural selection is a natural process in which some individuals have genetic basis linkages that may improve live survival or reproduction to adapt and have offspring that can survive in the environment. The existence of migration causes individuals to move from one area to another. If these individuals survive and reproduce in the new place, they will pass on their genes to their new environment. According Tarwiningsih (2009), the assumption is that the isolated population does not affect changes in genes with neighboring groups. Many populations that are not fully isolated from other populations of the same species, experience some normal changes in genes.

In addition, this research was also expected to reveal the genetic structure of the green peacock from the conservation and the wild ones, including the genetic structure of the kinship hierarchy between F1 and F2. It is important since the regulation stated that the flora and fauna of protected breeding parents from natural habitat (W) is declared belong to the state. Wildlife breeding parents first generation (F1) on conservation are declared as state property. Parent specimens of protected wildlife that come from the natural habitat, and/or captive breeding first generation (F1) of protected wildlife are illegal to be sold and must be submitted to the state at any time.

The research results showed that 10 (ten) samples from wild Java Green Peacock (*Pavo muticus*) feathers were genetically different from the green peacock reported in GenBank with code DQ 834 507 and from 6 (six) samples of conservation-bred Java Green Peacock (*Pavo muticus*). Compared with the conservation-bred ones, it was obvious that the Java Green Peacocks experienced mutation which is indicated by the results of phylogenetic analysis in which both species were not in the same cluster.

Therefore, it should be considered that the peacocks kept in conservation, despite sophisticated and modern facilities provided, experience a decrease in any inherited genetic characters which may eliminate Indonesian superior germs in relation to Java Green Peacock (*Pavo muticus*) in Indonesia.

Acknowledgement

Our gratitude is extended to the Rector of Universitas Airlangga who has granted the researchers BOPTN fund to support the excellent research universities in year 2013 and 2014.

References

- Awise, J.C. 1994. Molecular markers, Natural history and evolution. Chapman and Hall, New York.
- Carty, J., Ljungqvist, C., Prest, D., Segura, C. and Zimmering, H. 2009. How can molecular genetics help us to prioritize taxa for conservation. *Journal of Conservation Biology*. 3065. 1 : 38-45.
- Hartl, D. 2000. A primer of population genetics. 3rd ed. Sinauer Associates, Inc., Sunderland.
- Hernowo, J.B. 1995. Ecology and Behavior of the Green Peafowl (*Pavo muticus* Linnaeus 1766) In the Baluran National Park. East Java, Indonesia. Master Thesis Faculty of Forestry Science, Goerg August University of Gottingen. Germany.
- Hernowo, J.B. 2011. Ekologi merak hijau jawa (*Pavo muticus muticus*) Linnaeus 1758 pada beberapa tipe habitat di ujung timur penyebarannya di Jawa Timur, Indonesia. Disertasi. Program Studi Ilmu Pengetahuan Kehutanan Sekolah Pasca Sarjana Institut Pertanian Bogor
- Horai S., R. Kondo, Y. Nakagawa-Hattori, S. Hayashi, S. and K. Tajimaas Sonada. 1993. Peopling of the Americas, founded by four major lineages of mitochondrial DNA. *Mol Biol Evol*. 10 : 23-47.
- Holmes, D.A. 1989. Status report on Indonesian Galliformes. *Kukila* 4 : 133-143.
- IUCN. 2008. Red List - Home Page. <http://www.iucnredlist.org/> (Retrieved 9/2/2009)
- Kimball, R.T., Braun, E.L., Zwartjes, P.W., Crowe, T.M. and Ligon, D. 1999. A Molecular phylogeny of the pheasants and partridges Suggests that Reviews These lineages are not monophyletic. *Mol. Phylogenetics and Evol*. 11 (1) : 38-54
- Lamb, T. and Osentoski, M.F. 1995. Intraspecific phylogeography of the gopher tortoise, gopher polyphemus: RFLP analysis of amplified mtDNA segments. *Molecular Ecology* 4 : 709-718.
- Leeton, P., Christidis, L. and Westerman, M. 1993. Feathers from museum bird skins: a good source of DNA for phylogenetic studies. *The Condor*. 95 : 465-466.
- Li, W. and D. Graur. 1991. Fundamentals of molecular evolution. Sinauer Associates, Inc., Sunderland.
- Malago, W.Jr., Franco, HM, Matheucci, E.Jr., Medaglia, A. and Henrique-Silva, A.F. 2002. Large scale sex typing of ostriches using DNA extracted from feathers. *BMC Biotechnol*. 2 : 19.
- Moritz, C., Wilmer, J.W., Pope, L., Sherwin, W.B., Taylor, A.C. and Limpus, C.J. 1996. Applications of genetics to the conservation and management of the Australian fauna: Four case studies from Queensland. In: Smith, T.B. and R.K. Wayne. (Ed). Molecular genetic approaches in conservation. Oxford University Press, New York: pp. 442-456.
- Sativaningsih, D. 2005. Ekologi Perilaku Merak Hijau (*Pavo muticus* Linnaeus, 1766) Di Taman Nasional Alas Purwo Jawa Timur. Skripsi. Departemen Konservasi Sumberdaya Hutan dan Ekowisata, Fakultas Kehutanan IPB, Institut Pertanian Bogor, Unpublished.
- Sefc, K.M., Payne, R.B., Sorenson, M.D. 2003. Microsatellite amplification from museum feather samples: effects of fragment size and template concentration on genotyping errors. *The Auk*. 120 : 982-989.
- Sudoyo, H. 1995. Mutasi DNA mitokondria sebagai penyebab kelainan genetika. Bagian Biologi Fakultas Kedokteran Universitas Indonesia, Jakarta.
- Sulandari, S. and Zein, M.S.A. 2009. Analisis D-loop Mitokondria untuk memposisikan Ayam Hutan Merah dalam Domestikasi Ayam di Indonesia. *Media Peternakan*, Vol. 32 No. 1. things. 31-39.
- Taberlet, P. 1996. The use of mitochondrial DNA control region sequencing in conservation genetics. In: Smith, T.B. and R.K. Wayne (Eds). *Molecular Genetic Approaches in Conservation*. Oxford University Press, New York. pp. 125-142.
- Tarwiningsih, W. 2009. Analisis keragaman genetic kerbau lokal (*Bubalus bubalis*) berdasarkan haplotype DNA mitokondria. Skripsi. Major Ilmu Produksi dan Teknologi Peternakan, Fakultas Peternakan, Institut Pertanian Bogor.
- Wan, Q.H., Wu, H. and Fujihara, T. 2004. Which genetic marker for which conservation genetics issue? *Electrophoresis*. 25 : 2165-2176.
- Wiwegwean and Meckvichai, W. 2011. Genetic variation of Green Peafowl *Pavo muticus* captive in Thailand based on D-loop sequences. Short Communication. *International Journal of Galliformes Conservation, World Pheasant Association*. 2 : 38-42.



**KOMISI ETIK PENELITIAN
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA
*Animal Care and Use Committee (ACUC)***

**KETERANGAN KELAIKAN ETIK
“ ETHICAL CLEARENCE ”**

No : 213-KE

**KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE)
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA,
TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG
DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :**

**PENELITIAN BERJUDUL : CONSERVATION IMPACT ON MOLECULAR
GENETIC CHANGES ON JAVA GREEN PEACOCK
(PAVO MUTICUS) THROUGH MITOCHONDRIAL D-
LOOP MARKER**

PENELITI UTAMA : HERRY AGOES HERMADI

**UNIT/LEMBAGA/TEMPAT PENELITIAN : FAKULTAS KEDOKTERAN HEWAN,
UNIVERSITAS AIRLANGGA**

DINYATAKAN : LAIK ETIK

Surabaya, 14 Maret 2013

Ketua,

Dr. E. Bimo Aksono H, M.Kes.,Drh.
NIP. 196609201992031003

Mengetahui,
Dekan FKH-Unair

Prof. Bonziah Sidik, Ph.D.,Drh
NIP. 19531216198062001

