


Hunan Daxue Xuebao/Journal of Hunan University Natural Sciences

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
China	Multidisciplinary └ Multidisciplinary	Hunan Daxue/Hunan University Hunan University in Scimago Institutions Rankings	17
Journals	10002472, 16742974	1992-1993, 1996, 2001-2021	

SCOPE

Information not localized

Join the conversation about this journal



FIND SIMILAR JOURNALS ?options ⋮

1
Tumu Gongcheng
Xuebao/China Civil
CHN

25%
similarity

2
Xi'an Jianzhu Keji Daxue
Xuebao/Journal of Xi'an
CHN

24%
similarity

3
Tumu Jianzhu yu Huanjing
Gongcheng/Journal of Civil,
CHN









23%
similarity

4
Tumu yu Huanjing
Gongcheng Xuebao/Journal
CHN

23%
similarity

5
Dongnan Daxue Xuebao
(Ziran Kexue Ban)/Journal of
CHN

21%
similarity



← Show this widget in
your own website
Just copy the code below
and paste within your html
code:
[a href="https://www.scimagojr.com/widget.php?journal=HunanDaxueXuebao&year=2021&quartile=Q2&language=en&format=js">](https://www.scimagojr.com/widget.php?journal=HunanDaxueXuebao&year=2021&quartile=Q2&language=en&format=js)

G SCImago Graphica
Explore, visually
communicate and make
sense of data with our
[new data visualization
tool](#).



Metrics based on Scopus® data as of April 2022



jawad 3 weeks ago

Dear Sir MAdam

Is this journal still indexed in Scopus in the current year 2023 ?

Thank you so much



Source details

Hunan Daxue Xuebao/Journal of Hunan University Natural Sciences

Scopus coverage years: from 1992 to 1993, 1996, from 2001 to 2022

Publisher: Hunan Daxue/Hunan University

ISSN: 1674-2974

Subject area: Multidisciplinary

Source type: Journal

CiteScore 2021
0.9

SJR 2021
0.243

SNIP 2021
0.482

[View all documents >](#)[Set document alert](#) [Save to source list](#)[CiteScore](#)[CiteScore rank & trend](#)[Scopus content coverage](#)

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語版を表示する](#)
[查看简体中文版本](#)
[查看繁體中文版本](#)
[Просмотр версии на русском языке](#)

Customer Service

[Help](#)
[Tutorials](#)
[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗ All rights reserved. Scopus® is a registered trademark of Elsevier B.V.
We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

 RELX

Journal of Hunan University Natural Sciences

湖南大学学报

Journal of Hunan University



Home > Archives > Vol 48, No 6 (2021)

Vol 48, No 6 (2021)

Table of Contents

Articles

A New Approach to Detect COVID-19 in X-Ray Images of Indonesians

Anggraini Dwi Sensusiati, Aji Sapta Pramulen, Dewinda Julianensi Rumala, I Ketut Eddy Purnama, Alfian Nur Rosyid, Muhammad Amin, Rofida Lathifah

PDF



Analysis of the Effect of Green Organizational Culture on Organizational Performance and Competitive Advantages of Green through Green Innovation in Manufacturing Industries

Kristian Chandra, Willy Arafah, Yuzwar Z. Basri

PDF



Characteristics of Marine Heatwaves off West Sumatra Derived from High-Resolution Satellite Data

Mochamad Furqon Azis Ismail

PDF



Comparison of DEMs Spatial Resolution for Geomorphological Study in a Small Volcanic Island of Tidore, North Maluku, Indonesia

Bachtiar W. Mutaqin, Muh Aris Marfa, Danang Sri Hadmoko, Helvetia Wijayanti, Franck Lavigne, Audrey Faral

PDF



Date Palm Extract Effect on Quality of Mahseer Fish (Tor Soro) Spermatozoa after Frozen Storage

Abinawanto, Dyah Putri Alifiani, Anang Hari Kristanto, Astari Dwiranti, Fadhillah, Anom Bowolaksono, Retno Lestari

PDF



Determination of Low Level 137Cs in Environmental Water Sample Using AMP Method and a Review Comparing with Other Adsorbents

Deddy Irawan Permana Putra, Shinya Ochiai, Seiichi Tomihara, Wahyu Retno Prihatiningsih, Murdahayu Makmur, Seiya Nagao

PDF



Distribution Pattern Identification of Mineral using XRF and XRD Method in Jeneberang Watershed, Indonesia

Muhammad Altin Massinai, Wahyuningsih Mamudi, Muhammad Fawzy Ismullah Massinai

PDF



Enhancement of Eggplant Production in Intercropping Systems through Integrated Nutrient Management

Moch. Dawam Maghfoer, Mahayu Woro Lestari

PDF



High-Productivity Traditional Bali Palung Salt Method for Small Production Fields

Misri Gozan, Yeti Rochwulaningsih, Singgih Tri Sulistiyo, Makhfud Efendy, Nurul Intan, Andre Fahriz Perdana Harahap, Hafidzurrahman Suhairi, Muhammad Yusuf Arya Ramadhan, Nurul Hidayat, Siti Fauziah Rahman, Mahendra P. Utama, Noor Naelil Masruroh, Susilo Budiyanto, Muhammad Sahlan, Shigenori Kumazawa

PDF



Influence of Geomorphology and Siltstone Structure on the Land Vulnerability in Mamberamo Raya, Papua, Indonesia

Adi Tonggiroh

PDF



Integrated Family Development Intervention on Perceived Stress and Parental Self-Efficacy of Parents of Children with Autism Spectrum Disorder

Diah Curie Kharisma, Agus Suwandono, Annastasia Ediati

PDF





Krisna Damayanti, Noorlailie Soewarno



Not Privatization but Partnership: Government Policy in Providing Public Service Infrastructure in Indonesia



Sugeng Suharto, Sri Suharti, Untung Muhdiarto, Londa Very Yohanes, Sam Ratulangi

Optimal Machine Learning Models for Kitsune to Detect Mirai Botnet Malware Attack



Abdullah Alabdulatif, Syed Sajjad Hussain Rizvi, Manzoor Ahmed Hashmani

StegoBound: A Novel Image Steganography Technique Using Boundary-Based LSB Substitution



Syed Saad Ahmed, Mashal Memon, Rabeea Jaffari, Moazzam Jawaid

The Effect of Economic Growth, Investment on Local Indigenous Income in Jambi Province



M. Sudirman, M. S. Zahari, Willy Arafah

The Use of ICT in Thirukkural Teaching and Facilitation



Karthegees Ponniah, Ilangkumaran Sivanadhan, Muniswaran Kumar, Pooveneswaran Nadarajan, Aigul Akhmetova

Transplantation of Acropora in Different Depth: First Report on an Effort for Environmental Sustainability in Gili Ketapang, Probolinggo



Andys Dwianinda Rosaline, Mochamad Faisal Trisatrio, Muhammad Browijoyo Santanumurti, Fitri Budiyanto, Boedi Setya Rahardja, Putri Desi Wulan Sari

A Model for Prevention of Forest and Land Fires Crimes Using a Sustainable Plantation Approach



Aji Kurnia Dermawan, Sigid Suseno, Adrian E. Rompis, Dadang Epi Sukarsa

Comparison of CDC Bottle Bioassay Test with WHO Standard Method for Assessment of Aedes Aegypti Susceptibility to Carbamates and Organophosphates Insecticides in Semarang, Indonesia



Muhammad Choirul Hidajat, Martini Martini, Nur Endah Wahyuningish, Sayono, Ristiyanto, Triwibowo Ambar Garjito, Widiarti, Lasmiatyi, Sapto Prihasto Siswoko, Rima Tunjungsari Dyah Ayuningtyas, Ary Oktarsi Yanti, Suharyo Hadisaputro

Critical Reading Teaching Materials Show R-Tam-Based Malay Teaching Loaded with Humanist Literacy: Future Language Learning Innovations



Asnawi, Ida Zulaeha, Teguh Supriyanto, Hari Bakti Mardikantoro, Sri Wahyuni

Development of Junior Karate Physical Test Instruments in Kumite Category



Jujur Gunawan Manullang, Soegiyanto, Sulaiman, Oktia Woro Kasmini Handayani

Effect of Compensation and Discipline on Employee Performance: A Case Study Indonesia



Indra Prasetyo, Esa Wahyu Endarti, Budi Endarto, Nabilah Aliyyah, Rusdiyanto, Heru Tjaraka, Nawang Kalbuana, Arif Syafi'ur Rochman

Financial Decision-Making during the COVID-19 Pandemic in Population Administration of the Republic of Indonesia



Muhamad Arief Affandi, Etty Murwaningsari

Integrated Security System Implementation for Network Intrusion



Resevoa Moral Muhammad, Indrarini Dyah Irawati, Muhammad Iqbal

Isolation: The Root Cause of Indigenous Poverty in Lumoli Village



Elsina Titaley, Tony Donald Pariela

Pekanbaru Adolescents' Perceptions Concerning the Existence of Traditional Sports in the Millennial Era



M. Fransazeli Makorohim, Soegiyanto, Taufiq Hidayah, Setya Rahayu

Strategy for Strengthening Character Education in Physical Education Learning at Makassar City Elementary Education Level



Suastika Nurafiat, Tandiyo Rahayu, Sugiharto, Harry Pramono

The Cytotoxic Activity of Crude Aqueous Annona Plant-Derived Extracts on Spodoptera frugiperda, Cell Lines

PDF


 Jesús A. Polo Olivella, Marlinda Lobo de Souza, Ericsson Coy-Barrera, Jhon F. Betancur Pérez, Jorge W. Arboleda Valencia

The Effect of Company Size as a Mediating Variable on the Relationship of Working Capital to Profitability

PDF




Susanti Widhiastuti

The Effect of Economic Growth, Investment on Local Indigenous Income in Jambi Province

PDF




M. Sudirman, M. S. Zahari, Willy Arafah

The Effect of Handstand Exercises on Entry Results in Diving Athletes, South Sumatra

PDF




Kevin Octara, Soegiyanto, Tjepjep Rohendi Rohidi, Sulaiman

The Impact of Globalization and International Integration to Become a Global Citizenship in the Southeast – from the Practice of Ho Chi Minh City, Vietnam

PDF




Pham Ngoc Tram, Ngo Hong Diep

The Influence of Management Commitment on the Achievement of Lecturers through Work Motivation and Organizational Citizenship Behavior

PDF




Wahju A. Rini, Pamela H. Heng, Bobby Putrawan, Zeplin J. H. Tarigan, Sundring P. Djati, Ahmad Hidayat Sutawidjaya

The Religiosity Mechanism on Political Connections in the Indonesian Stock Exchange

PDF




Zunairoh, Liliana Inggrit Wijaya

The Role of Absorptive Capacity and Organizational Innovation in Culinary Business

PDF




Zoel Hutabarat, Willy Arafah, Farida Jasfar

The Role of ApOE, PSEN1, And Tnf- α as Predictor of COVID-19 Severity Evaluated Using Serial Thorax Radiograph

PDF




Widiana Ferriastuti, Aryati, Harianto Notopuro, Anggraini D. Sensusati, Paulus Sugianto, Anita Widyoningroem

Editorial Team

Editorial Policies

Focus & Scope

Author Guidelines

Abstracting and Indexing

Publication Ethics

Paper Submission

Article Processing Charge

Language Editing Services

Special Issues

INDEXED BY SCOPUS

Q2 Multidisciplinary

SJR 2021 best quartile
0.24

powered by scimagojr.com



WEB OF SCIENCE™

TEMPLATE



Journal
Template

USER

Username

Password

Remember me

Login

NOTIFICATIONS

› [View](#)

› [Subscribe](#)

JOURNAL CONTENT

Search

Search Scope

Search

Browse

▼ ▾

- [By Issue](#)
- [By Author](#)
- [By Title](#)

INFORMATION

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)

Journal of Hunan University Natural Sciences

湖南大学学报

Copyright 2012-2023
(e-ISSN: 1674-2974)

[Home](#)

[Search](#)

[Archives](#)

[Contact](#)

[About](#)

[Current](#)

[Sitemap](#)

[Paper](#)

[Submission](#)

+86-731-88823137

editorial-office@jonuns.com



Journal of Hunan University Natural Sciences

湖南大学学报

Journal of Hunan University



Home > Editorial Team

Editorial Team

Editor-in-Chief:

Prof. Yi Weijian

Advisors:

Deng Wei, Party secretary of the CPC Hunan University Committee; professor.

Duan Xianzhong, member of the CPC Hunan University Committee and its standing committee, president of Hunan University; professor and doctoral supervisor.

Yu Xiangcheng, vice secretary of the CPC Hunan University Committee.

Xie Chi, HNU Vice President, member of the Academic Research and Academic Degrees Committee and director of the Academic Committee of Department of Management of the HNU.

Wang Weibin, Vice president of Hunan University.

Editorial Team

Editorial Policies

Focus & Scope

Author Guidelines

Abstracting and Indexing

Publication Ethics

Paper Submission

Article Processing Charge

Language Editing Services

Special Issues

INDEXED BY SCOPUS





WEB OF SCIENCE™

TEMPLATE



Journal
Template

USER

Username

Password

Remember me

Login

NOTIFICATIONS

› [View](#)

› [Subscribe](#)

JOURNAL CONTENT

Search

Search Scope

 All ▾

Search

Browse

› [By Issue](#)

› [By Author](#)

› [By Title](#)

INFORMATION

- › [For Readers](#)
- › [For Authors](#)
- › [For Librarians](#)

Journal of Hunan University Natural Sciences

湖南大学学报

Copyright 2012-2023
(e-ISSN: 1674-2974)

[Home](#)

[Search](#)

[Archives](#)

[Contact](#)

[About](#)

[Current](#)

[Sitemap](#)

[Paper Submission](#)

+86-731-88823137

editorial-office@jonuns.com



Open Access Article

The Role of ApoE, PSEN1, And Tnf- α as Predictor of COVID-19 Severity Evaluated Using Serial Thorax Radiograph

Widiana Ferriastuti¹, Aryati^{2,*}, Hariant Notopuro³, Anggraini D. Sensusiat⁴, Paulus Sugianto⁵, Anita Widyoningroem¹

¹ Radiology Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia

² Clinical Pathology Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia

³ Biochemistry Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia

⁴ Radiology Department of the Universitas Airlangga Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia

⁵ Neurology Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia

Abstract: The high number of Covid-19 incidents is the main problem currently being faced, so how to reduce the number of incidents is a significant concern at this time. It is hoped that the risk factors and causes of mortality and how to detect the severity of Covid-19 can be a solution to reduce the morbidity of Covid-19 infection. This research is an analytical observational using the ApoE, Presenilin 1 (PSEN1), and TNF- α from the blood sample of Covid-19 patients; moreover, the thorax serial radiographs were evaluated by CARE score and were also used to determine the prognosis. All of the three indicators were examined using the ELISA Kit. Furthermore, the variables were analyzed using the SPSS 16.0 Ver. with Pearson and Spearman Rho data analysis. There is no correlation between ApoE, PSEN1, and TNF- α as Covid-19 severity predictors in this research regarding factors such as fewer geriatric samples, steroid usage as therapy, and fewer patients with comorbidities.

Keywords: apolipoprotein E, presenilin 1, TNF- α , COVID-19, severity, predictors.

载脂蛋白乙、早老素-1 和肿瘤坏死因子- 一种作为新冠肺炎严重程度预测因子的作用·使用连续胸片评估

摘要：新冠肺炎事件的大量发生是当前面临的主要问题，因此如何减少事件数量是此时的一个重要问题。希望死亡的风险因素和原因以及如何检测新冠肺炎的严重程度可以成为降低新冠肺炎感染发病率的解决方案。这项研究是使用新冠肺炎患者血液样本中的载脂蛋白乙、早老素1和肿瘤坏死因子-

一种进行的分析观察；此外，胸部系列射线照片片通过心脏麻醉风险评估评分进行评估，也用于判断预后。使用酶联免疫吸附测定试剂盒检查所有三个指标。此外，使用社会科学统计包16.0对变量进行分析。使用皮尔逊和斯皮尔曼罗数据分析。在本研究中，载脂蛋白乙、早

Received: March 14, 2021 / Revised: April 17, 2021 / Accepted: May 13, 2021 / Published: June 28, 2021

About the authors: Widiana Ferriastuti, M.D, Radiologists Consultant, Radiology Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia; Aryati, Clinical Pathology Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia; Hariant Notopuro, Biochemistry Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia; Anggraini D. Sensusiat, Radiology Department of the Universitas Airlangga Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia; Paulus Sugianto, Neurology Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia; Anita Widyoningroem, Radiology Department of the Dr. Soetomo General Academic Hospital, Medical Faculty, Universitas Airlangga, Surabaya, Indonesia

Corresponding author Aryati, aryati@fk.unair.ac.id

老素1和肿瘤坏死因子-

一种作为新冠肺炎严重程度预测因子与老年人样本较少、使用类固醇治疗和合并症患者较少等因素之间没有相关性。

关键词：载脂蛋白乙、早老素1、肿瘤坏死因子-

一种、新冠肺炎、严重程度、预测因子。

1. Introduction

SARS-CoV2, more popularly known as Coronavirus Disease 2019 (Covid-19), is a significant health problem that causes mortality, especially for the elderly. World Health Organization (WHO) has declared the Covid-19 as a pandemic on 11 March 2020, in which more than 1.8 million people were positive for Covid-19; moreover, there were more than 114.000 deaths globally [1].

The mortality for Covid-19 ARDS ranged between 26% and 61.5% if he had been treated in critical care. If he had received treatment using mechanical ventilation, the mortality could increase by about 65.7% to 94% [2]. Several risk factors could worsen the Covid-19 patients, which consist of older ages, comorbidity such as hypertension, diabetes, cardiovascular-related disease, kidney injury, lower lymphocyte counts, and raise D-dimer levels. Furthermore, 53% of deaths regarding Covid-19 ARDS were attributed to respiratory failure, 33% were caused by respiratory failure, combined with cardiac failure. In comparison, 7% of deaths were caused by myocardial damage and circulatory failure by unspecified causes [2].

In the brain and lungs infected with SARS-CoV2, an immune reaction occurs, which results in the activation of gold cells and the release of previously formed and newly formed mediators in the brain. This process then continues with the activation of glial cells, which causes neurons' degeneration when a cytokine storm occurs in the lungs, reaching the brain via the blood-brain barrier, possibly exacerbating neurological complications. In these circumstances, patients with previous neurological disorders (ADRD, PD, stroke, NMD) have a higher risk of developing comorbidities. Special attention and medical support systems are needed in these circumstances [3].

Adipocytes, hepatocytes, brain astrocytes, and arterial wall macrophages synthesize ApoE, which functions in lipids' transport, which plays a role in the brain, immune and vascular functions [4]. Compared with homozygous ApoE3, homozygous ApoE4 is 2.2 times the risk of positivity for Covid-19 and 4.3 times as much as death cases after Covid-19 [5], [6]. Covid-19 under linear dose dependence was associated with heterozygosity ($\epsilon 3/\epsilon 4$), whereas dementia, hypertension, coronary heart disease (CHD) or type II

diabetes did not reduce the strong association for the association between $\epsilon 4 / \epsilon 4$ and Covid-19 [4].

Cytokine storms play a central role in severity and mortality in SARS-CoV2 infection with increased plasma levels of IL-1 β , IL-7, IL-8, IL-9, IL-10, G-CSF, GM-CSF, IFN γ , IP-10, MCP-1, MIP-1 α , MIP-1 β , PDGF, TNF α , and VEGF in both ICU and non-ICU patients. Furthermore, patients with severe pneumonia who became ARDS admitted to the ICU and required oxygen compared to non-ICU patients with ARDS pneumonia found higher plasma levels of IL-2, IL-7, IL-10, G-CSF, IP-10, MCP-1, MIP-1 α , and TNF- α [7].

When Covid-19 infection meets the 2012 ARDS Berlin diagnostic criteria, the patient can be diagnosed with Covid-19 ARDS, with the assessment criteria (i) acute hypoxic respiratory failure, (ii) bad respiratory symptoms within one week, (iii) bilateral air space on radiographs not fully explained by effusion, lobar, pulmonary collapse or nodules, (iv) acute hypoxic respiratory failure not caused by heart failure [4].

The findings (CT) may be more sensitive than reverse transcription PCR (RT-PCR); chest radiography can be applied as the primary tool for diagnosis and monitoring. GG and Co on chest radiographs did not investigate their contribution; thus, chest radiographic scores (CASE) was used based on GG and Co's extension, considering the two subscores separately and evaluating their role in predicting patient outcomes [8]. There are numerous studies regarding Covid-19 that discuss practical and efficient methods to determine this disease as early as possible. The previous studies have tried ApOE and TNF- α as Covid-19 predictors with different results; however, the PSEN1 has never been used as the Covid-19 predictor. Hence, this study attempts to combine the ApOE and TNF- α with the PSEN1 to determine whether these indicators could be a proper Covid-19 predictor. Furthermore, we hypothesize that ApOE, PSEN1, and TNF- α might be a severity predictor for Covid-19 infection.

2. Methods

This research is quantitative and is included in the analytical observational study. It had 64 samples which then turned into 56 samples regarding the exclusion criteria of the samples. The instruments were used Human PS1 (Presiniilin 1) ELISA Kit, TNF- α (Human)

ELISA Kit, Human ApoE (Apolipoprotein E) ELISA Kit, and patient's thorax serial radiographs. The sampling procedure was first asked for the patient's informed consent, and then the blood sampling was taken due to the ApoE, PSEN1, and TNF- α examination. Second, the thorax serial radiographs were collected and examined by two radiographers in their fields. Furthermore, the ApoE, PSEN1, and TNF- α examinations were done using the ELISA Kit.

Moreover, the collected samples were processed using the SPSS Ver.16.0 application with Spearman rho data analysis to determine the correlation between the dependent and independent variables. In addition, this research had accepted the ethical clearance before it was started by the Health Research Ethics Committee of Dr. Soetomo General and Academic Hospital.

3. Results

There are several characteristics of Covid-19 patients collected as well, as the sampling process was done.

Table 1 Demography data of COVID-19 patients

Category	n = 65
Age	
(Adults = 20-60 years old)	48
(Elderly = > 60 years old)	8
Sex	
(Male)	40
(Female)	16
Comorbidity	12
Steroid Therapy	17
Severity	
(Remain)	19
(Better)	23
(Worsen)	14
Outcome	
(Heal)	41
(Passed away)	15

The table describes several Covid-19 patients' demography, which consists of Age, Sex, Comorbidity, Steroid therapy, Severity degree, and the outcome during treatments. It could be seen that the adults and the male sex have dominated the sample at 85.7% and 71.4 %, respectively. Moreover, a small proportion of respondents had the previous comorbidity at 21.42%, and about 30.3% of patients have received steroid therapy. The better prognosis was dominated the Covid-19 patients at 41.1%, followed by the better outcome at 73.2%.

The ApoE, PSEN1, and TNF- α were examined and processed by SPSS Ver.16 with Spearman Rho data analysis with the basis of decision making such as there is a significant correlation between variables if the value is sig. <0.05 and if the value is sig. > 0.05 , there is no significant correlation between variables. Interpretation of the coefficient correlation between variables ranges from 0.01 to 1.00 with the following description: almost no correlation with the range 0.00-0.20, low correlation range 0.21-0.40, moderate

correlation range 0.41-0.60, high correlation range 0.61-0.80, and perfect correlation in the range 0.81-1.00.

Table 2 Correlation between ApOE, PSEN1, and TNF- α with the severity score

Category	p-value	Coeficien Correlation
ApOE and severity score	0.440	-0.105
PSEN1 and severity score	0.338	0.118
TNF- α	0.776	0.039

The relation between severity score, which was tested with ApOE, PSEN1, and TNF- α , is mentioned in the table above. It could be seen that there is no correlation between all of the three indicators and severity score regarding the p-values of ApOE, PSEN1, and TNF- α (p-value = 0.440, 0.338, and 0.776 respectively; $\alpha = 0.05$). Furthermore, the ascendency of the indicators and severity score was tenuous, which was then proven by the result. The ApOE had the opposite direction of work; hence, the value was negative.

Table 3 Correlation between ApOE, PSEN1, and TNF- α as the severity predictors of COVID-19

Category	p-value	Coeficien Correlation
ApOE and severity	0.06	-0.253
PSEN1 and severity	0.730	0.047
TNF- α	0.183	0.180

The table describes the results of whether the ApoE, PSEN1, and TNF- α could become excellent severity predictors of Covid-19. It was found that neither the ApoE, PSEN1, nor TNF- α could become a Covid-19 severity predictor. There was no correlation between the three indicators as a Covid-19 severity predictor (p-value = 0.06, 0.730, and 0.183 respectively, which more than $\alpha = 0.05$). Each indicator had a poor relation as Covid-19 severity predictor; however, one thing that should be highlighted about the ApoE as the Covid-19 severity predictor is that it has controversy about whether the Variable ApoE increases the other variable will decrease.

4. Discussion

In previous studies, having one or two copies of ApoE4 predisposed the Severity of SARS-CoV2 disease, starting with a robust innate immune response, developing a cytokine storm, and ending with ARDS [9]. ApoE with e4e4 increases the risk of Severity of Covid-19 infection despite a pre-existing history of cardiovascular disease, type-2 diabetes, and dementia. Lipoprotein function and the pro-/anti-inflammatory macrophage phenotype were also influenced and moderated by ApoE e4. ACE2 and ApoE receptors are expressed in alveolar type II cells in the lungs, and as is known, SARS-CoV2 enters cells using ACE2 receptors. However, further study is needed to

understand better the biological mechanisms between the ApOE genotype and the Severity of Covid-19 [5].

Since the cytokines in this study consist of IL-6, IL-8, TNF- α , and IL-1 β as a whole, we can predict the patient's life expectancy by examining the patients when they first arrived. After adjusting for demographics and comorbidities, each cytokine could predict overall survival independently. IL-6 and IL-8 levels were known to have a strong association with severity (moderate, severe, and severe with end-organ damage), taking into account lung imaging, creatinine clearance (CrCl), vasoactivity, and use of ventilation. However, TNF- α was found to be can differentiate between moderate and severe Covid-19 presentation, use of mechanical ventilation, and only increase with end-organ damage [10].

On the contrary, Presenilin 1 (PSEN1) has not been found yet regarding its correlation to predict the Covid-19 Severity. Hence, the PSEN1 in this research is relatively new.

This study found that ApoE, PSEN1, and TNF- α could not predict the severity of Covid-19 disease due to several reasons: the patient's age where the elderly in this study were minor, then the use of steroids by some patients, and comorbidities suffered by the patient.

The high mortality rate belonged to Elderly patients due to the increase of CFR and symptomatic infection rates. Some studies also say that a significant risk factor for death from Covid-19 is old age [11], [12], [13]. Hence, this research could not prove the three indicators as the good predictors of Covid-19 Severity regarding the less sample of elderly.

Furthermore, steroid usage as therapy could affect the study's result. The timing of corticosteroid therapy was associated with the death of Covid-19 patients at the hospital. Bahl et al. mentioned in their research that Corticosteroid initiation will be triggered at age onset > 7 days. Corticosteroid administration should be withheld for 48-72 hours without invasive mechanical ventilation and can be given if the patient remains in hospital. Hypoxia with supplemental oxygen therapy cannot be the standard for corticosteroid administration unless at the right time [14].

However, in the other study by Gibson, the adjunctive therapies most frequently used in typical ARDS are continuous neuromuscular blocking agents, high-dose corticosteroids, and recruitment maneuvers. In Covid-19 ARDS, corticosteroid administration is only recommended in patients with concomitant shock who are unresponsive to vasopressors, so evidence of systemic steroids is scarce. It is feared that the release of the virus and a higher mortality rate could occur due to steroid administration [15]. Additionally, administration of corticosteroid therapy is associated with delayed viral clearance in other Novel Coronavirus outbreaks. Corticosteroid therapy in pneumonia caused by the influenza virus has a poor

outcome in patients' clinical work, including secondary bacterial infection and death [16].

In this research, 17 patients received the steroid therapy with results about three of 16 Covid-19 patients had worsened outcomes, and 5 of 15 patients who received the steroid were dead.

Factors that also influence this study are the comorbidities suffered by patients. It was found that 12 patients in this study had comorbidity that could exacerbate Covid-19 infections such as hypertension, type-2 diabetes mellitus, chronic kidney, and cardiovascular disease. Complementary conditions make it difficult for sufferers of SARS-CoV2 infection in a review with a broad scope, so patient management and proper medical care are essential steps to maintain patient survival [17]. Compared with the results of this study and previous research, there is an inequality in the results obtained, which may be due to the lack of sample size and time of the study and difficulties when collecting data with the Covid-19 situation is not flexible.

5. Conclusion

In this study, it was found that ApoE, PSEN1, and TNF- α were not good predictors for determining how severe the patient's Covid-19 infection was. This outcome is due to several factors that affect the results, such as the lack of a sample of the elderly, the use of steroid therapy, which still requires further research regarding the pros and cons of previous studies, and the small number of samples of patients who have comorbid that can exacerbate Covid-19 infection.

Acknowledgment

Gratitude is given to those who have helped in this research preparation process, namely, the Health Research Ethics Committee of Dr. Soetomo General and Academy Hospital, which funds this research. Other researchers and other related departments have helped this research process.

References

- [1] YUEN K. S., YE Z. W., FUNG S. Y., CHAN C. P., and JIN D. Y. Research Highlight SARS-CoV-2 and COVID-19: The Most Important Research Questions. *Cell & Bioscience*, 2020: 1–5. <https://doi.org/10.1186/s13578-020-00404-4>
- [2] WU C., CHEN X., CAI Y., XIA J., ZHOU X., XU S., HUANG H., ZHANG L., ZHOU X., DU C., ZHANG Y., SONG J., WANG S., CHAO Y., YANG Z., XU J., ZHOU X., CHEN D., XIONG W., XU L., ZHOU F., JIANG J., BAI C., ZHENG J., and SONG Y. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *Journal of the American Medical Association Internal Medicine*, 2020, 180(7): 934–943. <https://doi.org/10.1001/jamainternmed.2020.0994>
- [3] KUMAR D., JAHAN S., KHAN A., SIDDIQUI A. J., and REDHU N. S. Neurological Manifestation of SARS-

- CoV-2 Induced Inflammation and Possible Therapeutic Strategies Against COVID-19. *Molecular Neurobiology*, 2021, 14: 1–18. <https://doi.org/10.1007/s12035-021-02318-9>
- [4] FINCH C. E., & KULMINSKI A. M. Letter to the Editor The ApoE Locus and COVID-19 : Are We Going Where We Have Been ? *Journal of Gerontology: Series A*, 2021, 76(2): e1–e3. <https://doi.org/10.1093/gerona/glaa200>
- [5] KUO C., PILLING L. C., ATKINS J. L., MASOLI J. A. H., DELGADO J., KUCHEL G. A., and MELZER D. APOE e4 Genotype Predicts Severe COVID-19 in the UK Biobank Community Cohort. *Journal of Gerontology: Medical Sciences*, 2020, 75(11): 2231–2232. <https://doi.org/10.1093/gerona/glaa131>
- [6] KUO C., PILLING L. C., ATKINS J. L., MASOLI J. A. H., DELGADO J., KUCHEL G. A., and MELZER D. Letter to the Editor ApoE e4e4 Genotype and Mortality With COVID-19 in UK Biobank. *The Journals of Gerontology: Series A*, 2020, 75(9): 1801–1803. <https://doi.org/10.1093/gerona/glaa169>
- [7] KIRCHEIS R., HAASBACH E., LUEFTENEGGER D., and HEYKEN W. T. NF- κ B Pathway as a Potential Target for Treatment of Critical Stage COVID-19 Patients. *Frontiers in Immunology*, 2020, 11: 1–11. <https://dx.doi.org/10.3389%2Ffimmu.2020.598444>
- [8] GIRAUDO C., CAVALIERE A., FICHERA G., WEBER M., MOTTA R., PELLOSO M., TOSATO F., LUPI A., CALABRESE F., CARRETTA G., CATTELAN A. M., DE CONTI G., CIANCI V., NAVALESI P., PLEBANI M., REA F., VETTOR R., VIANELLO A., and STRAMARE R. Validation of a Composed COVID-19 Chest Radiography Score : The CARE Project. *European Respiratory Society Journal Open Research*, 2020, 6(4): 1–9. <http://dx.doi.org/10.1183/23120541.00359-2020>
- [9] GOLDSTEIN M. R., POLAND G. A., and GRAEBER C. W. Does Apolipoprotein E Genotype Predict COVID-19 Severity? *Quarterly Journal of Medicine*, 2020, 113(8): 529–530. <https://doi.org/10.1093/qjmed/hcaa142>
- [10] DEL VALLE D. M., KIM-SCHULZE S., HUANG H., BECKMANN N. D., NIRENBERG S., WANG B., LAVIN Y., SWARTZ T. H., MADDURI D., STOCK A., MARRON T. U., XIE H., PATEL M., TUBALLES K., VAN OEKELEN O., RAHMAN A., KOVATCH P., ABERG J. A., SCHADT E., JAGANNATH S., MAZUMDAR M., CHARNEY A. W., FIRPO-BETANCOURT A., RAO MENDU D., JHANG J., REICH D., SIGEL K., CORDON-CARDO C., FELDMANN M., PAREKH S., MERAD M., and GNJATIC S. An Inflammatory Cytokine Signature Predict COVID-19 Severity and Survival. *Nature Medicine*, 2020, 26: 1636–1643. <http://dx.doi.org/10.1038/s41591-020-1051-9>
- [11] KANG S. Age-Related Morbidity and Mortality among Patients with COVID-19. *Infection & Chemotherapy*, 2020, 52(2): 154–164. <https://dx.doi.org/10.3947%2Fic.2020.52.2.154>
- [12] HU X., XING Y., JIA J., NI W., LIANG J., ZHAO D., SONG X., GAO R., and JIANG F. Factors Associated with Negative Conversion of Viral RNA in Patients Hospitalized with COVID-19. *Scien Total Environment*, 2020, 728(175). <https://doi.org/10.1016/j.scitotenv.2020.138812>
- [13] ZHOU F., YU T., DU R., FANG G., LIU Y., LIU Z., XIANG J., WANG Y., SONG B., GU X., GUAN L., WEI Y., LI H., WU X., XU J., TU S., ZHANG Y., CHEN H., and CAO B. Clinical Course and Risk Factors for Mortality of Adult Inpatients with COVID-19 in Wuhan, China : A Retrospective Cohort Study. *The Lancet*, 2020, 395: 1054–1062. [http://dx.doi.org/10.1016/S0140-6736\(20\)30566-3](http://dx.doi.org/10.1016/S0140-6736(20)30566-3)
- [14] BAHL A., JOHNSON S., and WEI N. Timing of Corticosteroids Impacts Mortality in Hospitalized COVID-19 Patients. *Internal and Emergency Medicine*, 2021. <https://doi.org/10.1007/s11739-021-02655-6>
- [15] GIBSON P. G., QIN L., and PUAH S. H. COVID-19 Acute Respiratory Distress Syndrome (ARDS): Clinical Features and Differences from Typical Pre-COVID-19 ARDS. *Medical Journal of Australia*, 2020, 213(2): 54–56. <https://doi.org/10.5694/mja2.50674>
- [16] NATIONAL INSTITUTES OF HEALTH. *Coronavirus Disease 2019 (COVID-19) Treatment Guidelines*. National Institutes of Health, 2019. <https://www.covid19treatmentguidelines.nih.gov/>
- [17] EJAZ H., ALSRHANI A., ZAFAR A., JAVED H., JUNAID K., ABDALLA A. E., ABOSALIF K. O. A., AHMED Z., and YOUNAS S. COVID-19 and Comorbidities : Deleterious Impact on Infected Patients. *Journal of Infection and Public Health*, 2020, 13(12): 1833–1839. <https://doi.org/10.1016/j.jiph.2020.07.014>

参考文:

- [1] YUEN K. S., YE Z. W., FUNG S. Y., CHAN C. P., 和 JIN D. Y. 研究重点非典-冠状病毒-2 和新冠肺炎 : 最重要的研究问题。细胞与生物科学, 2020: 1–5. <https://doi.org/10.1186/s13578-020-00404-4>
- [2] WU C., CHEN X., CAI Y., XIA J., ZHOU X., XU S., HUANG H., ZHANG L., ZHOU X., DU C., ZHANG Y., SONG J., WANG S., CHAO Y., YANG Z., XU J., ZHOU X., CHEN D., XIONG W., XU L., ZHOU F., JIANG J., BAI C., ZHENG J., 和 SONG Y. 与中国武汉 2019 冠状病毒病肺炎患者急性呼吸窘迫综合征和死亡相关的危险因素。美国医学会内科杂志, 2020, 180(7): 934–943. <https://doi.org/10.1001/jamainternmed.2020.0994>
- [3] KUMAR D., JAHAN S., KHAN A., SIDDIQUI A. J., 和 REDHU N. S. 非典-冠状病毒-2 诱导炎症的神经学表现和针对冠状病毒病-19 的可能治疗策略。分子神经生物学, 2021, 14: 1–18. <https://doi.org/10.1007/s12035-021-02318-9>
- [4] FINCH C. E., 和 KULMINSKI A. M. 给编辑的信载脂蛋白乙位点和新冠肺炎 : 我们要去往过的地方吗 ? 老年学杂志 : 一种系列, 2021, 76(2): e1–e3. <https://doi.org/10.1093/gerona/glaa200>
- [5] KUO C., PILLING L. C., ATKINS J. L., MASOLI J. A. H., DELGADO J., KUCHEL G. A., 和 MELZER D. 载脂蛋白乙厄普西隆4基因型预测英国生物银行社区队列中的严重新冠肺炎。老年学杂志 : 医学科学, 2020, 75(11): 2231–2232. <https://doi.org/10.1093/gerona/glaa131>
- [6] KUO C., PILLING L. C., ATKINS J. L., MASOLI J. A. H., DELGADO J., KUCHEL G. A., 和 MELZER D. 给编辑的信载脂蛋白乙厄普西隆4厄普西隆4基因型和死亡率与英国生物银行的新冠肺炎。老年学杂志 : 一种系列, 2020, 75(9): 1801–1803. <https://doi.org/10.1093/gerona/glaa169>
- [7] KIRCHEIS R., HAASBACH E., LUEFTENEGGER D., 和 HEYKEN W. T. 活化乙细胞通路的核因子河童-轻链增强子作为治疗危重期新冠肺炎患者的潜在靶点。

- 免疫学前沿, 2020, 11: 1–11.
<https://dx.doi.org/10.3389%2Ffimmu.2020.598444>
- [8] GIRAUDO C., CAVALIERE A., FICHERA G., WEBER M., MOTTA R., PELLOSO M., TOSATO F., LUPI A., CALABRESE F., CARRETTA G., CATTELAN A. M., DE CONTI G., CIANCI V., NAVALESI P., PLEBANI M., REA F., VETTOR R., VIANELLO A., 和 STRAMARE R.
 合成的新冠肺炎胸部放射学评分的验证：心脏麻醉风险评估项目。欧洲呼吸学会期刊开放研究, 2020, 6(4): 1–9.
<http://dx.doi.org/10.1183/23120541.00359-2020>
- [9] GOLDSTEIN M. R., POLAND G. A., 和 GRAEBER C. W.
 载脂蛋白乙基因型能预测新冠肺炎的严重程度吗？医学季刊, 2020, 113(8): 529–530.
<https://doi.org/10.1093/qjmed/hcaa142>
- [10] DEL VALLE D. M., KIM-SCHULZE S., HUANG H., BECKMANN N. D., NIRENBERG S., WANG B., LAVIN Y., SWARTZ T. H., MADDURI D., STOCK A., MARRON T. U., XIE H., PATEL M., TUBALLES K., VAN OEKELEN O., RAHMAN A., KOVATCH P., ABERG J. A., SCHADT E., JAGANNATH S., MAZUMDAR M., CHARNEY A. W., FIRPO-BETANCOURT A., RAO MENDU D., JHANG J., REICH D., SIGEL K., CORDON-CARDO C., FELDMANN M., PAREKH S., MERAD M., 和 GNJATIC S.
 炎症细胞因子特征预测新冠肺炎的严重程度和存活率。
 自然医学, 2020, 26: 1636–1643.
<http://dx.doi.org/10.1038/s41591-020-1051-9>
- [11] KANG S.
 新冠肺炎患者的年龄相关发病率和死亡率。感染与化疗, 2020, 52(2): 154–164.
<https://dx.doi.org/10.3947%2Fic.2020.52.2.154>
- [12] HU X, XING Y, JIA J, NI W, LIANG J, ZHAO D, SONG X, GAO R, 和 JIANG F. 与新冠肺炎住院患者病毒核糖核酸负转化相关的因素。科学全面环境, 2020, 728(175).
<https://doi.org/10.1016/j.scitotenv.2020.138812>
- [13] ZHOU F, YU T, DU R, FAN G, LIU Y, LIU Z, XIANG J, WANG Y, SONG B, GU X, GUAN L, WEI Y, LI H, WU X, XU J, TU S, ZHANG Y, CHEN H, 和 CAO B. 中国武汉成人新冠肺炎住院患者死亡率的临床过程和危险因素：一项回顾性队列研究。柳叶刀, 2020, 395: 1054–1062. [http://dx.doi.org/10.1016/S0140-6736\(20\)30566-3](http://dx.doi.org/10.1016/S0140-6736(20)30566-3)
- [14] BAHL A., JOHNSON S., 和 WEI N. 皮质类固醇的时间影响住院新冠肺炎患者的死亡率。内科和急诊科, 2021. <https://doi.org/10.1007/s11739-021-02655-6>
- [15] GIBSON P. G., QIN L, 和 PUAH S. H. 新冠肺炎急性呼吸窘迫综合征：临床特征和与典型的新新冠肺炎前急性呼吸窘迫综合征的区别。澳大利亚医学杂志, 2020, 213(2): 54–56. <https://doi.org/10.5694/mja2.50674>
- [16] 国立卫生研究院. 2019 年冠状病毒病（新冠肺炎）治疗指南。美国国立卫生研究院, 2019.
<https://www.covid19treatmentguidelines.nih.gov/>
- [17] EJAZ H, ALSRHANI A, ZAFAR A, JAVED H, JUNAID K, ABDALLA A. E, ABOSALIF K. O. A, AHMED Z, 和 YOUNAS S. 新冠肺炎和合并症：对感染患者的有害影响。感染与公共卫生杂志, 2020, 13(12): 1833–1839.
<https://doi.org/10.1016/j.jiph.2020.07.014>