

Acute kidney injury after a coronary artery bypass graft surgery

by Kun Arifi Abbas

Submission date: 03-Jun-2022 09:02PM (UTC+0800)

Submission ID: 1849765902

File name: e_kidney_injury_after_a_coronary_artery_bypass_graft_surgery.pdf (1.08M)

Word count: 3131

Character count: 16238

Case Report

Acute Kidney Injury after a Coronary Artery Bypass Graft Surgery

Kun Arifi Abbas, Nancy Margarita Rehatta, Yan Efrata Sembiring¹, Ghuraba Adisurya, Prananda Surya Airlangga, Hamzah Hamzah, Hardiono Hardiono, Bambang Pujo Semedi, Christijogo Sumartono Waluejo

Department of Anesthesiology and Reanimation, Faculty of Medicine, Airlangga University - Dr. Soetomo General Hospital, ¹Department of Cardiovascular Thoracic Surgery, Faculty of Medicine, Airlangga University - RSUD Dr. Soetomo General Hospital Surabaya

Abstract

Coronary artery bypass graft (CABG) is performed on patients with blocked coronary arteries by creating a shortcut to resolve and improve blood flow to the heart muscle. The CABG procedure can be performed using a cardiopulmonary bypass (CPB) or an off-pump artery bypass graft technique. One complication in heart surgery is acute kidney injury (AKI) incidence. The risk factors in the incidence of AKI include CPB, complications from surgery (e.g., bleeding and blood transfusions), type of surgery, preoperative high-osmolality state, preoperative serum creatinine level, age >60 years, the presence of comorbid conditions (e.g., hypertension, diabetes, and dyslipidemia), impaired left ventricular function, hemodynamic disturbances, hypoperfusion, use of vasopressor drugs/inotropes, and preoperative hypoalbuminemia. We report an observation to six of our recent patients underwent CABG aged 36–69 years old. Five patients with coronary arterial diseases (CAD) underwent CABG, and one patient with CAD and mitral regurgitation underwent CABG surgery and mitral valve replacement. All patients had a preoperative albumin level of >3.5 g/dL. Two patients developed AKI <24 h postsurgery and three followed by the next day. Serum creatinine monitoring was carried out daily for 7 days, where two patients showed improvement in serum creatinine, and other two had their creatinine returned to normal. This case series found that AKI incidence after CABG was relatively high despite an excellent clinical outcome. Further research is needed with a larger number of cases to find the risk factors for AKI for perioperative prevention.

Keywords: Comorbid, complication, hemodynamic, hypoperfusion, vasopressor

INTRODUCTION

Coronary artery bypass graft (CABG) is performed on patients with blocked coronary arteries by creating a bypass to clear the blockage and improve blood flow to the heart muscle. This procedure uses the patient's autologous blood vessels (veins or arteries) as grafts to bypass the coronary arteries that are partially or wholly blocked due to the atherosclerotic plaque.^[1]

All CABG procedures performed in our institution, the Dr. Soetomo General Hospital, currently use the on-pump method. In 2018, our hospital had 82 cases of CABG procedures.^[1,2] The cardiopulmonary bypass (CPB) machine temporarily replaces the heart–lung function to provide oxygen, eliminates carbon dioxide in the blood during the surgical procedure, and creates a “bloodless” surgical field.^[3,4]

Acute kidney injury (AKI) is a syndrome of rapid abrupt in kidney function within hours to days and is often associated

with patients suffering from other acute diseases and appears in critically ill patients.^[5] The incidence of AKI is higher in heart surgery using CPB. The incidence of AKI in heart surgery ranges from 5% to 42% and even up to 70% depending on the type of surgery and the definition of AKI.^[5–8] Postoperative AKI can lead to prolonged stay, increased complication, increased mortality, and treatment costs.^[5,7,9,10]

CASE REPORT

Six patients underwent CABG aged 36–69 years old were observed. Five patients with coronary arterial

Address for correspondence: Dr. Kun Arifi Abbas, Jalan Mayjend. Prof. Dr. Moestopo no. 6-8, Surabaya, Indonesia. E-mail: kunarifi@gmail.com

Submitted: 24-Nov-2020

Revised: 26-Feb-2021

Accepted: 01-Mar-2021

Published: 09-Aug-2021

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Abbas KA, Rehatta NM, Sembiring YE, Adisurya G, Airlangga PS, Hamzah H, et al. Acute kidney injury after a coronary artery bypass graft surgery. *Bali J Anaesthesiol* 2021;5:204-7.

Access this article online

Quick Response Code:



Website:
www.bjoaonline.com

DOI:
10.4103/bjoa.bjoa_249_20

diseases (CAD) underwent CABG only, and one patient with CAD and mitral regurgitation underwent CABG surgery and mitral valve replacement. The comorbidities found in the patients were hypertension, diabetes mellitus, dyslipidemia, and no previous history of AKI. Two patients received diuretic therapy, and two patients received anticoagulant therapy. The lowest ejection fraction was 51%, and the highest was 76%. Chest X-ray examination showed no pulmonary edema, two patients presented with cardiomegaly [Figure 1].

All patients had a preoperative albumin level of >3.5 g/dL. The lowest preoperative hemoglobin was 9.8 g/dL, and the highest was 16.2 g/dL. An increase in serum creatinine was found in one patient, 1.6, with a BCR ratio of 20.6. Hypernatremia (150 mEq/L) was found in one patient. Blood sugar levels were within the normal limits.

The surgery duration lasts 6–7 h [Table 1], but one patient had an extended duration (10 h). To that particular patient, due to heart rhythm disturbances (atrial fibrillation) during surgery, the sternal closure is delayed until the next day.

Laboratory tests (hemoglobin, blood urea nitrogen, serum creatinine, and blood gas analysis) were performed after

anesthesia induction, during the CPB procedure, and after CPB. Albumin and random blood sugar test were performed after the first cardioplegia during CPB procedures [Table 2].

There was a decrease in albumin during CPB. The average value of albumin levels during CPB was 2.46 g/dL (2.2–2.6 g/dL). Only one patient experienced a decrease in hemoglobin levels <8.0 g/dL, so a 200 ml packed red cell (PRC) transfusion was performed. Hemofiltration of 1500–2000 mL ultrafiltration is performed during the surgery. Urine output during surgery was >1 ml/kg/hour. Serum creatinine levels during the CPB procedure tended to decrease in five patients but increased in one patient.

Postoperative care is carried out in the intensive care unit (ICU) with a length of stay of 3–5 days. Duration of the patients who use the ventilator until extubation is 1–2 days. The average length of treatment days postoperatively until discharge was 6.6 days (5–11 days). During ICU observation, five out of six patients required vasoactive/inotropic drugs to maintain hemodynamics.

Bleeding occurred at 60–700 ml drainage production on the 1st day, improved on the 2nd day, and stopped on the 3rd day. Replacement of blood volume loss using crystalloid,




Figure 1: Photo image of chest X-ray (CTR: Cardio Thorax Ratio)

Table 1: Surgery, cardiopulmonary bypass, and aortic cross-clamp duration

Patient	Diagnosis	Surgery	Surgery duration (min)	CPB duration (min)	Aorta cross-clamp duration (min)
1	CAD-3VD	CABG	395	116	71
2	CAD-3VD	CABG	365	128	67
3	CAD-2VD	CABG	440	68	51
4	CAD-3VD	CABG	390	129	75
5	CAD-3VD	CABG	410	106	70
6	CAD-3VD+MR	CABG+mitral replacement	600	286	151

CAD: Coronary arterial disease, 2VD: Two-vessels disease, 3VD: Three-vessels disease, CABG: Coronary artery bypass graft, CPB: Cardiopulmonary bypass, MR: Mitral regurgitation

Table 2: Investigation results

Patient	Sodium	BUN	SC	BUN to SC ratio	RBS	Osmolarity	pH	pCO2	BE
1	133	19	1	19	97	269	7.5	28	-1
2	139	11	1	11	154	280	7.48	30.6	0
3	133	8	0.8	10	74	263	7.39	37.7	-2
4	135	11	0.9	12.2	117	271	7.45	30.7	-2
5	140	14	1.3	10.8	123	281	7.54	31.4	5
6	138	15	1.2	12.5	99	277	7.43	31.3	-3

BUN: Blood urea nitrogen, SC: Serum creatinine, BS: Blood sugar, pCO2: Partial pressure of CO2, BE: Base excess, RBS: Random Blood Sugar

colloid, PRC, and fresh-frozen plasma. During ICU stay, hemoglobin levels were maintained >9 g/dL by blood transfusions. The level of albumin shortly after surgery (day 0) indicated a value of >3 g/dL. There was a decrease in base excess during ICU observation (day 0) in all patients compared to surgery.

Postoperative renal function was assessed by the level of urine production and changes in serum creatinine values after surgery. During postoperative care in the ICU, five out of six patients produced urine >0.5 ml/kg/hour, but there was one patient with urine output <0.5 ml/kg/hour that lasted 9 h.

Two patients developed AKI on day 0 and were followed by three other patients on day 1. Only one patient did not develop AKI at all. According to the kidney disease, Improving Global Outcomes (KDIGO), three patients had Stage-1 AKI, and two patients had Stage 2. Serum creatinine was monitored for 7 days, where two patients showed improvement in serum creatinine, and two other patients were fully recovered.

DISCUSSION

This case series report observed six patients with a length of stay in the ICU was 3–5 days, the length of stay on the ventilator was 2 days, and the length of hospital stay was 5–11 days. In general, the patient's clinical condition and postoperative outcome were satisfactory.

The diagnosis of AKI can generally be enforced with increased serum creatinine and decreased urine production. One of the criteria commonly used to assess AKI is KDIGO.^[11,12] Possible misdiagnosis of AKI can occur in the measurement of serum creatinine and urine production. In most CPB-utilized CABG cases, the blood creatinine levels may be decreased due to excess perioperative fluid (crystalloid, colloid, and transfusion) and hemodilution. Serum creatinine is a reliable test in diagnosing impaired kidney function because the body can still maintain urine production in the normal range even though the glomerular filtration rate has decreased.^[12]

Three patients were over 60 years of age and all developed AKI. With age, there is a decrease in the quality of kidney function so that the potential for kidney damage increases. This is due to the kidneys' structure and abilities that are worn out with age and are exacerbated by comorbidities.^[13] The BMI of the patients was still normal, so it is not a risk factor for AKI.^[14,15] Hypertension was found in four

patients, diabetes in two patients, and dyslipidemia in one. Hypertension, diabetes mellitus, and dyslipidemia are the risk factors for AKI.^[10,12]

The use of CPB is related to the incidence of AKI postcardiac surgery. The longer the CPB procedure will increase the probability of AKI. AKI occurs due to inflammation, reperfusion injury, pulsatile waves, hemolysis, micro embolism, and decreased cardiac output.^[5,16] Heart valve surgery is also a risk factor for AKI.^[17]

The target hemoglobin is always maintained by giving blood transfusions to avoid a decrease in oxygen supply.^[18] Postoperatively, there is an increase in albumin levels, possibly due to hemofiltration and rearrangement of fluid homeostasis. Hypoalbumin often occurs during the perioperative period and is associated with AKI events.^[19,20] An albumin level of <2.4 g/dL is associated with increased risk of AKI during hospitalization.^[20] Preoperative hypoalbumin is often associated with nutritional status and the possibility of edema.

Studies on hypoalbuminemia and perioperative AKI in cardiac surgery show different results. A serum albumin of <3 g/dL during CPB is less correlated with postoperative AKI, but the administration of perioperative albumin transfusion from cardiac surgery is associated with the occurrence of AKI.^[21,22] Administration of albumin during CPB priming does not provide any clinical benefit to patients but can prevent the decrease in platelet counts during CPB.^[23-25] In OPCAB surgery, preoperative administration of albumin to patients with albumin levels <4 g/dL reduces the risk of AKI and increases urine output during surgery but does not benefit patients who are converted to CPB.^[26]

CONCLUSION

This case series found that AKI incidence after CABG was relatively high despite an excellent clinical outcome. Most patients experience AKI only temporarily and recover from AKI. Further research is needed with a larger number of cases to find risk factors for AKI for perioperative prevention.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form, the patients have given their consent for their images and other clinical information

2

to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Alexander JH, Smith PK. Coronary-artery bypass grafting. *N Engl J Med* 2016;374:1954-64.
- Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, *et al.* Heart disease and stroke statistics-2019 update: A report from the American Heart Association. *Circulation* 2019;139:e56-528.
- Sarkar M, Prabh V. Basics of cardiopulmonary bypass. *Indian J Anesth* 2017;61:760-7.
- Hessel EA, Shann KG. Blood pumps, circuitry and cannulation techniques in cardiopulmonary bypass. In: Gravlee GP, Davis RF, Hammon J, eds. *Cardiopulmonary Bypass and Mechanical Support: Principles and Practice*. 4th ed. Philadelphia 2015.
- Nadim M, Forni L, Bihorac A, Hobson C, Koynier J, Shaw A, *et al.* Cardiac and Vascular Surgery-Associated Acute Kidney Injury: The 20th International Consensus Conference of the ADQI (Acute Disease Quality Initiative) Group. *Journal of the American Heart Association*. 2018;7(11).
- de Geus HR, Ronco C, Haase M, Jacob L, Lewington A, Vincent JL. The cardiac surgery-associated neutrophil gelatinase-associated lipocalin (CSA-NGAL) score: A potential tool to monitor acute tubular damage. *J Thorac Cardiovasc Surg* 2016;151:1476-81.
- Elmistekawy E, McDonald B, Hudson C, Ruel M, Mesana T, Chan V, *et al.* Clinical impact of mild acute kidney injury after cardiac surgery. *Ann Thorac Surg* 2014;98:815-22.
- Wu B, Chen J, Yang Y. Biomarkers of acute kidney injury after cardiac surgery: A narrative review. *Biomed Res Int* 2019;2019:1-11.
- Ostermann M, Joannidis M. Acute kidney injury 2016: Diagnosis and diagnostic workup. *Crit Care* 2016;20:299.
- Kumar AB, Suneja M. Cardiopulmonary bypass-associated acute kidney injury. *Anesthesiology* 2011;114:964-70.
- Kellum JA, Lameire N, Aspelin P, Barsoum R, Burdman E, Goldstein S, *et al.* Kidney Disease: Improving Global Outcomes (KDIGO) New Guideline for Diabetes Management in Chronic Kidney Disease: A Meet-the-Expert Session. *EMJ Diabetes*. 2020. Available from: <http://www.kidney-international.org>. [Last accessed on 2021 Jan 25].
- O'Neal JB, Shaw AD, Billings FT 4th. Acute kidney injury following cardiac surgery: Current understanding and future directions. *Crit Care* 2016;20:187.
- Wei Q, Liu H, Tu Y, Tung RN, Wang YL, Pan MM, *et al.* The characteristics and mortality risk factors for acute kidney injury in different age groups in China—A cross sectional study. *Renal Failure*. 2016;38:1413-7.
- WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2014;363:157-63.
- Liu AL, Wang J, Nikam M, Lai BC, Yeoh LY. Low, rather than High, Body Mass Index is a Risk Factor for Acute Kidney Injury in Multiethnic Asia Patients: A Retrospective Observational Study. *In J Nephrol*. 2018:1-17. DOI: 10.1155/2018/3284612.
- Xu S, Liu J, Li L, Wu Z, Li J, Liu Y, *et al.* Cardiopulmonary bypass time is an independent risk factor for acute kidney injury in emergent thoracic aortic surgery: A retrospective cohort study. *J Cardiothorac Surg* 2019;14:90.
- Yamauchi T, Miyagawa S, Yoshikawa Y, Toda K, Sawa Y, Osaka Cardiovascular Surgery Research (OSCAR) Group. Risk index for postoperative acute kidney injury after valvular surgery using cardiopulmonary bypass. *Ann Thorac Surg* 2017;104:868-75.
- Kementrian Kesehatan RI. Ministry of Health of the Republic of Indonesia. Decree of the Minister of Health of the Republic of Indonesia no HK.01.07/MENKES/659/2017 concerning the National Formulary. 2017. Jakarta: Ministry of Health RI. Jakarta: Kemenkes RI; 2017.
- Wiedermann CJ, Wiedermann W, Joannidis M. Causal relationship between hypoalbuminemia and acute kidney injury. *World J Nephrol* 2017;6:176-87.
- Thongprayoon C, Cheungpasitporn W, Mao MA, Sakhuja A, Kashani K. U-shape association of serum albumin level and acute kidney injury risk in hospitalized patients. *PLoS One* 2018;13:e0199153.
- Silva Ja V, Barra CR G, De Oca-sandoval ma Espinosa Blanco o M, Bazaldúa s F, Kc Casillas-Ramirez A T, Isaac F. [Internet]. *Caccv.org.ar*. 2021. Available from: http://www.caccv.org.ar/raccv-es-2014/RACCV_XII_nro2_2014_Art_pag21_ESP.pdf. [Last cited on 31 March 2021].
- Frenette AJ, Bouchard J, Bemier P, Charbonneau A, Nguyen LT, Rioux JP, *et al.* Albumin administration is associated with acute kidney injury in cardiac surgery: A propensity score analysis. *Crit Care* 2014;18:602.
- Russell JA, Navickis RJ, Wilkes MM. Albumin versus crystalloid for pump priming in cardiac surgery: Meta-analysis of controlled trials. *J Cardiothorac Vasc Anesth* 2004;18:429-37.
- Weiler P, Hamiko M, Mellert F, Roell W, Roell M, Welz C, *et al.* Impact of crystalloid or albumin priming of the heart-lung machine on in-hospital outcome after coronary artery bypass surgery. *Thorac Cardiovasc Surg* 2019;67:475-83.
- Tanaka A, Ota T, Uriel N, Asfaw Z, Onsager D, Lonchyna VA, *et al.* Cardiovascular surgery in Jehovah's Witness patients: The role of preoperative optimization. *J Thorac Cardiovasc Surg* 2015;150:976-830.
- Lee EH, Kim WJ, Kim JY, Chin JH, Choi DK, Sim JY, *et al.* Effect of exogenous albumin on the incidence of postoperative acute kidney injury in patients undergoing off-pump coronary artery bypass surgery with a preoperative albumin level of less than 4.0 g/dl. *Anesthesiology* 2016;124:1001-11.

Acute kidney injury after a coronary artery bypass graft surgery

ORIGINALITY REPORT

19%

SIMILARITY INDEX

16%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Mahasti Amoui, Reyhane Ahmadi, Mohsen Qutbi, Isa Neshandar Asli. "Somatostatin-receptor avidity of pancreatic neuroendocrine tumor thrombus in porto-caval venous systems on 99mTc-Octreotide and posttherapeutic 177Lu-DOTA-TATE scans", *World Journal of Nuclear Medicine*, 2022
Publication **3%**
- 2** www.thejhs.org
Internet Source **2%**
- 3** ijdvl.com
Internet Source **1%**
- 4** thejns.org
Internet Source **1%**
- 5** Mitra K. Nadim, Lui G. Forni, Azra Bihorac, Charles Hobson et al. "Cardiac and Vascular Surgery–Associated Acute Kidney Injury: The 20th International Consensus Conference of the ADQI (Acute Disease Quality Initiative) **1%**

Group", Journal of the American Heart Association, 2018

Publication

6	journals.lww.com Internet Source	1 %
7	worldwidescience.org Internet Source	1 %
8	academic.oup.com Internet Source	1 %
9	eprints.kaums.ac.ir Internet Source	1 %
10	Page, Karen, McKinney, Aidin. "EBOOK: Nursing the Acutely ill Adult: Case Book", EBOOK: Nursing the Acutely ill Adult: Case Book, 2012 Publication	1 %
11	www.wjgnet.com Internet Source	1 %
12	pulmo-ua.com Internet Source	1 %
13	link.springer.com Internet Source	<1 %
14	L. Abboud. "The slope of the end-systolic pressure-volume relationship compared with the global end-systolic pressure-volume ratio in humans", Clinical Cardiology, 01/1990	<1 %

15

gutpathogens.biomedcentral.com

Internet Source

<1 %

16

researchonline.jcu.edu.au

Internet Source

<1 %

17

Antonio Artigas, Jean Carlet, Ignacio Martin-Loeches, Michael Niederman, Antoni Torres. "23rd International Symposium on Infections in the Critically Ill Patient", Medical Sciences, 2018

Publication

<1 %

18

Kevin R. Regner. "Epidemiology of acute kidney injury after cardiac surgery: an update", Journal of Organ Dysfunction, 2009

Publication

<1 %

19

M. Ostermann, K. Weerapolchai, N. Lumlertgul. "Chapter 18 Prevention of Acute Kidney Injury After Cardiac Surgery", Springer Science and Business Media LLC, 2022

Publication

<1 %

20

O. Goren, I. Matot. "Perioperative acute kidney injury", British Journal of Anaesthesia, 2015

Publication

<1 %

21

Xiang Yu, Zhe Feng. "Analysis of Risk Factors for Perioperative Acute Kidney Injury and

<1 %

Management Strategies", Frontiers in Medicine, 2021

Publication

22

bmccresnotes.biomedcentral.com

Internet Source

<1 %

23

ccforum.biomedcentral.com

Internet Source

<1 %

24

doi.org

Internet Source

<1 %

25

www.science.gov

Internet Source

<1 %

26

Kyungmi Kim, Ji-Yeon Bang, Seon-Ok Kim, Saegyeol Kim, Joung Uk Kim, Jun-Gol Song. "Association of preoperative hypoalbuminemia with postoperative acute kidney injury in patients undergoing brain tumor surgery: a retrospective study", Journal of Neurosurgery, 2018

Publication

<1 %

27

Rezan Aksoy, Taylan Adademir, Ekrem Yilmaz, Deniz Cevirme, Mehmet Sengor, Cengiz Koksall, Murat Bulent Rabus. "Is Hypoalbuminemia a Predictor for Acute Kidney Injury after Coronary Bypass Grafting in Diabetes Mellitus Patients?", Brazilian Journal of Cardiovascular Surgery, 2019

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Acute kidney injury after a coronary artery bypass graft surgery

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
