



# Source details

## Cancer Control

Open Access ⓘ

Scopus coverage years: from 1995 to Present

Publisher: SAGE

ISSN: 1073-2748 E-ISSN: 1526-2359

Subject area: Medicine: Hematology Medicine: Oncology

Source type: Journal

CiteScore 2021

3.2 ⓘ

SJR 2021

0.586 ⓘ

SNIP 2021

0.972 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### i Improved CiteScore methodology ⓘ

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▾

$$3.2 = \frac{1,148 \text{ Citations } 2018 - 2021}{355 \text{ Documents } 2018 - 2021}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$3.0 = \frac{1,454 \text{ Citations to date}}{487 \text{ Documents to date}}$$

Last updated on 05 February, 2023 • Updated monthly

### CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine		
↳ Hematology	#69/126	45th
Medicine		
↳ Oncology	#215/360	40th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

---

## About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

## Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

## Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

---

## ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.





## Scientific Programming Papers

Open



## Cancer Control

### COUNTRY

United States

- Universities and research institutions in United States
- Media Ranking in United States

### SUBJECT AREA AND CATEGORY

- Medicine
  - Hematology
  - Medicine (miscellaneous)
  - Oncology



### PUBLISHER

H. Lee Moffitt Cancer Center and Research Institute

### H-INDEX

74

### PUBLICATION TYPE

Journals

### ISSN


10732748, 15262359

### COVERAGE

1995-2021

### INFORMATION

- [Homepage](#)
- [How to publish in this journal](#)
- [http://Jennifer.Lovick@sagepub.com/](mailto:Jennifer.Lovick@sagepub.com)



### SCOPE

Cancer Control is a JCR-ranked open access journal whose mission is to advance the prevention, detection, diagnosis, treatment, and palliative care of cancer by enabling researchers, doctors, policymakers, and other healthcare professionals to freely share research along the cancer control continuum. Our vision is a world where gold-standard cancer care is the norm, not the exception. The journal is organized into five topical sections, as well as a sixth curated by members of the Moffitt Cancer Center. The topical sections are inspired by the foci of the cancer control continuum (prevention, detection, diagnosis, treatment, survivorship). Article Types and Topics of Interest Cancer Control welcomes original research (both basic and clinical studies), reviews, and commentaries. We encourage submissions relating to advances in the following areas: Communications in cancer research Surveillance Social Determinants of Health Disparities Genetic Testing Decision-Making Dissemination of Evidence-Based Interventions Health Care Delivery and Quality of Care Epidemiology Measurement.

Join the conversation about this journal

## Complexity OA & Peer Reviewed







Open

Hindawi

FIND SIMILAR JOURNALS

options

<p>1 <b>BMC Cancer</b> GBR <b>74%</b> similarity</p>	<p>2 <b>ecancermedicalseience</b> GBR <b>74%</b> similarity</p>	<p>3 <b>Cancer Medicine</b> GBR <b>74%</b> similarity</p>	<p>4 <b>Wspolczesna Onkologia</b> POL <b>72%</b> similarity</p>	<p>5 <b>Voprosy Onkologii</b> RUS <b>71%</b> similarity</p>
--	---	---	---	---



Cancer Control

**Q2** Medicine (miscellaneous) best quartile

**SJR 2021** 0.59

powered by scimagojr.com


Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" >
```

SCImago Graphica


Explore, visually communicate and make sense of data with our **new data visualization tool.**



**Leave a comment**

Name

Email   
(will not be published)



**Submit**

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on [@ScimagoJR](#)

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)

[Edit Cookie Consent](#)

# Cancer Control

Impact Factor: **2.339**

5-Year Impact Factor: **3.671**

All articles

Filter



111-120 of 147

Sort by: Date




Select all

Export selected citations

## Original Research Article

 Open Access | Research article | First published Mar 29, 2022

[Correlation of Epidermal Growth Factor Receptor Mutation With Major Histologic Subtype of Lung Adenocarcinoma According to IASLC/ATS/ERS Classification](#)

Sara Boukansa PhD , Zineb Benbrahim MD, Sanaa Gamrani PhD, Sanae Bardai PhD, [...]


[View all](#) 

[Preview abstract](#) 


PDF / EPUB



## Original Research Article

 Open Access | Research article | First published Mar 26, 2022

## [Biweekly Raltitrexed Combined With Irinotecan as Second-Line Therapy for Patients With Metastatic Colorectal Cancer: A Phase II Trial](#)

Ke Cheng MD, PhD , Yu-Wen Zhou MD, PhD, Ye Chen MD, PhD, Zhi-Ping Li MD, PhD, [...]

[View all](#) 

[Preview abstract](#) 

Original Research Article



Open Access

Research article

First published Mar 26, 2022

PDF / EPUB



## [Clinical Characteristics and Prognosis of Small Cell Carcinoma in the Nasopharynx: A Population-Based Study](#)

Yu-ling Zhou, Ying-peng Peng, Qiao-dan Liu, Xian-zhen Chen, Jianzhong He, Wei Wei, [...]

[View all](#) 

[Preview abstract](#) 

PDF / EPUB



Original Research Article



Open Access

Research article

First published Mar 26, 2022

## [Efficacy of Drug-Eluting Beads Transarterial Chemoembolization Plus Camrelizumab Compared With Conventional Transarterial Chemoembolization Plus Camrelizumab for Unresectable Hepatocellular Carcinoma](#)

Yanqiao Ren, Yusheng Guo, Lei Chen, Tao Sun, Weihua Zhang, Bo Sun, Licheng Zhu, Fu Xiong, Chuansheng Zheng

[Preview abstract](#) 

PDF / EPUB



Research Article






Open Access

Research article

First published Mar 23, 2022

## [Active Immunotherapy for Glioblastoma Treatment: A Systematic Review and Meta-Analysis](#)

Joni Wahyuhadi , Irwan Barlian Immadoel Haq, Muhammad Reza Arifianto, Bagus Sulistyono , [...]  
[View all](#) 

[Preview abstract](#) 

Original Research Article

PDF / EPUB



Open Access

Research article

First published Mar 22, 2022

## [Knowledge, Attitude, and Practice Toward Cervical Cancer Screening Among Women Attending Health Facilities in Central Ethiopia](#)

Tulu Gebisa MPH, Elias Teferi Bala PhD, Berhanu Senbeta Deriba MPH 

[Preview abstract](#) 

PDF / EPUB



Original Research Article





Open Access

Research article

First published Mar 18, 2022

## [Profiling of Protein Adducts of Estrogen Quinones in 5-Year Survivors of Breast Cancer Without Recurrence](#)

Che Lin MD, PhD, Ding-Ru Chen MS, Shou-Jen Kuo MD, Chi-Yen Feng MD, Dar-Ren Chen MD , [...]  
[View all](#) 

[Preview abstract](#) 


PDF / EPUB



Perspective



## [The Role of CCN1 in Esophageal Adenocarcinoma: What We Have Learned From the Lab](#)



Zhiheng Chang MD, Tong Dang MD, Xianmei Meng MD, Jianyuan Chai PhD 

[Preview abstract](#) 

PDF / EPUB 

### Original Research Article

## [Clinicopathological Findings Associated With Cervical Lymph Node Metastasis in Papillary Thyroid Microcarcinoma: A Retrospective Study in China](#)

Yimeng Shi MD, Zheyu Yang MD, Yu Heng MS , Huijun Ju MD, Yu Pan MD, PhD, Yifan Zhang MD, PhD 

[Preview abstract](#) 

PDF / EPUB 

### Original Research Article

## [Health-Related Quality of Life of Omani Colorectal Cancer Survivors](#)

Maryam Al-Shandudi MD, MSc, Mansour Al-Mandhari MBChB, MRCP, FRACP, Moon Fai Chan PhD,

[...]

[View all](#) 

[Browse journal](#)  
[Preview abstract](#) 

All articles

Browse by year

Special collections index

< 1 ... 11 **12** 13 14 15 >

### Journal information

Journal description

 Privacy

Aims and scope

Editorial board

Submission guidelines

Journal indexing and metrics

Reprints

Journal permissions

Recommend to library

Advertising and promotion

## Keep up to date



Facebook



Twitter



LinkedIn



YouTube



RSS feed



Email alerts

[View all options](#)

## Also from SAGE Publishing

### CQ Library

American political resources

### Data Planet

A universe of data

### SAGE Business Cases

Real-world cases at your fingertips

### SAGE Campus

Online skills and methods courses

### SAGE Knowledge

### SAGE Research Methods

 Privacy

The ultimate social science library

The ultimate methods library

SAGE Video

---

Streaming video collections

Technology from SAGE

---

Make learning and research easier

# Cancer Control

Impact Factor: **2.339**

5-Year Impact Factor: **3.671**

## Editorial board

 [Hide all](#)

### Executive Editor

**Applonia Rose, PhD**

SAGE Publishing, Thousand Oaks, CA USA

### Deputy Editor

**Jennifer K. Lovick, PhD**

SAGE Publishing, Thousand Oaks, CA USA

### Managing Editor

**Priyanka Katyal, PhD**

SAGE Publications, New Delhi, India

**Garima Sharma, PhD**

SAGE Publications, New Delhi, India

**Pooja Sharma, MTech**

SAGE Publications, Dehradun, India

### Commissioning Editor

**Nirmala Bhoo Pathy, MD, PhD**

University of Malaya, Kuala Lumpur, Malaysia

Moffitt Section Editor



**Jonathan S. Zager, MD**

Moffitt Cancer Center, USA

Associate Editors - Moffitt Section



**Jhanelle Gray, PhD**

Moffitt Cancer Center, USA

**Arash Naghavi, PhD**

Moffitt Cancer Center, USA

Section Editor - Prevention



**Muhammad N. Aslam, MBBS, MD**

University of Michigan, Ann Arbor, USA

**Muni Balakrishnan Rubens, MD, PhD**

Miami Cancer Institute, USA

Section Editor - Survivorship



**Fernanda Machado Silva-Rodrigues, RN, PhD**

Santa Casa de Sao Paulo School of Medical Sciences (FCMSCSP) - Sao Paulo, Brazil

**Yingkun Xu, PhD**

Chongqing Medical University, China

Section Editor - Treatment



**Pietro Giorgio Calò, MD**

University of Cagliari, Italy

**Brian J. Czerniecki, MD, PhD**

Moffitt Cancer Center, USA

 Privacy

**Han-Yu Deng**

West Chin Hospital, Sichuan University, China

**Hanan Goldberg, MD**

SUNY Upstate Medical University, USA

**Yiqun Han, MD**

Chinese Academy of Medical Sciences and Peking Union Medical College, China

**Yong-cheng Hu, MD, PhD**

Tianjin Hospital, China

**Mohammad Yasin Karami, MD**

Shiraz University of Medical Sciences, Iran

**Xiangyi Kong, MD, PhD**

Chinese Academy of Medical Sciences and Peking Union Medical College, China

**Jinhui Liu, MD, PhD**

The First Affiliated Hospital of Nanjing Medical University, Nanjing, China

**Luigi Marano, MD, PhD**

University of Siena, Siena, Italy

**Jie Mei, MD**

Nanjing Medical University, China

**Carlos Eduardo Paiva, MD, PhD**

Barretos Cancer Hospital, Brazil

**Alessandro Rizzo, MD**

IRCCS Bari Hospital, Italy

**Ulrich Ronellenfitsch, MD**

University Hospital Halle, Halle, Germany

**Samer Salah, MD**

King Hussein Cancer Center, Amman, Jordan

**PengMing Sun, PhD**

Affiliated Hospital of Fujian Medical University, China

**Song Xu, MD, PhD**

Tianjin Medical University General Hospital, China

**Mohamed A. Yassin, MD**

Qatar University, Qatar

Section Editor – Detection



Privacy

**Nashwa El-Khazragy, MD**

Ain Shams University, Cairo, Egypt

**Abhishek Tyagi, PhD**

Wake Forest Baptist School of Medicine, USA

## Section Editor - Diagnosis

**Bernard Kwabi-Addo, PhD**

Howard University College of Medicine, USA

**Shengwen Calvin Li, PhD, M.Phil.,  
FRSM, FRSB**

CHOC Children's Hospital and University of California,  
Irvine, USA

**Haitao Wang, PhD**

Center for Cancer Research, NCI, Bethesda, USA

**Jian-Guo Zhou, MD**

The Second Affiliated Hospital of Zunyi Medical University,  
Zunyi, China

## Editorial Review Board

**Gulzhanat Aimagambetova, MD, PhD**

Nazarbayev University, Kazakhstan

**Massimiliano Bassi, MD**

Sapienza University of Rome, Italy

**Guntag Batihan, MD**

Kars Harakani State Hospital, Turkey

**Maria Rita Bianco, MD**

Magna Graecia University of Catanzaro, Italy

**Jessica Calderon-Mora, DrPh**

Texas Tech University Health Sciences Center El Paso,  
USA

**Deniz Can Güven, MD**

Hacettepe University Cancer Institute, Turkey

**Filippo Carannante, MD, PhD**

Fondazione Policlinico Campus Bio-Medico of Rome, Italy

**Roza Chaireti, MD, PhD**

Karolinska University Hospital, Sweden

**Timothy D Clay, MBBS (Hons) FRACP  
DMedSci**

St John of God Health Care, Australia

**Alejandro Zentella Dehesa, PhD, IIBO-  
UNAM**

National Autonomous University of Mexico, Mexico

**Durgesh K. Dwivedi, PhD**

King George Medical University, India

**Aydin Eresen, PhD**

University of California Irvine, USA

**Amit Gupta, MS, MBA, FRCS, FACS**

All India Institute of Medical Sciences, India

**Deniz Can Guven, MD**

Hacettepe University Cancer Institute, Turkey

**Ameneh Jafari, PhD**

Shahid Beheshti University of Medical Sciences, Tehran,  
Iran

**Ferda Kaleagasioglu, MD**

Istinye University, Turkey

**Sk. Kayum Alam, PhD**

University of Minnesota, USA

**Mohammad Amin Kerachian, MD, PhD,  
RMF**

Mashhad University of Medical Sciences, Iran

**Divya Khanna, MBBS, MD, PDCR**

TATA Memorial Cancer Center

**Arunkumar Krishnan, MBBS**

West Virginia University School of Medicine, USA

**Jew Win Kuan, MBBS, MRCP, PhD**

Universiti Malaysia Sarawak, Malaysia

**Chan-Yen Kuo, PhD**

Taipei Tzuchi Hospital, Taiwan

**Wang-Zhong Li, PhD**

First Affiliated Hospital of Guangzhou Medical University,  
Guangzhou, China

**Enoch Luis, PhD**

Instituto de Fisiología Celular, Mexico

**Ankit Madan, MD**

Medstar Georgetown Cancer Institute, USA



**Kate A. McBride, MPH, Grad Cert, PhD**

Western Sydney University, Australia

**Mona Osman, MD, MPH, MBA, Dipl  
IBLM**

American University of Beirut, Lebanon

**See-Hyoung Park, PhD**

Hongik University, Republic of Korea

**Alexandros Poutoglidis,  
MD,MSc,PhD(c),FEB(ORL-HNS)**

Genera Hospital of Thessaloniki, Greece

**Xiaonan Shao, MD**

The Third Affiliated Hospital of Soochow University,  
China

**Jie Shen, PhD, MS**

Massey Cancer Center, USA

**Micah A. Skeens, PhD, RN, APRN**

Abigail Research Institute at Nationwide Children's  
Hospital, USA

**Sumsum P. Sunny, MDS, MSc**

Mazumdar Shaw Cancer Center, India

**Ovidiu Tatar, MD, MSc**

Lady Davis Institute for Medical Research, Canada

**Clifton P. Thornton, PhD, RN, CPNP**

John Hopkins University, USA

**Georgios Tsoulfas, MD, PhD, FACS**

Aristotle University School of Medicine, Greece

**I-Shiang Tzeng, MBA, PhD**

Taipei Tzuchi Hospital, Taiwan

**Alessandro Uleri, PhD**

Humanitas University, Italy

**Suzanne Vang, PhD**

NYU Grossman School of Medicine, NY, USA

**Che Hsueh Yang, MD**

Tungs' Taichung MetroHarbor Hospital, Taiwan

**Chenlong Yang, MD, PhD**

Peking University Third Hospital, China

**Zhihao Yang, PhD**

Tianjin University, China

Editorial Board



**Matthew Zhang, PhD, MM, MB**

University Hospital, China

**Daniel A. Anaya, MD**

Moffitt Cancer Center, USA

**Aliyah Baluch, MD**

Moffitt Cancer Center, USA

**Idriss M. Bennani-Baiti, PhD,  
FRSM, CRSM**

Cancer Epigenetics Society, Austria

**Dung-Tsa Chen, PhD**

Moffitt Cancer Center, USA

**Hye Sook Chon, MD**

Moffitt Cancer Center, USA

**Michele L. Cote, PhD, MPH**

Wayne State University, USA

**Mahmoud Danaee, PhD**

University of Malaya, Kuala Lumpur, Malaysia

**Jasreman Dhillon, MD**

Moffitt Cancer Center, USA

**Ilana F. Gareen, PhD**

Brown University, USA

**Melissa J. Vilaro, PhD, MPH, CPH**

University of Florida, USA

**Clement K. Gwede, PhD, RN**

Moffitt Cancer Center, USA

**Sarah E. Hoffe, MD**

Moffitt Cancer Center, USA

**Jagdish Khubchandani, MBBS, PhD**

New Mexico State University, USA

**John V. Kiluk, MD**

Moffitt Cancer Center, USA

**Kyuwoong Kim, PhD**

National Cancer Control Institute, National Cancer Center,  
Republic of Korea

<b>Richard D. Kim, MD</b>	Moffitt Cancer Center, USA
<b>Bela Kis, MD, PhD</b>	Moffitt Cancer Center, USA
<b>Rami Komrokji, MD</b>	Moffitt Cancer Center, USA
<b>Marilyn L. Kwan, PhD</b>	Kaiser Permanente Northern California - Division of Research, USA
<b>Conor C. Lynch, PhD</b>	Moffitt Cancer Center, USA
<b>Tsutomu Nishida, MD, PhD</b>	Toyonaka Municipal Hospital, Osaka, Japan
<b>Catherine M. Olsen</b>	QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia
<b>Kristen J. Otto, MD</b>	Moffitt Cancer Center, USA
<b>Michael A. Poch, MD</b>	Moffitt Cancer Center, USA
<b>Elizabeth M. Sagatys, MD</b>	Moffitt Cancer Center, USA
<b>Saïd M. Sebti, PhD</b>	Moffitt Cancer Center, USA
<b>Bijal D. Shah, MD</b>	Moffitt Cancer Center, USA
<b>Lubomir Sokol, MD, PhD</b>	Moffitt Cancer Center, USA
<b>Hatem H. Soliman, MD</b>	Moffitt Cancer Center, USA
<b>Jonathan R. Strosberg, MD</b>	Moffitt Cancer Center, USA
<b>Jennifer Swank, PharmD</b>	Moffitt Cancer Center, USA
<b>Sarah W. Thirlwell, MSc, MSc(A), RN</b>	Moffitt Cancer Center, USA
<b>Jasmine A. Tiro, PhD</b>	University of Texas Southwestern Medical Center, USA

**Eric M. Toloza, MD, PhD**

Moffitt Cancer Center, USA

**Nam D. Tran, MD**

Moffitt Cancer Center, USA

**Yi Zhang, MD, PhD**

The First Affiliated Hospital of Zhengzhou University, China

## Browse journal

---

All articles

Browse by year

Special collections index

## Journal information

---

Journal description

Aims and scope

**Editorial board**

Submission guidelines

Journal indexing and metrics

Reprints

Journal permissions

Recommend to library

Advertising and promotion

## Keep up to date

---



Facebook



Twitter



LinkedIn



YouTube



RSS feed



Email alerts

[View all options](#)

## Also from SAGE Publishing

### CQ Library

American political resources

### Data Planet

A universe of data

### SAGE Business Cases

Real-world cases at your fingertips

### SAGE Campus

Online skills and methods courses

### SAGE Knowledge

The ultimate social science library

### SAGE Research Methods

The ultimate methods library

### SAGE Video

Streaming video collections

### Technology from SAGE

Make learning and research easier

# Active Immunotherapy for Glioblastoma Treatment: A Systematic Review and Meta-Analysis

Cancer Control  
Volume 29: 1–11  
© The Author(s) 2022  
Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
DOI: [10.1177/10732748221079474](https://doi.org/10.1177/10732748221079474)  
[journals.sagepub.com/home/ccx](https://journals.sagepub.com/home/ccx)

Joni Wahyuhadi<sup>1,3</sup> , Irwan Barlian Immadoel Haq<sup>1,3</sup>, Muhammad Reza Arifianto<sup>1,3</sup>, Bagus Sulistyono<sup>1,3</sup> , Rizki Meizikri<sup>1,3</sup> , Atika Rosada<sup>1,3</sup>, Cita Rosita Sigit Prakoeswa<sup>2,3</sup>, and Rahadian Indarto Susilo<sup>1,3</sup>

## Abstract

**Introduction:** Glioblastoma multiforme (GBM) makes 60–70% of gliomas and 15% of primary brain tumors. Despite the availability of standard multimodal therapy, 2 years, 3 years, and 5 years survival rate of GBM are still low. Active immunotherapy is a relatively new treatment option for GBM that seems promising.

**Methods:** An electronic database search on PubMed, Cochrane, Scopus, and [clinicaltrials.gov](https://clinicaltrials.gov) was performed to include all relevant studies. This study was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA). Reported parameters are OS, PFS, AEs, post treatment KPS, and 2 year mortality.

**Results:** Active immunotherapy provided better OS (HR = .85; 95% CI = .71–1.01;  $P = .06$ ) and PFS (HR = .83; 95% CI = .66 – 1.03;  $P = .11$ ) side albeit not statistically significant. Active immunotherapy reduces the risk of 2 year mortality as much as 2.5% compared to control group (NNT and RRR was 56.7078 and 0,0258, respectively).

**Conclusion:** Active immunotherapy might be beneficial in terms of survival rate in patients with GBM although not statistically significant. It could be a treatment option for GBM in the future.

## Keywords

active immunotherapy, vaccine, high-grade glioma, glioblastoma, glioblastoma multiforme

## Introduction

### Background

Glioblastoma Multiforme (GBM) is the most common form of primary brain malignancy.<sup>1</sup> GBM makes 60–70% of gliomas and 15% of primary brain tumor.<sup>2</sup> Optimal treatment with maximal tumor resection followed by adjuvant radiotherapy (RT) and temozolomide (TMZ), does not guarantee a good survival rate nor avoid recurrence of GBM,<sup>2,3</sup> as proven by the low 2 years, 3 years, and 5 years survival rate in population-based study (18%, 11% and 4% respectively).<sup>4</sup>

Researches on new therapeutic modalities for GBM have been done, among which is on targeted immunotherapy.<sup>5,6</sup> The immunosuppressive trait of its microenvironment, and its

ability to interfere microglial immune activity, laid basis for immunotherapy researches for GBM.<sup>7</sup> In general, immunotherapy is classified into active and passive immunotherapy.<sup>8</sup>

<sup>1</sup>Department of Neurosurgery, Dr Soetomo General Academic Hospital, Surabaya, Indonesia

<sup>2</sup>Department of Dermatology and Venereology, Dr Soetomo General Academic Hospital, Surabaya, Indonesia

<sup>3</sup>Faculty of Medicine - Universitas Airlangga, Surabaya, Indonesia

### Corresponding Author:

Joni Wahyuhadi, Department of Neurosurgery, Dr Soetomo General Academic Hospital, Surabaya, Indonesia. Jl. Mayjen Prof. Dr. Moestopo No.6-8, Gubeng, Surabaya, East Java 60286, Indonesia.  
Email: [joniwahyuhadi@fk.unair.ac.id](mailto:joniwahyuhadi@fk.unair.ac.id)



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE

and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

The idea of immunotherapy in GBM is to trigger immune system to be more reactive to GBM and therefore able to kill the tumor cells.<sup>9</sup> However, the impact of immunotherapeutic advances on improvement of patients' survival is still unknown. Therefore, a well-designed systematic review and data synthesis would be invaluable to conclude the overall efficacy of active immunotherapies for the treatment of GBM.

## Objective

This study was performed in order to draw conclusion about efficacy of active immunotherapy for the treatment of the patients with GBM.

## Materials and Methods

### Criteria for Considering Studies for This Review

*Type of Participants.* Eligible participants are adults aged  $\geq 18$  year-old with confirmed histopathological diagnosis of newly diagnosed or recurrent GBM (WHO Grade IV) who have received either active immunotherapy or conventional therapy.

### Type of Intervention

The intervention arms of eligible studies were active immunotherapy. Active immunotherapy included (1) dendritic cell vaccination, (2) peptide vaccination, (3) DNA vaccine, (4) viral vector-based vaccine, (5) antigen non-specific vaccine, and (6) autologous tumor cell therapy. Standard therapy stands for combination of surgical resection, radiotherapy or chemotherapy.

### Type of Comparators

Standard therapy (surgery, chemotherapy, and radiotherapy) with or without placebo.

### Type of Outcome Measures

Studies assessing both the primary and secondary outcomes were included in this study.

*Primary Outcome.* The primary outcomes of this review are:

- (1) Overall survival (OS)
- (2) Progression-free survival (PFS)

*Secondary Outcome.* The secondary outcomes of this review are:

- (1) Karnofsky performance scale (KPS)
- (2) Adverse events (AEs)
- (3) Two year mortality

### Type of Studies

Randomized controlled trials (RCTs) phase II or later on active immunotherapy

### Exclusion Criteria:

- (1) Patients younger than 18 or diagnosed with other type of tumor
- (2) Study in which non-active immunotherapy were used
- (3) Study in which treatment other than standard therapy is used as comparison
- (4) Studies in which neither OS nor PFS are measured
- (5) Case series, non-RCT, review studies, irrelevant articles
- (6) Studies published beyond the last 10 years or not written in English

### Search Methods for the Identification of Studies

This systematic review and meta-analysis was conducted according to the PRISMA.<sup>10</sup> Literature search was done on PubMed, Cochrane, Scopus, and [clinicaltrials.gov](http://clinicaltrials.gov). The keywords used on each electronic database in this review were ("active immunotherapy" OR "targeted therapy" OR "dendritic cell vaccine" OR "peptide vaccine" OR "DNA vaccine" OR "Viral vector-based vaccine" OR "antigen non-specific vaccines" OR "autologous tumour cell therapy") AND (glioblastoma OR GBM OR "high-grade glioma" OR HGG) AND (RCT OR "Randomized controlled trial"). The search result from all databases was imported to Mendeley. Duplicate results were detected using "check for duplicate" in Mendeley and were excluded. We have registered this systematic review and meta-analysis in PROSPERO with ID number 276332

### Data Collection and Analysis

*Selection of studies.* All articles' titles and abstracts were scanned independently by all authors. The results of independent searches were matched in order to find the common result; 2 physician reviewed unmatched findings once more, in order to check if they met the inclusion eligibility criteria.

### Data Appraisal and Extraction

IBIH, MRA, BS, RM, AR, and RIS independently reviewed the full texts in order to confirm their eligibility according to the predefined participants, intervention, comparison, outcome, and study type (PICOS). In case of disagreement between the authors, the issue was discussed until full agreement was reached. The required data were extracted and put into a table accordingly.

### Assessment of Risk of Bias in Included Studies

Risk of bias was assessed by IBIH, MRA, BS, RM, AR, and RIS. Should conclusion not be met, JW and CRSP would be asked to give their opinion. The biases assessed in this study were those mentioned in The Cochrane Collaboration Tool for Assessing Risk of Bias in Randomized Trials published in 2011.<sup>11</sup> Accordingly, all studies were classified as "low risk", "high risk," or "unclear risk" of bias.

## Measures of Treatment Effect

OS and PFS were collected as median and range, and presented as dichotomous outcome. KPS are presented in mean differences (MDs) while 2 Year Mortality are presented as Number-Needed to Treat (NNT). AEs are presented as pooled incidence. Only serious AEs are included in the calculation.

## Dealing With Missing Data

Data were extracted as they were reported in the studies. Missing data were noted and reported as bias or explained in the result and/or discussion.

## Assessment of Heterogeneity

Heterogeneity is addressed according to the  $I^2$  value reported on the forest plot made using RevMan 5.4.

## Data Synthesis

Data of OS and PFS were collected and presented in the form of forest plot.

## Result

### Description of Included Studies

**Search results.** A systematic search was done on July–August 2020. Initially, a total of 326 publications were identified. However, 44 studies were excluded due to articles duplication. From a total number of 282 studies, further 214 studies were excluded after abstract screening. Most of those excluded were either not a phase-II/above RCTs or did not report the outcome that we look upon. At final screening, 57 of 68 studies were excluded due to one or more of the following: (1) no available full text, (2) not written in English, and (3) meeting our exclusion criteria. Thirteen studies were included for qualitative synthesis, and six studies were included for quantitative synthesis. The PRISMA flow diagram was presented in [Figure 1](#).

### Included Studies

Six studies were included in this review. In detail, 2 evaluated DCV (Yao et al, 2018 and Buchroitner et al, 2018), 1 WLDCV (Cho et al, 2012), 1 CIK (Kong et al, 2017), 1 autologous dendritic cell (ICT-107) (Wen et al, 2019), and 1 Rindopepimut (Weller et al, 2017). All results of included studied were summarized in [Table 1](#).

### Risk of Bias in Included Studies

In general, the studies mostly (more than 50%) had low risk of selection bias as 4 of 6 studies reported the randomization process. However, in terms of detection bias, attrition bias,

reporting bias, and other bias, 50% or more of the studies were considered high risk ([Figures 2\(A\) and \(B\)](#)).

## Summary of Findings

Summary of findings of included studies and quality of evidence of this study are listed on [Table 1](#) and [Table 2](#), respectively. Difference of baseline characteristics between studies can be seen on [Table 3](#).

## Effects of Intervention

The effect of the intervention on OS and PFS can be seen on [Figures 3\(A\) and \(B\)](#). The median follow-up of the studies was 15.56 (10.26–26) months. Changes in KPS were unfortunately not reported in most studies, thus MDs was not possible to be analyzed.

## Overall Survival


Kong et al compared autologous CIK cells and standard therapy against standard therapy alone. The median OS in the treatment group was 22.47 (95% CI, 17.2–23.85) against 16.88 (95% CI, 13.91–21.94) in the control group. The median OS in Cho et al was 31.9 (95% CI, 20–56) for intervention group against 15 (95% CI, 5–27) in control group. Median OS in Yao's study was 13.7 in the treatment group against 10.7 in the control group. Median OS in intervention group and control group of Buchroitner's study was 18.8 (95% CI, 14.53–22.37) and 18.93 (95% CI, 11.63–22.67), respectively. Wen et al reported median OS for intervention group was 17 (95% CI, 13.68–20.61), and in controlled group was 15 (95% CI, 12.33–23.05). The median OS in Weller et al was 17.4 (95% CI, 16.1–19.4) for intervention group and 17.4 (95% CI, 16.2–18.1).

The cumulative hazard ratio (HR) for OS in this study was .85 (95% CI, .71–1.01). This indicated that active immunotherapy was favorable despite not statistically significant. The studies were heterogenous in terms of OS ( $I^2 = 54%$ ,  $P = .06$ ). Only one study had CI which did not cross or touch value of 1 ([Figure 3\(A\)](#)).

## Progression Free Survival

The median PFS in intervention and control group of Kong's study was 8.1 (95% CI, 5.8–8.5) and 5.4 (95% CI, 3.3–7.9), respectively. The median PFS in Cho et al was 8.5 (95% CI, 3–56) for intervention group and 8 (95% CI, 2–18) in control group. Yao et al compared DC vaccine for patients with new or recurrent GBM who had the tumor resected  $\geq 95%$  prior to study. Median PFS for intervention group in this study was 7.7, while for control group was 6.9. Wen et al reported median PFS for intervention group was 11.2 (95% CI, 8.22–13.05), and in controlled group was 9 (95% CI, 5.52–10.29). In Buchroitner's study, the median PFS in intervention group and control group was 6.8 (95% CI, 4.6–9.3) and 7 (95% CI, 5.97–9.53), respectively. The median OS in Weller et al was





**Figure 1.** PRISMA flowchart.

7.1 (95% CI, 5.4–7.9) for intervention group and 5.6 (95% CI, 5.1–7.1).

The cumulative hazard ratio (HR) for PFS was .83 (95% CI, .66–1.03). This indicated that active immunotherapy was favorable despite not statistically significant. The studies were heterogenous in terms of OS ( $I^2=44\%$ ,  $P = .11$ ). Only 2 studies had CI which did not cross value of 1, yet they were barely significant (Figure 3(B)).

### Two Year Mortality

We extracted the number of events of 2 year mortality from each study based on the given rate or inferred from the provided Kaplan–Meier graphs. One study<sup>14</sup> did not reach


2 year follow-up and thus was not included in the calculation. In total, there were 411 events of 2 year mortality out of 617 total subjects in the intervention arm. On the other hand, there were 400 similar events out of 585 total subjects in the control arm. We calculated the number-needed to treat (NNT), control event rate (CER), experimental event rate (EER), absolute risk reduction (ARR), and relative risk reduction (RRR) based on cumulative number of subjects from all included studies (Table 4).

The cumulative value of NNT was 56.7078, which means that 56–57 patients need to be treated with active immunotherapy to prevent 1 mortality. The RRR of .0257 901 signifies that patients gain a small benefit of reduction of 2 year mortality risk by 2.5%. Absolute risk reduction of .0176 343

**Table 1.** Included studies.

Author, year	Type of Study	Patients (n)	Patients Characteristic	Length of Follow-Up	Type of Intervention (n)	Outcome		
						OS*	PFS*	Ae KPS Post Treatment (n)
Kong, 2017 <sup>11</sup>	RCT, Phase III	180	53.3 ± 10.8 y.o for intervention 52.8 ± 10.5 y.o for control ND GBM KPS ≥ 60	10.26 (2.48–41.7)	Standard + CIK (91)	22.47 (17.2–23.85)	8.1 (5.8–8.5)	43 n/a
Cho, 2012 <sup>12</sup>	RCT, Phase II	34	53.8 ± 12.0 y.o for intervention 55.8 ± 11.1 y.o for control ND GBM KPS > 70	26 (17–56)	Standard (89) Standard + WLDCV (18)	16.88 (13.91–21.94) 31.9 (20–56)	5.4 (3.3–7.9) 8.5 (3–56)	32 n/a 0 70 (50–100)
Yao, 2018 <sup>13</sup>	RCT, Phase II	43	48 ± n/a y.o for intervention 50 ± n/a y.o for control ND or rGBM KPS ≥ 60	14 (9–23)	Standard (16) Standard + DCV (22)	15 (5–27) 13.7 (n/a)	8 (2–18) 7.7 (n/a)	n/a 50 (0–50) 0 n/a
Wen, 2019 <sup>14</sup>	RCT, Phase II	124	57.4 ± n/a y.o for intervention 57.4 ± n/a y.o for control ND GBM KPS ≥ 70	11.4 (1.4–28.4)	Standard (21) Standard + ICT-107 (81)	10.7 (n/a) 17 (13.68–20.61)	6.9 (n/a) 11.2 (8.22 – 13.05)	n/a n/a 47 n/a
Buchroiter, 2018 <sup>15</sup>	RCT, phase II	76	Median 54.6 (n/a) for intervention Median 54 (n/a) for control ND GBM KPS n/a	18.73 (1.4–53.3)	Standard (43) DCV + standard (34) Standard (42)	15 (12.33–23.05) 18.8 (14.53–22.37) 18.93 (11.63–22.67)	9 (5.52 – 10.29) 6.8 (4.6 – 9.3) 7 (5.97 – 9.53)	35 n/a 18 n/a 12 n/a
Weller, 2017 <sup>16</sup>	RCT, Phase III	745	Median age 59 (51–64) for intervention Median age 58 (52–64) for control ND GBM KPS n/a	17.12 (.5–48)	Standard therapy → rindopepimut + TMZ (371) Placebo + Standard (374)	17.4 (16.1–19.4) 17.4 (16.2–18.1)	7.1 (5.4 – 7.9) 5.6 (5.1–7.1)	337 n/a 343 n/a

AE: Adverse events; CIK: Cytokine-induced killer; DCV: Dendritic cell vaccine; \*Presented in median (range) month  
 KPS: Karnofsky performance score  
 ND GBM: Newly diagnosed GBM  
 rGBM: Recurrent GBM  
 WI: Whole-cell lysate OS and PFS presented in median (CI) Length of follow-up presented in median (range)



**Figure 2. Risk of bias (A) across all included studies, (B) each included study.**

signifies that there really is only 1.76% risk difference between control group and treatment group.

### Adverse Events

Kong et al reported 43 grade III–IV AEs in the intervention group and 32 in control group. However, there were only 3 AEs which were thought to be related to the CIK. No AEs were reported in Cho's and Yao's studies. Wen et al reported 47 and 35 serious AEs in intervention and control group, respectively. Serious AEs in Buchroitrner's study occurred in 18 and 12 subjects of intervention and control group, respectively. Weller et al reported 337 serious AEs in intervention group and 343 in control group

The pooled incidence of AEs among the treatment group was 37.87% (95% CI, 10.098–71.028). This finding, however, is severely heterogenous ( $I^2 = 98\%$ ,  $P < .0001$ ). It is also important to note that some studies did not clearly report if more than 2 AEs were present in a single subject. Two of the included studies did not report any AEs. (Figure 4)

### Discussion

Up to this point, standard therapy for GBM includes surgery, radiotherapy, and chemotherapy.<sup>18</sup> However, it is well known that GBM would still recur despite maximum effort with standard therapy.<sup>19–22</sup> Studies investigating the standard therapy of GBM reported that the survival rate of GBM was less than 2 years.<sup>23–25</sup> Patients with GBM was found to have a median survival time of 15–17 months.<sup>26</sup> Only  $\leq 5\%$  of patients were said to be alive at 5 year follow-up.<sup>25</sup>

Previous study shows that immunotherapy in general could be beneficial for glioblastoma treatment.<sup>8</sup> Here, we specifically review the efficacy of active immunotherapy for glioblastoma from recent RCTs phase II or above that met the

criteria. The principle of immunotherapy for cancer is amplifying the body's immune system to react towards the tumor through humoral and cytotoxic mechanism.<sup>6</sup> Active immunotherapy works by training the immune system to attack the tumor based on specific antigenicity<sup>6</sup> and also generates tumor-specific T-cells and immunological memory.<sup>27</sup>

Of all 6 studies reviewed, 2 evaluated DCV, 1 WLDCV, 1 CIK, 1 autologous dendritic cell (ICT-107), and 1 Rindopepimut. Active immunotherapy has been reported to be promising in other published studies.<sup>28–30</sup> Similarly, in our meta-analysis we found that the OS and PFS were in favor of active immunotherapy (.85 (95% CI, .71–1.01) and .83 (95% CI, .66–1.03), respectively). Our finding on OS and PFS of active immunotherapy indicated that this treatment modality is favorable despite not statistically significant.

This present study analyzed the mortality rate in the first 2 years since randomization in each study. Follow-up in Yao et al<sup>31</sup> did not reach 2 years so that it was excluded from 2 year mortality calculation. We found an NNT of 56.7078, which means that 56–57 patients need to be treated with active immunotherapy to prevent 1 mortality. The RRR of .0257 901 signifies that the patients gain a small benefit of reduction of 2 year mortality risk by 2.5%. Absolute risk reduction of .0176 343 signifies that there really is only 1.76% risk difference between the 2 groups. This calculation, however, was possibly biased as some studies did not provide censored events in their respective Kaplan–Meier graphs. In such cases, mortality rate was calculated based on either survival or mortality percentage provided by the authors from the related study.

Serious AEs from the included studies were 411 and 400 in immunotherapy and in control group, respectively. Two studies<sup>13,14</sup> did not provide data on AEs. It is important to note that this present study only gathered data on AEs of grade III or above. None of the studies made clear if the AEs were

**Table 2.** Quality of evidence.

Outcomes	Hazard Ratio (95% CI)	Number of Participants (studies)	Quality of Evidence (GRADE)	Comments
OS	HR .85 (.71 to 1.01)	1202 (6)	⊕⊕⊕⊖ Low	Presence of Bias, Imprecision Indicated by the Range of Confidence Interval (CI)
PFS	HR .83 (.66 to 1.03)	1202 (6)	⊕⊕⊕⊖ low	Presence of bias, imprecision indicated by the range of confidence interval (CI)

**Table 3.** Baseline characteristics of each study.

Author	Mean Age	Mutation (%)	GTR (%)	Mean Baseline KPS
Kong, 2017 <sup>11</sup>	53.3	Not checked	48.4%	84.4
Cho, 2012 <sup>12</sup>	52.11	MGMT 55.5%	53.9%	85.7
	55.81	MGMT 56.25%	77.7%	n/a
Yao, 2018 <sup>13</sup>	50	IDHI <sup>MT</sup> TERT <sup>WT</sup> 4.76%	n/a	n/a
		IDHI <sup>WT</sup> TERT <sup>WT</sup> 47.6%		79
Wen, 2019 <sup>14</sup>	48	IDHI <sup>WT</sup> TERT <sup>MT</sup> 47.6%		
		MGMT 57.14%		
Wen, 2019 <sup>14</sup>	57.4	IDHI <sup>MT</sup> TERT <sup>WT</sup> 13.63%	n/a	83
		IDHI <sup>WT</sup> TERT <sup>WT</sup> 45.45%		
Wen, 2019 <sup>14</sup>	57.5	IDHI <sup>WT</sup> TERT <sup>MT</sup> 40.9%		
		MGMT 40.9%		
Wen, 2019 <sup>14</sup>	57.4	MGMT 34.6%	71.6%	24.7% KPS <90
		HLA-A1 40.7%		
Wen, 2019 <sup>14</sup>	57.5	HLA-A2 51.9%		
		HLA-A1 and A2 7.4%		
Wen, 2019 <sup>14</sup>	57.5	MGMT 41.9%	74.4%	39.5% KPS <90
		HLA-A1 32.6%		
Wen, 2019 <sup>14</sup>	57.5	HLA-A2 51.2%		
		HLA-A1 and A2 16.3%		
Buchroitner, 2018 <sup>15</sup>	54.6	MGMT 20.58%	70.58%	n/a
	54	MGMT 14.28%	83.3%	n/a
Weller, 2017 <sup>16</sup>	59 <sup>§</sup>	MGMT 33%	n/a	84.4
	58 <sup>§</sup>	MGMT 35%	n/a	85.7


Row in green: intervention Row in yellow: control §Median

exclusively related to the immunotherapy agents. Study which best elaborated their AEs finding was that of Weller et al.<sup>17</sup> Most common grade III–IV AEs for all 369 treated patients in the rindopepimut group vs 372 treated patients in the control group were: thrombocytopenia (32 (9%) vs 23 (6%)), fatigue (6 (2%) vs 19 (5%)), brain edema (8 (2%) vs 11 (3%)), seizure (9 (2%) vs 8 (2%)), and headache (6 (2%) vs 10 (3%)). Serious AEs included seizure (18 (5%) vs 22 (6%)) and brain edema (7 (2%) vs 12 (3%)).<sup>32</sup> However, since the treatment group also received standard therapy in all studies, it is impossible to draw conclusion if the AEs were truly due to immunotherapy.

Reported AEs of immunotherapy for cancer treatment includes autoimmunity.<sup>33</sup> This phenomenon mainly occurred in immune-checkpoint types of immunotherapy. The induction of autoimmunity is thought to occur from enhancement of sub-clinical autoimmune disease which had presented previously in the patient, altering commensal microbiota in the GI tract and/or immune interactions with environmental microorganisms,

leading to pathology where there was previously tolerance, bystander tissue injury due to antitumor response, or aberrant response to the immunotherapy agents themselves.<sup>33</sup> In a retrospective study on patients with melanoma receiving DCV, Most common AEs were flu-like symptoms (67%) and injection site reactions (50%). Both of these AEs significantly correlated with the presence of tetramer-positive CD8 T-cells.<sup>34</sup>

Significant progress has been made in understanding how cancer cells elude the immune system by expressing immunological checkpoints that limit T cell activity and proliferation during the last several decades. The cytotoxic T lymphocyte antigen 4 (CTLA-4) receptor and its ligand (PD-L1) are currently the most clinically relevant immunological checkpoints. However, immune checkpoint inhibition is unable to demonstrate efficacy in GBM patients. A systematic review by Brahm et al, showed that majority of clinical trials involving GBM patients showed minimal clinical activity. Ongoing clinical trials by CheckMate studies with nivolumab have also failed to prove its efficacy. Thus, the role of



**Figure 3.** (A) The effect of intervention towards OS, (B) The effect of intervention towards PFS.

**Table 4.** Calculation of EER, CER, ARR, RRR, and NNT based on 2-year mortality.

Intervention Group Size	617
Control group size	585
Events in experimental group	411
Events in control group	400
EER (experimental event rate)	.666 126
CER (control event rate)	.683 761
ARR (absolute risk reduction)	.0176 343
RRR (relative risk reduction)	.0257 901
NNT (number-needed to treat)	56.7078

immune checkpoint inhibitors currently is not applicable in glioblastoma patients.

### Excluded Studies

A study by Buchroitner et al<sup>35</sup> was excluded as both JW and RIS deemed the authors did not report the study's method clearly enough. Akiyama et al published a phase I clinical trial.<sup>36</sup> Ardon et al published a clinical trial which was unfortunately not randomized.<sup>37</sup> We also identified trials which were still in pilot phase.<sup>38,39</sup> Bota et al compared ERC1671 (Gliovac), granulocyte-macrophage stimulating factor (GM-CSF), cyclophosphamide, and bevacizumab against placebo and bevacizumab for recurrent GBM. The study was excluded due to comparing patients not currently on standard therapy.<sup>40</sup> Similarly, study by Narita et al was excluded as they did not investigate patients on standard therapy.<sup>41</sup> Liau et al studied dendritic cell (DC) vaccine for 331 newly diagnosed GBM subjects aged. Despite amassing one of the biggest number of subjects for this topic, this study was a cross-over trial, which means that in the end all patients received the DC vaccine and, therefore,

was excluded.<sup>42</sup> A study by Inoges et al was excluded for not being an RCT.

### Quality, Bias, and Limitations of the Study

Quality of evidence for OS and PFS in this study was low grade as there is imprecision indicated by the range of CI and presence of bias.


**Bias.** Bias in each study was summarized in Supplementary Figure 1.

**Randomization and Allocation Concealment.** Kong et al, Cho et al, and Yao et al randomized the patients using a computer-based randomization, but the allocation concealment was not made clear in all 3.<sup>12-14</sup> Allocation concealment in Weller's study was prone to bias as there was MGMT methylation-based stratification.<sup>17</sup> Wen et al and Buchroitner et al did not mention the randomization and allocation concealment method.<sup>15,16</sup>

**Blinding.** Buchroitner et al<sup>16</sup> did not make mention of any blinding processes. Wen et al<sup>15</sup> and Yao et al<sup>14</sup> did not explain how blinding was kept throughout the study. Cho et al<sup>13</sup> and Kong et al<sup>12</sup> was an open-label study. Weller et al kept the participants and personnel blinded by assigning someone not involved in the study to prepare the drugs and label them only with patient's ID.<sup>17</sup>

**Incomplete Outcome Data.** There were some subjects who failed to complete the intervention in three studies.<sup>12,16,17</sup>

**Selective Reporting.** Wen et al,<sup>15</sup> Yao et al,<sup>14</sup> and Buchroitner et al<sup>16</sup> did not mark any censored events on the Kaplan–Meier graph.



**Figure 4.** Pooled incidence of grade III/IV AEs among treatment group.

*Other Potential Sources of Bias.* Five studies received findings.<sup>12,14-17</sup> Buchroitner et al<sup>16</sup> provided survival data in days instead of months. Therefore, we converted the given data to months by dividing them by 30. A patient in Cho et al did not complete the standard radiation therapy.

### Limitations

Other limitations in our study are different types of active immunotherapy, unspecified mutational characteristic, probable dissimilar baseline KPS between studies, and differences in extent of resection among subjects. Besides that, inconsistent reporting in most of the studies made conclusion regarding KPS impossible to infer. Our search of literature was limited to one language which was English. It is possible that other trials may be available in other languages or on other Websites. Not all RCTs included in this study discussed factors that related to immunity, so we could not analyze

those factors that might affect the patients immunity. More comprehensive RCTs about active immunotherapy for glioblastoma are needed.

### Conclusion

Active immunotherapy might be beneficial in terms of survival rate in patients with GBM although not statistically significant. Further high quality RCT with bigger number of samples are required to draw better conclusion. Severe AEs reported, yet none of the studies made clear if the AEs were exclusively related to the immunotherapy agents. As most of the studies did not report post-intervention KPS, we could not report the conclusion regarding KPS in this review. Randomized controlled trial on GBM with specific mutation characteristics will be beneficial for future researches and will direct future GBM treatments to be more patients-specific.

## Appendix

### Abbreviation

AA	Anaplastic astrocytoma;
ADCTA	Autologous dendritic cell/tumour antigen vaccine;
AE	Adverse events;
ARR	Absolute risk reduction;
CER	Control event rate;
CIK	Cytokine-induced killer;
DC	Dendritic cell;
DCV	dendritic cell vaccine;
EER	Experimental event rate;
GBM	Glioblastoma Multiforme;
GM-CSF	Granulocyte-macrophage stimulating factor;
HR	Hazard Ratio;
KPS	Karnofsky Performance Scale;
LAK	Lymphokine activated killer;
ND GBM	Newly diagnosed GBM;
NNT	Number-Needed to Treat;
OS	Overall Survival;
PD-1	Programmed Death-1;
PFS	Progression-free survival;
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analysis;
RCTs	Randomized controlled trials;
rGBM	Recurrent GBM;
RRR	Relative Risk Reduction;
RT	Adjuvant Radiotherapy;
TMZ	Temozolomide;
TILs	Tumour-infiltrating lymphocytes;
WL	Whole-cell Lysate.

### Acknowledgments

The Authors would like to acknowledge Dr Soetomo General Academic Hospital and Surabaya Neuroscience Institute for its support in this study.

### Authors' Contribution

All authors contributed to the study conception and writing manuscript. Should conclusion not be met, JW and CRSP would be asked to give their opinion. All authors read and approved the final manuscript.

### Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

### Disclaimers

The information and views presented in this review reflect the views of the authors of this review, not an official position of institution or funder.

### ORCID iDs

Joni Wahyuhadi  <https://orcid.org/0000-0003-2053-1805>

Bagus Sulistyono  <https://orcid.org/0000-0001-7776-8788>

Rizki Meizikri  <https://orcid.org/0000-0002-8243-9951>

### References

- Hawasli AH, Kim AH, Dunn GP, Tran DD, Leuthardt EC. Stereotactic laser ablation of high-grade gliomas. *Neurosurg Focus*. 2014;37:E1.
- Huang B. *Advances in Immunotherapy for Glioblastoma Multiforme*; 2017.
- Chen R, Deng X, Wu H, et al. Combined immunotherapy with dendritic cells and cytokine-induced killer cells for malignant tumors: A systematic review and meta-analysis. *Int Immunopharm*. 2014;22:451-464.
- Poon MTC, Sudlow CLM, Figueroa JD, Brennan PM. Longer-term ( $\geq 2$  years) survival in patients with glioblastoma in population-based studies pre- and post-2005: A systematic review and meta-analysis. *Sci Rep*. 2020;10:11622-11710.
- Vatu BI, Artene S-A, Staicu A-G, et al. Assessment of efficacy of dendritic cell therapy and viral therapy in high grade glioma clinical trials. A meta-analytic review. *J Immunoassay Immunochem*. 2019;40:70-80.
- Bregy A, Wong TM, Shah AH, Goldberg JM, Komotar RJ. Active immunotherapy using dendritic cells in the treatment of glioblastoma multiforme. *Cancer Treat Rev*. 2013;39:891-907.
- Safdari H, Hochberg FH, Richardson EP. Prognostic value of round cell (lymphocyte) infiltration in malignant gliomas. *Surg Neurol*. 1985;23:221-226.
- Hanaei S, Afshari K, Hirbod-Mobarakeh A, Mohajer B, Amir Dastmalchi D, Rezaei N. Therapeutic efficacy of specific immunotherapy for glioma: A systematic review and meta-analysis. *Rev Neurosci*. 2018;29:443-461.
- Weenink B, French PJ, Sillevius Smitt PAE, Debets R, Geurts M. Immunotherapy in glioblastoma: Current shortcomings and future perspectives. *Cancers*. 2020;12.
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Med*. 2009;6:e1000100.
- Higgins JPT, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928.
- Kong D-S, Nam D-H, Kang S-H, et al. Phase III randomized trial of autologous cytokine-induced killer cell immunotherapy for newly diagnosed glioblastoma in Korea. *Oncotarget*. 2017;8:7003-7013.
- Cho D-Y, Yang W-K, Lee H-C, et al. Adjuvant immunotherapy

- with whole-cell lysate dendritic cells vaccine for glioblastoma multiforme: A phase II clinical trial. *World Neurosurgery*. 2012; 77:736-744.
14. Yao Y, Luo F, Tang C, et al. Molecular subgroups and B7-H4 expression levels predict responses to dendritic cell vaccines in glioblastoma: An exploratory randomized phase II clinical trial. *Cancer Immunol Immunother*. 2018;67:1777-1788.
  15. Wen PY, Reardon DA, Armstrong TS, et al. A randomized double-blind placebo-controlled phase II trial of dendritic cell vaccine ICT-107 in newly diagnosed patients with glioblastoma. *Clin Cancer Res*. 2019;25:5799-5807.
  16. Buchroithner J, Erhart F, Pichler J, et al. Audenel immunotherapy based on dendritic cells has no effect on overall and progression-free survival in newly diagnosed glioblastoma: A phase II randomized trial. *Cancers*. 2018;10.
  17. Weller M, Butowski N, Tran DD, et al. Rindopepimut with temozolomide for patients with newly diagnosed, EGFRvIII-expressing glioblastoma (ACT IV): A randomised, double-blind, international phase 3 trial. *Lancet Oncol*. 2017;18:1373-1385.
  18. Komotar RJ, Otten ML, Moise G, Connolly ES. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma—a critical review. *Clin Med Oncol*. 2008;2:421-422.
  19. Alexander BM, Cloughesy TF. Adult glioblastoma. *J Clin Oncol*. 2019;35:2902-2909.
  20. Burri SH, Gondi V, Brown PD, Mehta MP. The evolving role of tumor treating fields in managing glioblastoma. *Am J Clin Oncol*. 2018;41:191-196.
  21. Ji X, Zhu H, Dai X, et al. Overexpression of GBP1 predicts poor prognosis and promotes tumor growth in human glioblastoma multiforme. *Cancer Biomarkers*. 2019;25:275-290.
  22. Tang X, Xu P, Wang B, et al. Identification of a specific gene module for predicting prognosis in glioblastoma patients. *Front Oncol*. 2019;9:812.
  23. Hegi ME, Diserens A-C, Gorlia T, et al. MGMT Gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med*. 2005;352:997-1003.
  24. Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med*. 2005;352:987-996.
  25. Stupp R, Hegi ME, Mason WP, et al. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol*. 2009;10:459-466.
  26. Ostrom QT, Gittleman H, Fulop J, et al. CBTRUS statistical report: Primary brain and central nervous system tumors diagnosed in the United States in 2008-2012. *Neuro Oncol*. 2015; 17(suppl 4):iv1-iv62
  27. Steinman RM. Decisions about dendritic cells: Past, present, and future. *Annu Rev Immunol*. 2012;30:1-22.
  28. Chodon T, Koya RC, Odunsi K. Active immunotherapy of cancer. *Immunol Invest*. 2015;44:817-836.
  29. Jin J, Joo KM, Lee SJ, et al. Synergistic therapeutic effects of cytokine-induced killer cells and temozolomide against glioblastoma. *Oncol Rep*. 2011;25:33-39.
  30. Rapp M, Grauer OM, Kamp M, et al. A randomized controlled phase II trial of vaccination with lysate-loaded, mature dendritic cells integrated into standard radiochemotherapy of newly diagnosed glioblastoma (GlioVax): Study protocol for a randomized controlled trial. *Trials*. 2018;19:293-314.
  31. Yao M, Li S, Wu X, et al. Cellular origin of glioblastoma and its implication in precision therapy. *Cell Mol Immunol*. 2018;15: 737-739.
  32. Pauken KE, Dougan M, Rose NR, Lichtman AH, Sharpe AH. Adverse events following cancer immunotherapy: Obstacles and opportunities. *Trends Immunol*. 2019;40:511-523.
  33. Boudewijns S, Westdorp H, Koomstra RHT, et al. Immune-related adverse events of dendritic cell vaccination correlate with immunologic and clinical outcome in stage III and IV melanoma patients. *J Immunother*. 2016;39:241-248.
  34. Buchroithner J, Pichler J, Marosi C, et al. Vascular endothelial growth factor targeted therapy may improve the effect of dendritic cell-based cancer immune therapy. *Int J Clin Pharm Ther*. 2014;52:76-77.
  35. Akiyama Y, Oshita C, Kume A, et al.  $\alpha$ -type-1 polarized dendritic cell-based vaccination in recurrent high-grade glioma: A phase I clinical trial. *BMC Cancer*. 2012;12:623-11.
  36. Ardon H, Van Gool SW, Verschuere T, et al. Integration of autologous dendritic cell-based immunotherapy in the standard of care treatment for patients with newly diagnosed glioblastoma: Results of the HGG-2006 phase I/II trial. *Cancer Immunol Immunother* 2012;61:2033-44.
  37. Sun J, Cai L, Zhang K, et al. A pilot study on EGFR-targeted molecular imaging of PET/CT With <sup>11</sup>C-PD153035 in human gliomas. *Clin Nucl Med*. 2014;39:e20-6.
  38. Reap EA, Suryadevara CM, Batich KA, et al. Dendritic cells enhance polyfunctionality of adoptively transferred t cells that target cytomegalovirus in glioblastoma. *Cancer Res*. 2018;78:256-264.
  39. Bota DA, Chung J, Dandekar M, et al. Phase II study of ERC1671 plus bevacizumab versus bevacizumab plus placebo in recurrent glioblastoma: Interim results and correlations with CD4+ T-lymphocyte counts. *CNS Oncology*. 2018;7:CNS22.
  40. Narita Y, Arakawa Y, Yamasaki F, et al. A randomized, double-blind, phase III trial of personalized peptide vaccination for recurrent glioblastoma. *Neuro Oncol*. 2019;21:348-359.
  41. Liao LM, Ashkan K, Tran DD, et al. Correction to: First results on survival from a large Phase 3 clinical trial of an autologous dendritic cell vaccine in newly diagnosed glioblastoma. *J Transl Med*. 2018;16: 179.
  42. Inogés S, Tejada S, de Cerio AL-D, et al. A phase II trial of autologous dendritic cell vaccination and radiochemotherapy following fluorescence-guided surgery in newly diagnosed glioblastoma patients. *J Transl Med*. 2017;15:1-12. doi:10.1186/s12967-017-1202-z.