Volume 11 Number 01 January 2020



Indian Journal of Public Health Research & Development

An International Journal

Website:

www.ijphrd.com

Vol. 11 No. 1 (2020): Indian Journal of Public Health Research & Development



DOI: https://doi.org/10.37506/ijphrd.v11i1

Published: 2020-05-21

Articles

Study Protocol: A Randomised Controlled Trial on Effectiveness of a Worksite Health Intervention on Common Musculoskeletal Problems and Work-Related Quality of Life (WRQoL) among Female Workers in the Garment Manufacturing Sectors

A. Santham Lilly Pet1, Timsi Jain2, Bobby Joseph3, Pethuru Devadason4, Gayathri M5 1-6



Health of the Elderly in India: A Socio-Legal Study

Arti1, J.K. Mittal2 7-11



General Awareness of Diabetes Mellitus among a Hospital Population in Chennai: A Survey

Sridhar M.1, Abilasha Ramasubramanian2 12-15





Identifying the Status of Menstrual Hygiene Management

Amrita Shilpi1, Rajasree Roy2, Gobina3, Spriha Roy4 20-26



Effect of Strengthening of Scapular Stabilizers in Treatment of Rounded Shoulder Posture in Dental Students

Ankita M. Patil1, Sayali Gijare2 27-30



Elder Abuse in Indian Setting-A Misconception or a Reality- A Deductive Analysis

Anusha Rashmi1, Linda Sequeira2, Prianka Shashi Kumar2, Rashmi3 31-34



A Study on Women Entrepreneurs Dealing Through Stress

Archana R.V.1, K. Vasanthi Kumari2 35-37



Effect of Bilateral Scapular Muscles Strengthening on Dynamic Balance in Post Stroke Individuals

Arpan Dhoka1, G. Varadharajulu2 38-44



In Vitro Antibacterial and Anticancer Study of Bioactive Compounds Isolated from Punica Grantum Peel

Arunava Das1, J. Bindhu, M. Bharath2, Nithin Johnson3, M. Jeevanantham3 45-52



A Study on Utilization of Primary Health Care Services among the People Residing in a Semi-Urban Area

B. Charumathi1, D. Jayashri1, S. Manisha2, Aadithya2, C. Hemanthkumar2, Timsi Jain3 53-56



Gonial Angle as a Determinant of Gender, a Panoramic Study in a Sample of Saudi Population

Ahmed Ali Alfawzan

1689-1693



Determinants of Under-Five Mortality in Southern Asia

Ankika Dutta

1694-1697



Can Better Infrastructure Ensure Better Healthcare? The Dialectic of Assam's Health Sector Scenario

Kasturi Goswami

1698-1701



Study the Link between Laparoscopic Cholecystectomy and Abdominal Wall Paraumbilical Hernia

Mohammed Mohammud Habash

1702-1704



Acute Head Injury with Pregnancy

Ajaydeep Singh1, Arvinpreet Kour2, Unmesh S. Santpur3 1705-1709



Effectiveness of Deep Brain Stimulation in Parkinsonism

Ajaydeep Singh1, Arvinpreet Kour

1710-1717



Significance of PD-L1 Expression and Tumor Infiltrating Lymphocytes in High Grade Serous Ovarian Cancer: Egyptian Experience

Mai Gad1, Amany Abou-Bakr2, Rasha Mahmoud Allam3, Hanan Ramdan Nassar4, Maher H. Ibraheem5, Soha Talima6, Ghada Mohamed7

1718-1724



Association between Jordanian Ostomates' Knowledge about Intestinal Ostomy Care and their Ostomy Health-Related Problems Rami A. Elshatarat

1725-1729



Comparison between Lord Dilitation Versus Lateral Internal Sphincterotomy for Management Post Hemorrhoidectomy Pain and Stenosis

Raisan Mahdi Shoramah Aljabery

1730-1734



Competency-based Assessment of Public Health Professionals in the Northeastern Region, Thailand: An Exploratory Factors Analysis

Wilawun Chada1, Songkramchai Leethongdee2, Supa Pengpid3, Sangud Chualinfa 1735-1740



Does Vertigo Predict Hypertension in Cervical Spondylosis Clinical Trail Study

Talib Kadhim Akar

1741-1744



Is there an Effect of Serotonin on Attention Deficit Hyperactivity Disorder

Yunias Setiawati1, H.J. Mukono2, Joni Wahyuhadi3, Endang Warsiki1, Sasanti Yuniar1 1745-1749



Different Grades of BMI is Correlated with Left Atrium and Ventricle Structure in Patients with Hypertensive Heart Disease

Meity Ardiana1, Rofida Lathifah2, Makhyan Jibril Al-Farabi1,3, Muhammad Satya Bhisma 1750-1754



Herbal Dental Products: The Impact of Social Media on Consumers' Behaviour

Manar E. Al-Samaray1, Humam M. Al-Somaiday1, Ali Mahmoud Al-Samydai2, Rudaina Othman Yousif 1755-1759



The Effect of Moringa Oleifera Flour Given for Mothers Breastfeeding Against Morbidity of Baby Ages 0-6 Months in Jeneponto District

Suhartatik1, Veni Hadju2, Masyita Muis3, Hasanuddin Ishak 4, Merryana Adriani 1760-1765



Evaluation of an Experimental Poly-Methyl Methacrylate/Nano Graphene Oxide Composite

Reem Gamal1, Yasser F. Gomaa2, Ashraf Mahroos3

1766-1771



Manipulative Movement Based on Information Technology Games for School Children Aged 10-12 Years

Nevi Hardika1, Moch. Asmawi2, James Tangkudung2, Firmansyah Dlis2, Abdul Sukur7, Widiastuti2, M.E. Winarno3

1772-1778



The Influence of Job Stressor on Organizational Loyalty and Intention to Quit among Health Care Staff

Mohammad Saipol Mohd Sukor1, Siti Aisyah Panatik2, Wan Mohd Azam Wan Mohd Yunus 1779-1783



Emotional Intelligence and Conflict Management Style among Staff in a Bank

Maisarah Mohd1, Halimah Mohd Yusof2

1784-1788



Trend on Drink Drive and Road Accident Across Asian Region: A Review Study

Siti Hawa Harith1,2, Norashikin Mahmud

1789-1796



Obesity and Job Performance among Teachers in Malaysia

Mohd Hakiki Md Tohid1, Zulkifli Khair2

1797-1802



Does Instagram's Like Affected Teenager Self-worth?

Hayinah Ipmawati1, Wiwien Dinar Pratisti2 1803-1806



The Application of MOPSI Module Forbreast Cancer Patients

Editorial Team

Editor

Dr Vidya Surwade

Professor, Dr Baba Saheb Ambedkar, Medical College & Hospital Rohini, Delhi

Associate Editor

Prof. (Dr.) Anil Gupta

Vice principal, Professor and Head Department of Biochemistry, Eklavya Dental College and Hospital, Kotputli, Affiliated with Rajasthan University of Health Sciences, Jaipur.

editor.ijphrd@gmail.com

Editorial Board

INTERNATIONAL BOARD

- Dr Abdul Rashid Khan B. Md Jagar Din, Associate Professor, Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
- Prof. Luca Testarelli, Department of Dentistry, Sapienza University of Rome, Italy
- Dr V Kumar, Consulting Physician, Mount View Hospital, Las Vegas, USA
- Basheer A. Al-Sum ,Botany and Microbiology Deptt , College of Science, King Saud University, Riyadh, Saudi Arabia
- Dr Ch Vijay Kumar, Associate Professor, Public Health and Community Medicine, University of Buraimi, Oman
- Dr VMC Ramaswamy, Senior Lecturer, Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
- Kartavya J. Vyas ,Clinical Researcher, Department of Deployment Health Research,Naval Health Research Center, San Diego, CA (USA)
- Hailemichael Mulugeta, Lecturer and Researcher inDebre Berhan University, Health science College, Debre Birhan Ethiopia
- Prof PK Pokharel, Community Medicine, BP Koirala Institute of Health Sciences, Nepal
- Sajjad Salim Issa AL-Musawi ,Supervisor for student of the Arab board in family medicine
- Prof. Dr. Ayad F. Alkaim, Professor of surface chemistry and applications of nanomaterials, Babylon University, Iraq
- Prof. Dr. Imad Hadi Hameed, Biomedical Science Department, College of Nursing, University of Babylon, Hillah city, Iraq
- Prof. Dr. Amean A Yasir, Department of Community, College of Nursing, University of Babylon, Hillah city, Iraq
- Dr Ammar Mohammed Hamood AL-Farga, Associate Professor, Department of Biochemistry, Faculty of Science, University of Jeddah, Saudi Arabia.
- Dr. Rajaa Ahmed Mahmoud University of Basrah, Al Zahraa College of Medicine, Iraq.

- Prof. Sushanta Kumar Mishra, Community Medicine, GSL Medical College Rajahmundry, Karnataka
- Prof D.K.Srivastava, Medical Biochemistry, Jamia Hamdard Medical College, New Delhi
- Dr Sunil Mehra, Paediatrician & Executive Director, MAMTA Health Institute of Mother & Child, New Delhi
- Prof M.Sriharibabu ,General Medicine ,GSL Medical College , Rajahmundry , Andhra Pradesh
- Prof Pankaj Datta, Principal & Prosthodentist, Indraprastha Dental College, Ghaziabad India
- Prof Samarendra Mahapatro, Pediatrician, Hi-Tech Medical College, Bhubaneswar, Orissa, India
- Dr Abhiruchi Galhotra, Additional Professor, Community and Family Medicine, AII India Institute of Medical Sciences, Raipur
- Prof Deepti Pruthvi,Pathologist,SS Institute of Medical Sciences & Research Center. Davangere, Karnataka
- Prof G S Meena, Director Professor, Maulana Azad Medical College, New Delhi
- Prof Pradeep Khanna, Community Medicine, Post Graduate Institute of Medical Sciences, Rohtak , Haryana
- Dr Shailendra Handu, Associate professor, Phrma, DM (Pharma, PGI Chandigarh)
- Dr. A.C. Dhariwal, Directorate of National Vector Borne Disease Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of India, Delhi.

SCIENTIFIC COMMITEE

- Dr Anju Ade, Professor, Community Medicine Department, SVIMS, Sri Padamavati Medical College, Tirupati. Andhra Pradesh.
- Dr.E.Venkata Rao, Associate Professor, Community Medicine,
 Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
- Dr. Amit K. Singh, Associate Professor, Community Medicine, VCSG Govt. Medical College,
 Srinagar Garhwal, Uttarakhand
- Dr R G Viveki, Prof n head, Dept of Community Medicine, BIMS Belgavi, India
- Dr. Santosh Kumar Mulage, Assistant Professor, Anatomy, Raichur Institute of Medical Sciences Raichur (RIMS), Karnataka, India
- Dr Gouri Ku. Padhy, Associate Professor, Community and Family Medicine,
 AII India Institute of Medical Sciences, Raipur, India
- Dr Ritu Goyal, Associate Professor, Anaesthesia , Sarswathi Institute of Medical Sciences, Panchsheel Nagar
- Dr. Anand Kalaskar, Associate Professor, Microbiology, Prathima Institute of Medical Sciences India
- Dr.Md.Amirul Hassan ,Associate Professor, Community Medicine Government Medical College, Ambedkar nagar, UP,India
- Dr.N.Girish, Associate Professor, Microbiology, VIMS&RC, Bangalore, India.
- Dr BR Hungund, Associate Professor Pathology, JNMC, Belgaum India
- Dr Sumeeta Soni, Associate Professor, Microbiology Department, B.J. Medical College, Ahmedabad ,Gujarat,India

Platform & workflow by OJS / PKP

DOI Number: 10.37506/v11/i1/2020/ijphrd/194103

Is there an Effect of Serotonin on Attention Deficit Hyperactivity Disorder

Yunias Setiawati¹, H.J. Mukono², Joni Wahyuhadi³, Endang Warsiki¹, Sasanti Yuniar¹

¹Department of Psychiatric, Faculty of Medicine Universitas Airlangga, Dr. Soetomo General Hospital, ²Department of Environmental Health, Faculty of Public Health Universitas Airlangga, ³Department of Neurosurgery, Faculty of Medicine Universitas Airlangga, Dr. Soetomo General Hospital

Abstract

Background: Attention deficit hyperactivity disorder (ADHD) is a neurobiological disorder with a prevalence of 5%-10% in the world that negatively impacts school behavior and achievement. There have been many studies reveals that attention disorders are caused by the decreased levels of dopamine and norepinephrine neurotransmitter. However, particular concern nowadays are the emotional and behavioral problems. The emotional and behavioral problem is suspected to be caused by the decrease in serotonin level. The aim of this study was to determine the difference in serotonin level between ADHD and non-ADHD children.

Method: This study was an observational case-control study design using random sampling method. Subject of this study was children in Bina Karya Elementary School Surabaya. ADHD severity was assessed using Conners Abreviated Rating Scale. Examination of serotonin levels in blood was done by ELISA method. Informed consent was signed by parents before the study. Data was analyzed using independent T-test.

Results: 44 (23 ADHD and 21 non-ADHD) children from grade 1 to 6 was included in the study. Based on the gender, 13 children were girls and 31 children were boys. Serotonin level in ADHD children was significantly higher than non-ADHD children (2.148 \pm 0.94 vs 2.006 \pm 0.115 μ mol/L; p = 0.0001).

Conclusion: Serotonin level in ADHD children was significantly higher than non-ADHD children.

Keyword: Serotonin, ADHD, Abbreviated Conners Teacher Rating Scale.

Introduction

Attention deficit hyperactivity disorder (ADHD) is a neurobiological disorder with a prevalence of 5%-10% in the world that negatively impacts school behavior and achievement. The onset of this disorder usually appears in the school-age⁽¹⁾. Many parents feel disconcerted in nurturingtheir children with ADHD since they often opposed, haphazardly behave,and disobey the rules^(2,3)

Corresponding Author: Yunias Setiawati

Department of Psychiatric, Faculty of Medicine Universitas Airlangga, Dr. Soetomo General Hospital e-mail: yunias.setiawati@gmail.com Parents, especially mothers, often have to lose their jobs because they have to take care of their ADHD children^(4,5). Teachers at school also often complain of being confused and discouraged in educating children with ADHD because they need extra attention while there are limited teachers available and a lot of students to be supervisedat one time^(6–8) ADHD often coexist with other disorders, such as anxiety, depression, and learning disorder due to short spans of attention, hyperactivity, and impulsive behavior^(9–12). Management of ADHD is multimodal, including pharmacotherapy and psychotherapy. Psychosocial therapy is also considered for parents that refuse pharmacotherapy^(6,9,13–15)

There have been many studies reveals that attention disorders are caused by the decreased levels of dopamine and norepinephrine neurotransmitter⁽¹⁶⁾ However,

particular concern nowadays are the emotional and behavioral problems. It is more severe than the communication difficulties in the family, as emotional and behavioral disorders will adversely affect the entire life span of an ADHD children⁽¹⁷⁾. The emotional and behavioral problem is suspected to be caused by the decrease in serotonin level. However, study about the effects of serotonin is very limited and the results are still controversial^(18,19).

Considering that the study about the effect of serotonin is very limited and the result is still controversial, authors conduct this study to determine the difference in serotonin level between ADHD and non-ADHD children.

Method

This study was an observational case-control study design using random sampling method. Subject of this study was children in Bina Karya Elementary School Surabaya. The inclusion criteria for the case group was ADHD children in class 1 to 6, the mothers were not experiencing any psychosocial stressor during the study period, and both the mothers and the children were able to communicate using Indonesian language and being cooperative during the study. The inclusion criteria for the control group was non-ADHD children in class 1

to 6, and both the mothers and the children were able to communicate using Indonesian language and being cooperative during the study. The exclusion criteria for the study was children with the others mental health condition.

ADHD scores was assessed using Abbreviated Conners Teacher Rating Scale in Indonesian version. Social Readjustment Rating Scale by Holmes and Rahe was used to assess psychosocial stressor of the mother (Holmes & Rahe, 1967). Serotonin serum level was measured using ELISA method.

Shappiro-wilk test was used for the normality test. Independent T-test was used to analyze the difference between serotonin level in ADHD and non-ADHD children. SPSS 17.0.0 for windows software, 2007, SPSS Inc. Chicago, IL, USA and Microsoft Office 2013 was used for the statistical analysis test.

This study follows the principles of the Declaration of Helsinki. This study has received recommendation from Surabaya city government office before the study begins (Recommendation Number 070/1574/436.8.5/2018). Parents were given information for consent and signed the informed consent regarding the study prior to their involvement in the study. Details that might disclose the identity of the subjects under study were omitted.

Results

Table 1: Characteristic of the Subjects' Family

		Non ADHD N (%)	ADHD N (%)	Total
Highest Education Background	Junior high school	11 (50)	11 (50)	22 (100)
	Senior high school	7 (41.2)	10 (58.8)	17 (100)
	university	3 (47.7)	2 (52.3)	5(100)
Occupation	Not work	16 (51.6)	15 (48.4)	31 (100)
	Work	5 (38.5)	8 (51.5)	13 (100)
Sufficiency of monthly income	Not sufficient	5 (50)	5 (50)	10 (100)
	Almostsufficient	6 (37.5)	10 (62.5)	16 (100)
	Sufficient	9 (56.2)	7 (43.8)	16 (100)
	Very sufficient	1 (47.7)	1 (52.3)	2 (100)
Parents relationship	Poor	0	3 (100)	3 (100)
	Good	21 (51.2)	20 (48.8)	41 (100)
Relationship among family member	Poor	0	5 (100)	5 (100)
	Good	21 (53.8)	18(46.2)	39 (100)
Family Member Social interaction	Poor	2 (66.7)	1 (33.3)	3 (100)
	Good	19 (46.3)	22 (53.7)	41 (100)

Characteristic of the subjects and the family: 21 non-ADHD children and 23 ADHD children was included in the study. Based on the gender, there were 13 girls (6 in non-ADHD group and 7 in ADHD group) and 31 boys (15 in non-ADHD group and 16 in ADHD group). From the mother highest education background, 22 parents had graduated from junior high school, 17 had graduated from senior high school education background, and 5 had graduated from university education background. Based on the occupation, 13 children had their mother worked, while 31 children had their mother not worked. From the sufficiency of the monthly income, only 18 family had sufficient monthly income. 3 family had a poor mother-father relationship, whilst 5 family had a poor family relationship (table 1).

Table 2. Serotonin Level in ADHD and Non-ADHD Children

Variable	Mean	P
ADHD	$2.148 \pm 0.94 \ \mu mol/L$	0.0001
Non-ADHD	$2.006 \pm 0.115 \ \mu mol/L$	

Results Table 2. The mean serotonin level in ADHD children was $2.148 \pm 0.94 \ \mu mol/L$, while the mean serotonin level in non-ADHD children was $2.006 \pm 0.115 \ \mu mol/L$. Normality test showed that the data distribution was normal. Using independent T-test, it was found that serotonin level in ADHD was significantly higher than non-ADHD children (p = 0.0001, 95%CI = 0.780-0.206).

Discussion

From the demographic data of the subjects, there were 16 boys and 7 girls with ADHD. This result is consistent with the results of previous studies which found that the prevalence of ADHD was greater in boys^(6,20,21). Psychostimulant methylphenidate (MPH) and amphetamine (AMPH) are the first-line ADHD drug choice^(22–25) Both of these drugs primarily work on dopamine (D2) and noradrenaline (NA) system, but not on the serotonin system^(8,26). More than half of the subjects did not respond positively to the administration of this psychostimulant^(6,27) Not all parents agree with methylphenidate that affects appetite and inhibits growth when given to their children and also the risk of drug dependence or abuse on long-term use with adolescent subject⁽²⁸⁾.

Serotonin selective reuptake inhibitors (SSRIs) that working in the serotonin system are the second-

line treatment in ADHD management(13,29-33). This fact becomes the basic etiologic hypothesis of serotonin to ADHD. Based on these results, hypotheses suggest that chronic serotonin deficiency causes hyperactivity and impulsivity symptoms in ADHD (19). The etiology of serotonin in ADHD up to now is not clearly known on brain imaging studies it was found that serotonin especially in the prefrontal area of the cortex contributes to the clinical symptoms of ADHD^(6,32) Prefrontal cortex plays a role in regulating attention, cognitive function, emotional regulation and motivation(34,35) Serotonin plays an important role in the development of the nervous system in early child development and has a different role with serotonin in adulthood Serotonin affects the symptoms of hyperactivity and impulsivity but has no effect on^(34,35). The statistical analysis of serotonin revealed a significant differences between ADHD and non-ADHD subjects. These new findings may provide input for alternative pharmacotherapy management other than methylphenidate because not all parents agree with the administration of these drugs that have side effects on decreased appetite and inhibit the growth of the child.

Conclusion

Serotonin level in ADHD children was significantly higher than in non-ADHD children.

Ethical Clearance: Taken from Health Research Ethics Committee Faculty of Medicine Airlangga University No. 212/EC/KEPK/FKUA/2018

Source of Funding: Self

Conflict of Interest: Nil

References

- Shin D, Ph D, Kim E, Oh K, Ph D. The Relationship between Hair Zinc and Lead Levels and Clinical Features of Attention-Deficit Hyperactivity Disorder. 2014;25(1):28–36.
- Shimabukuro S, Daley D, Thompson M, Laver-Bradbury C, Nakanishi E, Tripp G. Supporting Japanese Mothers of Children with ADHD: Cultural Adaptation of the New Forest Parent Training Programme. Jpn Psychol Res. 2017;59(1):35–48.
- 3. Scott S. Parenting programmes for attachment and conduct problems. Psychiatry. 2008;7(9):367–70.
- Derakhshanpoor F, Khaki S, Vakili A, Shahini
 N. Study of the Status of Mental Health in

- Mothers with Parenting Style in the Children with Attention Deficit and Hyperactivity Disorder (ADHD). Eur Psychiatry [Internet]. Elsevier Ltd; 28AD;30, Supple(0):578. Available from: http://www.sciencedirect.com/science/article/pii/S0924933815304594
- 5. Theule J, Wiener J, Tannock R, Jenkins JM. Parenting Stress in Families of Children With ADHD: A Meta-Analysis. J Emot Behav Disord. 2013;21(1):3–17.
- 6. Paule MG, Rowland AS, Ferguson SA, Chelonis JJ, Tannock R, Swanson JM, et al. Attention deficit/hyperactivity disorder: Characteristics, interventions and models. Neurotoxicol Teratol. 2000;22(5):631–51.
- 7. Nixon RDV. Treatment of behavior problems in preschoolers. Clin Psychol Rev. 2002;22(4):525–46.
- 8. Frei H, von Ammon K, Thurneysen A. Treatment of hyperactive children: Increased efficiency through modifications of homeopathic diagnostic procedure. Homeopathy. 2006;95(3):163–70.
- 9. Biederman J. Attention deficit hyperactivity disorder (ADHD). Ann Clin Psychiatry. 1991;3(1):9.
- 10. de Graaf R, Kessler RC, Fayyad J, ten Have M, Alonso J, Angermeyer M, et al. The prevalence and effects of adult attention-deficit/hyperactivity disorder (ADHD) on the performance of workers: results from the WHO World Mental Health Survey Initiative. Occup Environ Med [Internet]. 2008;65(12):835–42. Available from: http://oem.bmj.com/cgi/doi/10.1136/oem.2007.038448
- Chronis AM, Chacko A, Fabiano GA, Wymbs BT, Pelham, Jr. WE. Enhancements to the behavioral parenting training paradigm for families of children with ADHD: Review and future directions. Clin Child Fam Psychol Rev [Internet]. 2004;7(1):1–27. Available from: http://link.springer.com/article/10.1023/B:CCFP.0000020190.60808.a4
- 12. Motlagh MG, Katsovich L, Thompson N, Lin H, Kim YS, Scahill L, et al. Severe psychosocial stress and heavy cigarette smoking during pregnancy: An examination of the pre- and perinatal risk factors associated with ADHD and Tourette syndrome. Eur Child Adolesc Psychiatry. 2010;19(10):755–64.
- 13. Visser SN, Bitsko RH, Danielson ML, Ghandour RM, Blumberg SJ, Schieve LA, et al. Treatment of attention deficit/hyperactivity disorder among

- children with special health care needs. J Pediatr [Internet]. Elsevier Inc; 2015;166(6):1423–30. e2. Available from: http://dx.doi.org/10.1016/j. jpeds.2015.02.018
- Chronis AM, Jones HA, Raggi VL. Evidencebased psychosocial treatments for children and adolescents with attention-deficit/hyperactivity disorder. Clin Psychol Rev. 2006;26(4):486–502.
- 15. Kleinman RE, Brown RT, Cutter GR, Dupaul GJ, Clydesdale FM. A research model for investigating the effects of artificial food colorings on children with ADHD. Pediatrics. 2011;127(6):e1575–84.
- 16. Trent S, Davies W. The influence of sexlinked genetic mechanisms on attention and impulsivity. Biol Psychol [Internet]. Elsevier B.V.; 2012;89(1):1–13. Available from: http://dx.doi. org/10.1016/j.biopsycho.2011.09.011
- 17. Spencer TJ, Biederman J, Mick E. Attention-Deficit/Hyperactivity Disorder: Diagnosis, Lifespan, Comorbidities, and Neurobiology. 2007;32(6):631–42.
- Oades RD. The Role of Serotonin in Attention-Deficit Hyperactivity Disorder (ADHD). In: Muller C, Jacobs B, editors. Handbook of Behavioral Neurobiology of Serotonin [Internet]. 1st ed. London: Elsevier B.V.; 2010. p. 565–84. Available from: http://linkinghub.elsevier.com/retrieve/pii/ S1569733910701016
- 19. Oades RD. Role of the serotonin system in ADHD: treatment implications. 2007;1357–74.
- Marshall R, Neill P, Theodosiou L. Prevalence of attention deficit hyperactivity symptoms in parents of children diagnosed with the condition. Procedia Soc Behav Sci [Internet]. Elsevier B.V.; 2011;15:3056–8. Available from: http://dx.doi.org/10.1016/j.sbspro.2011.04.244
- 21. T.D. B, F. M. Environmental risk factors for attention-deficit hyperactivity disorder. Acta Paediatr Int J Paediatr [Internet]. 2007;96(9):1269–74. Available from: http://ovidsp.ovid.com/ovidweb.cgi?T=JS & PAGE=reference & D=emed8 & NEWS=N & AN=2007414048
- 22. Bolfer C, Pacheco SP, Tsunemi MH, Carreira WS, Casella BB, Casella EB. Attention-deficit/hyperactivity disorder: The impact of methylphenidate on working memory, inhibition capacity and mental flexibility. Arq Neuropsiquiatr [Internet]. 2017;75(4):204–8. Available from:

- http://www.scielo.br/pdf/anp/v75n4/0004-282X-anp-75-04-0204.pdf\nhttp://ovidsp.ovid.com/ovidweb.cgi?T=JS & PAGE=reference & D=emex & NEWS=N & AN=615927137
- 23. Graham J, Seth S, Coghill D. Adhd. Medicine (Baltimore). 2007;35(3):181–5.
- 24. Karabekiroglu K, Yazgan YM, Dedeoglu C. Can we predict short-term side effects of methylphenidate immediate-release? Int J Psychiatry Clin Pract. 2008;12(1):48–54.
- 25. Wigal SB, Gupta S, Greenhill L, Posner K, Lerner M, Steinhoff K, et al. Pharmacokinetics of Methylphenidate in Preschoolers with Attention-Deficit/Hyperactivity Disorder. J Child Adolesc Psychopharmacol [Internet]. 2007;17(2):153–64. Available from: http://www.liebertonline.com/doi/abs/10.1089/cap.2007.0043
- Southammakosane C, Schmitz K. Pediatric Psychopharmacology for Treatment of ADHD, Depression, and Anxiety. Pediatrics [Internet]. 2015;136(2):351–9. Available from: http://pediatrics.aappublications.org/cgi/doi/10.1542/peds.2014-1581
- 27. Kollins SH. ADHD, substance use disorders, and psychostimulant treatment: Current literature and treatment guidelines. J Atten Disord. 2008;12(2):115–25.
- 28. Iris RFÆ, Sam MÆ. ADHD, Temperament, and Parental Style as Predictors of the Child 's Attachment Patterns. 2006;103–14.
- 29. Biederman J, Spencer T, Wilens T. S P E C IA L Evidence-based pharmacotherapy for attention-deficit hyperactivity disorder. 2018;(July):77–97.
- 30. Li JJ, Lee SS. Negative emotionality mediates the

- association of 5-HTTLPR genotype and depression in children with and without ADHD. Psychiatry Res [Internet]. Elsevier; 2014;215(1):163–9. Available from: http://dx.doi.org/10.1016/j.psychres.2013.10.026
- 31. Cadoret RJ, Langbehn D, Caspers K, Troughton EP, Yucuis R, Sandhu HK, et al. Associations of the Serotonin Transporter Promoter Polymorphism With Aggressivity, Attention Deficit, and Conduct Disorder in an Adoptee Population. 2003;44(2):88–101.
- 32. Dalley JW, Roiser JP. Dopamine, serotonin and impulsivity. Neuroscience [Internet]. IBRO; 2012;215:42–58. Available from: http://dx.doi.org/10.1016/j.neuroscience.2012.03.065
- 33. Aktepe E, Ozkorumak E, Tanriover-Kandil S. Pregnancy and delivery complications and treatment approach in attention deficit hyperactivity disorder. Turk J Pediatr [Internet]. 2009;51(5):478–84. Available from: http://www.turkishjournalpediatrics.org/pediatrics/pdf/pdf_TJP_693.pdf\nhttp://ovidsp.ovid.com/ovidweb.cgi?T=JS & PAGE=reference & D=emed9 & NEWS=N & AN=2010096213
- Richardson JR, Taylor MM, Shalat SL, Guillot TS, Caudle WM, Hossain MM, et al. Developmental pesticide exposure reproduces features of attention deficit hyperactivity disorder. FASEB J. 2015;29(5):1960–72.
- 35. Villagomez A, Ramtekkar U. Iron, Magnesium, Vitamin D, and Zinc Deficiencies in Children Presenting with Symptoms of Attention-Deficit/ Hyperactivity Disorder. Children [Internet]. 2014;1(3):261–79. Available from: http://www.mdpi.com/2227-9067/1/3/261/