

Journal Profile

Berkala Ilmu Kesehatan Kulit dan Kelamin (Periodical of Dermatology and Venerology)

eISSN : 25494082 | pISSN : 25494082

Health Science

[Universitas Airlangga](#)



S2

Sinta Score



Indexed by GARUDA

9

H-Index

8

H5-Index

518

Citations

466


5 Year Citations

p-ISSN 1978-4279

e-ISSN 2549-4082

Vol. 33 / No.3 / December 2021

Berkala Ilmu Kesehatan Kulit dan Kelamin



Periodical of Dermatology and Venereology



Vol. 33 No. 3 (2021): DECEMBER

Current Issue



Vol. 33 No. 3 (2021): DECEMBER

Published: 2021-11-30

Research / Retro

A Five-Year Review of Adverse Cutaneous Drug Reaction in a Tertiary Care Hospital in Yogyakarta, Indonesia

Dyah Ayu Mira Oktarina , Maria Sophiati , Erinda Maharani Rambu Moha

150-155

Abstract : 589

PDF : 268

PDF

DOI :
10.20473/bikk.V33.3.2021.150-155

Mucocutaneous Manifestations in HIV/AIDS Patients

Afif Nurul Hidayati , Citra Dwi Harningtyas , Damayanti Damayanti

156-161

Abstract : 393

PDF : 197

PDF

DOI :
10.20473/bikk.V33.3.2021.156-161

Prevention of Contact Dermatitis Due to Hand Hygiene in The Era of COVID-19

Damayanti Damayanti , Cita Rosita Sigit Prakoeswa , Sylvia Anggraeni

162-167

Abstract : 359

PDF : 172

PDF

DOI :
10.20473/bikk.V33.3.2021.162-167

The Profile of Type 1 Leprosy Reaction at Leprosy Division of Dermatology and Venereology

Outpatient Clinic of Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

Brigita Ika Rosdiana , Linda Astari , Astindari Astindari

168-172

Abstract : 380

PDF : 129

PDF

DOI :

10.20473/bikk.V33.3.2021.168-172

The Profile of Psoriasis Vulgaris Patients: A Descriptive Study

Cita Rosita Sigit Prakoeswa , Afif Nurul Hidayati , Made Putri Hendaria

173-181

Abstract : 617

PDF : 272

PDF

DOI :

10.20473/bikk.V33.3.2021.173-181

Susstance P Levels in Children with Atopic Dermatitis

Khairina Nasution , Deryne Anggia Paramita , Nova Zairina Lubis

182-186

Abstract : 283

PDF : 99

PDF

DOI :

10.20473/bikk.V33.3.2021.182-186

The Demography, Clinical Characteristics, and White Blood Analysis of Leprosy Reactions in Multibacillary Leprosy: A Retrospective Study

Natalia Tanojo , Damayanti Damayanti , Budi Utomo

187-193

Abstract : 310

PDF : 146

PDF

DOI :

10.20473/bikk.V33.3.2021.187-193

The Effect of Corncob (Zea mays) Extract Cream on the Number of Melanin Pigments of Guinea Pig Exposed to Ultraviolet

Pasid Harlisa , Sita Mahardika , Suryani Yuliyanti

194-199

Abstract : 456

PDF : 93

PDF

DOI :

10.20473/bikk.V33.3.2021.194-199

The Effect of Systemic Methotrexate and Cyclosporine Combination Therapy in Psoriasis Vulgaris Patients in Bandung, Indonesia

Okí Suwarsa , Fatima Aulia Khairani , Syawalika Ulya Isneny

200-204

Abstract : 343

PDF : 102

PDF

DOI :

10.20473/bikk.V33.3.2021.200-204

Literature Review

Phototherapy in Pediatric Dermatology

 Citra Dwi Harningtyas , Yuri Widia

 205-212

 Abstract : 424

 PDF : 207

 PDF

 DOI :
10.20473/bikk.V33.3.2021.205-212

Case Report

Chickenpox Mimicking Monkeypox in Adult with Diabetes Mellitus and Acute Kidney Injury: Diagnosis and Management

 Maya Wardiana , Rahmadewi Rahmadewi , Dwi Murtiastutik

 213-223

 Abstract : 259

 PDF : 88

 PDF

 DOI :
10.20473/bikk.V33.3.2021.213-223

The Confusion between Pustular Psoriasis and Acute Generalized Exanthematous Pustulosis as a Cause of Exfoliative Dermatitis: A Case Report

 Desiana Widityaning Sari , Cita Rosita Sigit Prakoeswa , Damayanti Damayanti

 224-231

 Abstract : 312

 PDF : 128

 PDF

 DOI :
10.20473/bikk.V33.3.2021.224-231

Focus and Scope	Publication Ethics
Peer Review Process	Article Processing Charge
Editorial Team	Open Access Policy
Archiving	Plagiarism
Copyright	Contact
Old Website	



Editorial Team

Editor in Chief

Prof. Dr. Cita Rosita Sigit Prakoeswa, dr., Sp.KK(K), FINS DV, FAADV, Faculty of Medicine, Airlangga University Surabaya, Indonesia, Indonesia

Editorial Manager

dr. Damayanti Damayanti, Sp.KK(K), FINS DV, Faculty of Medicine, Airlangga University Surabaya, Indonesia

Editorial Handling

dr. Irmadita Citrashanty, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Maylita Sari, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Septiana Widyantari, Sp.KK, FINS DV, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Medhi Denisa Alinda, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Hasnikmah Mappamasing, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Sylvia Anggraeni, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

Editorial Board

Dr. Afif Nurul Hidayati, dr., Sp.KK(K), FINS DV, FAADV, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Evy Ervianti, Sp.KK(K), FINS DV, FAADV, Faculty of Medicine, Airlangga University Surabaya, Indonesia, Indonesia, Indonesia

dr. Linda Astari, Sp.KK(K), FINS DV, FAADV, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Astindari Astindari, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

Prof. Dr. med. Isaak Effendy, Academic Teaching Hospital Bielefeld Klinikum Bielefeld, Germany
Dermatology and Allergology, Germany

Prof. dr. Taruna Ikrar, M.Pharm, MD., Ph.D, Biomedical Science, The National Health University,
California, United States

dr. Sri Manovita Pateda, S.Ked., M.Kes, Ehime University, Japan, Japan

dr. Jimmy Rusdian Masjkur, M.Sc., Ph.D, University Hospital Carl Gustav Carus Dresden, Germany,
Germany

Robert Tungadi, S.Si., M.Si., Apt, Aachen University, Germany, Germany

dr. David Sudarto Oeiria, Sp.KK, FINSVD, FAADV, Division of Dermatologic Surgery, Dept. of
Dermatology, Huashan Hospital - Fudan University, Shanghai, China

Assistant Editor

dr. Yuri Widia, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Menul Ayu Umborowati, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Bagus Haryo Kusumaputra, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Putri Hendria Wardhani, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Regitta Indira Agusni, Sp.KK, Faculty of Medicine, Airlangga University Surabaya, Indonesia

Reviewer

dr. Lili Legiwati, Sp.KK(K), Faculty of Medicine, Indonesia University, Jakarta, Indonesia

Prof. Hari Sukanto, Faculty of Medicine, Airlangga University, Surabaya, Indonesia

dr. Reti Hindritiani, Faculty of Medicine, Padjajaran University, Bandung, Indonesia

dr. Lilik Norawati, Sp.KK, RSPAD Gatot Soebroto, Jakarta, Indonesia

Robert Tungadi, S.Si., M.Si., Apt, Aachen University, Germany, Germany

dr. Jimmy Rusdian Masjkur, M.Sc., Ph.D, University Hospital Carl Gustav Carus Dresden, Germany,
Germany

dr. Sri Manovita Pateda, S.Ked., M.Kes, Ehime University, Japan, Japan

Prof. dr. Taruna Ikrar, M.Pharm, MD., Ph.D, Biomedical Science, The National Health University, California, United States

Prof. Dr. med. Isaak Effendy, Academic Teaching Hospital Bielefeld Klinikum Bielefeld, Germany
Dermatology and Allergology, Germany

Dr. Afif Nurul Hidayati, dr., Sp.KK(K), FINS DV, FAADV, Faculty of Medicine, Airlangga University Surabaya, Indonesia

dr. Diah Mira Indramaya, Departemen/Staf Medik Fungsional Ilmu Kesehatan Kulit dan Kelamin Fakultas Kedokteran Universitas Airlangga/Rumah Sakit Umum Daerah Dr. Soetomo Surabaya, Indonesia

dr. Abraham Arimurko, Sp.KK, RSPAD Gatot Soebroto, Jakarta, Indonesia

dr. Asmaja D Soedarwoto, Sp.KK(K), RS dr. Hasan Sadikin, Bandung, Indonesia

dr. Lis Surachmiati Suseno, Sp.KK, Faculty of Medicine, Indonesia University/RS Premier Jatinegara, Jakarta, Indonesia

dr. Dhiana Ernawati, Sp.KK(K), Faculty of Medicine, Diponegoro University/RS Dr. Kariadi, Semarang, Indonesia


trisniartami setyaningrum, Indonesia

rahmadewi rahmadewi, Departemen Staf Medik Fungsional Ilmu Kesehatan Kulit dan Kelamin Fakultas Kedokteran Universitas Airlangga, Rumah Sakit Umum Daerah Dr. Soetomo, Indonesia

Dr. Nelva Karmila Jusuf, dr., Sp.KK(K), Faculty of Medicine, Sumatera Utara University, Medan, Indonesia

Focus and Scope	Publication Ethics
Peer Review Process	Article Processing Charge
Editorial Team	Open Access Policy
Archiving	Plagiarism
Copyright	Contact

The Profile of Psoriasis Vulgaris Patients: A Descriptive Study

Cita Rosita Sigit Prakoeswa¹, Afif Nurul Hidayati¹, Made Putri Hendaria¹ , M. Yulianto Listiawan¹, Budi Utomo², Damayanti¹, Irmadita Citrashanty¹, Sylvia Anggraeni¹, Menul Ayu Umborowati¹, Evy Ervianti¹

¹Departement of Dermatology and Venereology, Faculty of Medicine Universitas Airlangga/Dr. Soetomo General Academic Hospital, Universitas Airlangga Teaching Hospital, Surabaya, Indonesia

²Department of Public Health Sciences, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

ABSTRACT

Background: Psoriasis is a skin disease that occurs due to proliferation and differentiation disturbances of the epidermis. This chronic condition often reduces the human quality of life. There have been studies investigating psoriatic patient profiles and treatments, but few are conducted in Surabaya. Hence, those studies cannot fully represent the population demographic in Surabaya. **Purpose:** To investigate the prevalence of psoriasis vulgaris and the sociodemographic profile of the patients in Dr Soetomo General Academic Hospital. **Methods:** This was a Retrospective study psoriasis vulgaris patients in January 2016–December 2018. The data were collected by retrieving Electronic Medical Records (EMR) as secondary data. **Result:** The psoriasis vulgaris prevalence of outpatients was 0.46% and 4.59% in inpatients. Most study subjects were adults aged 18 and above (98.1%), most of the patients were obese (46.3%), the most common provoking factor was focal tooth infection (33.8%) and stress (28.8%), and none reported family history of psoriasis. **Conclusion:** Some results are similar to other studies conducted in Surabaya, which were the obesity and Type 2 Diabetes Mellitus (DM) being strong a factor to psoriasis development. However, the result on psoriasis family history was not in line with other studies.

Keywords: psoriasis vulgaris, patient profile, severity, treatment, human and disease.

Correspondence: Afif Nurul Hidayati, Department of Dermatology and Venereology Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital/Universitas Airlangga, Surabaya, Jl. Mayjen Prof. Dr. Moestopo No. 6-8 Surabaya 60131, Indonesia. Phone: (031) 5501609, e-mail: afif_nurulhidayati@fk.unair.ac.id.

Article info | Submitted: 2-6-2021, Accepted: 20-7-2021, Published: 30-11-2021

BACKGROUND

Psoriasis is an inflammatory skin disease with chronic immunologic mediation, and it is often found with characteristics such as epidermal hyperplasia, increased arthritis risk, cardiovascular morbidity, and psychosocial problems.¹ It is marked by erythematous plaque covered with thick white squamous layers that occur due to proliferation and differentiation disturbances of the epidermis.^{1,2} It is a multifactorial disease in which pathogenesis is not fully understood but is thought to relate to genetic predisposition, autoimmune disease, and other environmental factors.^{1,3} It is also thought to arise from abnormal interactions of the innate immune system in which the cells involved are keratinocytes, dendritic cells, T lymphocytes, neutrophils, and mast cells.^{1,4}

The estimated prevalence of adult psoriasis ranges from 0.51% to 11.43%, increasing globally.^{5,6} In Indonesia, a retrospective study by Gayatri shows an equal predilection between males and females. The disease could occur in various stages of life but

mainly occurred in 15 to 40 years old, rarely found in ages less than ten years old.⁷ The many types of psoriasis, psoriasis vulgaris is the most common and is observed in 90% of patients.^{1,8} Surabaya is the second-largest city in Indonesia, second to the capital city of Jakarta. As of January 2019, it is home to 3 million people. Despite being populous, there are still established epidemiological data regarding psoriasis vulgaris that represent the demographic of people in Surabaya.⁹

A thorough physical examination is required for diagnosis because laboratory examination does not usually present specific outcomes. Additionally, histopathologic studies are needed in diagnosing complex cases.^{1,10} In choosing a therapeutic regiment, it is important to consider the scale and severity of the lesion.^{1,11,12} Severity of psoriasis can be scored with Psoriasis Area and Severity Index (PASI), which combines severity and area of lesions in a single index score.¹³ Successful treatment means 75% lower of the initial PASI score, known as PASI 75.¹⁰ A number of

topical that are available such as topical corticosteroid, vitamin D analogue, tazarotene, and calcineurin inhibitor.^{1,14} Furthermore, systemic agents include cyclosporine A, methotrexate, acitretin fumarate, hydroxyurea, acid esters, 6-thioguanine, sulfasalazine, mycophenolate mofetil, apremilast, and tofacitinib.¹ Another therapeutic mechanism that can be combined with the former includes phototherapy, which thins T cells selectively, especially those on the epidermis, prompting apoptosis and shifting of response from Th1 to Th2.^{1,8}

This retrospective study is aimed to evaluate the profiles of psoriasis vulgaris patients of Dr Soetomo General Academic Hospital in Surabaya, ranging from their general profiling, disease severity, and prescribed treatment.

RESULT

Table 1. Characteristics of psoriasis vulgaris patients

Variables	2016 n(%)	2017 n(%)	2018 n(%)	Total n(%)
Outpatient				
Sex				
Male	1 (33.3)	4 (33.3)	22 (55.0)	27 (50.0)
Female	2 (66.7)	8 (66.7)	17 (45.0)	27 (50.0)
Age				
< 18 years	0 (0)	1 (7.7)	0 (0)	1 (1.9)
≥ 18 years	3 (100)	12 (92.3)	39 (100)	54 (98.1)
Job				
Housewife	1 (100)	5 (41.7)	10 (25.6)	16 (29.6)
Civil servants	1 (100)	0 (0)	9 (23.1)	10 (18.5)
Private employees	1 (100)	6 (50)	16 (41)	23 (42.6)
Student	0 (0)	1 (8.3)	0 (0)	1 (1.9)
Entrepreneur	0 (0)	0 (0)	4 (10,3)	4 (7.4)
BMI				
<18.5 kg/m ²	0 (0)	3 (25)	0 (0)	3 (5.6)
18.5–22.9 kg/m ²	0 (0)	5 (41.7)	23 (59)	28 (51.9)
≥ 23.0 kg/m ²	3 (100)	4 (33.3)	16 (41)	23 (42.6)
Predilection areas				
Scalp	0 (0)	2 (16.7)	5 (12.9)	7 (13.0)
Upper extremities	3 (100)	11 (91.7)	36 (92.3)	50 (92.6)
Lower extremities	3 (100)	12 (100)	35 (89.7)	50 (92.6)
Back and chest	3 (100)	12 (100)	34 (87.1)	49 (90.7)
Provoking factors				
Tooth focal infection	2 (66.7)	2 (16.7)	16 (41)	20 (37)
ENT focal infection	0 (0)	5 (41.6)	4 (10.3)	9 (16.7)
Stress	1 (33.3)	4 (33.3)	13 (33.3)	18 (33.3)
Low compliance	0 (0)	1 (8,3)	4 (10.3)	5 (9.3)
Unknown	0 (0)	0 (0)	2 (5.1)	2 (3.7)
Family history				
Yes	0 (0)	0 (0)	0 (0)	0 (0)
No	3 (100)	12 (100)	39 (100)	54 (100)
Biopsy result				

METHODS

This research was a retrospective descriptive study of psoriasis vulgaris involving inpatients and outpatients of Dr Soetomo General Academic Hospital, Departement of Dermatology and Venereology. The secondary data were obtained through the electric medical record (EMR) of psoriasis vulgaris patients admitted from January 2016 to December 2018.

The inclusion criterion was medical records containing PASI scores before and after treatment. Data collected include general sex, age, body mass index (BMI), predilected areas, provoking factors, family history, and biopsy results. Ethical clearance was obtained from the Ethical Committee of Dr Soetomo General Academic Hospital Surabaya with reference number: 1979/KEPK/IV/2020.

Fitting psoriasis	0 (0)	7 (58.3)	31 (79.4)	38 (70.4)
Not fitting	2 (66.7)	5 (41.7)	6 (15.6)	13 (24)
No result	1 (33.3)	0 (0)	2 (5)	3 (5.6)
Inpatient				
Sex				
Male	4 (80)	9 (56.3)	3 (60)	16 (61.5)
Female	1 (20)	7 (43.8)	2 (40)	10 (38.5)
Age				
< 18 years	0 (0)	2 (12.5)	1 (20)	3 (11.5)
≥ 18 years	5 (100)	14 (87.5)	4 (80)	23 (88.5)
Job				
Housewife	1 (20)	5 (31.3)	1 (20)	7 (26.9)
Civil servants	1 (20)	1 (6.3)	0 (0)	2 (7.7)
Private employees	2 (40)	6 (37.5)	2 (40)	10 (48.5)
Student	0 (0)	1 (6.3)	0 (0)	2 (7.7)
Entrepreneur	1 (20)	2 (12.5)	0 (0)	3 (11.5)
Others	0 (0)	0 (0)	1 (20)	1 (3.8)
Unemployed	0 (0)	1 (6.3)	0 (0)	1 (3.8)
BMI				
<18.5 kg/m ²	0 (0)	1 (6.3)	1 (20)	2 (7.7)
18.5–22.9 kg/m ²	1 (20)	5 (31.3)	3 (60)	9 (34.6)
≥ 23.0 kg/m ²	4 (80)	10 (62.3)	1 (20)	15 (57.7)
Predilection areas				
Scalp	3 (60)	5 (31.3)	3 (60)	11 (42.3)
Upper extremities	4 (80)	16 (100)	5 (100)	25 (96.2)
Lower extremities	5 (100)	16 (100)	5 (100)	26 (100)
Back and chest	5 (100)	16 (100)	5 (100)	26 (100)
Behing	1 (20)	1 (6.3)	0 (0)	2 (7.7)
Provoking factors				
Tooth focal infection	2 (40)	4 (25)	1 (20)	7 (26.7)
ENT focal infection	1 (20)	3 (18.8)	0 (0)	4 (15.4)
Stress	1 (20)	3 (18.8)	1 (20)	5 (19.2)
Low compliance	0 (0)	3 (18.8)	1 (20)	4 (15.4)
Unknown	2 (40)	3 (18.8)	2 (40)	7 (26.9)
Family history				
Yes	0 (0)	0 (0)	0 (0)	0 (0)
No	5 (100)	16 (100)	5 (100)	26 (100)
Biopsy result				
Fitting psoriasis	4 (80)	13 (81.3)	3 (60)	20 (76.9)
Not fitting	1 (20)	3 (18.8)	2 (40)	6 (23.1)
No result	0 (0)	0 (0)	0 (0)	0 (0)

BMI: Body Mass Index

ENT : Ear, Nose, Throat

The table above shows that the distribution of sex of outpatient was balanced. Most of them were older than 18 years old (98.1%), and most of them worked as private employees (42.6%) and housewives (29.6%). More than half of the outpatients have a normal BMI. The majority of the patients have predilections in the upper extremities, lower extremities, and back and chest, with 92.6%, 92.6%, and 90.7%, respectively. The most often provoking


factors are tooth focal infection and stress. All outpatients have no family history of psoriasis.

For inpatients, 88.5% of them were 18 years old and above. Most inpatients were males (61.5%), and most of their occupation was private employees, followed by housewives. More than half have excessive weight. The lesions mainly occurred around the lower extremities, back, and chest (100% each), followed by upper extremities (96,2%). The most provoking factors were tooth focal infection and

stress. None have a family history of psoriasis. A small portion of psoriatic outpatient (24%) and inpatient (23.1%) have biopsy result that does not fit with psoriasis pictures. The diagnosis is based upon clinical exam, while biopsy functions assist the diagnosis in atypical or doubtful cases. Three biopsy

results of outpatients are not found on the record, which may be suspected to them refusing biopsy or having biopsied somewhere else.

Type 2 DM is the most common comorbidity inpatients and outpatients, presenting 30.8% and 29.6%, respectively, followed by dyslipidemia



DM: Diabetes Mellitus

Figure 1. Comorbidity of in- and outpatients of psoriasis vulgaris in Dermatology and Venereology Department of Dr. Soetomo General Academic Hospital Period January 2016 – December 2018.




Figure 2. In- and outpatient psoriasis vulgaris degree of severity before and after treatment in Dermatology and Venereology Department of Dr. Soetomo General Academic Hospital Period January 2016 – December 2018.

Almost all (96.3%) outpatient subjects have severe psoriasis, and no patient has mild psoriasis at the beginning of the regimen. By the end, most outpatient issues have moderate psoriasis (48.1%). On the other hand, 100% of inpatient subjects had severe psoriasis at the beginning of the treatment which 19.2% revert to moderate psoriasis, leaving 80.8% of severe psoriasis inpatients at the end. Thirty-five outpatients out of 54 achieved PASI 75, success treatment, by the end of the treatment, while only one

inpatient out of twenty-six could accomplish the same.

For outpatients subjects, 92.6% received a combination of topical and systemic therapy because all outpatients have moderate and severe psoriasis and no mild psoriasis. Four severe psoriasis patients have topical, systemic, and phototherapy regimens. Since all inpatients started with severe psoriasis (PASI>10), all of them were treated with a combination of topical and systemic therapy.

Table 2. Success rate of PASI 75 on psoriasis vulgaris treatment

PASI 75	2016 n(%)	2017 n(%)	2018 n(%)	Total n(%)
Outpatient				
Achieved	2 (66.7)	11(91.7)	22 (56.4)	35(64.8)
Unachieved	1 (33.3)	1 (98.3)	17(43.6)	19(35.2)
Inpatient				
Achieved	0(0)	0(0)	1(20)	1(3.8)
Unachieved	5(100)	16(100)	4(80)	25(96.2)

PASI : Psoriasis Area and Severity Index

Table 3. Type of therapy for psoriasis vulgaris

Type	2016 n(%)	2017 n(%)	2018 n(%)	Total n(%)
Outpatient				
Topical and systemic	3 (100)	12 (100)	35 (89.7)	50 (92.6)
Topical, systemic, and phototherapy	0 (0)	0 (0)	4 (10.3)	4 (7.4)
Inpatient				
Topical and systemic	5(100)	16(100)	5(100)	26(100)

Table 4. Type of topical therapy of psoriasis vulgaris

Type	2016 n(%)	2017 n(%)	2018 n(%)	Total n(%)
Outpatient				
Vaseline album	0 (0)	6 (50)	14(61.5)	30(55.6)
Urea 10% cream	1(33.3)	9 (75)	19(48.7)	29(53.7)
Hydrocortisone 2.5% cream	0 (0)	1 (8.3)	5 (12.8)	6 (11.1)
Momethasone furoate 0.1% cream	1(33.3)	9(75)	28(71.8)	38(70.4)
Desoximethasone 0.25% cream	3(100)	12(100)	36(92.3)	51(94.4)
Clobetasol proprionate 0.05% cream	0 (0)	1 (8.3)	1 (2.6)	2 (3.7)
Desonide Lotion	0 (0)	2(16.7)	11(28.2)	13(24.1)
Inpatient				
Oleum coccus	0 (0)	0 (0)	2 (40)	2 (7.7)
Vaseline album	2 (40)	14(87.5)	2 (40)	18(69.2)
Urea 10% cream	1 (20)	1 (6.3)	0 (0)	2 (7.7)
Hydrocortisone 2.5% cream	0 (0)	0 (0)	1 (20)	1 (3.8)
Momethasone furoate 0.1% cream	4 (80)	8 (50)	1 (20)	13 (50)
Desoximethasone 0.25% cream	5(100)	14(87.5)	4 (80)	23(88.5)
Clobetasol proprionate 0.05% cream	0(0)	1 (6.3)	0 (0)	1 (3.8)
Desonide Lotion	2 (40)	3(18.8)	0 (0)	5 (19.2)

Table 5. Type of systemic therapy of psoriasis vulgaris

Type	2016 n(%)	2017 n(%)	2018 n(%)	Total n(%)
Outpatient				
Methotrexate	2(66.7)	8(66.7)	29(74.4)	39(72.2)
Cyclosporine	0 (0)	1 (8.3)	0 (0)	1 (1.9)
Folic acid	3 (100)	11(91.7)	32(82.1)	46(85.2)
CTM	0 (0)	0 (0)	2 (5.1)	2 (3.7)
Loratadin	0 (0)	0 (0)	14(35.9)	14(25.9)
Cetirizine	2 (66.7)	12(100)	35(89.7)	49(90.7)
Inpatient				
Methotrexate	5 (100)	10(62.5)	5 (100)	20(76.9)
Cyclosporine	0 (0)	4 (25)	0 (0)	4 (15.4)
Folic acid	4 (80)	9(56.3)	2 (40)	15(57.7)
CTM	3 (60)	2(12.5)	1 (20)	6 (23.1)
Cetirizine	4 (80)	13(81.3)	4 (80)	21(80.8)

CTM : Chlorphenamine maleate

Table 6. Methotrexate doses on psoriasis vulgaris treatment

MTX (/week)	2016 n(%)	2017 n(%)	2018 n(%)	Total n(%)
Outpatient				
2.5 mg	0 (0)	0 (0)	3 (10.3)	3 (7.7)
5 mg	1 (50)	0 (0)	1 (3.4)	2 (5.1)
7.5 mg	0 (0)	0 (0)	1 (3.3)	1 (2.6)
10 mg	0 (0)	0 (0)	3 (10.3)	3 (2.6)
12.5 mg	0 (0)	2 (25)	3 (10.3)	5 (12.8)
15 mg	1 (50)	6 (75)	7 (58.6)	24(61.5)
25 mg	0 (0)	0 (0)	1 (3.4)	1 (2.6)
Inpatient				
2.5 mg	0 (0)	0 (0)	1 (20)	1 (5)
7.5 mg	0 (0)	0 (0)	1 (20)	1 (5)
10 mg	0 (0)	1 (10)	0 (0)	1 (5)
15 mg	5 (100)	8 (80)	3 (60)	16 (80)
17.5 mg	0 (0)	1 (10)	0 (0)	1 (5)

MTX : Methotrexate

DISCUSSION

The average psoriasis vulgaris prevalence in the outpatient Department of Dermatology and Venerology of Dr Soetomo General Academic Hospital, Surabaya, between January 2016-December 2018 was 0.19%, with an increasing trend of prevalence from 2016 to 2018. Additionally, the prevalence rate of psoriasis vulgaris in the inpatient department was 3.98%, corresponding to the global prevalence of 0.09%–11.4%.¹⁵ The number has not been compared with the prevalence in Indonesia due to a lack of data. In 2007-2008, studies conducted in Jakarta, Indonesia, reported that the psoriasis prevalence was 1.73%, with most (82.98%) were psoriasis vulgaris.¹⁶ The distribution of sex in outpatient subjects was balanced, while in inpatient

subjects, it was dominated by males. Globally, studies found that the sex distribution of psoriasis patients is similar.¹⁵

Most of the outpatients were 18 years old and above. It also corresponds to numerous findings that psoriasis is more prevalent in the adult population, which increases with age. The average onset of psoriasis is 33 years old. Three-fourths of them occur before the age of 46.¹⁵ The predilection location in this study shows to be within the upper and lower extremities and back and chest. These predilection locations are typical of the psoriasis type. Psoriasis vulgaris is typically localized in the extensors extremities, head, lumbosacral, buttocks, and genitalia.¹

Factors that provoke psoriasis in outpatients and inpatients the most are focal tooth infection, stress, and ENT infections. Several studies show that endogenous and exogenous factors could provoke psoriasis. Those factors include skin aggression, infection, alcohol and smoking, stress, drugs, hormonal changes, and allergy.^{1,15,17}

All study subjects have no family history of psoriasis. It is contradictory to many other previous studies which reported that the percentage of psoriasis history in the family is relatively high amongst psoriasis patients; 17.5% in Egypt, 45.9% in Italy, 40.7% in Spain, 28.6% in the Maghreb, 23% in China, and 23.1% in Malaysia.¹⁸ The inheritance pattern of psoriasis is best described as multifactorial. The psoriasis concordance rate ranges from 35% to 73%, which decreases nearer to the equator, suggesting the possible involvement of UV light as an environmental factor.¹ The absence of family history in the study may be caused by an undetected case in the patient's family or other sociocultural factors deeming sickness a private matter or may be due to Indonesia being along the equatorial line.

Some of the subjects had unfitting histopathological results to psoriasis. It might be caused by the diagnosis firstly being made based on clinical examination. Still, the histopathology exam was done to assist the diagnosis of atypical and doubtful cases.¹⁹

Type 2 DM and dyslipidemia were the most common comorbidity of psoriasis patients. Others were obesity, spondyloarthritis/arthropathy, hypertensive heart disease, and depression. Often, it is for psoriasis to be accompanied by comorbidities. Some classic examples are psoriatic arthritis, inflammatory bowel disease, psychiatric and psychological problems, as well as uveitis. Some of these may be caused by the genetic fragility of the psoriatic patients, while the condition itself greatly affects the quality of life and person confidence. This disease is related to anxiety and depression, with 30% and 60%, respectively.²⁰

Obesity, hypertriglyceridemia, and hyperglycemia also increase with psoriasis degree of severity, independent of other components of metabolic syndrome.²⁰

In a cohort study for a systemic therapy for psoriasis patients that involves non-psoriatic patients, the hazard ratio is 1.53 (after adjustment to age, sex, diabetes, hypertension, dyslipidemia, and smoking). Chronic inflammation of psoriasis produces Th1 and Th17 that promote systemic inflammation. Proinflammatory cytokines stimulate the hypothalamus-hypophysis axis, which relates to

central obesity, hypertension, and insulin resistance. In other words, psoriasis worsens obesity, diabetes, thrombosis, and atherosclerosis.²⁰ Obesity is an independent risk factor for psoriasis as the risk increases with increasing BMI. Social isolation, poor dietary habits, depression, alcoholism, and sedentary lifestyle in psoriasis patients may explain why psoriasis can cause obesity.²¹ Further supporting evidence is shown by a cross-sectional study in the UK. Numerous epidemiological studies have concrete proof that obesity is a predisposing factor for psoriasis and worsens psoriasis inflammation.²² Some studies show positive effects of reducing body weight with reduced severity of psoriasis.²¹

At the time of admission, almost all psoriatic outpatients have severe psoriasis (PASI>10). By the time of therapy, 48.1% have moderate psoriasis (PASI 5–10), and 25.9% have mild psoriasis (PASI<5). PASI 75 success is achieved by 64.8% of outpatients. Although, the target of the therapy is based on several parameters, not only skin symptoms but also the quality of life. However, PASI 75 is currently used as a basis to measure primary outcomes of clinical studies for psoriasis. The goal remains at the complete recovery of skin lesions and 90% improvement or PASI 90, especially amongst severe psoriatic patients. Patients who do not improve at least 50% in week 12 or 16 from the initial PASI score is called non-responder.²³

All psoriatic inpatients had a PASI score of more than 10, indicating a severe condition. However, after the therapy, only a small portion whose severity decreased to moderate with PASI 5-10 (19.2%). The percentage of subjects achieving PASI 75 is only 3.8%. Inpatient subjects were more severe than outpatients at the beginning of their care, and the management of their condition was 'emerging'. Hence once they show improvement, they are discharged for home treatment. All inpatients subjects got systemic therapy of methotrexate or cyclosporin. PASI score evaluation in the systemic treatment of psoriasis needs to be done after at least 12 weeks, but their length of stay is shorter. It may explain why inpatient subjects do not achieve PASI 75. It was due to the limited time window.²³

Three therapy modalities for psoriasis vulgaris are topical, systemic, and phototherapy. Topical medicine is for mild patients. For moderate to severe cases, a combination of topical and systemic are used. The majority of the outpatients received combination therapy (92.6%). A small population gets a variety of the three modalities (7.4%). All inpatients subjects received combination therapy. Methotrexate and cyclosporin were the most used topical therapy for

both inpatients and outpatients. Cyclosporine was the alternative if treatment with methotrexate did not show clinical improvement. Methotrexate is a folic acid analogue that hinders the formation of nucleic acid in activated T cells and keratinocytes, hence becoming an antiproliferative agent and an immunomodulator in psoriasis.¹ The drug is prescribed in divided doses to increase efficacy and minimize side effects.²⁴

There are patterns in the profiles of psoriatic patients. Some are similar to other findings, such as the proportional relationship between obesity and psoriasis prevalence, as explained in an international study by de Oliveira, Kunz, and Takeshita. They reported that obesity is related to the inflammation process. The lack of family history in all subjects in this study contradicts other findings suggesting a strong familial relationship to the disease. Despite so, this may be due to a lack of testing or lack of awareness. The success of a treatment regimen depends on several factors such as degree of lesion severity, genetics, environment, pharmacokinetics, and pharmacodynamics.

REFERENCES

- Gudjonsson J, Elder J. Psoriasis. In: Kang S, Amagai M, Bruckner A, Enk A, Margolis D, editors. *Fitzpatrick Dermatology in General Medicine*. 9th ed. New York: McGraw Hill Education 2019. p. 457–97.
- Luo Y, Ru Y, Sun X, Zhou Y, Yang Y, Ma T. Characteristics of psoriasis vulgaris in China: a prospective cohort study protocol. *Am Transl Med* 2019;7(22):694.
- Kerkhof P, Nestle F. Psoriasis. In: Bologna J, Jorizzo J, Schagger J, editors. *Dermatology* 2. 3rd ed. Philadelphia: Elsevier 2012. p. 135–55.
- Michalek I, Loring B, John S. A systematic review of worldwide epidemiology of psoriasis. *J Eur Acad Dermatol Venereol* 2017;31(5):205–12.
- Sinniah B, Saraswathy S, Prashant S. Epidemiology of psoriasis in Malaysia: a hospital-based study. *Med J Malaysia* 2010;65(1):112–4.
- Zhang J. Epidemiology and risk factors of psoriasis. *Pr J Clin Med* 2013;1(2):4–6.
- Gayatri L. Studi retrospektif: Psoriasis pustulosa generalisata. *Berkala Ilmu Kesehatan Kulit dan Kelamin* 2014;26(1):49–55.
- Widaty S, Soebono H, Nilasari H, Listiawan M, Siswati A, Triwahyudi D, et al. Psoriasis. In: Iidaty S, Soebono H, Nilasari H, Listiawan M, Siswati A, Triwahyudi D, et al., editors. *Panduan praktik klinis bagi dokter spesialis kulit dan kelamin di Indonesia*. 2nd ed. Jakarta: Perhimpunan Dokter Spesialis Kulit dan Kelamin Indonesia 2017. p. 230–40.
- Dispenduk capil. Jumlah penduduk kota Surabaya [Internet] 2019. Available from: <http://dispendukcapil.surabaya.go.id/berita/483-jumlah-penduduk-kota-Surabaya>.
- Novianto E, Irawan A, Wiryadi B, Prakoeswa C, Kariosentono H, Roesyani I, et al. Alur tata laksana psoriasis (tipe plak). In: Novianto E, Fitri E, Buadanti W, editors. *Alur tata laksana psoriasis vulgaris (tipe plak) di Indonesia*. 4th ed. Jakarta: Perhimpunan Dokter Spesialis Kulit dan Kelamin Indonesia 2019. p. 1–3.
- Wong HR, Freishtat RJ, Monaco M, Odoms K, Shanley TP. Leukocyte subset-derived genome-wide expression profiles in pediatric septic shock. *Pediatr Crit Care Med* 2010;11(3):349–55.
- Palfreeman AC, McNamee KE, McCann FE. New developments in the management of psoriasis and psoriatic arthritis: A focus on apremilast. *Drug Des Dev Ther* 2013;7(2):201–10.
- Haag D, Sundstrom A, Eriksson M, Egenolf M. Severity of psoriasis differs between men and women: a study of the clinical outcome measure psoriasis area and severity index (PASI) In 5438 Swedish Register Patients. *Am J Clin Dermatol* 2017;1(1):1–5.
- Jacob T. Psoriasis. In: Menaldi S, Bramono K, Indriatmi W, editors. *Ilmu penyakit kulit dan kelamin*. Jakarta: Badan Penerbit Fakultas Kedokteran Universitas Indonesia 2015. p. 213–21.
- World Health Organization. Global report on psoriasis 2016; 1(1): 1-10
- Setyorini, M Triestianawati W, Wiryadi B, Jacob T. Proporsi sindrom metabolik pada pasien psoriasis vulgaris berdasarkan kriteria national cholesterol education program adult treatment panel III di RS Dr. Cipto Mangunkusumo dan sebuah klinik swasta di Jakarta. *MDVI* 2012;39(1):2–9.
- Xhaja A, Shkodrani E, Frangaj S, Kuneshka L, Vasili E. An epidemiological study on trigger factors and quality of life in psoriatic patients. *Mater Socio Medica* 2014;26(3):168–71.
- El-Komy M, Mashaly H, Sayed K, Hafez V, El-Mesidy M, Said E et al. Clinical and epidemiologic features of psoriasis patients in an Egyptian medical centre. *JAAD Int*

- 2020;1(2):81–90.
19. Ferreli C, Pinna A, Piloni L, Tomasini C, Rongioletti F. Histopathological aspects of psoriasis and its uncommon variants. *Genital Di Dermatologia e Venereol* 2018;153(2):173–84.
 20. Rocha B de O, Duarte G. Psoriasis: classical and emerging comorbidities. *A Bras Dermatol* 2015;90(1):9–20.
 21. Kunz M, Simon J, Saalback A. Psoriasis: obesity and fatty acids. *Front Immunol* 2019;10(5):180–7.
 22. Takeshita J, Grewal S, Langan S, Mehta N, Ogdie A, Voorhes A. Psoriasis and comorbid diseases epidemiology. *J Am Acad Dermatology* 2016; 12(1): 377–91.
 23. Gisondi P, Del Giglio M, Girolomoni G. Treatment approaches moderate to severe psoriasis. *Int J Mol Sci* 2017;18(11):2427.
 24. Panduan Pelayanan Klinis (PPK). Psoriasis vulgaris. SMF kesehatan kulit dan kelamin RSUD Dr. Soetomo Surabaya 2016;1(1): 1-5