



**HASIL VALIDASI DAN PENILAIAN**  
**KARYA ILMIAH DOSEN FAKULTAS KEDOKTERAN UNIVERSITAS AIRLANGGA**  
**(MEDIA PUBLIKASI KARYA ILMIAH : JURNAL ILMIAH INTERNASIONAL)**  
**NOMOR : 32 /UN3.1.1/BPRE/ II/2023**

**FORM 15**

**A. Identitas Karya Ilmiah**

- Judul Jurnal Ilmiah (Artikel) : **Combination of platelet rich plasma and stromal vascular fraction on the level of vascular endothelial growth factor in rat subjects experiencing deep dermal burn injury .**
- Jumlah penulis : 10 (sepuluh) orang, SachraswatyRachmanLaiding, FonnyJosh, SartianBattung, AgussalimBukhari, Warsinggih, IlhamJayaPatellongi, MuhNassrumMassi, AndiAsadullIslam, **IswinarnoDososaputro**, MuhammadFaruk
- Status Pengusul : Penulis Ke. 9
- Identitas Jurnal Ilmiah
- a. Nama Jurnal : Annals of Medicine and Surgery
  - b. Nomor ISSN : 20490801
  - c. Volume, Nomor, bulan, tahun : 64 /2021,
  - d. Penerbit : Elsevier
  - e. Doi artikel : doi.org/10.1016/j.amsu.2021.102254
  - f. Alamat web Jurnal :  
<https://reader.elsevier.com/reader/sd/pii/S2049080121002041?token=CCB0514DF6E077F3F4F0A3BD4845B39BB19396A920A2E790535A906502A0BA4970DAF6A5148C283BEBBE914AEF20CBDB&originRegion=eu-west-1&originCreation=20210405022525>
  - g. Terindeks di Scimagojr/Thomson Reuter ISI Knowledge atau di :  
**Q3 (2021) 0,37 H.Index.: 30, Coverage : 2012-2021,**

**B. Kategori Publikasi Karya Ilmiah (beri lipada kategori yang tepat)**

- :  Jurnal Ilmiah Internasional Bereputasi (SJR > 0,10)  
 Jurnal Ilmiah Internasional Bereputasi (SJR < 0,10)  
 Jurnal Ilmiah Internasional terindeks di Web of Science clarivate analytics / kelompok emerging sources citation indeks (tidak terindeks SJR)

**C. Hasil Validasi Ketua Departemen**

Telah diperiksa dan divalidasi dengan baik, dan sampai pernyataan ini dibuat sebagai karya ilmiah **original / plagiat\***, sehingga kami turut bertanggung jawab bahwa karya ilmiah tersebut telah memenuhi syarat kaidah ilmiah, norma akademik, dan norma hukum, sesuai dengan Peraturan Menteri Pendidikan Nasional Nomor 17 Tahun 2010 tanggal 16 Agustus 2010 tentang Pencegahan dan Pananggulangan Plagiat di Perguruan Tinggi. Namun demikian, apabila di kemudian hari ternyata terbukti bahwa karya ilmiah tersebut merupakan karya Ilmiah Plagiat, maka akan menjadi tanggung jawab mutlak penulis tersebut di atas, baik secara perdata maupun pidana. Demikian surat pernyataan ini saya buat untuk dipergunakan sebagaimana mestinya.

Surabaya, 10 Pebruari 2023

Ketua Departemen  
Bedah Plastik Rekonstruksi dan Estetik

Prof. Dr. David S. Perdanakusuma, dr., SpBP-RE(K)  
NIP. 19600305 198901 1 002

## Form Penilaian Kualitas Karil dan Kesesuaian Bidang Ilmu

Profil Sinta : Link Sinta <https://sinta.kemdikbud.go.id/profile>

A		Identitas Karya Ilmiah
1	Judul :	“Combination of platelet rich plasma and stromal vascular fraction on the level of vascular endothelial growth factor in rat subjects experiencing deep dermal burn injury.”
2	Nama Penulis :	SachraswatyRachmanLaidding, FonnyJosh, SartianBattung, AgussalimBukhari, Warsinggih, IlhamJayaPatellongi, MuhNassrumMassi,AndiAsadulIslam, IswinarnoDososaputro, MuhammadFaruk
3	Nama Jurnal :	Annals of Medicine and Surgery
B		Peng-index :
		<b>Q3 (2021) 0,37 H.Index. 30, Coverage : 2012- to present</b> ISSN. 20490801 , Publisher : Elsevier Vol.64 Tahun 2021, 102254 Doi : //doi.org/10.1016/j.amsu.2021.102254
C		Relevansi kompetensi dosen dengan substansi karya ilmiah
		1. Tujuan dari penelitian ini adalah untuk mengetahui pengaruh kombinasi PRP dan SVF pada tingkat VEGF dalam model tikus penyembuhan luka bakar dermal dalam. 2. Artikel ini sesuai dengan bidang keahlian pengusul yaitu <b>Luka Bakar</b> 3. Tidak ada keterkaitan dengan naskah Disertasi pengusul yang berjudul : <b>Mekanisme hambatan Deposisi Koleagen Tipe 1 Pada Luka Bakar Derajat 2 Dalam yang diberi Ekstrak Buah Delima Terstandar pad tikus Putih (<i>Rattus Norvegicus</i>)</b>
D		Kesesuaian antara lingkup / subjek area jurnal dengan karya ilmiah yang diusulkan
		1. Alamat Web Jurnal : <a href="https://www.sciencedirect.com/science/article/pii/S2049080121002041">https://www.sciencedirect.com/science/article/pii/S2049080121002041</a> 2. Kebenaran ISSN : 20490801 3. Termasuk "Predatory" tidak (jurnal; penerbit) : <b>tidak masuk pada predatory</b> 4. Syarat komposisi Editor Board : <b>lebih dari 4 negara</b> 5. Syarat kontributor penulis artikel : <b>Penulis anggota</b> 6. Keberkalaan penerbitan : 12 terbitan pertahun 7. <b>Subjek area dan katagori jurnal</b> : Medicine, Medicine (Miscellaneous), Surgery
E		Kepastian tidak ada pelanggaran integritas akademik
		1. Indikasi plagiasi (lihat check similarity) : Similarity 22 % artikel tidak ada indikasi plagiasi. 2. Fabrikasi : Tidak ada

		3. Falsifikasi : Tidak ada 4. Praktek kepalsuan : Tidak ada
	Nilai pengusul (penulis pertama dan corespondensi 60% )	
	Nilai pengusul (penulis pertama / penulis corespondensi masing - masing 40%)	
	Nilai lainnya sesuai PO PAK 2019 dan suplemennya	

Surabaya,

Penilai Angka Kredit I

Nama :

NIP :

Bidang Ilmu :

Unit Kerja :



## Experimental Research

## Combination of platelet rich plasma and stromal vascular fraction on the level of vascular endothelial growth factor in rat subjects experiencing deep dermal burn injury

Sachraswaty Rachman Laididing<sup>a</sup>, Fonny Josh<sup>a</sup>, Sartian Battung<sup>b</sup>, Agussalim Bukhari<sup>c</sup>, Warsinggih<sup>b</sup>, Ilham Jaya Patellongi<sup>d</sup>, Muh Nassrum Massi<sup>e</sup>, Andi Asadul Islam<sup>f</sup>, Iswinarno Dososaputro<sup>g</sup>, Muhammad Faruk<sup>b,\*</sup>

<sup>a</sup> Division of Plastic and Reconstructive Surgery, Department of Surgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>b</sup> Department of Surgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>c</sup> Department of Clinical Nutrition, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>d</sup> Department of Biostatistics, Faculty of Public Health, Hasanuddin University, Makassar, Indonesia

<sup>e</sup> Departement of Microbiology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>f</sup> Department of Neurosurgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>g</sup> Department of Plastic and Reconstructive Surgery, Airlangga University, Dr. Soetomo Hospital, Surabaya, Indonesia



## ARTICLE INFO

## Keywords:

Vascular endothelial growth factor

Platelet-rich plasma

Stromal vascular fraction

Rats

Deep dermal burn

## ABSTRACT

**Background:** The healing process of burns includes coagulation, inflammation, and remodeling. Vascular endothelial growth factor (VEGF) is involved throughout this healing process. Stem cells from the platelet-rich plasma (PRP) with stromal vascular fraction (SVF) can increase concentrations of growth factors, including VEGF. This is expected to accelerate burn healing. The aim of this study was to determine the effect of a combination of PRP and SVF on VEGF levels in a rats model of deep dermal burn wound healing.

**Materials and methods:** This is an experimental research study in rats using a post-test control group design with 4 groups: A) control, B) Vaseline, C) topical PRP and SVF, and D) PRP and SVF injection. Burn wounds were induced according to the modified Guo method.

**Results:** In a rats model of deep dermal wound healing, topical Vaseline significantly increased serum VEGF compared to control. Topical application and injection of stem cells also significantly increased serum VEGF compared to control and Vaseline. The VEGF concentration was significantly higher following injection of PRP and SVF, suggesting that the injection route is more effective at increasing VEGF levels compared to the topical application of stem cells.

**Conclusion:** The combination of PRP and SVF, either by injection or topical application, can increase VEGF levels during the healing process from deep dermal burns.

## 1. Introduction

A burn is defined by injury on the skin or underlying tissue caused by radiation, heat, cold, chemicals, or electricity [1–3]. Many researches have been performed to discover the best method in solving the problem of burns and accelerating the burn healing process. One of them is stem cell therapy [4]. Stem cells are undifferentiated primitive cells that have

the ability to differentiate from just one type of cell (unipotent), or into several types of cells (multipotent). They can even become various types of cells (totipotent). This ability can be utilized to repair body cells damaged by disease or trauma [5].

The stromal vascular fraction (SVF) is a collection of lipoaspirate components derived from fat tissue liposuction [6–9]. Lipoaspirate contains a large number of adipose-derived stem cells (ASCs). SVFs from

\* Corresponding author. Department of Surgery, Faculty of Medicine, Hasanuddin University, Jalan Perintis Kemerdekaan KM 11, Makassar, 90245, Indonesia.

E-mail addresses: [sachras1276@gmail.com](mailto:sachras1276@gmail.com), [sachras1276@gmail.com](mailto:sachras1276@gmail.com) (S.R. Laididing), [fonnyjosh2003@yahoo.com](mailto:fonnyjosh2003@yahoo.com), [fonnyjosh2003@yahoo.com](mailto:fonnyjosh2003@yahoo.com) (F. Josh), [battungsartian@gmail.com](mailto:battungsartian@gmail.com) (S. Battung), [agussalimbukhari@yahoo.com](mailto:agussalimbukhari@yahoo.com) (A. Bukhari), [kbd.warsinggih@gmail.com](mailto:kbd.warsinggih@gmail.com) (Warsinggih), [ilham\\_pt@yahoo.com](mailto:ilham_pt@yahoo.com) (I.J. Patellongi), [nasrum2000@yahoo.com](mailto:nasrum2000@yahoo.com) (M.N. Massi), [andiasadul@yahoo.com](mailto:andiasadul@yahoo.com) (A.A. Islam), [iswinarno@yahoo.com](mailto:iswinarno@yahoo.com) (I. Dososaputro), [farox8283@gmail.com](mailto:farox8283@gmail.com) (M. Faruk).

<https://doi.org/10.1016/j.amsu.2021.102254>

Received 2 March 2021; Received in revised form 21 March 2021; Accepted 22 March 2021

Available online 26 March 2021

2049-0801/© 2021 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

fat tissue contain T cell regulators, endothelial precursor cells, pre-adipocytes, anti-inflammation macrophages, superoxide dismutase (SOD), IGF, TGF, FGF, hepatocyte growth factor (HGF), interleukin (IL), hematopoietic stem and progenitor cells, erythrocytes, fibroblasts, lymphocytes, monocyte/macrophages and pericytes are found in the SVF [10–13]. SVFs are known to promote burn healing by increasing cell proliferation and vascularization, enhancing inflammation, and increasing fibroblast activity [14,15].

Platelet-rich plasma (PRP) is a concentrate of thrombocytes in a small volume of plasma. It include growth factors, vascular endothelial growth factor (VEGF), hepatocyte growth factor (HGF), epidermal growth factor (EGF), basic fibroblast growth factor (bFGF), platelet-derived growth factor (PDGF), transforming growth factor- $\beta$  (TGF- $\beta$ ), and insulin-like growth factor-1 (IGF-1), which are secreted after thrombocyte activation [13,15–21].

VEGF is key signal used by oxygen-hungry cells to stimulate blood vessel growth [13]. VEGF is produced in large amounts by keratinocytes, macrophages, endothelial cells, thrombocytes, and fibroblasts during wound healing. Cell disruption and hypoxia, signs of tissue damage, are strong initial inducers of angiogenic factors at the wound site, such as VEGF and its receptor [22,23]. VEGF-A promotes the initial phase of angiogenesis and is important for wound healing. It binds the surface receptors tyrosine kinase Flt-1 (VEGF-1 receptor, or VEGFR-1) and KDR (VEGF-2 receptor, or VEGFR-2). Flt-1 is needed for blood vessel organization, whereas KDR is essential for chemotaxis, proliferation, and endothelial cell differentiation [24]. We aimed to investigate the combination of PRP and SVF can increase the level of VEGF compared to control and Vaseline-treated rats with deep dermal burns.

## 2. Methods

This was an experimental research study using rats with a post-test control group design comprise a control group and 3 experimental groups (sacrificed on Day 1, 4, 7, 10 and 14 post experiment) conducted in the animal laboratory in our institutions over 2 weeks. The subjects were adult male rats (*Rattus norvegicus*), aged 10 weeks and weighing 150–250 g, obtained from the animal laboratory in our institution. A total of 64 rats were divided into 4 groups. Group A was the negative control in which samples were collected from the blood of healthy rats without burns or treatment (negative control). The 3 experimental groups were: Group B (Vaseline, topical), Group C (PRP and SVF, topical), and Group D (PRP and SVF, injected). Groups B, C, and D consisted of 5 groups of 4 rats, respectively. Group A was a group of 4 rats. This study was conducted in the animal laboratory Faculty of Medicine, Hasanuddin University, and received approval from our local Ethics Commission registration number: 216/UN4.6.4.5.31/PP36/2020. The work was also carried out in line with the ARRIVE guidelines for reporting animal research [25,26].

**Table 1**  
Average change in VEGF level in each group.

Termination Day	Group A (mean $\pm$ SD) (pg/ml)	Group B (mean $\pm$ SD) (pg/ml)	Group C (mean $\pm$ SD) (pg/ml)	Group D (mean $\pm$ SD) (pg/ml)	*p value	**p value
1	65.2 $\pm$ 0.7	70.1 $\pm$ 1.5	75.5 $\pm$ 4.0	83.7 $\pm$ 5.0	0.086	<0.001
4	65.2 $\pm$ 0.7	75.7 $\pm$ 5.1	86.3 $\pm$ 4.3	96.1 $\pm$ 5.6	0.051	<0.001
7	65.2 $\pm$ 0.7	75.8 $\pm$ 4.3	87.0 $\pm$ 5.2	96.4 $\pm$ 4.0	0.046	<0.001
10	65.2 $\pm$ 0.7	77.8 $\pm$ 4.1	89.8 $\pm$ 4.8	98.7 $\pm$ 3.9	0.166	<0.001
14	65.2 $\pm$ 0.7	78.5 $\pm$ 2.4	88.4 $\pm$ 7.2	98.6 $\pm$ 3.7	0.016	<0.001

\*Levene test \*\*One-way Anova test.

### 2.1. PRP and SVF preparations

For PRP preparations, rats were first shaved on their backs and then anesthetized using ether inhalation. The stem cell donor group underwent a thoracotomy until the heart was exposed. The cardiac apex identification was then conducted by aspirating blood from the cardiac. The blood drawn was transferred to a tube containing EDTA. Briefly, blood was centrifuged for 10 min at 2400 rpm (450 g) for the first centrifugation. The supernatant plasma with a buffy coat was mixed then for 15 min and again centrifuged at a speed of 3600 rpm (850 g). The infranatant buffy coat was used to prepare the final PRP product [6, 27].

SVF preparations were derived from fat tissue of donor rats collected by incising the groin area of the four limbs using a scalpel. Then, the fat was washed with phosphate buffer salt and chopped until smooth and transferred to a 15 cc tube. Then, 0.15% collagenase was added to the tube containing fat and incubated for 30 min at 37 °C. To neutralize collagenase activation, Dulbecco Modified Eagle Media (DMEM) with 10% FBS and 1% antibiotic-aticimotic was added and then samples were centrifuged at 1,500 rpm for 5 min. Cell pellets were resuspended with aquadest then treated with Trypan blue dye to count the number of SVF cells in the Neubauer counting room [27,28]. A total of 50,000 SVF cells mixed with 0.5 aquadest were transferred to the Eppendorf tube for the final SVF product.

The combination of PRP and SVF was obtained by mixing 0.5 cc PRP with 50,000 SVF cells [27].

### 2.2. Deep dermal burn model

In each treatment group, hair was removed in the treatment area using the chemical Jolen (Veet®). Prior to burn injury, rats were anesthetized in a special box using ether inhalation until the state of awareness decreased. The wound area was disinfected with 1% povidone-iodine. Burn wounds were induced according to the modified Guo method of exposure to a hot aluminum metal which its size 10  $\times$  10 mm previously heated in 100 °C water [29]. After modeling the burns, the experimental animals were orally administered an analgesic (sodium dipiron/metamizole, 50 mg/kg BW) and antibiotic (amoxycillin, 15 mg/kg BW) for three consecutive days [29,30].

### 2.3. Wound analysis

Skin tissue was subjected to histopathological examination to analyzed deep dermal burns using hematoxylin and eosin (H&E) staining. The histopathological slides were analyzed using a light microscope (Leica Microsystems).

### 2.4. Vaseline and combination stem cell treatment

Vaseline (petroleum jelly) was used as a moisturizer in the burn wound area [31]. In group B, topical Vaseline was applied to the burn area. In group C, the topical combination of PRP and SVFs was applied to the middle of the wound and the four edges of the wound at 6, 9, 12, 3 for a 50,000 SVF cells and total volume of 0.5 cc PRP per experimental rats. In group D, rats received the combination of PRP and SVFs subcutaneously injected into the middle of the wound and the four edges of the wound at 6, 9, 12, 3 for a 50,000 SVF cells and total volume of 0.5 cc PRP per experimental rats.

### 2.5. VEGF measurements

ELISAs were performed on days 1, 4, 7, 10, and 14 days following treatment. All blood samples were examined using the VEGF ELISA from MyBioSource, Inc. (catalog No: MBS724516) [6,32].

**Table 2**  
Comparison of VEGF levels between groups A and B.

Termination Day-	Group A (Mean ± SD) (pg/ml)	Group B (Mean ± SD) (pg/ml)	*p value
1		70.1 ± 1.5	0.001
4		75.7 ± 5.1	0.026
7	65.2 ± 0.7	75.8 ± 4.3	0.003
10		77.8 ± 4.1	0.008
14		78.5 ± 2.4	<0.001

Independent t-test.

### 2.6. Statistical analysis

A Levene's test, One-way ANOVA test and paired t-test were employed by using a combination of SPSS statistics 24 (IBM Corp., Armonk, NY) and Excel (Microsoft Corp, Redmond, WA). A p-value of less than 0.05 was considered statistically significant.

### 3. Results

In this study, 64 rats were divided into 4 experimental groups: A) negative control, B) topical Vaseline, C) topical PRP and SVF, and D) subcutaneously injected PRP and SVF. Groups B, C, and D consisted of 5 subgroups of 4 rats, respectively. Group A was a group of 4 rats.

Based on Table 1, we can conclude the mean value of changes in VEGF levels for 14 days in 4 treatment groups. The group treated with vaseline, topical PRP and SVFs, and injection PRP and SVFs showed an increase in VEGF levels from day one to day 10 and then decreased on day 14. This indicated that the peak increase in VEGF levels was on day 10. In the PRP and SVFs group, injection in each treatment day had the highest mean value among all groups. In the group treated with vaseline, PRP and SVFs topical, PRP and SVFs injection, and the control group, there were significant differences in changes of VEGF levels with each

value < 0.001.

#### 3.1. Comparison between group A (negative control) and group B (topical vaseline)

As shown in Table 2, VEGF levels in the Vaseline group were significantly different from the control group ( $p < 0.05$ ). Based on this result, H0 is rejected and H1 is accepted, suggesting there is a significant effect of Vaseline on VEGF levels in a rat model of deep dermal burns on days 1, 4, 7, 10, and 14. The VEGF levels with Vaseline were highest on day 14 (Fig. 1). This result is similar to a study that reported VEGF serum levels with Vaseline were lower than control on day 7, but higher than control on days 14 and 21 [16]. In another study comparing Vaseline to controls with an angiogenesis parameter, the researchers found that on days 7, 14, and 21, Vaseline was better than the control group [17]. In another study comparing VEGF levels in Pelnac, Vaseline, and a control group found that on day 7, Vaseline was better than control and Pelnac was significantly better than Vaseline [18].

**Table 3**  
Comparison of VEGF levels between groups B and C.

Termination Day-	Group B (Mean ± SD) (pg/ml)	Group C (Mean ± SD) (pg/ml)	*p value
1	70.1 ± 1.5	75.5 ± 4.0	0.047
4	75.7 ± 5.1	86.3 ± 4.3	0.020
7	75.8 ± 4.3	87.0 ± 5.2	0.017
10	77.8 ± 4.1	89.8 ± 4.8	0.010
14	78.5 ± 2.4	88.4 ± 7.2	0.041

\* Independent t-test.




Fig. 1. Comparison result of VEGF levels between groups A and B.




Fig. 2. Comparison of VEGF levels between groups B and C.

Table 4

Comparison of VEGF levels between groups B and D.

Termination Day-	Group B (Mean $\pm$ SD) (pg/ml)	Group D (Mean $\pm$ SD) (pg/ml)	<sup>a</sup> p value
1	70.1 $\pm$ 1.5	83.7 $\pm$ 5.0	0.002
4	79.0 $\pm$ 2.1	96.1 $\pm$ 5.6	0.002
7	77.8 $\pm$ 4.1	96.4 $\pm$ 4.0	<0.001
10	75.8 $\pm$ 4.3	98.7 $\pm$ 3.9	<0.001
14	75.7 $\pm$ 5.1	98.6 $\pm$ 3.7	<0.001

<sup>a</sup> Independent t-test.

### 3.2. Comparison between group B (topical vaseline) and group C (PRP and SVF, topical)

As shown in Table 3, the comparison of VEGF levels after topical PRP and SVF to Vaseline on each day was significantly different ( $p < 0.05$ ) (Fig. 2). Based on this result, H0 is rejected and H1 is accepted, suggesting that topical application of stem cells is more effective than Vaseline for increasing VEGF levels in a rat model of deep dermal burns on days 1, 4, 7, 10 and 14.

### 3.3. Comparison between group B (topical vaseline) and group D (PRP and SVF, injected)

As shown in Table 4, VEGF levels after stem cell injections were significantly increased compared with Vaseline on each day ( $p < 0.05$ ) (Fig. 3). Based on this result, H0 is rejected and H1 is accepted, suggesting that injections of stem cells are better able to increase VEGF levels compared with Vaseline in a rat model of deep dermal burns on days 1, 4, 7, 10 and 14.

### 3.4. Comparison between group C (PRP and SVF, topical) and group D (PRP and SVF, injected)

As shown in Table 5, VEGF levels were significantly increased after injection of stem cells and topical stem cells ( $p < 0.05$ ), and there was a significant difference between the two groups (Fig. 4). Based on this result, H0 is rejected and H1 is accepted, suggesting the injection of stem cells is better than topical stem cells in increasing VEGF levels in a rat model of deep dermal burns on days 1, 4, 7, 10 and 14.

## 4. Discussion

One of the factors that plays an important role in the wound healing process is angiogenesis, which is the process of forming new blood vessels [33,34]. One of the most important and widely studied angiogenic factors is vascular endothelial growth factor (VEGF) [34]. VEGF is one of the most potent mediators of vascular permeability and vascular regulation in angiogenesis. VEGF is also commonly called "vascular permeability factor," which refers to a specific endothelial cell mitogen secreted as a 45 kDa protein consisting of two subunits that do not induce cell proliferation in other cell types [24,34].

PRP contains many growth factors, including VEGF, HGF, bFGF, EGF, TGF- $\beta$ , IGF-1, and PDGF [35]. Angiogenesis is particularly stimulated by VEGF, PDGF and basic fibroblast growth factor (bFGF). Angiogenesis is also assisted by pericytes, which in turn depend on the availability of PDGF and VEGF [36,37]. Likewise, ADSC/SVF that was found contain many growth factors and cytokines, which in turn influence angiogenesis, as well as mononuclear infiltrates, fibroblast and production on collagen. VEGF, a potent angiogenic agents, migration of fibroblast and activity stimulators such as TGF, and macrophage chemotaxis agents are regulated by ADSCs [38].

The administration of PRP and SVF, either topically or by injection enhances VEGF levels up to day 10. This result was significant compared




Fig. 3. Comparison of VEGF levels between groups B and D.

Table 5

Comparison of VEGF levels between groups C and D.

Termination Day-	Group C (Mean ± SD) (pg/ml)	Group D (Mean ± SD) (pg/ml)	p value
1	75.5 ± 4.0	83.7 ± 5.0	0.045
4	86.3 ± 4.3	96.1 ± 5.6	0.034
7	87.0 ± 5.2	96.4 ± 4.0	0.030
10	89.8 ± 4.8	98.7 ± 3.9	0.029
14	88.4 ± 7.2	98.6 ± 3.7	0.047

<sup>a</sup> Independent *t*-test.

to Vaseline and control ( $p < 0.05$ ). Our findings are similar to a previous study that reported the benefits of topical PRP gel in treating soft tissue wounds, in which the recovery proceeded faster compared to Vaseline. PRP gel is not only beneficial in chronic wound treatment, but is also effective in acute injury [39]. Another study regarding the administration of the SVF and PRF combination showed significantly better results in the concentration of VEGF levels compared to SVF or PRP alone with VEGF levels peaking on day 7 and slowly declining on days 14 and 28 [40]. Additionally, another study showed that PRP and SVF injection increased VEGF levels on day 1 to day 7 then decreased on day 10 [41].

In our study, we found a significant difference between injection and topical PRP and SVF administration. PRP and SVF by injection is more effective at increasing VEGF levels compared with topical PRP and SVF. The main objective of PRP therapy is to centralize growth factors at the injury site. These factors are considered beneficial in the healing process in soft and hard tissue and are also considered the most effective way to provide a biological stimulus to the tissue [42]. In a previous *in vitro* study, it was stated that freezing and diluting PRP does not significantly affect VEGF release. There was no significant difference in VEGF levels after freezing for 1 h and 7 days. This study also concluded that kinetic differences in VEGF release are not significantly affected by PRP

preparation; further clinical studies could explore and clarify the PRP effect [35]. Based on our results, it can be concluded that the injection route is significantly more effective than topical application for increasing VEGF release.

In a study analyzing the musculoskeletal effect of PRP and SVF, the authors stated that direct injection of PRP to the joints can control the inflammatory response and promote long-term healing. The growth factor components in PRP have special anti-inflammatory effects and can also reduce pain. Anti-inflammation, immunomodulation, and analgesia are also influenced by soluble factors excreted by SVF or ADSCs, including HGF, VEGF, NGF, EGF, FGF and TGF- $\beta$  [43].

Inflammation regulation, proliferation of cells, remodeling, deposition of extracellular matrix, angiogenesis and epithelization play important role in skin wound healing. In the mouse skin, PRP and SVFs can reduce inflammatory process, increase collagen deposition, and promote angiogenesis and neurogenesis, thus promote wound healing [44]. PRP and adipose-derived stem cells combination is potent in stimulating increase of growth factor concentration (IGF-1, TGF- $\beta$  1, HGF, and VEGF) [27]. PRP is an accessible cell therapy to help treat chronic wounds and severe burns [16,19]. PRP is an autologous cell therapy containing a large number of bioactive factors involved in wound healing and tissue repair. PRP can be used for treating and stimulating wound healing in burns and split-thickness skin grafts, which are often used for the management of burns [45].

Based on this study, further research is needed regarding the effect of the combination of PRP and SVF on VEGF levels in dermal burn healing. Future studies could involve determining the quantity of neo-vascularization in burn healing through histopathological examination.

In addition, our results may be of interest to other researchers investigating the additive and synergistic effects of therapeutic strategies involving stem cells. The fact that the PRP and SVF efficacy data for burn injuries are limited highlights the need for phase I–II clinical trials. We hope that this research can serve as a fundamental resource and an






Fig. 4. Comparison of VEGF levels between groups C and D.

additional reference when making decisions about the application of PRP and SVF as a new breakthrough in the management of burn injuries.

## 5. Conclusion

In our study, we can conclude that the administration of PRP and SVF by injection can significantly increase VEGF levels more effectively than Vaseline in rats with induced deep dermal burns. Further, the administration of topical PRP and SVF significantly increased VEGF levels compared to the Vaseline group. There is a significant difference between PRP and SVF administration by injection and topical application. The injection of PRP and SVF is more effective at increasing VEGF levels compared to topical application.

## Provenance and peer review

Our study was non-commissioned and externally peer-reviewed.

## Ethical approval

All procedure for Animal experiment has been approved by Ethics Commission Faculty of Medicine, Hasanuddin University Number: 216/UN4.6.4.5.31/PP36/2020.

## Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Author contribution

SRL, FJ, SB, and MF wrote the manuscript and participated in the study design. SRL, FJ, SB, MF, ASB, WS, MNM, IDS, and AAI drafted and

revised the manuscript. SRL, FJ, SB, and MF performed deep dermal burn model. SRL, FJ, MF, and LJP performed bioinformatics analyses and revised the manuscript. All authors read and approved the final manuscript.

## Registration of research studies

None.

## Guarantor

Sachraswaty Rachman Laidding.

## Consent

This manuscript does not involve human participants, human data, or human tissue.

## Declaration of competing interest

The authors declare that they have no conflict of interests.

## Acknowledgment

A higher appreciation to all staff from the Hasanuddin University Medical Research Center (HUMRC), Makassar, Indonesia.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.amsu.2021.102254>.

## References

- [1] ANZBA, Emergency Management of Severe Burns (EMSB), eighteenth ed., THE EDUCATION COMMITTEE OF AUSTRALIA AND NEW ZEALAND BURN ASSOCIATION LTD, Albany Creek, 2016.
- [2] I. Kaddoura, G. Abu-Sittah, A. Ibrahim, R. Karamanoukian, N. Papazian, Burn injury: review of pathophysiology and therapeutic modalities in major burns, *Ann. Burns Fire Disasters* 30 (2017) 95–102, <http://www.ncbi.nlm.nih.gov/pubmed/29921720>.
- [3] M.A.R. Stokes, W.D. Johnson, Burns in the third world: an unmet need, *Ann. Burns Fire Disasters* 30 (2017) 243–246, <https://pubmed.ncbi.nlm.nih.gov/29983673>.
- [4] F. Ghieh, R. Jurjus, A. Ibrahim, A.G. Geagea, H. Daouk, B. El Baba, S. Chams, M. Matar, W. Zein, A. Jurjus, The use of stem cells in burn wound healing: a review, *BioMed Res. Int.* 2015 (2015) 684084, <https://doi.org/10.1155/2015/684084>.
- [5] V.K. Singh, A. Saini, M. Kalsan, N. Kumar, R. Chandra, Describing the stem cell potency: the various methods of functional assessment and in silico diagnostics, *Front. Cell Dev. Biol.* 4 (2016), <https://doi.org/10.3389/fcell.2016.00134>.
- [6] S.R. Laidling, F. Josh, Francisca, M. Faruk, A.S. Palissei, B. Satria, Warsinggih, A. Bukhari, M.N. Massi, A.A. Islam, Combination of platelet-rich plasma and stromal vascular fraction on the level of transforming growth factor- $\beta$  in rat subjects experiencing deep dermal burn injury, *Ann. Med. Surg.* 60 (2020) 737–742, <https://doi.org/10.1016/j.amsu.2020.11.085>.
- [7] P. Foubert, A.D. Gonzalez, S. Teodosescu, F. Berard, M. Doyle-Eisele, K. Yekkala, M. Tenenhaus, J.K. Fraser, Adipose-derived regenerative cell therapy for burn wound healing: a comparison of two delivery methods, *Adv. Wound Care* 5 (2016) 288–298, <https://doi.org/10.1089/wound.2015.0672>.
- [8] S. Atalay, A. Coruh, K. Deniz, Stromal vascular fraction improves deep partial thickness burn wound healing, *Burns* 40 (2014) 1375–1383, <https://doi.org/10.1016/j.burns.2014.01.023>.
- [9] J. Han, Y.J. Koh, H.R. Moon, H.G. Ryou, C.-H. Cho, I. Kim, G.Y. Koh, Adipose tissue is an extramedullary reservoir for functional hematopoietic stem and progenitor cells, *Blood* 115 (2010) 957–964, <https://doi.org/10.1182/blood-2009-05-219923>.
- [10] V. Tantuway, P. Bhambani, A. Nagla, P. Mantry, R. Sharma, P. Mehto, S. Bansal, Autologous grafting of non manipulated freshly isolated - adipose tissue derived stromal vascular fraction in single surgical sitting for treatment of knee osteoarthritis, *Int. J. Res. Orthop.* 3 (2016) 107, <https://doi.org/10.18203/issn.2455-4510.IntJResOrthop20164834>.
- [11] A. Darinskias, M. Paskevicius, G. Apanavicius, G. Vilkevicius, L. Labanauskas, T. E. Ichim, R. Rimdeika, Stromal vascular fraction cells for the treatment of critical limb ischemia: a pilot study, *J. Transl. Med.* 15 (2017) 143, <https://doi.org/10.1186/s12967-017-1243-3>.
- [12] K. Comella, R. Silbert, M. Parlo, Effects of the intradiscal implantation of stromal vascular fraction plus platelet rich plasma in patients with degenerative disc disease, *J. Transl. Med.* 15 (2017) 12, <https://doi.org/10.1186/s12967-016-1109-0>.
- [13] P. Gentile, M.G. Scioli, A. Bielli, A. Orlandi, V. Cervelli, Concise review: the use of adipose-derived stromal vascular fraction cells and platelet rich plasma in regenerative plastic surgery, *Stem Cell.* 35 (2017) 117–134, <https://doi.org/10.1002/stem.2498>.
- [14] P. Bourin, B.A. Bunnell, L. Casteilla, M. Dominici, A.J. Katz, K.L. March, H. Redl, J. P. Rubin, K. Yoshimura, J.M. Gimble, Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cytotherapy 15 (2013) 641–648, <https://doi.org/10.1016/j.jcyt.2013.02.006>.
- [15] J. Choi, K.W. Minn, H. Chang, The efficacy and safety of platelet-rich plasma and adipose-derived stem cells: an update, *Arch. Plast. Surg.* 39 (2012) 585–592, <https://doi.org/10.5999/aps.2012.39.6.585>.
- [16] E. Rapisio, N. Bertuzzi, S. Bonamini, G. Bernuzzi, A. Formentini, E. Grignattini, M. Pio Grieco, Adipose-derived stem cells added to platelet-rich plasma for chronic skin ulcer therapy: wounds a compend, *Clin. Res. Pract.* 28 (2016) 126–131.
- [17] P. Borriore, A. Di Gianfrancesco, M.T. Pereira, F. Pigozzi, Platelet-rich plasma in muscle healing, *Am. J. Phys. Med. Rehabil.* 89 (2010) 854–861, <https://doi.org/10.1097/PHM.0b013e3181f1c1c7>.
- [18] A. Duran, S. Yasar, P. Gunes, S. Aytelkin, A. Duran, Clinical and Histopathological Evaluation of the Effects of Platelet Rich Plasma, Platelet Poor Plasma and Topical Serum Physiologic Treatment on Wound Healing Caused by Radiofrequency Electrosurgery in Rats, 2016.
- [19] T. Stessuk, M.B. Puzzi, E.A. Chaim, P.C.M. Alves, E.V. de Paula, A. Forte, J. M. Izumizawa, C.C. Oliveira, F. Frei, J.T. Ribeiro-Paes, Platelet-rich plasma (PRP) and adipose-derived mesenchymal stem cells: stimulatory effects on proliferation and migration of fibroblasts and keratinocytes in vitro, *Arch. Dermatol. Res.* 308 (2016) 511–520, <https://doi.org/10.1007/s00403-016-1676-1>.
- [20] M. Tohidnezhad, D. Varoga, C.J. Wruck, L.O. Brandenburg, A. Seekamp, M. Shakibaei, T.T. Sönmez, T. Pufe, S. Lippross, Platelet-released growth factors can accelerate tenocyte proliferation and activate the anti-oxidant response element, *Histochem. Cell Biol.* 135 (2011) 453–460, <https://doi.org/10.1007/s00418-011-0808-0>.
- [21] R.P. Martins, D.D. Hartmann, J.P. de Moraes, F.A.A. Soares, G.O. Puntel, Platelet-rich plasma reduces the oxidative damage determined by a skeletal muscle contusion in rats, *Platelets* 27 (2016) 784–790, <https://doi.org/10.1080/09537104.2016.1184752>.
- [22] N.H. Mansoub, M. Gürdal, E. Karadadas, H. Kabadayi, S. Vatansver, G. Ercan, The role of PRP and adipose tissue-derived keratinocytes on burn wound healing in diabetic rats, *Bioimpacts* 8 (2018) 5–12, <https://doi.org/10.15171/bi.2018.02>.
- [23] N. Subramaniam, J.J. Petrik, M.K. Vickaryous, VEGF, FGF-2 and TGF $\beta$  expression in the normal and regenerating epidermis of geckos: implications for epidermal homeostasis and wound healing in reptiles, *J. Anat.* 232 (2018) 768–782, <https://doi.org/10.1111/joa.12784>.
- [24] C.S. Melincovici, A.B. Bosca, S. Şuşman, M. Murginean, C. Mihu, M. Istrate, I. M. Moldovan, A.L. Roman, C.M. Mihu, Vascular endothelial growth factor (VEGF)-key factor in normal and pathological angiogenesis, *Rom. J. Morphol. Embryol. - Rev. Roum. Morphol. Embryol.* 59 (2018) 455–467.
- [25] C. Kilkenny, W.J. Browne, I.C. Cuthill, M. Emerson, D.G. Altman, Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research, *PLoS Biol.* 8 (2010), e1000412, <https://doi.org/10.1371/journal.pbio.1000412>.
- [26] K.F. Schulz, D.G. Altman, D. Moher, CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials, *BMJ* 340 (2010) c332, <https://doi.org/10.1136/bmj.c332>, c332.
- [27] S. Tajima, M. Tobita, H. Orbay, H. Hyakusoku, H. Mizuno, Direct and indirect effects of a combination of adipose-derived stem cells and platelet-rich plasma on bone regeneration, *Tissue Eng.* 21 (2014), <https://doi.org/10.1089/ten.tea.2014.0336>.
- [28] F. Josh, K. Kobe, M. Tobita, R. Tanaka, K. Suzuki, K. Ono, H. Hyakusoku, H. Mizuno, Accelerated and safe proliferation of human adipose-derived stem cells in medium supplemented with human serum, *J. Nippon Med. Sch.* 79 (2012) 444–452, <https://doi.org/10.1272/jnms.79.444>.
- [29] H.-F. Guo, R.M. Abi, R.A. Hamid, A.A. Zaini, H. Khaza' ai, A new model for studying deep partial-thickness burns in rats, *Int. J. Burns Trauma.* 7 (2017) 107–114.
- [30] D. dos, S. Tavares Pereira, M.H.M. Lima-Ribeiro, N.T. de Pontes-Filho, A.M. dos, A. Carneiro-Leão, M.T. dos, S. Correia, Development of animal model for studying deep second-degree thermal burns, *J. Biomed. Biotechnol.* (2012) 460841, <https://doi.org/10.1155/2012/460841>, 2012.
- [31] A. Sethi, T. Kaur, S.K. Malhotra, M.L. Gambhir, Moisturizers: the slippery road, *Indian J. Dermatol.* 61 (2016) 279–287, <https://doi.org/10.4103/0019-5154.182427>.
- [32] Warsinggih, B. Irawan, I. Labeled, R.E. Lusikooy, S. Sampetoding, M.I. Kusuma, J. A. Uwuratuw, E. Syarifuddin, Prihantono, M. Faruk, Association of superoxide dismutase enzyme with staging and grade of differentiation colorectal cancer: a cross-sectional study, *Ann. Med. Surg.* 58 (2020) 194–199, <https://doi.org/10.1016/j.amsu.2020.08.032>.
- [33] R.J. Bodnar, Epidermal growth factor and epidermal growth factor receptor: the yin and yang in the treatment of cutaneous wounds and cancer, *Adv. Wound Care* 2 (2013) 24–29, <https://doi.org/10.1089/wound.2011.0326>.
- [34] K.E. Johnson, T.A. Wilgus, Vascular endothelial growth factor and angiogenesis in the regulation of cutaneous wound repair, *Adv. Wound Care* 3 (2014) 647–661, <https://doi.org/10.1089/wound.2013.0517>.
- [35] N. Hosny, F. Goubran, B. BadrEldin Hasan, N. Kamel, Assessment of vascular endothelial growth factor in fresh versus frozen platelet rich plasma, *J. Blood Transfus.* 2015 (2015) 1–5, <https://doi.org/10.1155/2015/706903>.
- [36] J.I. Greenberg, D.J. Shields, S.G. Barillas, L.M. Acevedo, E. Murphy, J. Huang, L. Schepke, C. Stockmann, R.S. Johnson, N. Angle, D.A. Cheresch, A role for VEGF as a negative regulator of pericyte function and vessel maturation, *Nature* 456 (2008) 809–813, <https://doi.org/10.1038/nature07424>.
- [37] N. Caporarello, F. D'Angeli, M.T. Cambria, S. Candido, C. Giallongo, M. Salmeri, C. Lombardo, A. Longo, G. Giurdanella, C.D. Anfufo, G. Lupo, Pericytes in microvessels: from "mural" function to brain and retina regeneration, *Int. J. Mol. Sci.* 20 (2019) 6351, <https://doi.org/10.3390/ijms20246351>.
- [38] A.L. Cardoso, M.M. Bachion, J. de M. Moraes, M.S. Fantinati, V.L.L. de Almeida, R. S. Lino Júnior, Adipose tissue stromal vascular fraction in the treatment of full thickness burns in rats, *Acta Cir. Bras.* 31 (2016) 578–585, <https://doi.org/10.1590/S0102-865020160090000032>.
- [39] K. Kazakos, D.N. Lyras, D. Verettas, K. Tilkeridis, M. Tryfonidis, The use of autologous PRP gel as an aid in the management of acute trauma wounds, *Injury* 40 (2009) 801–805, <https://doi.org/10.1016/j.injury.2008.05.002>.
- [40] Z.-J. Liang, X. Lu, D.-Q. Li, Y.-D. Liang, D.-D. Zhu, F.-X. Wu, X.-L. Yi, N. He, Y.-Q. Huang, C. Tang, H. Li, Precise intradermal injection of nanofat-derived stromal cells combined with platelet-rich fibrin improves the efficacy of facial skin rejuvenation, *Cell. Physiol. Biochem.* 47 (2018) 316–329, <https://doi.org/10.1159/000489809>.
- [41] N. Seyhan, D. Alhan, A.U. Ural, A. Gunal, M.C. Avunduk, N. Savaci, The effect of combined use of platelet-rich plasma and adipose-derived stem cells on fat graft survival, *Ann. Plast. Surg.* 74 (2015) 615–620, <https://doi.org/10.1097/SAP.0000000000000480>.
- [42] A.D. Mazzocca, M.B.R. McCarthy, D.M. Chowaniec, M.P. Cote, A.A. Romeo, J. P. Bradley, R.A. Arciero, K. Beitzel, Platelet-rich plasma differs according to preparation method and human variability, *J. Bone Jt. Surg.* 94 (2012) 308–316, <https://doi.org/10.2106/JBJS.K.00430>.
- [43] A. Fotouhi, A. Maleki, S. Dolati, A. Aghebati-Maleki, L. Aghebati-Maleki, Platelet rich plasma, stromal vascular fraction and autologous conditioned serum in treatment of knee osteoarthritis, *Biomed. Pharmacother.* 104 (2018) 652–660, <https://doi.org/10.1016/j.biopha.2018.05.019>.
- [44] T. Liu, C. Qiu, C. Ben, H. Li, S. Zhu, One-step approach for full-thickness skin defect reconstruction in rats using minced split-thickness skin grafts with Pelnac overlay, *Burn, Trauma* 7 (2019), <https://doi.org/10.1186/s41038-019-0157-0>.
- [45] K. Akhundov, G. Pietramaggiore, L. Waselle, S. Darwiche, S. Guerid, C. Scaletta, N. Hirt-Burri, L.A. Applegate, W. V Raffouf, Development of a cost-effective method for platelet-rich plasma (PRP) preparation for topical wound healing, *Ann. Burns Fire Disasters* 25 (2012) 207–213.

# Combination of platelet rich plasma and stromal vascular fraction on the level of vascular endothelial growth factor in rat subjects experiencing deep dermal burn injury

*by Sachraswaty Rachman Laiding*

---

**Submission date:** 01-May-2021 03:15PM (UTC+0800)

**Submission ID:** 1575197860

**File name:** factor\_in\_rat\_subjects\_experiencing\_deep\_dermal\_burn\_injury.pdf (1.41M)

**Word count:** 6002


**Character count:** 30479



ELSEVIER

Contents lists available at ScienceDirect

Annals of Medicine and Surgery

journal homepage: [www.elsevier.com/locate/amsu](http://www.elsevier.com/locate/amsu)

Experimental Research

## Combination of platelet rich plasma and stromal vascular fraction on the level of vascular endothelial growth factor in rat subjects experiencing deep dermal burn injury

Sachraswaty Rachman Laidding<sup>a</sup>, Fonny Josh<sup>a</sup>, Sartian Battung<sup>b</sup>, Agussalim Bukhari<sup>c</sup>, Warsinggih<sup>d</sup>, Ilham Jaya Patellongi<sup>d</sup>, Muh Nassrum Massi<sup>e</sup>, Andi Asadul Islam<sup>f</sup>, Iswinarno Dososaputro<sup>g</sup>, Muhammad Faruk<sup>b, \*</sup><sup>a</sup> Division of Plastic and Reconstructive Surgery, Department of Surgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia<sup>b</sup> Department of Surgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia<sup>c</sup> Department of Clinical Nutrition, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia<sup>d</sup> Department of Biostatistics, Faculty of Public Health, Hasanuddin University, Makassar, Indonesia<sup>e</sup> Department of Microbiology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia<sup>f</sup> Department of Neurosurgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia<sup>g</sup> Department of Plastic and Reconstructive Surgery, Airlangga University, Dr. Soetomo Hospital, Surabaya, Indonesia

### ARTICLE INFO

#### Keywords:

Vascular endothelial growth factor  
Platelet-rich plasma  
Stromal vascular fraction  
Rats  
Deep dermal burn

### ABSTRACT

**Background:** The healing process of burns includes coagulation, inflammation, and remodeling. Vascular endothelial growth factor (VEGF) is involved throughout this healing process. Stem cells from the platelet-rich plasma (PRP) with stromal vascular fraction (SVF) can increase concentrations of growth factors, including VEGF. This is expected to accelerate burn healing. The aim of this study was to determine the effect of a combination of PRP and SVF on VEGF levels in a rats model of deep dermal burn wound healing.

**Materials and methods:** This is an experimental research study in rats using a post-test control group design with 4 groups: A) control, B) Vaseline, C) topical PRP and SVF, and D) PRP and SVF injection. Burn wounds were induced according to the modified Guo method.

**Results:** In a rats model of deep dermal wound healing, topical Vaseline significantly increased serum VEGF compared to control. Topical application and injection of stem cells also significantly increased serum VEGF compared to control and Vaseline. The VEGF concentration was significantly higher following injection of PRP and SVF, suggesting that the injection route is more effective at increasing VEGF levels compared to the topical application of stem cells.

**Conclusion:** The combination of PRP and SVF, either by injection or topical application, can increase VEGF levels during the healing process from deep dermal burns.

### 1. Introduction

A burn is defined by injury on the skin or underlying tissue caused by radiation, heat, cold, chemicals, or electricity [1–3]. Many researches have been performed to discover the best method in solving the problem of burns and accelerating the burn healing process. One of them is stem cell therapy [4]. Stem cells are undifferentiated primitive cells that have

the ability to differentiate from just one type of cell (unipotent), or into several types of cells (multipotent). They can even become various types of cells (totipotent). This ability can be utilized to repair body cells damaged by disease or trauma [5].

The stromal vascular fraction (SVF) is a collection of lipoaspirate components derived from fat tissue liposuction [6–9]. Lipoaspirate contains a large number of adipose-derived stem cells (ASCs). SVFs from

\* Corresponding author. Department of Surgery, Faculty of Medicine, Hasanuddin University, Jalan Perintis Kemerdekaan KM 11, Makassar, 90245, Indonesia.

E-mail addresses: [sachras1276@gmail.com](mailto:sachras1276@gmail.com), [sachras1276@gmail.com](mailto:sachras1276@gmail.com) (S.R. Laidding), [fonnyjosh2003@yahoo.com](mailto:fonnyjosh2003@yahoo.com), [fonnyjosh2003@yahoo.com](mailto:fonnyjosh2003@yahoo.com) (F. Josh), [battungsartian@gmail.com](mailto:battungsartian@gmail.com) (S. Battung), [agussalimbukhari@yahoo.com](mailto:agussalimbukhari@yahoo.com) (A. Bukhari), [kbd.warsinggih@gmail.com](mailto:kbd.warsinggih@gmail.com) (Warsinggih), [ilham.pti@yahoo.com](mailto:ilham.pti@yahoo.com) (I.J. Patellongi), [nasrum2000@yahoo.com](mailto:nasrum2000@yahoo.com) (M.N. Massi), [andiasadul@yahoo.com](mailto:andiasadul@yahoo.com) (A.A. Islam), [iswinarno@yahoo.com](mailto:iswinarno@yahoo.com) (I. Dososaputro), [farox8283@gmail.com](mailto:farox8283@gmail.com), [farox8283@gmail.com](mailto:farox8283@gmail.com) (M. Faruk).

<https://doi.org/10.1016/j.amsu.2021.102254>

Received 2 March 2021; Received in revised form 21 March 2021; Accepted 22 March 2021

Available online 26 March 2021

2049-0801/© 2021 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license

<http://dx.doi.org/10.1016/j.amsu.2021.102254>

fat tissue contain T cell regulators, endothelial precursor cells, pre-adipocytes, anti-inflammation macrophages, superoxide dismutase (SOD), IGF, TGF, FGF, hepatocyte growth factor (HGF), interleukin (IL), hematopoietic stem and progenitor cells, erythrocytes, fibroblasts, lymphocytes, monocyte/macrophages and pericytes are found in the SVF [10–13]. SVFs are known to promote burn healing by increasing cell proliferation and vascularization, enhancing inflammation, and increasing fibroblast activity [14,15].

Platelet-rich plasma (PRP) is a concentrate of thrombocytes in a small volume of plasma. It include growth factors, vascular endothelial growth factor (VEGF), hepatocyte growth factor (HGF), epidermal growth factor (EGF), basic fibroblast growth factor (bFGF), platelet-derived growth factor (PDGF), transforming growth factor- $\beta$  (TGF- $\beta$ ), and insulin-like growth factor-1 (IGF-1), which are secreted after thrombocyte activation [13,15–21].

VEGF is key signal used by oxygen-hungry cells to stimulate blood vessel growth [13]. VEGF is produced in large amounts by keratinocytes, macrophages, endothelial cells, thrombocytes, and fibroblasts during wound healing. Cell disruption and hypoxia, signs of tissue damage, are strong initial inducers of angiogenic factors at the wound site, such as VEGF and its receptor [22,23]. VEGF-A promotes the initial phase of angiogenesis and is important for wound healing. It binds the surface receptors tyrosine kinase Flt-1 (VEGF-1 receptor, or VEGFR-1) and KDR (VEGF-2 receptor, or VEGFR-2). Flt-1 is needed for blood vessel organization, whereas KDR is essential for chemotaxis, proliferation, and endothelial cell differentiation [24]. We aimed to investigate the combination of PRP and SVF can increase the level of VEGF compared to control and Vaseline-treated rats with deep dermal burns.

## 2. Methods

This was an experimental research study using rats with a post-test control group design comprise a control group and 3 experimental groups (sacrificed on Day 1, 4, 7, 10 and 14 post experiment) conducted in the animal laboratory in our institutions over 2 weeks. The subjects were adult male rats (*Rattus norvegicus*), aged 10 weeks and weighing 150–250 g, obtained from the animal laboratory in our institution. A total of 64 rats were divided into 4 groups. Group A was the negative control in which samples were collected from the blood of healthy rats without burns or treatment (negative control). The 3 experimental groups were: Group B (Vaseline, topical), Group C (PRP and SVF, topical), and Group D (PRP and SVF, injected). Groups B, C, and D consisted of 5 groups of 4 rats, respectively. Group A was a group of 4 rats. This study was conducted in the animal laboratory Faculty of Medicine, Hasanuddin University, and received approval from our local Ethics Commission registration number: 216/UN4.6.4.5.31/PP36/2020. The work was also carried out in line with the ARRIVE guidelines for reporting animal research [25,26].

**Table 1**  
Average change in VEGF level in each group.

Termination Day.	Group A (mean ± SD) (pg/ml)	Group B (mean ± SD) (pg/ml)	Group C (mean ± SD) (pg/ml)	Group D (mean ± SD) (pg/ml)	*p value	**p value
1	65.2 ± 0.7	70.1 ± 1.5	75.5 ± 4.0	83.7 ± 5.0	0.096	<0.001
4	65.2 ± 0.7	75.7 ± 5.1	86.3 ± 4.3	96.1 ± 5.6	0.051	<0.001
7	65.2 ± 0.7	75.8 ± 4.3	87.0 ± 5.2	96.4 ± 4.0	0.046	<0.001
10	65.2 ± 0.7	77.8 ± 4.1	89.8 ± 4.8	98.7 ± 3.9	0.166	<0.001
14	65.2 ± 0.7	78.5 ± 2.4	88.4 ± 7.2	98.6 ± 3.7	0.016	<0.001

\*Levene test \*\*One-way Anova test.

### 2.1. PRP and SVF preparations

For PRP preparations, rats were first shaved on their backs and then anesthetized using ether inhalation. The stem cell donor group underwent a thoracotomy until the heart was exposed. The cardiac apex identification was then conducted by aspirating blood from the cardiac. The blood drawn was transferred to a tube containing EDTA. Briefly, blood was centrifuged for 10 min at 2400 rpm (450 g) for the first centrifugation. The supernatant plasma with a buffy coat was mixed then for 15 min and again centrifuged at a speed of 3600 rpm (850 g). The infranant buffy coat was used to prepare the final PRP product [6, 27].

SVF preparations were derived from fat tissue of donor rats collected by incising the groin area of the four limbs using a scalpel. Then, the fat was washed with phosphate buffer salt and chopped until smooth and transferred to a 15 cc tube. Then, 0.15% collagenase was added to the tube containing fat and incubated for 30 min at 37 °C. To neutralize collagenase activation, Dulbecco Modified Eagle Media (DMEM) with 10% FBS and 1% antibiotic-antimycotic was added and then samples were centrifuged at 1,500 rpm for 5 min. Cell pellets were resuspended with aquadest then treated with Trypan blue dye to count the number of SVF cells in the Neubauer counting room [27,28]. A total of 50,000 SVF cells mixed with 0.5 aquadest were transferred to the Eppendorf tube for the final SVF product.

The combination of PRP and SVF was obtained by mixing 0.5 cc PRP with 50,000 SVF cells [27].

### 2.2. Deep dermal burn model

In each treatment group, hair was removed in the treatment area using the chemical Jolen (Veet®). Prior to burn injury, rats were anesthetized in a special box using ether inhalation until the state of awareness decreased. The wound area was disinfected with 1% povidone-iodine. Burn wounds were induced according to the modified Guo method of exposure to a hot aluminum metal which its size 10 × 10 mm previously heated in 100 °C water [29]. After modeling the burns, the experimental animals were orally administered an analgesic (sodium dipiron/metamizole, 50 mg/kg BW) and antibiotic (amoxycillin, 15 mg/kg BW) for three consecutive days [29,30].

### 2.3. Wound analysis

Skin tissue was subjected to histopathological examination to analyzed deep dermal burns using hematoxylin and eosin (H&E) staining. The histopathological slides were analyzed using a light microscope (Leica Microsystems).

### 2.4. Vaseline and combination stem cell treatment

Vaseline (petroleum jelly) was used as a moisturizer in the burn wound area [31]. In group B, topical Vaseline was applied to the burn area. In group C, the topical combination of PRP and SVFs was applied to the middle of the wound and the four edges of the wound at 6, 9, 12, 3 for a 50,000 SVF cells and total volume of 0.5 cc PRP per experimental rats. In group D, rats received the combination of PRP and SVFs subcutaneously injected into the middle of the wound and the four edges of the wound at 6, 9, 12, 3 for a 50,000 SVF cells and total volume of 0.5 cc PRP per experimental rats.

### 2.5. VEGF measurements

ELISAs were performed on days 1, 4, 7, 10, and 14 days following treatment. All blood samples were examined using the VEGF ELISA from MyBioSource, Inc. (catalog No: MBS724516) [6,32].

**Table 2**  
Comparison of VEGF levels between groups A and B.

Termination Day-	Group A (Mean ± SD) (pg/ml)	Group B (Mean ± SD) (pg/ml)	*p value
1		70.1 ± 1.5	0.001
4		75.7 ± 5.1	0.026
7	65.2 ± 0.7	75.8 ± 4.3	0.003
10		77.8 ± 4.1	0.008
14		78.5 ± 2.4	<0.001

Independent t-test.

### 2.6. Statistical analysis

A Levene's test, One-way ANOVA test and paired t-test were employed by using a combination of SPSS statistics 24 (IBM Corp., Armonk, NY) and Excel (Microsoft Corp, Redmond, WA). A p-value of less than 0.05 was considered statistically significant.

### 3. Results

In this study, 64 rats were divided into 4 experimental groups: A) negative control, B) topical Vaseline, C) topical PRP and SVF, and D) subcutaneously injected PRP and SVF. Groups B, C, and D consisted of 5 groups of 4 rats, respectively. Group A was a group of 4 rats.

Based on Table 1, we can conclude the mean value of changes in VEGF levels for 14 days in 4 treatment groups. The group treated with vaseline, topical PRP and SVFs, and injection PRP and SVFs showed an increase in VEGF levels from day one to day 10 and then decreased on day 14. This indicated that the peak increase in VEGF levels was on day 10. In the PRP and SVFs group, injection in each treatment day had the highest mean value among all groups. In the group treated with vaseline, PRP and SVFs topical, PRP and SVFs injection, and the control group, there were significant differences in changes of VEGF levels with each p

value < 0.001.


### 3.1. Comparison between group A (negative control) and group B (topical vaseline)

As shown in Table 2, VEGF levels in the Vaseline group were significantly different from the control group ( $p < 0.05$ ). Based on this result,  $H_0$  is rejected and  $H_1$  is accepted, suggesting there is a significant effect of Vaseline on VEGF levels in a rat model of deep dermal burns on days 1, 4, 7, 10, and 14. The VEGF levels with Vaseline were highest on day 14 (Fig. 1). This result is similar to a study that reported VEGF serum levels with Vaseline were lower than control on day 7, but higher than control on days 14 and 21 [16]. In another study comparing Vaseline to controls with an angiogenesis parameter, the researchers found that on days 7, 14, and 21, Vaseline was better than the control group [17]. In another study comparing VEGF levels in Pelnae, Vaseline, and a control group found that on day 7, Vaseline was better than control and Pelnae was significantly better than Vaseline [18].

**Table 3**  
Comparison of VEGF levels between groups B and C.

Termination Day-	Group B (Mean ± SD) (pg/ml)	Group C (Mean ± SD) (pg/ml)	*p value
1	70.1 ± 1.5	75.5 ± 4.0	0.047
4	75.7 ± 5.1	86.3 ± 4.3	0.020
7	75.8 ± 4.3	87.0 ± 5.2	0.017
10	77.8 ± 4.1	89.8 ± 4.8	0.030
14	78.5 ± 2.4	88.4 ± 7.2	0.041

\* Independent t-test.



**Fig. 1.** Comparison result of VEGF levels between groups A and B.




Fig. 2. Comparison of VEGF levels between groups B and C.

Table 4

Comparison of VEGF levels between groups B and D.

Termination Day-	Group B (Mean $\pm$ SD) (pg/ml)	Group D (Mean $\pm$ SD) (pg/ml)	p value
1	70.1 $\pm$ 1.5	83.7 $\pm$ 5.0	0.002
4	79.0 $\pm$ 2.1	96.1 $\pm$ 5.6	0.002
7	77.8 $\pm$ 4.1	96.4 $\pm$ 4.0	<0.001
10	75.8 $\pm$ 4.3	98.7 $\pm$ 3.9	<0.001
14	75.7 $\pm$ 5.1	98.6 $\pm$ 3.7	<0.001

<sup>a</sup> Independent t-test.

### 3.2. Comparison between group B (topical vaseline) and group C (PRP and SVF, topical)

As shown in Table 3, the comparison of VEGF levels after topical PRP and SVF to Vaseline on each day was significantly different ( $p < 0.05$ ) (Fig. 2). Based on this result, H<sub>0</sub> is rejected and H<sub>1</sub> is accepted, suggesting that topical application of stem cells is more effective than Vaseline for increasing VEGF levels in a rat model of deep dermal burns on days 1, 4, 7, 10 and 14.

### 3.3. Comparison between group B (topical vaseline) and group D (PRP and SVF, injected)

As shown in Table 4, VEGF levels after stem cell injections were significantly increased compared with Vaseline on each day ( $p < 0.05$ ) (Fig. 3). Based on this result, H<sub>0</sub> is rejected and H<sub>1</sub> is accepted, suggesting that injections of stem cells are better able to increase VEGF levels compared with Vaseline in a rat model of deep dermal burns on days 1, 4, 7, 10 and 14.

### 3.4. Comparison between group C (PRP and SVF, topical) and group D (PRP and SVF, injected)

As shown in Table 5, VEGF levels were significantly increased after injection of stem cells and topical stem cells ( $p < 0.05$ ), and there was a significant difference between the two groups (Fig. 4). Based on this result, H<sub>0</sub> is rejected and H<sub>1</sub> is accepted, suggesting the injection of stem cells is better than topical stem cells in increasing VEGF levels in a rat model of deep dermal burns on days 1, 4, 7, 10 and 14.

## 4. Discussion

One of the factors that plays an important role in the wound healing process is angiogenesis, which is the process of forming new blood vessels [33,34]. One of the most important and widely studied angiogenic factors is vascular endothelial growth factor (VEGF) [34]. VEGF is one of the most potent mediators of vascular permeability and vascular regulation in angiogenesis. VEGF is also commonly called "vascular permeability factor," which refers to a specific endothelial cell mitogen secreted as a 45 kDa protein consisting of two subunits that do not induce cell proliferation in other cell types [24,34].

PRP contains many growth factors, including VEGF, HGF, bFGF, EGF, TGF- $\beta$ , IGF-1, and PDGF [35]. Angiogenesis is particularly stimulated by VEGF, PDGF and basic fibroblast growth factor (bFGF). Angiogenesis is also assisted by pericytes, which in turn depend on the availability of PDGF and VEGF [36,37]. Likewise, ADSC/SVF that was found contain many growth factors and cytokines, which in turn influence angiogenesis, as well as mononuclear infiltrates, fibroblast and production on collagen. VEGF, a potent angiogenic agents, migration of fibroblast and activity stimulators such as TGF, and macrophage chemotaxis agents are regulated by ADSCs [38].

The administration of PRP and SVF, either topically or by injection enhances VEGF levels up to day 10. This result was significant compared




Fig. 3. Comparison of VEGF levels between groups B and D.

Table 5

Comparison of VEGF levels between groups C and D.

Termination Day-	Group C (Mean ± SD) (pg/ml)	Group D (Mean ± SD) (pg/ml)	p value
1	75.5 ± 4.0	83.7 ± 5.0	0.045
4	86.3 ± 4.3	96.1 ± 5.6	0.034
7	87.0 ± 5.2	96.4 ± 4.0	0.030
10	89.8 ± 4.8	98.7 ± 3.9	0.029
14	88.4 ± 7.2	98.6 ± 3.7	0.047

<sup>a</sup> Independent t-test.

to Vaseline and control ( $p < 0.05$ ). Our findings are similar to a previous study that reported the benefits of topical PRP gel in treating soft tissue wounds, in which the recovery proceeded faster compared to Vaseline. PRP gel is not only beneficial in chronic wound treatment, but is also effective in acute injury [39]. Another study regarding the administration of the SVF and PRF combination showed significantly better results in the concentration of VEGF levels compared to SVF or PRP alone with VEGF levels peaking on day 7 and slowly declining on days 14 and 28 [40]. Additionally, another study showed that PRP and SVF injection increased VEGF levels on day 1 to day 7 then decreased on day 10 [41].

In our study, we found a significant difference between injection and topical PRP and SVF administration. PRP and SVF by injection is more effective at increasing VEGF levels compared with topical PRP and SVF. The main objective of PRP therapy is to centralize growth factors at the injury site. These factors are considered beneficial in the healing process in soft and hard tissue and are also considered the most effective way to provide a biological stimulus to the tissue [42]. In a previous in vitro study, it was stated that freezing and diluting PRP does not significantly affect VEGF release. There was no significant difference in VEGF levels after freezing for 1 h and 7 days. This study also concluded that kinetic differences in VEGF release are not significantly affected by PRP

preparation; further clinical studies could explore and clarify the PRP effect [35]. Based on our results, it can be concluded that the injection route is significantly more effective than topical application for increasing VEGF release.

In a study analyzing the musculoskeletal effect of PRP and SVF, the authors stated that direct injection of PRP to the joints can control the inflammatory response and promote long-term healing. The growth factor components in PRP have special anti-inflammatory effects and can also reduce pain. Anti-inflammation, immunomodulation, and analgesia are also influenced by soluble factors excreted by SVF or ADSCs, including HGF, VEGF, NGF, EGF, FGF and TGF- $\beta$  [43].

Inflammation regulation, proliferation of cells, remodeling, deposition of extracellular matrix, angiogenesis and epithelization play important role in skin wound healing. In the mouse skin, PRP and SVFs can reduce inflammatory process, increase collagen deposition, and promote angiogenesis and neurogenesis, thus promote wound healing [44]. PRP and adipose-derived stem cells combination is potent in stimulating increase of growth factor concentration (IGF-1, TGF- $\beta$  1, HGF, and VEGF) [27]. PRP is an accessible cell therapy to helps treat chronic wounds and severe burns [16,19]. PRP is an autologous cell therapy containing a large number of bioactive factors involved in wound healing and tissue repair. PRP can be used for treating and stimulating wound healing in burns and split-thickness skin grafts, which are often used for the management of burns [45].

Based on this study, further research is needed regarding the effect of the combination of PRP and SVF on VEGF levels in dermal burn healing. Future studies could involve determining the quantity of neo-vascularization in burn healing through histopathological examination.

In addition, our results may be of interest to other researchers investigating the additive and synergistic effects of therapeutic strategies involving stem cells. The fact that the PRP and SVF efficacy data for burn injuries are limited highlights the need for phase I-II clinical trials. We hope that this research can serve as a fundamental resource and an






Fig. 4. Comparison of VEGF levels between groups C and D.

additional reference when making decisions about the application of PRP and SVF as a new breakthrough in the management of burn injuries.

## 5. Conclusion

In our study, we can conclude that the administration of PRP and SVF by injection can significantly increase VEGF levels more effectively than Vaseline in rats with induced deep dermal burns. Further, the administration of topical PRP and SVF significantly increased VEGF levels compared to the Vaseline group. There is a significant difference between PRP and SVF administration by injection and topical application. The injection of PRP and SVF is more effective at increasing VEGF levels compared to topical application.

## Provenance and peer review

Our study was non-commissioned and externally peer-reviewed.

## Ethical approval

All procedure for Animal experiment has been approved by Ethics Commission Faculty of Medicine, Hasanuddin University Number: 216/UN4.6.4.5.31/PP36/2020.

## Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Author contribution

SRL, FJ, SB, and MF wrote the manuscript and participated in the study design. SRL, FJ, SB, MF, ASB, WS, MNM, IDS, and AAI drafted and

revised the manuscript. SRL, FJ, SB, and MF performed deep dermal burn model, SRL, FJ, MF, and IJP performed bioinformatics analyses and revised the manuscript. All authors read and approved the final manuscript.

## Registration of research studies

None.

## Guarantor

Sachraswaty Rachman Laidding.

## Consent

This manuscript does not involve human participants, human data, or human tissue.

## Declaration of competing interest

The authors declare that they have no conflict of interests.

## Acknowledgment

A higher appreciation to all staff from the Hasanuddin University Medical Research Center (HUMRC), Makassar, Indonesia.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.amsu.2021.102254>.

## References

- [1] ANZBA, Emergency Management of Severe Burns (EMSB), eighteenth ed., THE EDUCATION COMMITTEE OF AUSTRALIA AND NEW ZEALAND BURN ASSOCIATION LTD, Albany Creek, 2016.
- [2] I. Kaddoura, G. Abu-Sittah, A. Ibrahim, R. Karamanoukian, N. Pipazian, Burn injury: review of pathophysiology and therapeutic modalities in major burns, *Ann. Burns Fire Disasters* 30 (2017) 95–102. <http://www.ncbi.nlm.nih.gov/pubmed/29021720>.
- [3] M.A.R. Stokes, W.D. Johnson, Burns in the third world: an unmet need, *Ann. Burns Fire Disasters* 30 (2017) 243–246. <https://pubmed.ncbi.nlm.nih.gov/29963673/>.
- [4] F. Ghieh, R. Jurjus, A. Ibrahim, A.G. Geagea, H. Daouk, B. El Baba, S. Chama, M. Matar, W. Zein, A. Juajus, The use of stem cells in burn wound healing: a review, *BioMed Res. Int.* 2015 (2015) 684084. <https://doi.org/10.1155/2015/684084>.
- [5] V.K. Singh, A. Saini, M. Kalsan, N. Kumar, R. Chandra, Describing the stem cell potency: the various methods of functional assessment and in silico diagnostics, *Front. Cell Dev. Biol.* 4 (2016). <https://doi.org/10.3389/fcell.2016.00134>.
- [6] S.R. Laidling, F. Josh, Francisca, M. Faruk, A.S. Palissey, B. Satrin, Warsinggh, A. Bukhari, M.N. Massi, A.A. Islam, Combination of platelet-rich plasma and stromal vascular fraction on the level of transforming growth factor- $\beta$  in rat subjects experiencing deep dermal burn injury, *Ann. Med. Surg.* 60 (2020) 737–742. <https://doi.org/10.1016/j.amss.2020.11.084>.
- [7] P. Foubert, A.D. Gonzalez, S. Teodorescu, F. Berard, M. Doyle-Eisele, K. Yekkala, M. Teunhaus, J.K. Fraser, Adipose-derived regenerative cell therapy for burn wound healing: a comparison of two delivery methods, *Adv. Wound Care* 5 (2016) 288–298. <https://doi.org/10.1089/wound.2015.0672>.
- [8] S. Atalay, A. Coruh, K. Deniz, Stromal vascular fraction improves deep partial thickness burn wound healing, *Burns* 40 (2014) 1375–1383. <https://doi.org/10.1016/j.burns.2014.01.023>.
- [9] J. Han, Y.-I. Koh, H.R. Moon, H.G. Ryoo, C.-H. Cho, I. Kim, G.-Y. Koh, Adipose tissue is an extramedullary reservoir for functional hematopoietic stem and progenitor cells, *Blood* 115 (2010) 957–964. <https://doi.org/10.1182/blood-2009-05-219923>.
- [10] V. Tantiway, P. Bhamhani, A. Nagla, P. Mantry, R. Sharma, P. Mehto, S. Bansal, Autologous grafting of non manipulated freshly isolated - adipose tissue derived stromal vascular fraction in single surgical sitting for treatment of knee osteoarthritis, *Int. J. Res. Orthop.* 3 (2016) 107. <https://doi.org/10.18203/issn.2455-4510.ijro.orthop.20160484>.
- [11] A. Darinskas, M. Paskevicius, G. Apanavicius, G. Vilkevicius, L. Labanavskas, T. E. Ichim, R. Rimdeika, Stromal vascular fraction cells for the treatment of critical limb ischemia: a pilot study, *J. Transl. Med.* 15 (2017) 143. <https://doi.org/10.1186/s12967-017-1243-3>.
- [12] K. Comella, R. Silbert, M. Parlo, Effects of the intradiscal implantation of stromal vascular fraction plus platelet rich plasma in patients with degenerative disc disease, *J. Transl. Med.* 15 (2017) 12. <https://doi.org/10.1186/s12967-016-1109-8>.
- [13] P. Gentile, M.G. Scioli, A. Bielefi, A. Olandi, V. Cervelli, Concise review: the use of adipose-derived stromal vascular fraction cells and platelet rich plasma in regenerative plastic surgery, *Stem Cell.* 35 (2017) 117–134. <https://doi.org/10.1002/stem.2498>.
- [14] P. Bourin, B.A. Burnell, L. Castella, M. Dominici, A.J. Katz, K.L. March, H. Redl, J. P. Rubin, K. Yoshimura, J.M. Gimble, Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society of Cytotherapy 15 (2013) 641–648. <https://doi.org/10.1016/j.jcyt.2013.02.006>.
- [15] J. Choi, K.W. Min, H. Chang, The efficacy and safety of platelet-rich plasma and adipose-derived stem cells: an update, *Arch. Plast. Surg.* 39 (2012) 585–592. <https://doi.org/10.5999/aps.2012.39.4.585>.
- [16] E. Raposo, N. Bertozzi, S. Bonasini, G. Bernuzzi, A. Formentini, E. Grignaffini, M. Pio Grieco, Adipose-derived stem cells added to platelet-rich plasma for chronic skin ulcer therapy: wounds a compound, *Clin. Res. Pract.* 28 (2016) 126–131.
- [17] P. Borrione, A. Di Gianfrancesco, M.T. Pereira, F. Pigozzi, Platelet-rich plasma in muscle healing, *Am. J. Phys. Med. Rehabil.* 89 (2010) 854–861. <https://doi.org/10.1097/PHM.0b013e3181f1c1e7>.
- [18] A. Duran, S. Yasar, P. Gomes, S. Aytekin, A. Duran, Clinical and Histopathological Evaluation of the Effects of Platelet Rich Plasma, Platelet Poor Plasma and Topical Serum Physiologic Treatment on Wound Healing Caused by Radiofrequency Electrosurgery in Rats, 2016.
- [19] T. Stesuk, M.B. Puzzi, E.A. Chaim, P.C.M. Alves, E.V. de Paula, A. Forte, J. M. Izumizawa, C.C. Oliveira, F. Frei, J.T. Ribeiro-Paes, Platelet-rich plasma (PRP) and adipose-derived mesenchymal stem cells: stimulatory effects on proliferation and migration of fibroblasts and keratinocytes in vitro, *Arch. Dermatol. Res.* 308 (2016) 511–520. <https://doi.org/10.1007/s00403-016-1674-1>.
- [20] M. Tohidnezhad, D. Varoga, C.J. Wruck, L.O. Brandenburg, A. Seekamp, M. Shakibaei, T.T. Sozaev, T. Pufe, S. Lippross, Platelet-released growth factors can accelerate tenocyte proliferation and activate the anti-oxidant response element, *Histochem. Cell Biol.* 135 (2011) 453–460. <https://doi.org/10.1007/s00418-011-0808-6>.
- [21] R.P. Martins, D.D. Hartmann, J.P. de Moraes, F.A.A. Soares, G.O. Puntel, Platelet-rich plasma reduces the oxidative damage determined by a skeletal muscle contusion in rats, *Platelets* 27 (2016) 784–790. <https://doi.org/10.1080/09537104.2016.1184752>.
- [22] N.H. Mansoub, M. Girdal, E. Karadadas, H. Kabadayi, S. Yatansever, G. Ercan, The role of PRP and adipose tissue-derived keratinocytes on burn wound healing in diabetic rats, *Bioimpacts* 5 (2018) 5–12. <https://doi.org/10.15371/bi.201802>.
- [23] N. Subramaniam, J.J. Petrlik, M.K. Vickaryous, VEGF, FGF-2 and TGF $\beta$  expression in the normal and regenerating epidermis of geckos: implications for epidermal homeostasis and wound healing in reptiles, *J. Anat.* 232 (2018) 768–782. <https://doi.org/10.1111/joa.12784>.
- [24] C.S. Melincovic, A.B. Bojca, S. Sujanon, M. Margison, C. Mihu, M. Istaitie, I. M. Moldovan, A.L. Roman, C.M. Mihu, Vascular endothelial growth factor (VEGF)-like factor in normal and pathological angiogenesis, *Roum. J. Morphol. Embryol. Rev. Roum. Morphol. Embryol.* 59 (2018) 455–467.
- [25] C. Kilkenny, W.J. Browne, L.C. Cuthill, M. Emerson, D.G. Altman, Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research, *PLoS Biol.* 8 (2010), e1000412. <https://doi.org/10.1371/journal.pbio.1000412>.
- [26] K.F. Schulz, D.G. Altman, D. Moher, CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials, *BMJ* 340 (2010) e332. <https://doi.org/10.1136/bmj.e332>.
- [27] S. Tajima, M. Tobita, H. Orbay, H. Hyakusoku, H. Mizuno, Direct and indirect effects of a combination of adipose-derived stem cells and platelet-rich plasma on bone regeneration, *Tissue Eng.* 21 (2014). <https://doi.org/10.1089/ten.tea.2014.00326>.
- [28] F. Josh, K. Kobe, M. Tobita, R. Tanaka, K. Suzuki, K. Ono, H. Hyakusoku, H. Mizuno, Accelerated and safe proliferation of human adipose-derived stem cells in medium supplemented with human serum, *J. Nippon Med. Sch.* 79 (2012) 444–452. <https://doi.org/10.1272/jnms.79.444>.
- [29] H.-F. Guo, R.M. Ali, B.A. Hamid, A.A. Zaini, H. Khayat, A new model for studying deep partial-thickness burns in rats, *Int. J. Burns Trauma* 7 (2017) 107–114.
- [30] D. dos, S. Tavares Pereira, M.H.M. Lima Ribeiro, N.T. de Pontes-Filho, A.M. dos, A. Carneiro-Leão, M.T. dos, S. Correia, Development of animal model for studying deep second-degree thermal burns, *J. Biomed. Biotechnol.* (2012) 460841. <https://doi.org/10.1155/2012/460841>.
- [31] A. Sethi, T. Kaur, S.K. Malhotra, M.L. Gambhir, Moisturizer: the slippery road, *Indian J. Dermatol.* 61 (2016) 279–287. <https://doi.org/10.4103/0019-5154.182427>.
- [32] Warsinggh, B. Irawan, I. Labeda, R.E. Luskiyov, S. Sampetodong, M.I. Kusuma, J. A. Uwaruwat, E. Syarifuddin, Prihantono, M. Faruk, Association of superoxide dismutase enzyme with staging and grade of differentiation colorectal cancer: a cross-sectional study, *Ann. Med. Surg.* 58 (2020) 194–199. <https://doi.org/10.1016/j.amss.2020.08.032>.
- [33] R.J. Bodnar, Epidermal growth factor and epidermal growth factor receptor: the yin and yang in the treatment of cutaneous wounds and cancer, *Adv. Wound Care* 2 (2013) 24–29. <https://doi.org/10.1089/wound.2011.0326>.
- [34] K.E. Johnson, T.A. Wilgus, Vascular endothelial growth factor and angiogenesis in the regulation of cutaneous wound repair, *Adv. Wound Care* 3 (2014) 647–661. <https://doi.org/10.1089/wound.2013.0517>.
- [35] N. Hosny, F. Goubran, B. BadrEldin Husan, N. Kamel, Assessment of vascular endothelial growth factor in fresh versus frozen platelet rich plasma, *J. Blood Transfus.* 2015 (2015) 1–5. <https://doi.org/10.1155/2015/706903>.
- [36] J.I. Greenberg, D.J. Shields, S.G. Barillas, L.M. Acevedo, E. Murphy, J. Huang, L. Schepke, C. Stockmann, R.S. Johnson, N. Angle, D.A. Cheresh, A role for VEGF as a negative regulator of pericyte function and vessel maturation, *Nature* 456 (2008) 809–813. <https://doi.org/10.1038/nature07424>.
- [37] N. Caporarello, F. D'Angeli, M.T. Cambria, S. Candido, C. Gallongo, M. Salmeri, C. Lombardo, A. Longo, G. Giordanello, C.D. Anfuso, G. Lupo, Pericytes in microvessels from "murine" function to brain and retina regeneration, *Int. J. Mol. Sci.* 20 (2019) 6351. <https://doi.org/10.3390/ijms20246351>.
- [38] A.L. Cardoso, M.M. Bachion, J. de M. Morais, M.S. Fantinati, V.L.L. de Almeida, R. S. Lino Júnior, Adipose tissue stromal vascular fraction in the treatment of full thickness burns in rats, *Acta Cir. Bras.* 31 (2016) 578–585. <https://doi.org/10.1590/s0102-86502016000000002>.
- [39] K. Kazakos, D.N. Lyras, D. Verettas, K. Tilkeridis, M. Tryfonidis, The use of autologous PRP gel as an aid in the management of acute trauma wounds, *Injury* 40 (2009) 801–805. <https://doi.org/10.1016/j.injury.2008.05.002>.
- [40] Z.-J. Liang, X. Lu, D.-Q. Li, Y.-D. Liang, D.-D. Zhu, F.-X. Wu, X.-L. Yi, N. He, Y.-Q. Huang, C. Tang, H. Li, Precise intradermal injection of manofact-derived stromal cells combined with platelet-rich fibrin improves the efficacy of facial skin rejuvenation, *Cell. Physiol. Biochem.* 47 (2018) 316–329. <https://doi.org/10.1159/000489869>.
- [41] N. Seyhan, D. Altan, A.U. Ural, A. Gural, M.C. Avunduk, N. Savaci, The effect of combined use of platelet-rich plasma and adipose-derived stem cells on fat graft survival, *Ann. Plast. Surg.* 74 (2015) 615–620. <https://doi.org/10.1097/SAP.0000000000000480>.
- [42] A.D. Mazzecca, M.B.R. McCarthy, D.M. Chowanicek, M.P. Cote, A.A. Roman, J. P. Bradley, R.A. Arciero, K. Betzel, Platelet-rich plasma differs according to preparation method and human variability, *J. Bone Jt. Surg.* 94 (2012) 308–316. <https://doi.org/10.2106/JBJS.E.00420>.
- [43] A. Fotouhi, A. Maleki, S. Dolati, A. Aghebati-Maleki, L. Aghebati-Maleki, Platelet rich plasma, stromal vascular fraction and autologous conditioned serum in treatment of knee osteoarthritis, *Biomed. Pharmacother.* 104 (2018) 652–660. <https://doi.org/10.1016/j.biopha.2018.05.019>.
- [44] T. Liu, C. Qu, C. Ben, H. Li, S. Zhu, One-step approach for full-thickness skin defect reconstruction in rats using minced split-thickness skin grafts with Pelnae overlay, *Burn. Trauma* 7 (2019). <https://doi.org/10.1186/s13087-019-0157-0>.
- [45] K. Akhond, G. Pietramaggioli, L. Waselle, S. Darwiche, S. Guerci, C. Scatena, N. Hirt-Burn, L.A. Applegate, W. V. Rafford, Development of a cost-effective method for platelet-rich plasma (PRP) preparation for topical wound healing, *Ann. Burns Fire Disasters* 25 (2012) 207–213.

# Combination of platelet rich plasma and stromal vascular fraction on the level of vascular endothelial growth factor in rat subjects experiencing deep dermal burn injury

## ORIGINALITY REPORT

22%

SIMILARITY INDEX

14%

INTERNET SOURCES

19%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1

Yusuf Emes, Buket Aybar, Pervin Vural, Neslihan Yildirim Saral et al. "Effects of Bone Morphogenetic Proteins on Osteoblast Cells: Vascular Endothelial Growth Factor, Calcium, Inorganic Phosphate, and Nitric Oxide Levels", *Implant Dentistry*, 2010

Publication

1%

2

Widodo, Bambang Pristiwanto, Muhaimin Rifa'i, Irfan Mustafa, Fahrul Zaman Huyop. "A single epitope of Epstein-Barr Virus stimulate IgG production in mice", *Annals of Medicine and Surgery*, 2018

Publication

1%

3

Manfred Infanger, Oliver Schmidt, Peter Kossmehl, Sibylle Grad, Wolfgang Ertel, Daniela Grimm. "Vascular endothelial growth factor serum level is strongly enhanced after burn injury and correlated with local and general tissue edema", *Burns*, 2004

Publication

1%

Leung, Jerry Shapiro, Kevin J. McElwee.  
"Growth factor concentrations in platelet -  
rich plasma for androgenetic alopecia: An  
intra - subject, randomized, blinded,  
placebo - controlled, pilot study",  
Experimental Dermatology, 2020  
Publication

- 
- |           |   |     |
|-----------|---|-----|
| <b>10</b> | <a href="http://www.meredithvanstone.com">www.meredithvanstone.com</a><br>Internet Source | 1 % |
|-----------|---|-----|
- 
- |           |   |     |
|-----------|---|-----|
| <b>11</b> | Dimitris Lyras. "Immunohistochemical study<br>of angiogenesis after local administration of<br>platelet-rich plasma in a patellar tendon<br>defect", International Orthopaedics,<br>02/11/2009<br>Publication | 1 % |
|-----------|---|-----|
- 
- |           |  |     |
|-----------|--|-----|
| <b>12</b> | Yi Sun, Xiaoqi Chi, Haoye Meng, Mengjiao Ma<br>et al. "Polylysine-decorated macroporous<br>microcarriers laden with adipose-derived<br>stem cells promote nerve regeneration in<br>vivo", Bioactive Materials, 2021<br>Publication | 1 % |
|-----------|--|-----|
- 
- |           |   |     |
|-----------|---|-----|
| <b>13</b> | <a href="http://tehj.springeropen.com">tehj.springeropen.com</a><br>Internet Source | 1 % |
|-----------|---|-----|
- 
- |           |  |      |
|-----------|--|------|
| <b>14</b> | Cagri A. Uysal, Morikuni Tobita, Hiko<br>Hyakusoku, Hiroshi Mizuno. "Adipose-derived<br>stem cells enhance primary tendon repair:<br>Biomechanical and immunohistochemical | <1 % |
|-----------|--|------|