



Source details

F1000Research

Open Access ⓘ

Scopus coverage years: from 2012 to 2022

Publisher: Taylor & Francis

E-ISSN: 2046-1402

Subject area: Pharmacology, Toxicology and Pharmaceutics: General Pharmacology, Toxicology and Pharmaceutics

Biochemistry, Genetics and Molecular Biology: General Biochemistry, Genetics and Molecular Biology View all ▾

Source type: Journal

CiteScore 2021

5.0 ⓘ

SJR 2021

0.939 ⓘ

SNIP 2021

1.010 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology ⓘ

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▾

$$5.0 = \frac{15,170 \text{ Citations 2018 - 2021}}{3,056 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$5.1 = \frac{13,399 \text{ Citations to date}}{2,618 \text{ Documents to date}}$$

Last updated on 05 January, 2023 • Updated monthly

CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#11/74	85th
General Pharmacology, Toxicology and Pharmaceutics		
Biochemistry, Genetics and Molecular Biology	#65/204	68th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER



[Terms and conditions](#) ↗ [Privacy policy](#) ↗


Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.





<p>COUNTRY</p> <p>United Kingdom</p> <div data-bbox="119 264 354 309">  Universities and research institutions in United Kingdom </div> <div data-bbox="119 331 354 358">  Media Ranking in United Kingdom </div>	<p>SUBJECT AREA AND CATEGORY</p> <p>Biochemistry, Genetics and Molecular Biology</p> <ul style="list-style-type: none"> └ Biochemistry, Genetics and Molecular Biology (miscellaneous) <p>Immunology and Microbiology</p> <ul style="list-style-type: none"> └ Immunology and Microbiology (miscellaneous) <p>Medicine</p> <ul style="list-style-type: none"> └ Medicine (miscellaneous) <p>Pharmacology, Toxicology and Pharmaceutics</p> <ul style="list-style-type: none"> └ Pharmacology, Toxicology and Pharmaceutics (miscellaneous) 	<p>PUBLISHER</p>	<p>H-INDEX</p> <p>72</p>
<p>PUBLICATION TYPE</p> <p>Journals</p>	<p>ISSN</p> <p>20461402</p>	<p>COVERAGE</p> <p>2012-2021</p>	<p>INFORMATION</p> <p>Homepage</p> <p>How to publish in this journal</p> <p>research@f1000.com</p>



SCOPE








F1000Research publishes articles and other research outputs reporting basic scientific, scholarly, translational and clinical research across the physical and life sciences, engineering, medicine, social sciences and humanities. F1000Research is a scholarly publication platform set up for the scientific, scholarly and medical research community; each article has at least one author who is a qualified researcher, scholar or clinician actively working in their speciality and who has made a key contribution to the article. Articles must be original (not duplications). All research is suitable irrespective of the perceived level of interest or novelty; we welcome confirmatory and negative results, as well as null studies. F1000Research publishes different type of research, including clinical trials, systematic reviews, software tools, method articles, and many others. Reviews and Opinion articles providing a balanced and comprehensive overview of the latest discoveries in a particular field, or presenting a personal perspective on recent developments, are also welcome. See the full list of article types we accept for more information.

 Join the conversation about this journal

FIND SIMILAR JOURNALS

options

<p>1 Advances in Experimental Medicine and Biology USA</p> <p>79% similarity</p>	<p>2 EBioMedicine NLD</p> <p>78% similarity</p>	<p>3 Science Translational Medicine USA</p> <p>73% similarity</p>	<p>4 Frontiers of Medicine CHN</p> <p>73% similarity</p>	<p>5 Journal of Biomedical Science GBR</p> <p>71% similarity</p>
--	---	---	--	--



F1000Research

Q1
Biochemistry, Genetics and Molecular Biology...
best quartile

SJR 2021
0.94
powered by scimagojr.com


Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" >
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool.**



D

deivasigamani 4 months ago

What is the impact factor of this journal?

← reply

**Melanie Ortiz** 4 months ago

SCImago Team

Dear Deivasigamani, thank you very much for your comment. SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR (Check it on our website). We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team

H

hany akeel naji 2 years ago

I have been waiting about one year for my article to publish or review but till now nothing happened

← reply

**Melanie Ortiz** 2 years ago

SCImago Team

Dear Hany,
Thank you for contacting us.
Unfortunately, we cannot help you with your request, we suggest you contact the journal's editorial staff, so they could inform you more deeply.
Best Regards, SCImago Team

A

alemu 2 years ago

Background: Nowadays, people die due to metabolic aging than chronological age worldwide. Unhealthy lifestyle behaviors may be long-term risk factors for chronic circumstances and its effect on quality of life is not routinely examined.
Objective: To investigate the association between healths related quality of life and unhealthy lifestyles behaviors among middle aged urban residents in West Ethiopia.
Methods: This cross-sectional study was conducted on 266 adults. Data were collected using health promoting questionnaires for lifestyle behaviors and EQ_5D of health related quality of life. Data were analyzed using descriptive statistics, T-test and a multiple regression analysis with SPSS version24.
Results: Of a total surveyed a mean of age was 52.5 years. The proportion of poor health related quality of life (HRQoL), low dietary diversity score and physical inactivity were 24.4%, 68% and 91% respectively. Individuals with poor HRQoL, had high unhealthy lifestyle behaviors scores and significant mean differences were observed ($P < 0.001$). On multiple linear regression models, significant association was seen between physical activity, alcohol consumption, smoking, diet and HRQoL. Being physically inactive ($\beta = 2.972$; 95.0% CI: 2.32, 3.62; $P < 0.001$) and unhealthy diet ($\beta = 0.043$; 95.0% CI: 0.015, 0.070; $P < 0.002$) were showed positive significant association with HRQoL. In contrast, smoking, alcohol drinking, and khat ($p = 0.042$) were negatively associated with HRQoL. Combined poor lifestyle behaviors and poor health related quality of life were significantly associated ($P = 0.023$) as seen independently.
Conclusion: This study revealed that physical inactivity, unhealthy diet and combined poor lifestyle factors were significantly associated to poor HRQOL and prevalent. Community based healthy lifestyles education is recommended to improve adult health and quality of life.

← reply

**Melanie Ortiz** 2 years ago

SCImago Team

Dear Alemu,
thank you for contacting us.
We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.
Unfortunately, we cannot help you with your request, we suggest you visit the journal's homepage (See submission/author guidelines) or contact the journal's editorial staff, so they could inform you more deeply.
Best Regards, SCImago Team

Y

Yasmeen 2 years ago

What is the impact factor og this journal in 2020

← reply



Melanie Ortiz 2 years ago

SCImago Team

Dear Yasmeen, thank you very much for your comment. SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR. Check out our web to localize the journal. We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team



AHMAD 3 years ago

i wanna ask if your journal is SCI?

← reply



Melanie Ortiz 3 years ago

SCImago Team

Dear Ahmad, SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR. Check our web to locate the journal. We suggest you to consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImag



Seyed 4 years ago

Hello please help me. sending manuscripts online is complicated for me because of IP protocols. Can you tell me an electronic mailing way to sending or submitting manuscripts to F1000?

← reply



Dr Ramesh 4 years ago

F1000 Research is indexed ISI or web of Science OR not

← reply



Elena Corera 4 years ago

SCImago Team

Dear Dr Ramesh, SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR. Check our page to locate the journal. We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team

Leave a comment

Name

Email
(will not be published)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:

Powered by:



Follow us on [@ScimagoJR](#)

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS
Horatio (Satire 1,1,106)

[Edit Cookie Consent](#)

Advisory Board

The Advisory Board of F1000Research comprises a large group of leading experts across biology and medicine. They do not act as Editors in the traditional sense (they do not handle manuscripts or make decisions to accept or reject a paper), but they provide strategic input on the direction we should take with F1000Research. They occasionally advise us on issues arising with specific articles, and many members of the board also review for us.

[A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#)

B

Ian Beales
Nelson Bennett
Avri Ben-Ze'ev
Benedikt Berninger
Eric Beyer
Azra Bihorac
Daniel Bikle
Kevin J Black
Chellakkan Selvanesan Blesson
Erin Aiello Bowles
Bruce Brew

C

David Catcheside
Andrew Chalmers
Tak Mao Chan
Karen Chapman
Declan Chard
Walter Chazin
Jonathan Chernoff
Cheng-Ming Chiang
Ryan Chisholm
Wei-Sheng Chong
Sandra Citi
Vitaly Citovsky
Tim Clark
James Coker
Giuseppe Colloca
William Colmers
Jason Crawford
David Criddle

D

Ira Daar
Linda Dagi
Blossom Damania
Eric Dannaoui
Vinicio de Jesus Perez
Sharon DeMorrow
Gonzalo G de Polavieja
Saskia de Wildt
Harriet de Wit



Annette Dolphin
Lucy Donaldson
Sylvie Doublé
Paschalis-Thoma Doulias
Crislyn D'Souza-Schorey
James Duffin
Janice Du Mont

E
Sharyn Endow
Markus Engstler
Sam Enna
Erim Erdem

F
Alastair Ferguson
Gerardo Ferrara
Richard Festenstein
Thomas Finger
Céline Fiset
Heike Fölsch
Steven Frank
Bernd Fritsch

G
Gus Gazzard
Jozef Gécz
Robert Gerlai
Ivan Gerling
Carole Goble
Richard Gomer
Andrew Goryachev
John Greenspan
Guy Griebel
W Sue Griffin
Elizabeth Grove
Jaime Grutzendler
Wei Guo

H
Adam Hartman
Johannes Hell
Winston Hide
Stephen Hoffman
Stephen Holgate
Thorsten Hoppe
Wolfgang Huber
Arthur Hurwitz

I
Radu Iliescu
Robert Insall
Harry Ischiropoulos

J
Jan Jakobsson
Guilhem Janbon



Wael Kariemah
Chaya Kalcheim
Lynn Kamerlin
Mikhail Kazachkov
Johannes S Kern
Jean-Pierre Kinet
Edward Kipreos
Fenella Kirkham
Gordon Klein
Alisa Koch
Amos Korczyn
Benoit Kornmann
Jan Kucera
Anuj Kumar
Saravana Kumar

L

Eileen Lafer
Hans Lassmann
Mario Lebediker
John Lee
Laurel Lenz
Simon Levin
Stefan Linder
Ton Lisman
Creighton M Litton
Hartmut Lode
Theresa Lu
Robyn Lucas
Ben Lugtenberg
Paul Lyons

M

Roberto Maggi
Martin Marinus
M Rashad Massoud
Jocelyn McDonald
Robert McPeck
Anthony Means
Julien Mendlewicz
Arthur Mercurio
Ralph Mistlberger
Ali Mobasher
David Moher
Randall Moon
Carlos Morel
Dimitrios Morikis
Nicola Mulder

N

Corey Nislow

O

Chiadi Onyike



Stephen Pinner
Michel Pohl
Simon Portsmouth
David Potter
Chaim Putterman

R

Adam Ratner
Ana Recober
Victor Reus
José Luis Riechmann
Karin Romisch
Vincent Rotello
Barry Rouse
Gloria Rudenko
James Russell

S

Philippe Saas
Paul R Sanberg
Alan Schechter
Werner Scheithauer
Tamar Schlick
Thomas Schnider
Alfons Schnitzler
Irene Schulz
Michael Sendtner
Andrew D Sharrocks
Nilabh Shastri
Kazim Sheikh
Andrew Shennan
Xiao Shifu
Chiara Simonelli
Helmy Siragy
Cassian Sitaru
Richard Smith
H Peter Soyer
Pamela Stanley
Christoph Stein
Carly Stevens
Charles Stevens
Bruno Stieger

T

Paul-Peter Tak
Paul Terry
Igor Tetko
Jacques Thibodeau
Jakub Tolar
Peter Tonellato
Francis Tsai
Takeshi Tsubata
Tom Tullius
Burkhard Tümmler



Hans van Bokhoven

Martin van den Berg

Peter Van Endert

Dirk van Helden

Chandra Verma

Jan Vermorken

David Voehringer

W Claire Walczak

Nick Ward

Peter Wark

Stephen Waxman

Alan Wein

Tom Woodcock

Long-Jun Wu

Jeremy C Wyatt

Kevan Wylie

X Yongbiao Xue

Y Michael B Yaffe

Kenneth Yamada

Helen Yap

Dorothy Yuan

Z Yunde Zhao

Deyou Zheng

Guy Zimmerman

Christos Zouboulis

F1000Research

An innovative open access publishing platform offering rapid publication and open peer review, whilst supporting data deposition and sharing.

[BROWSE](#)

[GATEWAYS](#)

[COLLECTIONS](#)

[HOW IT WORKS](#)

[BLOG](#)

[CONTACT](#)

[PRIVACY NOTICE](#)

[RSS](#)



[Home](#) » [Browse Articles](#)

You searched for Johanes Nugroho

[ARTICLES](#) [FACULTY REVIEWS](#) [DOCUMENTS](#) [POSTERS](#) [SLIDES](#)

FILTERS ▾

1-1 of 1 ARTICLE

SYSTEMATIC REVIEW  metrics 

Factor Xa inhibitor for venous thromboembolism management in patient with cancer: a systematic review and meta-analysis [version 1; peer review: 2 approved]

Johanes Nugroho Eko Putranto, Ardyan Wardhana, Yoga Alfian Noor, Pirhot Lambok Marnala Yosua Siahaan, Makhyan Jibril Al Farabi

 **PEER REVIEWERS** Suko Adiarto; Irene Terrenato

FUNDER Indonesian Endowment Fund for Education (LPDP-Lembaga Pengelola Dana Pendidikan RI)

PUBLISHED 08 Dec 2021

PUBLISH YOUR RESEARCH



ARTICLES

We publish a wide range of article types in science, engineering, medicine, social sciences and humanities, with no editorial biases.

SUBMIT AN ARTICLE (/FOR-AUTHORS/PUBLISH-YOUR-RESEARCH)

See [guidelines](#) and [policies](#).



BROWSE
GATEWAYS
COLLECTIONS
HOW IT WORKS

BLOG
CONTACT
PRIVACY NOTICE
RSS



Follow us



© 2012-2023 F1000 Research Ltd. ISSN 2046-1402 | [Legal](#) | Partner of [Research4Life](#) · [CrossRef](#) · [ORCID](#) · [FAIRSharing](#)



The F1000Research website uses cookies. By continuing to browse the site, you are agreeing to our use of cookies. [Find out more »](#)



SYSTEMATIC REVIEW

Factor Xa inhibitor for venous thromboembolism management in patient with cancer: a systematic review and meta-analysis [version 1; peer review: awaiting peer review]

Johanes Nugroho Eko Putranto ^{1,2}, Ardyan Wardhana ³, Yoga Alfian Noor^{1,2}, Pirhot Lambok Marnala Yosua Siahaan^{2,4}, Makhyan Jibril Al Farabi^{1,4}

¹Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

²Dr. Soetomo General Hospital, Surabaya, Indonesia

³Faculty of Medicine, Universitas Surabaya, Surabaya, Indonesia

⁴Department of Anesthesia and Reanimation, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

V1 First published: 08 Dec 2021, 10:1257
<https://doi.org/10.12688/f1000research.73883.1>
Latest published: 08 Dec 2021, 10:1257
<https://doi.org/10.12688/f1000research.73883.1>

Abstract

Background: An earlier systematic review reported no differences in the incidence of recurrent venous thromboembolism and major bleeding between factor Xa inhibitors and standard anticoagulation. The present meta-analysis aimed to assess the effectiveness of factor Xa inhibitors for the management of venous thromboembolism (VTE), specifically in patients with cancer, as there were more randomized clinical trials (RCTs) available.

Methods: The PubMed and Cochrane Library databases were systematically screened for all RCTs assessing factor Xa inhibitor efficacy for VTE management in cancer patients. Using RevMan 5.3, we performed a Mantel-Haenszel fixed-effects meta-analysis of the following outcomes: recurrent VTE, VTE events, and major bleeding rates.

Results: We identified 11 studies involving 7,965 patients. Factor Xa inhibitors were superior in preventing VTE recurrence, compared to low-molecular-weight heparin (LMWH) (OR 0.60; 95% CI 0.45–0.80; $P < 0.01$) and vitamin K antagonists (VKA) (OR 0.51; 95% CI 0.33–0.78; $P < 0.01$). As prophylaxis, factor Xa inhibitors had a similar rate of VTE compared to VKAs (OR 1.08 [95% CI 0.31–3.77]; $P = 0.90$) and a lower rate compared to placebo (OR 0.54 [95% CI 0.35–0.81]; $P < 0.01$). Major bleeding rates were higher with factor Xa inhibitors than with LMWHs (OR 1.34 [95% CI 0.83–2.18]; $P = 0.23$), but significantly lower than VKAs (OR 0.71 [95% CI 0.55–0.92]; $P < 0.01$).

Conclusions: Factor Xa inhibitors are effective for VTE management in patients with cancer; however, they are also associated with an increased bleeding risk compared to LMWH, but decreased when compared to VKA.

Open Peer Review

Reviewer Status AWAITING PEER REVIEW

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

bleeding, cancer, factor Xa inhibitor, oral anticoagulant, venous thromboembolism.

Corresponding author: Johannes Nugroho Eko Putranto (j.nugroho.eko@fk.unair.ac.id)

Author roles: **Nugroho Eko Putranto J:** Conceptualization, Investigation, Resources, Supervision, Validation, Writing – Review & Editing; **Wardhana A:** Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation; **Noor YA:** Investigation, Methodology, Project Administration, Resources, Software, Supervision, Writing – Original Draft Preparation; **Lambok Marnala Yosua Siahhaan P:** Formal Analysis, Funding Acquisition, Investigation, Resources, Validation; **Al Farabi MJ:** Data Curation, Investigation, Project Administration, Resources, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Indonesian Endowment Fund for Education (LPDP) Republic of Indonesia *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2021 Nugroho Eko Putranto J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Nugroho Eko Putranto J, Wardhana A, Noor YA *et al.* **Factor Xa inhibitor for venous thromboembolism management in patient with cancer: a systematic review and meta-analysis [version 1; peer review: awaiting peer review]** F1000Research 2021, **10**:1257 <https://doi.org/10.12688/f1000research.73883.1>

First published: 08 Dec 2021, **10**:1257 <https://doi.org/10.12688/f1000research.73883.1>

Introduction

Cancer patients are five times more likely to experience venous thromboembolism (VTE) than the general population.¹ Second only to cancer itself, VTE is the second most common cause of mortality in cancer patients.² According to previous clinical management recommendations, the typical VTE treatment in cancer patients involves the initial use of parenteral low-molecular-weight heparin (LMWH) followed by long-term use of oral vitamin K antagonists (VKA).³ However, recent recommendations proposed factor Xa inhibitors as one of the options of the main initial treatment for VTE.⁴

Factor Xa inhibitors are preferred over LMWH and VKA because they conveniently do not require injections every day compared to LMWH, their more predictable effects, lack of monitoring or frequent repeat doses, and fewer drug interactions compared to VKA.⁵ An earlier systematic review reported differences between factor Xa inhibitors and standard anticoagulation drugs in the incidence of recurrent VTE and major bleeding.⁶ Based on this research, the present meta-analysis aims to evaluate the effectiveness of factor Xa inhibitors for the management of venous thromboembolism, particularly in patients with cancer.

Ethical considerations

Ethical approval for this research was obtained from the Dr. Soetomo General Hospital Surabaya Ethical Committee in Health Research (1964/KEPK/IV/2020).

Trial registry

UMIN Clinical Trial Registry (UMIN ID 000040346).

Methods

We adopted the Preferred Reporting Items for Reviews and Meta-Analyses guidelines for analysis reporting.⁷ Any RCTs that studied VTE rates or major bleeding, as primary or secondary outcomes, in cancer patients who received an oral factor Xa inhibitor were included. Phase II trials, trials with an antiplatelet control group, and trials using an anticoagulant as VTE post-procedure prophylaxis were excluded.

We conducted a systematic search using the PubMed and Cochrane Library databases on April 24, 2020, after gaining approval from the Institutional Review Board. As for the title, abstract, and medical subject heading, we used search terms like “cancer,” “factor Xa inhibitor,” “oral anticoagulant,” “venous thromboembolism,” “apixaban,” “rivaroxaban,” “edoxaban,” “prophylaxis,” “bleeding,” “thromboembolism,” “thromboprophylaxis,” “randomized,” and “rct.”

We screened more studies by looking at the references in the included articles. Two investigators independently selected studies, with disagreements resolved through discussion and a third investigator's opinion. Thereafter, for each report, two investigators independently extracted the following information: authors, year of publication, trial name, cancer status, sample size, dose and duration of anticoagulation, duration of patient follow-up, and outcomes for the two treatment groups where available.

We determined four comparison groups: (1) factor Xa inhibitor versus LMWH as treatment for VTE; (2) factor Xa inhibitor versus VKA as treatment for VTE; (3) factor Xa inhibitor versus placebo as prophylaxis for VTE; (4) factor Xa inhibitor versus VKA as prophylaxis for VTE. The outcomes of our meta-analysis were recurrent VTE or new VTE event rates and incidence of major bleeding. VTE events were confirmed by leg vein ultrasound scanning, D-dimer testing, or both; alternatively, clinically overt pulmonary embolism was confirmed by imaging. Major bleeding was defined as in Schulman *et al.*⁸

The Cochrane Collaboration Risk of Bias Tool was used by two independent investigators to assess the methodological quality of included studies, and the GRADE approach was employed to grade each outcome.^{9,10} Any disputes were settled through discussion with a third investigator. We calculated odds ratios (ORs) for all outcomes at the longest follow-up period and used Review Manager (RevMan v5.3 2014) to apply the Mantel–Haenszel fixed-effects method. We conducted a modified intention-to-treat analysis including patients who had received ≥ 1 medication dose. We planned to conduct sensitivity analysis by removing studies likely to be biased. The I² statistic was used to assess statistical heterogeneity between studies. If the heterogeneity was $> 50\%$, we applied a random-effects model for analysis.¹¹

Results

The search identified 202 citations in PubMed and 41 in the Cochrane Library, among which 43 were duplicates (Figure 1). We found 22 more studies of which we evaluated the full text. Four studies were post-procedure prophylaxis




Figure 1. PRISMA flow diagram.

trials, three lacked a control, two were phase II trials, and two were extensions of included trials, so 11 were omitted. As a result, we could include 11 studies in our analysis.¹²⁻²²

Table 1 lists the characteristics of the included studies. There were four trials on apixaban, four on rivaroxaban, and three on edoxaban. The study size ranged from 300 to 1,170 patients. Five studies were subgroup analyses of patients with cancer from larger primary trials.¹²⁻¹⁶ We pooled their data only from the subgroup of patients with cancer, not all study population. One study was a pooled analysis of the subgroup of patients with cancer in “sister” trials.¹⁷ Four trials^{13,18-20} compared factor Xa inhibitors with LMWH, and three^{12,14,17} compared factor Xa inhibitors with VKA as a VTE treatment. Two trials^{15,16} compared factor Xa inhibitors with placebo and two^{21,22} compared factor Xa inhibitors with VKA as prophylaxis of VTE. We included one trial that investigated two doses of edoxaban for VTE prophylaxis, where the outcomes of both groups were combined and analyzed as one intervention group.¹⁶

The risk of bias across domains is presented in **Figure 2**. In most studies, the randomization process, adherence to the intervention, assessment, missing outcome results, and reporting were deemed adequate. In four trials, participants were blinded. The percentage of patients not followed up ranged from 0.2% to 5.6%. All trials reported the results from modified intention-to-treat analysis.

Table 1. The characteristics of the included trials.

Author	Blinding to subjects	Population	Randomized patients	Intervention	Dose	Control	Follow up period	Death	Lost to follow up
Prins et al., 2013; EINSTEIN-DVT and PE	No	Cancer patients with VTE (100% active cancer)	597	Rivaroxaban	15 mg bid for 3 wk followed by 20 mg qd	Heparin/VKA	3-12 months	30% vs 35%	N/A
Agnelli et al., 2015; AMPLIFY	Yes	Cancer patients with VTE (31.6% active cancer)	534	Apixaban	10 mg bid for 7 d followed by 5 mg bid	Heparin/VKA	6 months	N/A	N/A
Raskob et al., 2016; HOKUSAI-VTE	Yes	Cancer patients with VTE (48% active cancer)	771	Edoxaban	60 mg once daily	Heparin/VKA	3-12 months	N/A	N/A
Raskob et al., 2017; HOKUSAI-VTE	No	Cancer patients with VTE (97.9% active cancer)	1050	Edoxaban	60 mg once daily	Dalteparin (200 UI/kg/d during 30 days, then 150 UI/kg/d)	12 months	39% vs 36%	0.8% (3 vs 5)
Young et al., 2017; SELECT-D	No	Cancer patients with VTE (100% active cancer)	406	Rivaroxaban	15 mg bid for 3 wk followed by 20 mg qd	Dalteparin (200 UI/kg/d during 30 days, then 150 UI/kg/d)	6 months	75% vs 70%	0.2% (0 vs 1)
McBane et al., 2018; ADAM VTE	No	Cancer patients with VTE (100% active cancer)	300	Apixaban	10 mg bid for 7 d followed by 5 mg bid	Dalteparin (200 UI/kg/d during 30 days, then 150 UI/kg/d)	6 months	15% vs 10%	5.6% (9 vs 7)
Fanola et al., 2018; ENGAGE AF-TIMI	No	Cancer patients with AF (100% active cancer)	1153	Edoxaban	60 mg once daily or 30 mg once daily	VKA	> 2 years	32% vs 30%	N/A
Chen et al., 2019; ROCKET AF	No	Cancer patients with AF (7.8% active cancer)	640	Rivaroxaban	20 mg qd	VKA	2 years	10% vs 15%	N/A
Carrier et al., 2019; AVERT	Yes	Ambulatory patients with risk of VTE	574	Apixaban	2.5 mg bid	Placebo	6 months	12% vs 10%	4.3% (13 vs 11)
Khorana et al., 2019; CASSINI	Yes	Ambulatory patients with risk of VTE	841	Rivaroxaban	10 mg qd	Placebo	6 months	20% vs 25%	N/A
Agnelli et al., 2020; CARAVAGGIO	No	Cancer patients with VTE (97.3% active cancer)	1170	Apixaban	10 mg bid for 7 d followed by 5 mg bid	Dalteparin (200 UI/kg/d during 30 days, then 150 UI/kg/d)	6 months	23% vs 25%	1.7% (12 vs 8)

	Randomization	Adhering to intervention	Missing outcome data	Measurement	Reporting	Overall
Agnelli et al 2020	○	○	○	○	○	Low
McBane et al 2018	○	⊘	○	○	○	Some
Raskop et al 2017	○	○	○	○	○	Low
Young et al 2018	○	⊘	○	○	○	Some
Agnelli et al 2015	○	○	○	○	○	Low
Prins et al 2013	○	○	○	○	○	Low
Raskop et al 2016	○	○	○	○	○	Low
Chen et al 2019	○	○	○	○	○	Low
Fanola et al 2018	○	○	○	○	○	Low
Carrier et al 2019	○	⊘	○	○	○	Some
Khorana et al 2019	○	○	○	○	○	Low

○ : LOW RISK ⊘ : HIGH RISK

Figure 2. Risk of bias assessment.

The quality of evidence for each outcome analyzed using the GRADE approach is presented in [Table 2](#). We did not downgrade from the risk of bias, inconsistency, indirectness, and imprecision aspect of all outcomes, because of a low risk of bias, no substantial heterogeneity, a large enough sample size, and narrow confidence interval (CI). We downgraded one level for the major bleeding outcome because the funnel plot of major bleeding outcome suggested publication bias ([Figure 3](#)).

Seven studies involving 4,771 patients reported VTE recurrence ([Table 2](#)). Recurrence occurred in 4.9% (117/2,399) of patients allocated to factor Xa inhibitors, 9.1% (132/1,445) allocated to LMWHs, and 6.9% (64/927) of those allocated to VKAs. In comparison ([Figure 4](#)), the reduction of the risk of VTE recurrence with factor Xa inhibitors compared to LMWH was acceptable (four trials; OR 0.60; 95% CI 0.45–0.80; $P < 0.01$), without substantial heterogeneity ($I^2 = 26%$; $P = 0.26$). VTE recurrence rates were lower in patients treated with factor Xa inhibitors compared to patients treated using VKAs (three trials; OR 0.51; 95% CI 0.33–0.78; $P < 0.01$), without substantial heterogeneity ($I^2 = 0%$; $P = 0.37$).

Three studies, including 2,056 patients, reported the incidence of new VTE after anticoagulant prophylaxis. The factor Xa inhibitor group had a 4.1% (42/1,021) VTE occurrence rate, while the VKA and placebo groups each had 1.45% (5/355) and 9.6% (65/680), respectively. According to the meta-analysis shown in [Figure 5](#), there were similar VTE incidences in the factor Xa inhibitor and the VKA groups (one trial; OR = 1.08 [95% CI, 0.31–3.77]; $P = 0.90$); however, the heterogeneity analysis could not be applied. The estimated effect of factor Xa inhibitors on VTE incidence compared to placebo showed a statistically significant reduction (two trials; OR = 0.54 [95% CI, 0.35–0.81]; $P < 0.01$), without substantial heterogeneity ($I^2 = 31%$; $P = 0.23$).

Table 2. Summary of findings.

	No of studies	Total participants	Pooled OR (95% CI)	P	I2 (P)	GRADE
Recurrence						High
vs LMWH	4	2890	0.60 (0.45, 0.80)	0.0004	26% (0.26)	
vs VKA	3	1881	0.51 (0.33, 0.78)	0.002	0% (0.37)	
New VTE						High
vs VKA	1	684	1.08 (0.31, 3.77)	0.90	N/A	
vs Placebo	2	1372	0.54 (0.35, 0.81)	0.003	31% (0.23)	
Major bleeding						Moderate
vs LMWH	4	2890	1.34 (0.83, 2.18)	0.23	28% (0.25)	
vs VKA	5	3703	0.71 (0.55, 0.92)	0.009	0% (0.72)	
vs Placebo	2	1372	1.98 (0.88, 4.44)	0.10	0% (0.96)	




Figure 3. Funnel plot of (A) recurrent VTE outcome; (B) new VTE outcome; (C) major bleeding outcome.




Figure 4. Forest plot of recurrent VTE outcome.

Eleven studies, including 7,965 patients, reported major bleeding (Table 2). Major bleeding occurred in 5.5% (231/4,178) of patients allocated to factor Xa inhibitors, 3.6% (52/1445) to LMWHs, 8.1% (134/1,662) to VKAs and 1.3% (9/680) to placebo. According to the meta-analysis shown in Figure 6, the acceptable increase of risk cannot be confirmed from the description of major bleeding with factor Xa inhibitors compared to LMWH, as based on an OR of 1.34 (95% CI, 0.83–2.18) with a P = 0.23, which is not statistically significant. However, factor Xa inhibitors significantly reduced the risk of major bleeding compared to VKAs (five trials; OR = 0.71 [95% CI, 0.55–0.92]; P = 0.009), without substantial heterogeneity (I2 = 0%; P = 0.72). The risk of major bleeding was higher with factor Xa inhibitors versus placebo




Figure 5. Forest plot of new VTE outcome.




Figure 6. Forest plot of major bleeding outcome.

(two trials; OR = 1.98 [95% CI, 0.88–4.44]; P = 0.10) but not statistically significant, without substantial heterogeneity (I² = 0%; P = 0.96).

Discussion

The aim of this meta-analysis was to determine the efficacy and safety of factor Xa inhibitors for VTE treatment in cancer patients. Recurrence was 4.9%, 9.1%, and 6.9% for the factor Xa inhibitor, LMWH, and VKA groups, respectively. All were lower than the findings of a retrospective cohort study which reported an incidence of 13.1%, 17.6%, and 17.9%, respectively.²³ Our review of four studies involving over 4,771 patients found that factor Xa inhibitors were associated with a lower risk of VTE recurrence when compared to LMWH, and even lower when compared to VKA. This result was consistent with a recent meta-analysis which combined data from RCTs and retrospective cohort studies.²⁴

Another finding in our meta-analysis in terms of safety profiles was that factor Xa inhibitors were associated with an increased risk of bleeding when compared to LMWH, but a lower risk when compared to VKA. This result is in line with the findings of other systematic reviews.^{24–26} However, another meta-analysis found a significantly higher incidence of bleeding (two trials, OR= 2.72 [95% CI: 1.05–7.01]; P= 0.039) with factor Xa inhibitors, relative to LMWH.²⁷ Importantly, the bleeding outcome in comparison to LMWH was the result of pooled data from nonspecific cancer patients. The results of the analysis of major bleeding in comparison to LMWH were mainly influenced by those of the HOKUSAI VTE Cancer trial and the recent CARAVAGGIO trial.^{28,29} Both had different results: the former showed significantly higher bleeding in the edoxaban group while the second showed similar major bleeding events between groups.

Our meta-analysis also provided information about the efficacy of factor Xa inhibitors as prophylaxis, which suggested that, compared to placebo, it can significantly reduce VTE incidence. According to a recent clinical practice guideline, high-risk cancer outpatients can receive thromboprophylaxis with a factor Xa inhibitor or LMWH, in the absence of major risk factors for bleeding.³⁰ The high cost and the pain of daily LMWH injections was avoided with the factor Xa inhibitor regimen.

With respect to factor Xa inhibitors and LMWH, the inclusion of the CARAVAGGIO trial, with highly rigorous evidence, increased the accuracy of the estimated outcomes. There are a number of limitations to the current meta-analysis: the majority of the data corresponded to subgroup or post-hoc analyses. Further, the following variables were not controlled for: cancer stage, type of cancer, follow-up period. While most of the included studies evaluated patients for six months, the optimal duration of anticoagulation treatment was not evaluated to achieve an agreement. Finally, despite our systematic electronic database search and our investigation of the references in the included studies, we may have missed relevant studies.

Conclusion

Factor Xa inhibitors are effective for VTE management in patients with cancer; however, they are also associated with an increased bleeding risk compared to LMWH, but decreased when compared to VKA.

Data availability statement

Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

Reporting guidelines⁸

Figshare: PRISMA checklist for 'Factor Xa inhibitor for venous thromboembolism management in Patients with cancer: a systematic review and meta-analysis'. <https://doi.org/10.6084/m9.figshare.16590086.v3>³¹

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/) (CC-BY 4.0).

Acknowledgments

This work was performed as part of Johanes Nugroho employment at the Department of Cardiology and Vascular Medicine, Universitas Airlangga/Dr. Soetomo General Hospital, East Java, Indonesia. The authors also would like to thank Enago (www.enago.com) for the English language review and this work was supported by the Indonesian Endowment Fund for Education (Lembaga Pengelola Dana Pendidikan).

References

1. Elyamany G, Alzahrani AM, Bukhary E: **Cancer-associated thrombosis: an overview.** *Clin Med Insights Oncol.* 2014; **8**: 129–137. [PubMed Abstract](#) | [Publisher Full Text](#)
2. Khorana AA, Francis CW, Culakova E, et al.: **Thromboembolism is a leading cause of death in cancer patients receiving outpatient chemotherapy.** *J. Thromb. Haemost.* 2007; **5**: 632–634. [PubMed Abstract](#) | [Publisher Full Text](#)
3. Kearon C, Akl EA, Comerota AJ, et al.: **Antithrombotic therapy for VTE disease: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines.** *Chest.* 2012; **141**: e419S–e496S. [Publisher Full Text](#)
4. Lyman GH, Carrier M, Ay C, et al.: **American Society of Hematology 2021 guidelines for management of venous thromboembolism: prevention and treatment in patients with cancer.** *Blood Adv.* 2021; **5**(4): 927–974. [PubMed Abstract](#) | [Publisher Full Text](#)
5. Fox BD, Kahn SR, Langleben D, et al.: **Efficacy and safety of novel oral anticoagulants for treatment of acute venous thromboembolism: direct and adjusted indirect meta-analysis of randomised controlled trials.** *BMJ.* 2012; **345**: e7498. [PubMed Abstract](#) | [Publisher Full Text](#)
6. Robertson L, Kesteven P, McCaslin JE: **Oral direct thrombin inhibitors or oral factor Xa inhibitors for the treatment of pulmonary embolism.** *Cochrane Database Syst. Rev.* 2015;

- 2016(Issue 12): Art. No.: CD010957.
[Publisher Full Text](#)
7. Moher D, Liberati A, Tetzlaff J, *et al.*: **The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement.** *Open Med.* 2009; **3**: 123–130.
 8. Schulman S, Kearon C: **Definition of major bleeding in clinical investigations of antithrombotic medicinal products in non-surgical patients.** *J. Thromb. Haemost.* 2005; **3**: 692–694.
[PubMed Abstract](#) | [Publisher Full Text](#)
 9. Sterne JAC, Savović J, Page MJ, *et al.*: **RoB 2: a revised tool for assessing risk of bias in randomised trials.** *BMJ.* 2019 Aug 28; **366**: l4898.
[Publisher Full Text](#)
 10. Ryan R, Hill S: *How to GRADE the quality of the evidence Version 3.0.* Cochrane Consumers and Communication Group; 2018 Dec [cited 2019 July 13]; [about 25 p.].
[Reference Source](#)
 11. Higgins JPT, Green S: *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0.* The Cochrane Collaboration; 2011 [cited 2019 July 10].
[Reference Source](#)
 12. Raskob GE, van Es N, Segers A, *et al.*: **Edoxaban for venous thromboembolism in patients with cancer: results from a non-inferiority subgroup analysis of the Hokusai-VTE randomised, double-blind, double-dummy trial.** *Lancet Haematol.* 2016; **3**: e379–e387.
[PubMed Abstract](#) | [Publisher Full Text](#)
 13. Raskob GE, van Es N, Verhamme P, *et al.*: **on behalf of the Hokusai VTE Cancer Investigators. Edoxaban for the treatment of cancer-associated venous thromboembolism.** *N. Engl. J. Med.* 2018; **378**(7): 615–624.
[Publisher Full Text](#)
 14. Agnelli G, Buller HR, Cohen A, *et al.*: **Oral apixaban for the treatment of venous thromboembolism in cancer patients: Results from the AMPLIFY trial.** *J. Thromb. Haemost.* 2015; **13**(12): 2187–2191.
[PubMed Abstract](#) | [Publisher Full Text](#)
 15. Chen ST, Hellkamp AS, Becker RC, *et al.*: **Efficacy and safety of rivaroxaban vs. warfarin in patients with non-valvular atrial fibrillation and a history of cancer: observations from ROCKET AF.** *Eur Heart J Qual Care Clin Outcomes.* 2019; **5**(2): 145–152.
[PubMed Abstract](#) | [Publisher Full Text](#)
 16. Fanola CL, Ruff CT, Murphy SA, *et al.*: **Efficacy and safety of edoxaban in patients with active malignancy and atrial fibrillation: analysis of the ENGAGE AF - TIMI 48 trial.** *J Am Heart Assoc.* 2018; **7**(16): e008987.
[PubMed Abstract](#) | [Publisher Full Text](#)
 17. Prins MH, Lensing AW, Bauersachs R, *et al.*: **Oral rivaroxaban versus standard therapy for the treatment of symptomatic venous thromboembolism: a pooled analysis of the EINSTEIN-DVT and PE randomized studies.** *Thromb J.* 2013; **11**(1): 21.
[PubMed Abstract](#) | [Publisher Full Text](#)
 18. Agnelli G, Becattini C, Meyer G, *et al.*: **Apixaban for the treatment of venous thromboembolism associated with cancer.** *N Engl J Med.* 2020; **382**: 1599–1607.
[Publisher Full Text](#)
 19. Young AM, Marshall A, Thirlwall J, *et al.*: **Comparison of an oral factor Xa inhibitor with low molecular weight heparin in patients with cancer with venous thromboembolism: Results of a randomized trial (SELECT-D).** *J Clin Oncol.* 2018; **36**(20): 2017–2023.
[PubMed Abstract](#) | [Publisher Full Text](#)
 20. McBane RD 2nd, Wysokinski WE, Le-Rademacher JG, *et al.*: **Apixaban, dalteparin, in active cancer associated venous thromboembolism, the ADAM VTE trial.** *Blood.* 2020; **18**(2): 411–421.
 21. Khorana AA, Soff GA, Kakkar AK, *et al.*: **Rivaroxaban for thromboprophylaxis in high-risk ambulatory patients with cancer.** *N Engl J Med.* 2019; **380**: 720–728.
[PubMed Abstract](#) | [Publisher Full Text](#)
 22. Carrier M, Abou-Nassar K, Mallick R, *et al.*: **Apixaban to prevent venous thromboembolism in patients with cancer.** *N Engl J Med.* 2019; **380**: 711–19.
[Publisher Full Text](#) | [PubMed Abstract](#)
 23. Streiff MB, Mccrae K, Yannicelli D, *et al.*: **Effectiveness and safety of anticoagulants for the treatment of venous thromboembolism in patients with cancer.** *Am J Hematol.* 2018; **93**(5): 664–671.
[PubMed Abstract](#) | [Publisher Full Text](#)
 24. Yang M, Li J, Sun R, *et al.*: **Comparison between direct factor Xa inhibitors and low-molecular-weight heparin for efficacy and safety in the treatment of cancer-associated venous thromboembolism: A meta-analysis.** *J Can Res Ther.* 2019; **15**: 1541–1546.
[PubMed Abstract](#) | [Publisher Full Text](#)
 25. Fuentes HE, McBane RD 2nd, Wysokinski WE, *et al.*: **Direct Oral Factor Xa Inhibitors for the Treatment of Acute Cancer-Associated Venous Thromboembolism: A Systematic Review and Network Meta-analysis.** *Mayo Clin Proc.* 2019; **94**(12): 2444–2454.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. Li A, Garcia DA, Lyman GH, *et al.*: **Direct oral anticoagulant (DOAC) versus low-molecular-weight heparin (LMWH) for treatment of cancer associated thrombosis (CAT): A systematic review and meta-analysis.** *Thromb Res.* 2019; **173**: 158–163.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. Brunetti ND, Gesuete E, De Gennaro L, *et al.*: **Direct oral anti-coagulants compared with vitamin-K inhibitors and low-molecular-weight-heparin for the prevention of venous thromboembolism in patients with cancer: A meta-analysis study.** *Int J Cardiol.* 2017; **230**: 214–221.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Hokusai VTE Cancer Investigators: **Edoxaban for the Treatment of Cancer-Associated Venous Thromboembolism.** *N Engl J Med.* 2018; **378**: 673–674.
[Publisher Full Text](#)
 29. Caravaggio Investigators: **Apixaban for the Treatment of Venous Thromboembolism Associated with Cancer.** *N Engl J Med.* 2020; **382**: 1599–1607.
[Publisher Full Text](#)
 30. Key NS, Khorana AA, Kuderer NM, *et al.*: **Venous thromboembolism prophylaxis and treatment in patients with cancer: ASCO clinical practice guideline update.** *J Clin Oncol.* 2020; **38**: 496–520.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Al Farabi, Jibril M: **PRISMA XA Checklist and Diagram.** figshare. Dataset. 2021.
[Publisher Full Text](#)

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research