

Dexamethasone and olmesartan as potential antiremodelling agents of valvular interstitial cell into myofibroblast: In vitro study on α - smooth muscle actin expression

by Ardianto Nandiwardhana

Submission date: 22-Jan-2023 03:04PM (UTC+0800)

Submission ID: 1996838964

File name: 26DEXA_1.PDF (854.05K)

Word count: 5156

Character count: 29944

12

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/363081875>

Dexamethasone and olmesartan as potential antiremodelling agents of valvular interstitial cell into myofibroblast

Article in *International Journal of Health Sciences* · August 2022

DOI: 10.53730/ijhs.v6i59.12283

CITATIONS

0

READS

37

7 authors, including:



Ardianto Nandiwardhana

3 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



Eka Prasetya Budi Mulia

Airlangga University

59 PUBLICATIONS 74 CITATIONS

[SEE PROFILE](#)



David Nugraha

Airlangga University

22 PUBLICATIONS 75 CITATIONS

[SEE PROFILE](#)



Achmad Lefi

Dr Soetomo General Hospital

25 PUBLICATIONS 35 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



[COVID-19 View project](#)



[Polymorphism of CT49, CT1822, and PTPN22 in Type 1 Diabetes Mellitus in Pediatric Patients View project](#)

How to Cite:

Nandiwardhana, A., Suwanto, D., Mulia, E. P. B., Nugraha, D., Lefi, A., Budiarto, M., & Nugroho, J. (2022). Dexamethasone and olmesartan as potential antiremodelling agents of valvular interstitial cell into myofibroblast. *International Journal of Health Sciences*, 6(S9), 379–391. <https://doi.org/10.53730/ijhs.v6nS9.12283>

Dexamethasone and olmesartan as potential antiremodelling agents of valvular interstitial cell into myofibroblast: *In vitro* study on α -smooth muscle actin expression

Ardianto Nandiwardhana

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga - Dr. Soetomo General Hospital, Surabaya, Indonesia
Corresponding author email: wardhanaaddo@gmail.com

Denny Suwanto

22

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga - Dr. Soetomo General Hospital, Surabaya, Indonesia

Eka Prasetya Budi Mulia

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga - Dr. Soetomo General Hospital, Surabaya, Indonesia

David Nugraha

Faculty of Medicine, Universitas Airlangga - Dr. Soetomo General Hospital, Surabaya, Indonesia

Achmad Lefi

1

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga - Dr. Soetomo General Hospital, Surabaya, Indonesia

Mohammad Budiarto

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga - Dr. Soetomo General Hospital, Surabaya, Indonesia

Johanes Nugroho

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga - Dr. Soetomo General Hospital, Surabaya, Indonesia

Abstract--Rheumatic heart disease is a late complication of valvular inflammation caused by rheumatic fever. Studies have shown that the differentiation of valvular interstitial cells (VIC) into fibroblasts plays an important role in valvular remodeling and fibrosis. Various strategies to minimize valvular fibrosis has increased recently. This study aims to analyze the effect of dexamethasone, olmesartan, and its

combination in inhibiting TGF- β 1-induced VIC differentiation into myofibroblast. In vitro laboratory experimental-posttest only control group design was conducted. Isolated VIC of *Oryctolagus cuniculus* was pretreated using 2,5 ng/mL of TGF- β 1 and divided into groups of dexamethasone (0.1 μ M/L), olmesartan (100 nmol/L), and its combination. Inhibition of myofibroblast differentiation was quantified by the expression of α -SMA levels detected by immunofluorescence. Dexamethasone, olmesartan, and its combination administration were significantly reduced TGF- β 1-induced VIC differentiation into myofibroblast expressed by α -SMA levels (dexamethasone 6823 \pm 1735.3, olmesartan 6683.7 \pm 2795.05). Combination of dexamethasone and olmesartan exhibit the most potent inhibition compared to control (5051.87 \pm 1612.210 vs 22286.73 \pm 2780.2; $p < 0.000$). In conclusion, dexamethasone, olmesartan, and the combination can significantly reduce the differentiation of VIC into myofibroblasts. The greatest potential is the combined effect of dexamethasone and olmesartan, while dexamethasone and olmesartan have the same potential.

Keywords--Valvular interstitial cell, myofibroblast, α -Smooth muscle actin, dexamethasone, Olmesartan

27

Introduction

Rheumatic heart disease (RHD) is a sequela of acute rheumatic fever with serious long term complication and morbidity which still a significant public health problem with 33.4 million cases and mortality rate up to 319,400 in 2015.(Guilherme & Kalil, 2020; Ordunez et al., 2019; Watkins et al., 2017) One of the serious sequela of RHD is the progression of mitral valve damage in 50-60% of cases, multivalvular and mixed lesions are found in 20%, with the most common pathology being stenosis (34% mitral; 9% aortic). The long-term effectiveness of mitral valve surgery, both mitral valve replacement and mitral valve repair, has not shown satisfactory results with 10-year survival rate of 62% in the mitral valve replacement group and 46% in the mitral valve repair group.(Thourani et al., 2003) Meanwhile, in individuals undergoing percutaneous mitral commissurotomy (PMC), prior prospective cohort study showed a survival rate of 68% at 20 years and only 16% of patients did not require reinvention.(Bouleti et al., 2013)

The use of PMC and valve replacement or repair surgery as the currently available standard therapy, shows the low long-term effectiveness of valve intervention measures. Interventions using surgery and minimally invasive procedures are practically only focused on the valve leaflets and annulus, even though the pathological process also occurs at the cellular level, especially fibrosis in the sub-valvular apparatus, ventricular, and atrial myocardial remodeling due to pressure/volume overload exposure and chronic inflammation.(Yankah et al., 2011) Intervention in the early inflammatory phase using antifibrotic agents is a new hope to break the bridge of acute rheumatic fever into valvular pathology in RHD. Chronic inflammatory exposure involving TGF- β 1 in valve interstitial cells

(VIC) due to cross-reactivity of group A streptococcal M protein, induces activation of the *Smad* signaling pathway, mitogen-activated protein kinase (MAPK), and extracellular signal-regulated kinase 1/2 (ERK1/2) will cause myofibroblastic differentiation, induction of extracellular matrix synthesis resulting in fibrosis and distortion of valve architecture.(Vaideeswar & Butany, 2016)

Valve stenosis is a terminal process of the acute phase of valvular disease inflammatory response. Various methods of treating valve stenosis surgically and minimally invasive methods have been developed, but limitations in terms of long-term benefit, survival, patient selection, and complications limit the application of this method. This issue has recently attracted interest to find novel antiremodelling agents. The antifibrotic effect of angiotensin receptor blockers (ARBs) are demonstrated through antagonism of the angiotensin II type I receptor (AT1R) on local Ang II exposure in inflamed tissues, resulting in inhibition of myofibroblast growth and induction of extracellular matrix synthesis.(Yang et al., 2005) In addition, early studies revealed that glucocorticoids also may exert some anti-atherosclerosis effects with reduced atherosclerosis severity in animal model.(Asai et al., 1993; Makheja et al., 1989) The onset of atherosclerosis is triggered by proinflammatory mediators, which induce adhesion molecules in endothelial cells by activating MAPK, especially p38 MAPK.

Furthermore, inhibition of fibrogenesis via transforming growth factor (TGF- β 1) pathway using various agents has been investigated in pulmonary alveolar cells (Guzy, 2020), renal glomerular cells (Isaka, 2018), hepatocytes(Masuda et al., 2020), myocytes (Parichatikanond et al., 2020), and fibroblasts ventricles(Peng et al., 2010; Zhou et al., 2015). However, to the best of our knowledge, there have been no studies examining the antifibrotic properties of ARBs and steroids on VIC. Based on these considerations, this study aimed to assess the effectiveness of dexamethasone and olmesartan in inhibiting fibrogenesis of VIC in the early inflammatory phase, so that the process of valve fibrosis can be prevented in the acute phase of the disease and patients can avoid long-term sequelae.

Method

1.1 Study design and animal model

This in vitro post-test control group only study was conducted in Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia. Some of the interventions given were (1) Dexamethasone; (2) Olmesartan; (3) Dexamethasone-Olmesartan combination. The experimental subject used in this study was valvular interstitial cell (VIC) induced by TGF- β 1. The experimental unit was obtained from VIC cultures taken from the valves of New Zealand rabbits (*Oryctolagus cuniculus*), aged 12-13 weeks with weight of 2.5-3 kg. The model then divided into three groups, namely the dexamethasone group, olmesartan group, and a combination group. The sample size calculated using the Higgins and Klinbaum formula was 6 replications.

1.2 Isolation, culture, and characterization of VIC

Isolation of rabbit VIC was performed as previously described by Linn et al with some modification on animal model selection. (Lin et al., 2017) Rabbits which met the standardized inclusion criteria were terminated by lethal injection using midazolam 5 mg/kg and phenytoin 125 mg/kg intraperitoneally. Surgically extracted rabbit leaflets were then separated from myocytes and valvular endothelial cells by digestion method using collagenase II enzyme followed by centrifugation and seeding of cell suspensions. Adhered cells after centrifugation at 100g were transferred to culture media.

Disintegrated VIC cells were cultured in Dulbecco's Modified Eagle Medium (DMEM; Gibco BRL, Grand Island, NY, USA) with supplementation of 10% foetal bovine serum (FBS), and 1% penicillin/streptomycin solution at 37°C with 95% humidity and 5% CO₂. After reaching 80% confluency, cells from the second passage were then transferred to non-coated six wells with DMEM and 10% FBS at a density of 2600 cells/dish. After 24 hours, the media was replaced with new DMEM containing 10% FBS and TGF- β 1 (2.5 ng/ml) and then incubated for one to seven days.

Characterization of VIC was done through measuring vimentin expression. After washing with phosphate buffer saline (PBS), cells were fixed with 3% formaldehyde for 15 minutes. In order to inhibit non-specific binding, PBS containing 1% serum was added for 15 minutes and washed again using PBS. Subsequently, the fluorescein isothiocyanate (FITC)-labeled anti-vimentin from Thermo Fisher Scientific (Schwerte, Germany) reagent was added, and washed again with PBS. Vimentin expression was documented by fluorescent microscopy.

1.3 Intervention and myofibroblast differentiation assessment protocol

After the cells reached 80-90% confluency, the medium was replaced with serum free DMEM. Afterwards, pre-treatment condition was carried out with Olmesartan 100 nM, Dexamethasone 1000 nM, along with antifibrotic combination for 3 hours followed by incubation with TGF β 1 2.5 ng/ml for 24 hours. Myofibroblast characterization was carried out by fixation with 2% paraformaldehyde/PBS and 1% Triton X-100/PBS for 30 minutes. Then the cells were blocked with 2% BSA/PBS and incubated with mouse monoclonal antibody against FITC-labeled α -SMA, then washed again with PBS. α -SMA expression was documented by fluorescent microscopy.

1.4 Statistical analysis

After each experiment was carried out six times, the data were then collected and tabulated using Microsoft Office Excel 2016 and were further processed using IBM SPSS ver.26 for Windows (SPSS Inc, Chicago, IL). After the data cleaning, descriptive analysis was performed to calculate the mean, standard deviation, and box-plot diagrams for each group. In addition, unpaired numerical comparative statistical analysis in both groups was carried out using unpaired T-test for normal distributed data, otherwise the Mann-Whitney test was used. For comparison among three or more groups, one way analysis of variance (ANOVA)

with post hoc Bonferroni was conducted for the same variance, otherwise ANOVA with post hoc Games Howell was performed, but if the data not normally distributions then Kruskal-Wallis with post hoc Mann-Whitney will used. p-value <0.05 was considered as statistically significant.

Discussion

The results of the cultured and isolated rabbit *Oryctolagus cuniculus* valvular' cells in this study showed islands of cuboidal-spindle shape elongated cells with regular cell membranes that visually under light microscope, as demonstrated in figure 1. The characterization of VIC was carried out with positive markers of vimentin through immunocytochemical examination. Immunocytochemical analysis showed that there was a positive expression (luminescence above 95% of cell luminescence per field of view) of vimentin. Immunocytochemical staining results showed positive markers for vimentin which is a marker of valve interstitial cells (figure 2). The vimentin marker expression in the process of VIC isolation after incubation on the 9th day was detected with median of 6,585.4 (384 – 16,909).




Figure 1 Valve Interstitial Cell Morphology A) Magnification 4 x 10; B) Magnification 10x10. The spindle-elongated cells are seen as valve interstitial cells.




Figure 2 Vimentin Expression in Valve Interstitial Cells A) Magnification 4 x 10; B) Magnification 10 x 10. Valve interstitial cells appeared evenly distributed across the visual field tightly with strong luminescence

Subsequently, after exposure to TGF- β 1, VIC differentiated into myofibroblasts with characterized as irregular surfaces and cytoplasm filled with stress fibers. The Wilcoxon test results showed in the group exposed to TGF- β 1 were significantly different ($p < 0.001$) compared to the control group. This suggests TGF- β 1 as a potent cytokine that induce VIC differentiation into myofibroblasts. Furthermore, immunocytochemical staining after exposure to TGF- β 1 showed a significant increase in α -SMA expression (fluorescence above 95% luminescence of cells per field of view) when compared to controls. This indicates the process of differentiation of VIC into myofibroblasts (figure 3). The median quantity value of α -SMA expression in VIC exposed to TGF- β 1 as an inductor of myofibroblastic differentiation obtained a median of 23,474 (18,452 – 25,671), whereas in the control group the median expression of α -SMA was only 1,075 (118 – 7,696).




Figure 3 A) Myofibroblast Cell Morphology and α -SMA Expression in Immunocytochemistry with TGF- β 1 Induction B) Myofibroblast Cell Morphology and α -SMA Expression in Control Group Immunocytochemistry

Quantity of α -SMA expression in VIC exposed to TGF- β 1 and treated with dexamethasone, olmesartan, and combination showed that the lowest α -SMA expression was observed in the combination group of dexamethasone and olmesartan with a mean of $5,051.8 \pm 1,612.2$. The α -SMA expression in the dexamethasone group was $6,823 \pm 1735.3$ and the α -SMA expression in the olmesartan group was $6,942.7 \pm 2,455.9$. Statistical analysis using Kruskal-Wallis showed significant difference in these groups (p -value < 0.001).

Visual expression of α -SMA were shown in figure 4. These data were further analyzed by the Mann-Whitney post hoc test, and it was found that there were significant differences between the dexamethasone and control groups ($p < 0.001$), dexamethasone in combination ($p < 0.001$), olmesartan with control ($p < 0.001$), olmesartan with combination ($p < 0.004$) and combination with control ($p < 0.001$), but there was no difference between the dexamethasone and olmesartan groups ($p < 0.705$) (figure 5). According to these results, it demonstrated that the combination of dexamethasone and olmesartan can significantly reduce the

differentiation of rabbit VIC into myofibroblasts based on the expression of α -SMA, while dexamethasone and olmesartan alone have same potential




Figure 5 (A) Dexamethasone Group in Phase Contrast. (B) α -SMA Immunofluorescence Expression Group in the Dexamethasone Group. (C) Olmesartan Group in Phase Contrast. (D) Expression of α -SMA Immunofluorescence in the Olmesartan Group. (E) Combination Group in Phase Contrast. (F) Expression of α -SMA Immunofluorescence in the Combination Group. (G) Control Group on Contrast Phase, (H) Expression of α -SMA Immunofluorescence on contrast phase microscope.




Figure 4 Comparison of each intervention treatment group on α -SMA expression. It can be seen that there is a significant difference between the groups against the control.

To the best of our knowledge, this is the first research that used rabbit VIC to make in vitro fibrosis models with single induction using TGF- β 1. This study proved the role of TGF- β 1 as an inducer of VIC differentiation as evidenced by changes in cell morphology and the presence of significant quantity of α -SMA luminescence. TGF- β 1 can activate the small-mothers-against-decapentaplegic (Smad) transduction pathway and initiate transcription of myofibroblastic pro-differentiation genes.(Cucoranu et al., 2005) TGF- β 1 has been shown to act as a key profibrogenic cytokine in vitro and in vivo studies(Bujak, 2001; Calderone et al., 2001; Frangogiannis et al., 2000), as well as an increase in TGF- β 1 levels in fibrotic heart organs.(Kim et al., 1995; Kupfahl et al., 2000) Prior study also revealed the effect of TGF-1 exposure on ventricular fibroblast cells to induce the fibrosis process through induction of collagen production by fibroblasts, myofibroblasts, 50 and induction of fibroblast differentiation into myofibroblasts.(Lijnen et al., 2000; Petrov et al., 2002) Our study was successfully able to produce VIC isolates similar to the method used by Lin et al (2017), but with a lower potential for contamination of ventricular or atrial myocytes and fibroblasts through rational selection of experimental animals.(Lin et al., 2017)

TGF- β 1 as one of the main profibrotic cytokines is widely involved in the pathogenesis of RHD. This cytokine has the ability to stimulate myofibroblastic differentiation which will functionally increase the upregulation of α -SMA, to produce inflammatory mediators, growth factors, and increase the production of extracellular matrix such as the secretion of collagen I, III, and fibronectin which amplify the fibrosis process. Induction of TGF- β 1 will initiate signaling cascades through cell membrane and intracellular protein receptors Smad 2/3, the Mitogen-Activated Protein Kinase (MAPK) pathway, and the c-Jun N-terminal kinase (JNK) pathway.(Du et al., 2020; Heldin et al., 1997) The characteristics of myofibroblast cells that we found were similar to previous result reported with high content of exocytic vesicles, irregular cell membranes, and stress fibers in the cell cytoplasm.(Baum & Duffy, 2011; Petrov et al., 2002)

Myofibroblastic differentiation is a physiological response of VIC after exposure to pathological injury. This induces TGF- β 1 expression and initiates smooth muscle actin transcription. Myofibroblasts will increase the ability of collagen production and molecular adhesion (paxilin, tensin, ED-A fibronectin).(Frangogiannis et al., 2000; Gabbiani, 2003) Myofibroblasts also have high levels of microfilaments with abundant extracellular fibronectin, which can cause traction on histological tissue structures.(Gabbiani, 2003) The transformation of VIC into myofibroblasts will then cause excessive collagen deposition, thereby increasing the distance between functional VIC and disrupting cell-cell adhesion junctions (Connexin 43 and Connexin 45), as well as the effect of tissue contraction, which will bind to impaired valve function at the macroscopic level.(Baum & Duffy, 2011)

This study demonstrated the potential antiremodelling properties of dexamethasone and olmesartan as well as their combination in successfully inhibit VIC differentiation into myofibroblastic. Previous study has shown that dexamethasone treatment resulted in high extracellular matrix deposition.(Lo et al., 2003) It was also described that several components of the myofibroblast process could be inhibited by dexamethasone as well as inhibit α -SMA

amplification in endothelial injury.(Gu et al., 2010) Actin has been implicated as a major element in cellular movement and is associated with increased cellular migration.(Christian et al., 2001) Consequently, it is reasonable to assume that decreased expression of one of its isoforms would lead to decreased cell motility. In the study by Tsai et.al, cell migration decreased in parallel with decreased α -SMA expression after administration with dexamethasone.(Tsai et al., 2003) These findings demonstrate a direct relationship between cell migration and α -SMA expression and suggest that this protein plays a crucial role in driving cell migration. Moreover, the antifibrotic properties of dexamethasone can also be achieved through the induction of the MKP-1 pathway through increased levels of 15-deoxy-delta -12,14-PGJ2 (15DPGJ2), an endogenous MKP-1 ligand, via the p38-MAPK pathway. Activation of the MKP-1 pathway then act as an inhibitor of NF- κ B pathway activation.(Almawi et al., 1996)

Angiotensin receptor blocker (ARB) drugs, one of which is olmesartan, has an antiremodelling effect through inhibition of the RAAS system which suppresses the production of AngII. AngII has the role in enhancing the process of fibrogenesis in various organs has been demonstrated in various studies. Likewise, inhibition of the RAAS hormone using ARB has been shown to provide benefits in the process of organ fibrosis.(Kim et al., 1995) This Inhibition effect can be an intervention target to suppress the process of fibrosis and inflammation in VIC. Our results also in accordance with a previous study revealed that administration of 10 nM olmesartan was shown to reduce the levels of TGF- β 1 and α -SMA in cardiac fibroblasts.(Zhou et al., 2015) In addition, olmesartan administration 5 mg/kg/day in rats showed changes in profibrotic parameters, including: decreased activation of the TGF-1/Smad signaling pathway, increased levels of ACE2 and TNF- Converting Enzyme (TACE), and inhibition of apoptotic effects characterized by decreased levels of Caspase 3, BAX, Bcl-2.(Suh et al., 2019) The study using ARB as an inhibitory agent of fibroblast cell differentiation and its equivalent, is similar to the findings in this study, which showed that administration of olmesartan at a dose of 10 nanomol/L could significantly inhibit the differentiation of VIC.

In this study, the strongest potential in inhibiting myofibroblastic differentiation of VIC was found in the dexamethasone-olmesartan combination group. This is because the combination mechanism of action through different pathways that mutually potentiate each other. Olmesartan inhibits the interaction of ATII molecules at the ATII type 1 receptor, thereby reducing the synthesis of Ras molecules and activation of the JAK/STAT pathway, while dexamethasone acts as a negative inhibitor and regulator of Smad 2/3, TNF α , IL-1 and downstream NF- κ B signaling pathways. pro-myofibroblast differentiation. The combination of these different working points is mutually potentiating so that it provides the most potent inhibition effect of myofibroblast differentiation compared to the use of a single agent.

To date, there is no combination tested in the context of in vitro VIC model with TGF- β 1 induction. The finding of this potential inhibitory effect of VIC differentiation into myofibroblasts, especially the dexamethasone-olmesartan combination that we chose with in-depth consideration of its simultaneous and potentiating effects through various cell differentiation pathways, can be a

reference for further research in exploring more deeply the antifibrotic potential of dexamethasone, olmesartan, and dexamethasone. The combination in order to be an alternative to pharmacological therapy to inhibit the fibrosis process in RHD.

Conclusion

Dexamethasone and olmesartan has inhibitory effects on the differentiation of rabbit VIC induced by TGF- β 1 into myofibroblasts based on the expression of α -SMA. The combination of both drugs has a better potentiating effect in the ability to inhibit VIC differentiation into myofibroblasts. The combination of these two agents has great potential as an antiremodelling agent, especially in valvular disease due to RHD.

References

- Almawi, V., Beyhum, N., & Riedert, J. (1996). *of cytokine and cytokine receptor expression by glucocorticoids*. 60(November), 563–572. <https://doi.org/10.1002/jlb.60.5.563>
- Asai, K., Funaki, C., Hayashi, T., Yamada, K., Naito, M., Kuzuya, M., Yoshida, F., Yoshimine, N., & Kuzuya, F. (1993). Dexamethasone-induced suppression of aortic atherosclerosis in cholesterol-fed rabbits: Possible mechanisms. In *Arteriosclerosis and Thrombosis* (Vol. 13, Issue 6, pp. 892–899). <https://doi.org/10.1161/01.atv.13.6.892>
- Baum, J., & Duffy, H. S. (2011). Fibroblasts and myofibroblasts: What are we talking about? *Journal of Cardiovascular Pharmacology*, 57(4), 376–379. <https://doi.org/10.1097/FJC.0b013e3182116e39>
- Bouleti, C., Jung, B., Himbert, D., Brochet, E., Messika-Zeitoun, D., D etaint, D., Garbarz, E., Cormier, B., & Vahanian, A. (2013). Long-term efficacy of percutaneous mitral commissurotomy for restenosis after previous mitral commissurotomy. *Heart*, 99(18), 1336–1341. <https://doi.org/10.1136/heartjnl-2013-303944>
- Bujak. (2001). *doi_10.1016_j.cardiores.2006.10.002_Enhanced Reader.pdf*.
- Calderone, A., Murphy, R. J. L., Lavoie, J., Colombo, F., & B eliveau, L. (2001). TGF- β 1 and prepro-ANP mRNAs are differentially regulated in exercise-induced cardiac hypertrophy. In *Journal of Applied Physiology* (Vol. 91, Issue 2, pp. 771–776). <https://doi.org/10.1152/jappl.2001.91.2.771>
- Christian, M. M., Moy, R. L., Wagner, R. F., & Yen-Moore, A. (2001). A correlation of alpha-smooth muscle actin and invasion in micronodular basal cell carcinoma. *Dermatologic Surgery: Official Publication for American Society for Dermatologic Surgery [et Al.]*, 27(5), 441–445. <https://doi.org/10.1046/J.1524-4725.2001.00200.X>
- Cucoranu, I., Clempus, R., Dikalova, A., Phelan, P. J., Ariyan, S., Dikalov, S., & Sorescu, D. (2005). NAD(P)H oxidase 4 mediates transforming growth factor- β 1-induced differentiation of cardiac fibroblasts into myofibroblasts. *Circulation Research*, 97(9), 900–907. <https://doi.org/10.1161/01.RES.0000187457.24338.3D>
- Du, Y., Xiao, H., Wan, J., Wang, X., Li, T., Zheng, S., Feng, J., Ye, Q., Li, J., Li, G., & Fan, Z. (2020). Atorvastatin attenuates TGF- β 1-induced fibrogenesis by inhibiting Smad3 and MAPK signaling in human ventricular fibroblasts. *International Journal of Molecular Medicine*.

- <https://doi.org/10.3892/ijmm.2020.4607>
- Frangogiannis, N. G., Michael, L. H., & Entman, M. L. (2000). Myofibroblasts in reperfused myocardial infarcts express the embryonic form of smooth muscle myosin heavy chain (SMemb). *Cardiovascular Research*, 48(1), 89–100. [https://doi.org/10.1016/S0008-6363\(00\)00158-9](https://doi.org/10.1016/S0008-6363(00)00158-9)
- Gabbiani, G. (2003). The myofibroblast in wound healing and fibrocontractive diseases. In *Journal of Pathology*. <https://doi.org/10.1002/path.1427>
- Gu, Z., Rolfe, B. E., Xu, Z. P., Thomas, A. C., Campbell, J. H., & Lu, G. Q. M. (2010). Enhanced effects of low molecular weight heparin intercalated with layered double hydroxide nanoparticles on rat vascular smooth muscle cells. *Biomaterials*, 31(20), 5455–5462. <https://doi.org/10.1016/J.BIOMATERIALS.2010.03.050>
- Guilherme, L., & Kalil, J. (2020). Rheumatic Fever and Rheumatic Heart Disease. In *The Autoimmune Diseases* (Issue May, pp. 1255–1268). Elsevier. <https://doi.org/10.1016/B978-0-12-812102-3.00063-4>
- Guzy, R. (2020). Fibroblast Growth Factor Inhibitors in Lung Fibrosis: Friends or Foes? In *American Journal of Respiratory Cell and Molecular Biology*. <https://doi.org/10.1165/rcmb.2020-0156ED>
- Heldin, C. H., Miyazono, K., & Ten Dijke, P. (1997). TGF- β signalling from cell membrane to nucleus through SMAD proteins. In *Nature*. <https://doi.org/10.1038/37284>
- Isaka, Y. (2018). Targeting TGF- β signaling in kidney fibrosis. In *International Journal of Molecular Sciences*. <https://doi.org/10.3390/ijms19092532>
- Kim, S., Ohta, K., Hamaguchi, A., Yukimura, T., Miura, K., & Iwao, H. (1995). Angiotensin II induces cardiac phenotypic modulation and remodeling in vivo in rats. *Hypertension*, 25(6), 1252–1259. <https://doi.org/10.1161/01.HYP.25.6.1252/FORMAT/EPUB>
- Khidoyatova, M. R., Kayumov, U. K., Inoyatova, F. K., Fozilov, K. G., Khamidullaeva, G. A., & Eshpulatov, A. S. (2022). Clinical status of patients with coronary artery disease post COVID-19. *International Journal of Health & Medical Sciences*, 5(1), 137–144. <https://doi.org/10.21744/ijhms.v5n1.1858>
- Kupfahl, C., Pink, D., Friedrich, K., Zurbrugg, H. R., Neuss, M., Warnecke, C., Fielitz, J., Graf, K., Fleck, E., & Regitz-Zagrosek, V. (2000). Angiotensin II directly increases transforming growth factor beta1 and osteopontin and indirectly affects collagen mRNA expression in the human heart. *Cardiovascular Research*, 46(3), 463–475. [https://doi.org/10.1016/S0008-6363\(00\)00037-7](https://doi.org/10.1016/S0008-6363(00)00037-7)
- Lijnen, P. J., Petrov, V. V., & Fagard, R. H. (2000). Induction of cardiac fibrosis by transforming growth factor- β 1. *Molecular Genetics and Metabolism*. <https://doi.org/10.1006/mgme.2000.3032>
- Lin, C., Zhu, D., Markby, G., Corcoran, B. M., Farquharson, C., & Macrae, V. E. (2017). Isolation and characterization of primary rat valve interstitial cells: A new model to study aortic valve calcification. *Journal of Visualized Experiments*, 2017(129), 3–8. <https://doi.org/10.3791/56126>
- Lo, W. R., Rowlette, L. L., Caballero, M., Yang, P., Hernandez, M. R., & Borrás, T. (2003). Tissue differential microarray analysis of dexamethasone induction reveals potential mechanisms of steroid glaucoma. *Investigative Ophthalmology & Visual Science*, 44(2), 473–485. <https://doi.org/10.1167/IOVS.02-0444>
- Makheja, A. N., Bloom, S., Muesing, R., Simon, T., & Bailey, J. M. (1989). Anti-inflammatory drugs in experimental atherosclerosis. *Atherosclerosis*, 76(2–3),

- 155–161. [https://doi.org/10.1016/0021-9150\(89\)90099-3](https://doi.org/10.1016/0021-9150(89)90099-3)
- Masuda, A., Nakamura, T., Abe, M., Iwamoto, H., Sakaue, T., Tanaka, T., Suzuki, H., Koga, H., & Torimura, T. (2020). Promotion of liver regeneration and anti-fibrotic effects of the TGF- β receptor kinase inhibitor galunisertib in CCl₄-treated mice. *International Journal of Molecular Medicine*. <https://doi.org/10.3892/ijmm.2020.4594>
- Ordunez, P., Martinez, R., Soliz, P., Giraldo, G., Mujica, O. J., & Nordet, P. (2019). Rheumatic heart disease burden, trends, and inequalities in the Americas, 1990–2017: a population-based study. *The Lancet Global Health*, 7(10), e1388–e1397. [https://doi.org/10.1016/S2214-109X\(19\)30360-2](https://doi.org/10.1016/S2214-109X(19)30360-2)
- Parichatikanond, W., Luangmonkong, T., Mangmool, S., & Kurose, H. (2020). Therapeutic targets for the treatment of cardiac fibrosis and cancer: Focusing on tgf- β Signaling. In *Frontiers in Cardiovascular Medicine*. <https://doi.org/10.3389/fcvm.2020.00034>
- Peng, H., Carretero, O. A., Peterson, E. L., & Rhaleb, N. E. (2010). Ac-SDKP inhibits transforming growth factor- β 1-induced differentiation of human cardiac fibroblasts into myofibroblasts. *American Journal of Physiology - Heart and Circulatory Physiology*. <https://doi.org/10.1152/ajpheart.00464.2009>
- Petrov, V. V., Fagard, R. H., & Lijnen, P. J. (2002). Stimulation of collagen production by transforming growth factor- β 1 during differentiation of cardiac fibroblasts to myofibroblasts. *Hypertension*, 39(2 1), 258–263. <https://doi.org/10.1161/hy0202.103268>
- Suh, S. H., Choi, H. S., Kim, C. S., Kim, I. J., Ma, S. K., Scholey, J. W., Kim, S. W., & Bae, E. H. (2019). Olmesartan Attenuates Kidney Fibrosis in a Murine Model of Alport Syndrome by Suppressing Tubular Expression of TGF β . *International Journal of Molecular Sciences*, 20(15). <https://doi.org/10.3390/IJMS20153843>
- Suryasa, I. W., Rodríguez-Gámez, M., & Koldoris, T. (2022). Post-pandemic health and its sustainability: Educational situation. *International Journal of Health Sciences*, 6(1), i-v. <https://doi.org/10.53730/ijhs.v6n1.5949>
- Thourani, V. H., Weintraub, W. S., Guyton, R. A., Jones, E. L., Williams, W. H., Elkabbani, S., & Craver, J. M. (2003). Outcomes and long-term survival for patients undergoing mitral valve repair versus replacement: Effect of age and concomitant coronary artery bypass grafting. *Circulation*. <https://doi.org/10.1161/01.CIR.0000079169.15862.13>
- Tsai, W. C., Tang, F. T., Wong, M. K., & Pang, J. H. S. (2003). Inhibition of tendon cell migration by dexamethasone is correlated with reduced alpha-smooth muscle actin gene expression: a potential mechanism of delayed tendon healing. *Journal of Orthopaedic Research : Official Publication of the Orthopaedic Research Society*, 21(2), 265–271. [https://doi.org/10.1016/S0736-0266\(02\)00151-1](https://doi.org/10.1016/S0736-0266(02)00151-1)
- Vaideeswar, P., & Butany, J. (2016). Valvular Heart Disease. In *Cardiovascular Pathology: Fourth Edition* (4th ed.). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-420219-1.00012-4>
- Watkins, D. A., Johnson, C. O., Colquhoun, S. M., Karthikeyan, G., Beaton, A., Bukhman, G., Forouzanfar, M. H., Longenecker, C. T., Mayosi, B. M., Mensah, G. A., Nascimento, B. R., Ribeiro, A. L. P., Sable, C. A., Steer, A. C., Naghavi, M., Mokdad, A. H., Murray, C. J. L., Vos, T., Carapetis, J. R., & Roth, G. A. (2017). Global, Regional, and National Burden of Rheumatic Heart Disease, 1990–2015. *New England Journal of Medicine*.

- <https://doi.org/10.1056/nejmoa1603693>
- Yang, L., Bataller, R., Dulyx, J., Coffman, T. M., Ginès, P., Rippe, R. A., & Brenner, D. A. (2005). Attenuated hepatic inflammation and fibrosis in angiotensin type 1a receptor deficient mice. *Journal of Hepatology*, 43(2), 317–323. <https://doi.org/10.1016/j.jhep.2005.02.034>
- Yankah, C. A., Siniawski, H., Detschades, C., Stein, J., & Hetzer, R. (2011). Rheumatic mitral valve repair: 22-Year clinical results. *Journal of Heart Valve Disease*.
- Zhou, J., Jiang, K., Ding, X., Fu, M., Wang, S., Zhu, L., He, T., Wang, J., Sun, A., Hu, K., Chen, L., Zou, Y., & Ge, J. (2015). Qiliqiangxin inhibits angiotensin II-induced transdifferentiation of rat cardiac fibroblasts through suppressing interleukin-6. *Journal of Cellular and Molecular Medicine*. <https://doi.org/10.1111/jcmm.12512>

Dexamethasone and olmesartan as potential antiremodelling agents of valvular interstitial cell into myofibroblast: In vitro study on α - smooth muscle actin expression

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to iGroup Student Paper	1%
2	journals.lww.com Internet Source	1%
3	erepository.mkuit.ac.rw Internet Source	1%
4	etheses.whiterose.ac.uk Internet Source	1%
5	Jessica Hoppstädter, Alaina J. Ammit. "Role of Dual-Specificity Phosphatase 1 in Glucocorticoid-Driven Anti-inflammatory Responses", <i>Frontiers in Immunology</i> , 2019 Publication	1%
6	www.journalijar.com Internet Source	1%
7	"43rd Annual AEPC Meeting 18th Spring Meeting of ECHSA Palazzo del Casinò, Venice Lido (Italy), 21–24 May, 2008", <i>Cardiology in the Young</i> , 2008	<1%

- 8 Toshiyuki Maezawa, Masayuki Tanaka, Miho Kanazashi, Noriaki Maeshige, Hiroyo Kondo, Akihiko Ishihara, Hidemi Fujino. "Astaxanthin supplementation attenuates immobilization-induced skeletal muscle fibrosis via suppression of oxidative stress", The Journal of Physiological Sciences, 2016
Publication <1 %
-
- 9 Yuqi Huang, Wenbin Guo, Heping Kan. "TPX2 Is a Prognostic Marker and Contributes to Growth and Metastasis of Human Hepatocellular Carcinoma", International Journal of Molecular Sciences, 2014
Publication <1 %
-
- 10 Submitted to Aberystwyth University
Student Paper <1 %
-
- 11 www.ncbi.nlm.nih.gov
Internet Source <1 %
-
- 12 erl.ucc.edu.gh:8080
Internet Source <1 %
-
- 13 skeletaljournal.biomedcentral.com
Internet Source <1 %
-
- 14 Juan He, Jian Ding, Qihua Lai, Xinke Wang, Aimin Li, Side Liu. "Irbesartan Ameliorates Lipid Deposition by Enhancing Autophagy via PKC/AMPK/ULK1 Axis in Free Fatty Acid <1 %

Induced Hepatocytes", Frontiers in Physiology, 2019

Publication

15

Litao Yang, Yinlu Liu, Cuicui Bi, Bo Zhang. " Effects of Kütz polysaccharide on renal fibrosis in high - fat mice ", Food Science & Nutrition, 2022

Publication

<1 %

16

assets.researchsquare.com

Internet Source

<1 %

17

www.pjps.pk

Internet Source

<1 %

18

Brian Mendel, Christianto Christianto, Phoniex Angellia, Indra Holiyono, Radityo Prakoso, Sisca Natalia Siagian. "Reversed Potts Shunt Outcome in Suprasystemic Pulmonary Arterial Hypertension: A Systematic Review and Meta-Analysis", Current Cardiology Reviews, 2022

Publication

<1 %

19

iovs.arvojournals.org

Internet Source

<1 %

20

mdpi-res.com

Internet Source

<1 %

21

Cohn, L.H.. "The effect of pathophysiology on the surgical treatment of ischemic mitral regurgitation: operative and late risks of repair versus replacement", European Journal of Cardio-Thoracic Surgery, 1995

<1 %

22 Submitted to Universitas Airlangga <1 %
Student Paper

23 biosciencetrends.com <1 %
Internet Source

24 docshare.tips <1 %
Internet Source

25 ediss.uni-goettingen.de <1 %
Internet Source

26 link.springer.com <1 %
Internet Source

27 pubmed.ncbi.nlm.nih.gov <1 %
Internet Source

28 "46th Annual Meeting of the Association for <1 %
European Paediatric and Congenital
Cardiology, AEPC with joint sessions with
the Japanese Society of Pediatric Cardiology
and Cardiac Surgery Istanbul, Turkey, May
23–26, 2012", Cardiology in the Young, 2012
Publication

29 V V Petrov. "Transforming growth factor- <1 %
beta(1) induces angiotensin-converting
enzyme synthesis in rat cardiac fibroblasts
during their differentiation to
myofibroblasts.", Journal of Renin-
Angiotensin-Aldosterone System, 2000
Publication

30	Internet Source	<1 %
31	livrepository.liverpool.ac.uk Internet Source	<1 %
32	orca.cardiff.ac.uk Internet Source	<1 %
33	research.kums.ac.ir Internet Source	<1 %
34	researchexperts.utmb.edu Internet Source	<1 %
35	rjme.ro Internet Source	<1 %
36	s-space.snu.ac.kr Internet Source	<1 %
37	s3.eu-west-1.amazonaws.com Internet Source	<1 %
38	theses.ncl.ac.uk Internet Source	<1 %
39	Linshen Xie, Ye Zeng. "Therapeutic Potential of Exosomes in Pulmonary Fibrosis", Frontiers in Pharmacology, 2020 Publication	<1 %
40	ebin.pub Internet Source	<1 %
41	jaha.ahajournals.org Internet Source	<1 %

- | | | |
|----|--|------|
| 42 | pure.rug.nl
Internet Source | <1 % |
| 43 | www.science.gov
Internet Source | <1 % |
| 44 | Alexey Fedulov. "Serum TGF-Beta 1 and TNF-Alpha Levels and Cardiac Fibrosis in Experimental Chronic Renal Failure", Immunological Investigations, 5/1/2005
Publication | <1 % |
| 45 | Anbiao wu, Lihong Zhang, Jingyang Chen, Hekai Li, Pingzhen Yang, Minsheng Chen, Qicai Liu. "Limb-bud and Heart (LBH) mediates proliferation, fibroblast-to-myofibroblast transition and EMT-like processes in cardiac fibroblasts", Molecular and Cellular Biochemistry, 2021
Publication | <1 % |
| 46 | Terunao Iwanaga, Tetsuhiro Chiba, Masato Nakamura, Tatsuya Kaneko et al. "Miglustat, a glucosylceramide synthase inhibitor, mitigates liver fibrosis through TGF- β /Smad pathway suppression in hepatic stellate cells", Biochemical and Biophysical Research Communications, 2023
Publication | <1 % |
| 47 | Tian, Xiao, Xue, Zhang, Jia, Luo, Wang, Zhu, Cao. "The Expression of BNP, ET-1, and TGF- β 1 in Myocardium of Rats with Ventricular | <1 % |

Arrhythmias", International Journal of Molecular Sciences, 2019

Publication

48	arro.anglia.ac.uk Internet Source	<1 %
49	ecommons.luc.edu Internet Source	<1 %
50	elifesciences.org Internet Source	<1 %
51	wlv.openrepository.com Internet Source	<1 %
52	www.pubmedcentral.nih.gov Internet Source	<1 %
53	www.scielo.br Internet Source	<1 %
54	Handbook of Experimental Pharmacology, 2004. Publication	<1 %
55	Sadashiva S. Karnik, Hamiyet Unal, Jacqueline R. Kemp, Kalyan C. Tirupula et al. "International Union of Basic and Clinical Pharmacology. XCIX. Angiotensin Receptors: Interpreters of Pathophysiological Angiotensinergic Stimuli", Pharmacological Reviews, 2015 Publication	<1 %
56	Zhengri Lu, Chunshu Hao, Hao Qian, Yuanyuan Zhao, Xiangwei Bo, Yuyu Yao,	<1 %

Genshan Ma, Lijuan Chen. "Tripartite motif 38 attenuates cardiac fibrosis after myocardial infarction by suppressing TAK1 activation via TAB2/3 degradation", iScience, 2022

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Dexamethasone and olmesartan as potential antiremodelling agents of valvular interstitial cell into myofibroblast: In vitro study on α - smooth muscle actin expression

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14
