

# Can oral microbiome dysbiosis affect the behavior of children with Autism Spectrum Disorders (ASD)? : Narrative review

*by Udijanto Tedjosongko*

---

**Submission date:** 06-Jan-2023 03:41PM (UTC+0800)

**Submission ID:** 1989123573

**File name:** WJARR-2022-1461.pdf (731.27K)

**Word count:** 3709

**Character count:** 21019

## Can oral microbiome dysbiosis affect the behavior of children with Autism Spectrum Disorders (ASD)? : Narrative review

Udijanto Tedjosongko \*, Paramita Devi Oktaviani, Salma Nadia, Dimas Prasetyanto Wicaksono and Seno Pradopo

*Department of Pediatric Dentistry, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.*

World Journal of Advanced Research and Reviews, 2023, 17(01), 051–056

Publication history: Received on 17 November 2022; revised on 28 December 2022; accepted on 31 December 2022

Article DOI: <https://doi.org/10.30574/wjarr.2023.17.1.1461>

### Abstract

Background: Autism spectrum disorders (ASD) are complex developmental disorders characterised by several levels of social interaction and communication difficulties. The prevalence of children with ASD worldwide is around 1:160, and the number of cases of autism has increased significantly. The oral microbiome is a diverse microbial biofilm with the second most complex microbiota in the human body after the gut. Several factors, such as lifestyle and health, can influence the oral microbiome's composition and metabolism. Children with ASD generally have poorer oral hygiene and suffer from periodontal disease and dental caries more than typical children; this is associated with behavioural disturbances, such as communication limitations, motoric limitations, eating habits and the side effects of drugs. Continued oral microbiome dysbiosis can lead to gut microbiome dysbiosis. There are two hypotheses for the transmission of oral bacteria to the gut, via the hematogenous route or via the enteral route. Gut microbiome dysbiosis may regulate social behaviour, such as emotional dissonance and anxiety, through blood-brain-barrier and gut-brain-axis pathways. When there is inflammation of the periodontal tissue in ASD children, it can cause lipopolysaccharide (LPS) leakage to the blood-brain barrier (BBB) that can cause an inflammatory response and disrupt metabolic activity in the central nervous system. Oral dysbiosis, which continues into gut dysbiosis, can affect behaviour through the gut-brain axis. This review aims to determine the impact of oral microbiome dysbiosis on the behaviour of children with ASD.

**Keywords:** Autism spectrum disorder (ASD); Oral microbiome; Dysbiosis; Behaviour; Children;

### 1. Introduction

Autism spectrum disorders (ASD) are complex developmental disorders classified as very diverse in exceptional children and characterised by some difficulty in social interaction and communication. Other characteristics include atypical behaviours and habits such as difficulty switching from one activity to another, attention to detail and unusual reactions to sensations. The three main symptoms of ASD according to the classification of mental illness in ICD-10 and DSM-IV (1), are:

- Impaired social relationships,
- Impaired speech and language, and
- Repetitive actions and interests (2–4).

Most children with ASD have poor oral hygiene and many suffer from gingivitis and dental caries. An imbalance in the commensal microbiota ecology can cause dental caries. These changes could be related to irregular tooth-brushing habits because caregivers and parents have difficulty brushing their children's teeth. The pooled prevalence of periodontal disease in ASD children was 69.4% (5–7).

\* Corresponding author: Udijanto Tedjosongko

Children with ASD often experience problems in maintaining healthy teeth and mouths. Dental care is the most frequently unmet need to preserve the health of special needs children, especially children with ASD. This is a problem because dental and oral health is very important in the growth and development of children (8).

The oral microbiome can be described as a diverse group of microbial biofilms and has the second most complex microbiota in the human body after the gut. The microbiota can be considered more stable over time, for example, there may be only short-term disturbances in microbiota composition following external stress, such as the use of antibiotics. This stability is called resilience and can be divided into two explanations: resistance, namely the ability to deal with disturbances, and recovery, namely the capability to recover from the effects of disorders. Overall, resilience is the capacity of an ecosystem to deal with disruption without changing its symbiotic state (9).

Several factors can affect the oral microbiome's composition and metabolism, such as the host's lifestyle and health status. Each anatomic surface of the oral cavity has a specific microbiota organised as a biofilm. Even within the same tooth surface environment (either supragingival or subgingival), the composition of the bacterial community varies greatly. When the microbiota balance is broken, it will cause dysbiosis. When oral dysbiosis occurs, the microbiome will cause pathogens to develop and disease will occur (9).

Continued oral microbiome dysbiosis can lead to gut microbiome dysbiosis. There are two hypotheses for the transmission of oral bacteria to the gut. The first is via the hematogenous route, whereby oral bacteria enter the lesion and systemically circulate and colonise the gastrointestinal mucosa. The second is through the enteral route, in which oral bacteria move through the digestive tract to the intestine. The human body has several defence mechanisms against microbes, including neutralisation via gastric acid and resistance to colonisation by the enteral route. However, this defence mechanism may not work optimally. Some microbes, such as *P. gingivalis*, are known to be acid resistant, especially at higher inoculation doses. Regardless of the route, evidence suggests that more than half of the bacterial species in the gastrointestinal system undergo oral-gut translocation, even in the absence of pathology. Among the many oral bacteria found in the intestines of patients with gastrointestinal disease are members of the genera *Staphylococcus*, *Porphyromonas*, *Veillonella*, *Fusobacterium*, *Actinomyces* and *Parvimonas* (10).

Oral microbiome dysbiosis can influence complex behaviours such as emotional behaviour and anxiety in two ways. First, when there is inflammation of the periodontal tissue in ASD children, it can cause lipopolysaccharide (LPS) leakage to the blood-brain barrier (BBB) that can cause an inflammatory response and disrupt metabolic activity in the central nervous system. Second, oral dysbiosis, which continues into gut dysbiosis, can affect behaviour through the gut-brain axis. Therefore, oral health must be maintained because it can affect behaviour and emotions (11,12). This review aims to determine the effect of oral microbiome dysbiosis on the behaviour of children with ASD.

---

## 2. Literature review

### 2.1. Oral microbiome dysbiosis

Behaviours in children with ASD, such as communication limitations, motor limitations, eating habits, drug side effects and infrequent dental treatment, can cause oral microbiome dysbiosis. Alternative therapies that are carried out for children with ASD, such as a gluten/casein-free diet, can limit protein intake and indirectly cause an increase in carbohydrate content in saliva. Protein plays a role in increasing the buffer capacity of saliva. This can affect the condition of the oral microbiome (7).

There are significant differences in the bacterial content of ASD children and typical children. In children with ASD, significantly higher colonisation of *Proteobacteria*, *Actinobacteria* and *Bacteroidetes* was observed. At the genus level, increased colonisation by *Streptococcus* and *Haemophilus* and reduced colonisation by *Prevotella*, *Actinomyces*, *Porphyromonas* and *Fusobacterium* in tooth samples of children with ASD compared to typical children indicates dysbiosis of the oral microbiome of ASD children (5,9,13-15). Dysbiosis can be characterised by three different scenarios and can occur simultaneously, namely: a) loss of overall microbial diversity, b) loss of beneficial microbes, and c) increase in pathogenic microbes (16).

### 2.1.1. Loss of microbial diversity

The general ecological concept is the loss of biodiversity within a community, denoting a decrease in the number, genetic variability and variety of species of a biological community in a specified location. This loss of biodiversity can cause damage to the ecosystem. In the context of caries, several reports indicate a loss of diversity with increasing disease severity. This suggests that increased acidification of the oral microenvironment accompanied by loss of diversity and decreased level and metabolic activity of beneficial bacteria leads to the emergence of cariogenic bacteria (16-18).

### 2.1.2. Loss of beneficial microbes

One of the main features of dysbiosis is the loss of some of the benefits derived from a healthy oral microbiome. The oral microbiome is important for the maturation and development of an appropriate oral immune response; protects the host from oral pathogens and carcinogenic metabolites; and is part of the nitrate-nitrite-nitric oxide pathway, so there are several benefits. Loss of beneficial microbes can reduce the host's ability to fight pathogenic bacteria, responding to an exaggerated immune response against the host's tissues and exposing the host to carcinogenic metabolites and adverse vascular changes. This loss is especially important in periodontal disease, where excessive chronic inflammation leads to loss of supporting tissue around the teeth, including loss of alveolar bone, which can lead to tooth loss over time (16,19).

### 2.1.3. Increase in pathogenic microbes

In a healthy environment, the oral microbiome contains opportunistic pathogens at such low levels that they do not cause any problems to the host. However, increasing microbial pathogens can increase the risk of dental caries, periodontal disease and systemic disease (16,20).

## 2.2. Connection between oral microbiome dysbiosis and behaviour in children with ASD

The oral microbiome can influence the behaviour of children with ASD in two ways. First, changes in transcription results in microglia, leading to impaired microglia function, so the permeability of the BBB is higher (Figure 1). This causes the brain to be more easily exposed to bacterial metabolites. When inflammation of the periodontal tissue occurs in ASD children, it can cause leakage of LPS to the BBB. It can cause an inflammatory response and disrupt metabolic activity in the central nervous system. Prolonged disruption of energy metabolism in neurons, oligodendrocytes and glia can cause structural changes in the cortex, hippocampus, amygdala, or cerebellum, thus affecting behaviour disorders in ASD children. The increase in LPS in ASD children correlates with the increase in Interleukin 6 (IL-6) which acts as both a pro-inflammatory cytokine and an anti-inflammatory myokine. In addition, inflammation in the developing brain causes synapse malfunction. Synaptic malfunction resulting in vasopressin secretion can also affect social behaviour (21-23).




Figure 1 Direct and indirect mechanisms of bacteria infecting the brain (22).

In the direct mechanism, the oral cavity infects the olfactory tract and the olfactory nerves transfer the bacteria to the brain. In an indirect mechanism, bacteria in the mouth infect the blood and find their way through the blood, blood-brain barrier (BBB), perivascular space and circumventricular organs to the brain (22).

The second way is through the gut-brain axis, based on two-way physiological connections where information is exchanged between the host microbiome, the gut and the brain. This involves a link between the central nervous system and microbes in the gastrointestinal tract through direct neural activation. Prolonged oral microbiome dysbiosis can lead to microbiota dysbiosis in the gastrointestinal tract. Dysbiosis of the intestinal microbiota can affect several complex behaviours, such as emotional behaviour and anxiety (12,22,24,25). The microbiome is important for the immune system, brain and genome. A series of recent studies have demonstrated that the microbiome influences the central nervous system in multiple ways and influences the psychological state of individuals by mediating reactions to stress and anxiety (26-30).

The enteric nervous system (ENS), composed of two layers of approximately 100 million nerve cells that line the digestive tract from the oesophagus to the rectum, is an independent centre for neural processing and integration. The ENS can trigger nausea or malaise, build stress and evoke emotions. The microbiome, among other things, achieves both the recognition and synthesis of neuroendocrine hormones and produces neuroactive factors capable of communicating not only with the ENS but also with the central nervous system. The gut microbiome can influence synaptogenesis, regulation of neurotransmitters and neurotropic factors in the cerebral hemispheres. Administering probiotics (e.g. *Lactobacillus*) and faecal microbiome transplantation to treat conditions associated with depression and anxiety are no longer experimental but are good therapeutic avenues. The link between the microbiome and depression has been proven by research on bacterial abnormalities in depressed patients. It has also been shown that feeding multiple microbiome populations (*L. rhamnosus*, *B. infantis*, *B. longum*) can improve mood even in healthy subjects (26, 31-33).

In ASD children, compared to typical children, the adverse gut microbiome is more numerous than the beneficial microbiome. Behaviour, oral hygiene and the gut microbiome in ASD are interrelated factors. In general, gut microbiome dysbiosis often causes illnesses such as diarrhoea and abdominal pain, which can affect the psychology and behaviour of ASD children (11). The immune system is produced on the epithelial surface of the digestive system, which is inhabited by many bacteria, comprising both systemic and mucosal immune systems. Beneficial bacteria living in the intestinal wall epithelium play a major role in immunomodulation. When the healthy gut flora is disrupted, the number of cells that produce Immunoglobulin A (IgA) drops dramatically, reducing the immune system's ability and causing digestive disorders (14,27,34).

Digestive problems that children with ASD often experience include bloating, diarrhoea, constipation, difficulty eating and malnutrition in various degrees of severity. These digestive problems cause discomfort and pain. However, due to the inability of ASD children to communicate, most of them cannot tell their parents about it, so they express their feelings in other ways, such as self-harm, tantrums, refusing to eat, etc. (24,34,35).

13  
The mainstay of therapy for ASD is based on behavioural therapy, applied behavioural analysis, and speech and occupational therapy. Given that ASD is so complex and diverse, additional treatments exist to improve cognitive, linguistic and adaptive abilities. Probiotic supplementation may improve ASD symptoms (25). Probiotics can intervene in the gut microbiome, thereby increasing the good bacteria in the intestines of children with ASD. The most effective probiotics are probiotics which include four lactobacilli strains, three bifidobacteria strains and one *S. thermophiles* strain. Such probiotics significantly reduce gastrointestinal distress, especially constipation, and substantially improve certain aspects of behaviour, such as repetitive and rebellious behaviour (36-38).

From this literature review, it can be seen that the condition of the oral microbiome needs more attention from dental practitioners because oral microbiome dysbiosis can progress to gut microbiome dysbiosis, which can affect emotional and anxiety behaviour in children with ASD through the blood-brain-barrier and gut-brain-axis pathways. It does not rule out the possibility that direct access from the oral microbiome to the brain can influence behaviour, which requires further research.

---

### 3. Conclusion

Oral microbiome dysbiosis may affect the behaviour of children with ASD. Oral microbiome dysbiosis can progress to gut microbiome dysbiosis, which may affect emotional and anxiety behaviour in children with ASD through the blood-brain-barrier and gut-brain-axis pathways.

## Compliance with ethical standards

### Acknowledgments

The authors thank the reviewers for their insightful suggestions.

### Disclosure of conflict of interest

The authors declare that there is no conflict of interest regarding the publication of this document.

## References

- [1] Ali EM, Al- Adwan F eid ziad, Al-Naimat YM. Autism Spectrum Disorder (ASD); Symptoms, Causes, Diagnosis, Intervention, and Counseling Needs of the Families in Jordan. *Mod Appl Sci.* 2019;13(5):48.
- [2] Primasari IFND, Supena A. The Parents' Role of Children with Special Needs (Autism) in Elementary Schools During the Covid-19 {Peran Orang Tua Anak Berkebutuhan Khusus (Autis) Sekolah Dasar Selama Masa Covid-19}. *Jurnal Sekolah PGSD FIP UNIMED* 2020;5(1):133–42.
- [3] Firdaus F, Santy WH. Analysis of the Factors Causing Tantrum in Children with Autism {Analisis Faktor Penyebab Perilaku Tantrum Pada Anak Autis}. *Med Technol Public Heal J.* 2020;4(1):55–60.
- [4] WHO. Autism spectrum disorders [Internet]. 2021. Available from: <https://www.who.int/news-room/fact-sheets/detail/autism-spectrum-disorders>
- [5] Qiao Y, Wu M, Feng Y, Zhou Z, Chen L, Chen F. Alterations of oral microbiota distinguish children with autism spectrum disorders from healthy controls. *Sci Rep* [Internet]. 2018;8(1):1–12. Available from: <http://dx.doi.org/10.1038/s41598-018-19982-y>
- [6] Rachmawati D, Ermawati T. Oral Hygiene Status and Caries in Students with Special Needs at Autism SLB and TPA B SLB Branjangan, Jember Regency {Status Kebersihan Mulut dan Karies Pada Siswa Berkebutuhan Khusus di SLB Autis dan TPA B SLB Branjangan Kabupaten Jember}. *Warta Pengabdian.* 2019;13(3):74–9.
- [7] Ferrazzano GF, Salerno C, Bravaccio C, Ingenito A, Sangianantoni G, Cantile T. Autism spectrum disorders and oral health status: Review of the literature. *Eur J Paediatr Dent.* 2020;21(1):9–12.
- [8] Dwiputra S, Damayanti L, Sasmita IS. Dental fear in children with autism spectrum disorders {Dental fear anak dengan gangguan spektrum autisme terhadap perawatan gigi}. *J Kedokt Gigi Univ Padjadjaran.* 2021;33(3):213.
- [9] Barboza-Solís DDS, MSc, PhD C, Acuña-Amador PhD LA. The Oral Microbiota: A Literature Review for Updating Professionals in Dentistry. Part I. *Odovtos - Int J Dent Sci.* 2019;3(22):143–52.
- [10] Khor B, Snow M, Herrman E, Ray N, Mansukhani K, Patel KA, et al. Interconnections between the oral and gut microbiomes: Reversal of microbial dysbiosis and the balance between systemic health and disease. *Microorganisms.* 2021;9(3):1–22.
- [11] Lombardi M, Troisi J. Gut reactions: How far are we from understanding and manipulating the microbiota complexity and the interaction with its host? lessons from autism spectrum disorder studies. *Nutrients.* 2021;13(10):1–15.
- [12] Hemalakshmi R, Pavithra Amritkumar. Critical review on role of Microbiota transfer therapy in Autism spectrum disorder. *Int J Res Pharm Sci.* 2021;12(3):2095–114.
- [13] Forsyth A, Raslan K, Lyashenko C, Bona S, Snow M, Khor B, et al. Children with autism spectrum disorder: Pilot studies examining the salivary microbiome and implications for gut metabolism and social behavior. *Hum Microbiome J* [Internet]. 2020;15:100066. Available from: <https://doi.org/10.1016/j.humic.2019.100066>
- [14] Berding K, Donovan SM. Microbiome and nutrition in autism spectrum disorder: Current knowledge and research needs. *Nutr Rev.* 2016;74(12):723–36.
- [15] Strati F, Cavalieri D, Albanese D, De Felice C, Donati C, Hayek J, et al. New evidences on the altered gut microbiota in autism spectrum disorders. *Microbiome.* 2017;5(1):1–11.
- [16] Radaic A, Kapila YL. The oralome and its dysbiosis: New insights into oral microbiome-host interactions. *Comput Struct Biotechnol J* [Internet]. 2021;19:1335–60. Available from: <https://doi.org/10.1016/j.csbj.2021.02.010>

- [17] Lamont RJ, Koo H, Hajishengallis G. The oral microbiota: dynamic communities and host interactions. *Nat Rev Microbiol* [Internet]. 2018;16(12):745–59. Available from: <http://dx.doi.org/10.1038/s41579-018-0089-x>
- [18] Lu M, Xuan S, Wang Z. Oral microbiota: A new view of body health. *Food Sci Hum Wellness* [Internet]. 2019;8(1):8–15. Available from: <https://doi.org/10.1016/j.fshw.2018.12.001>
- [19] Gao L, Xu T, Huang G, Jiang S, Gu Y, Chen F. Oral microbiomes: more and more importance in oral cavity and whole body. *Protein Cell* [Internet]. 2018;9(5):488–500. Available from: <https://doi.org/10.1007/s13238-018-0548-1>
- [20] Degroot AK, Low D, Mizoguchi A, Mizoguchi E. Current understanding of dysbiosis in disease in human and animal models. *Inflamm Bowel Dis*. 2016;22(5):1137–50.
- [21] Johnson D, Letchumanan V, Thurairajasingam S, Lee LH. A revolutionizing approach to autism spectrum disorder using the microbiome. *Nutrients*. 2020;12(7):1–25.
- [22] Olsen I, Hicks SD. Oral microbiota and autism spectrum disorder (ASD). *J Oral Microbiol* [Internet]. 2020;12(1). Available from: <https://doi.org/10.1080/20002297.2019.1702806>
- [23] Fouquier J, Moreno Huizar N, Donnelly J, Glickman C, Kang D-W, Maldonado J, et al. The Gut Microbiome in Autism: Study-Site Effects and Longitudinal Analysis of Behavior Change. *mSystems*. 2021;6(2):1–17.
- [24] Hicks SD, Uhlig R, Afshari P, Williams J, Chronos M, Tierney-Aves C, et al. Oral microbiome activity in children with autism spectrum disorder. *Autism Res*. 2018;11(9):1286–99.
- [25] Sivamaruthi BS, Suganthy N, Kesika P, Chaiyasut C. The role of microbiome, dietary supplements, and probiotics in autism spectrum disorder. *Int J Environ Res Public Human and Health*. 2020;17(8).
- [26] Lavazza A, Sironi VA. Are we ready for a “microbiome-guided behaviour” approach? *Cambridge Q Healthc Ethics*. 2019;28(4):708–24.
- [27] Mitu De, Bhattacharya M, Saha AC, Basu I, Ganguli S, Dey SR. Influence of Oral Microbiome on Human Health: an Overview. *Int J Adv Life Sci Res*. 2019;2(1):16–21.
- [28] Rivera-Iñiguez I, Roman S, Ojeda-Granados C, Arturo P. Impact of the gut microbiome on behavior and emotions. *Microbiome Metabolome Diagnosis, Ther other Strateg Appl*. 2019;(January):379–90.
- [29] Ignatova V. Influence of Gut Microbiota on Behavior and Its Disturbances. *Behav Neurosci*. 2019;1–27.
- [30] Góralczyk-Bińkowska A, Szmajda-Krygier D, Kozłowska E. The Microbiota–Gut–Brain Axis in Psychiatric Disorders. *Int J Mol Sci*. 2022;23(19):1–23.
- [31] DiCarlo GE, Mabry SJ, Cao X, McMillan C, Woynarowski TG, Harrison FE, et al. Autism-Associated Variant in the SLC6A3 Gene Alters the Oral Microbiome and Metabolism in a Murine Model. *Front Psychiatry*. 2021;12(April):1–10.
- [32] Guest PC. Role of the Gut Microbiome in Autism Spectrum Disorders. *Adv Exp Med Biol*. 2019;1178(August):v–vi.
- [33] Sarkar A, Harty S, Johnson KVA, Moeller AH, Carmody RN, Lehto SM, et al. The role of the microbiome in the neurobiology of social behaviour. *Biol Rev*. 2020;95(5):1131–66.
- [34] McBride-Campbell N. Gut and Psychology Syndrome [Internet]. Vol. 23, *Journal of Orthomolecular Medicine*. 2008. Available from: <http://www.orthomolecular.org/library/jom/2008/pdf/2008-v23n02-p090.pdf>
- [35] James DM, Davidson EA, Yanes J, Moshiree B, Dallman JE. The Gut-Brain-Microbiome Axis and Its Link to Autism: Emerging Insights and the Potential of Zebrafish Models. *Front Cell Dev Biol*. 2021;9(April).
- [36] Davies C, Mishra D, Eshraghi RS, Mittal J, Sinha R, Bulut E, et al. Altering the gut microbiome to potentially modulate behavioral manifestations in autism spectrum disorders: A systematic review. *Neurosci Biobehav Rev* [Internet]. 2021;128(July):549–57. Available from: <https://doi.org/10.1016/j.neubiorev.2021.07.001>
- [37] Banerjee A, Dey SR, Basu I, De M. Autism Spectrum Disorder and Gut Microbiome: A Brief Review. *Int J Adv Life Sci Res*. 2021;4(1).
- [38] Singla K, Baruah A, Chapadgaonkar SS. Probiotics and Management of Autism Spectrum Disorders. *J Microbiol Biotechnol Food Sci*. 2021;10(5):5.

# Can oral microbiome dysbiosis affect the behavior of children with Autism Spectrum Disorders (ASD)? : Narrative review

## ORIGINALITY REPORT

**20%**  
SIMILARITY INDEX

**13%**  
INTERNET SOURCES

**13%**  
PUBLICATIONS

**0%**  
STUDENT PAPERS

## PRIMARY SOURCES

- 1** [www.scielo.sa.cr](http://www.scielo.sa.cr)  
Internet Source **3%**
- 2** [mdpi-res.com](http://mdpi-res.com)  
Internet Source **2%**
- 3** Han Qin, Guangyue Li, Xiaohui Xu, Chuangwei Zhang, Wenjie Zhong, Shihan Xu, Yuanyuan Yin, Jinlin Song. "The role of oral microbiome in periodontitis under diabetes mellitus", *Journal of Oral Microbiology*, 2022  
Publication **2%**
- 4** Eleonora Lacorte, Giuseppe Gervasi, Ilaria Bacigalupo, Nicola Vanacore, Umberto Raucci, Pasquale Parisi. "A Systematic Review of the Microbiome in Children With Neurodevelopmental Disorders", *Frontiers in Neurology*, 2019  
Publication **1%**
- 5** Christopher Busayo Olowosoke, Otitoola Shobi Gbemisola, Adebola Abosede Alaba, Oluwadamilola Hope Adepoju et al. "Insilico **1%**



Validation of Selected Natural Products as Multi-regulator of EZH2-PPAR Therapeutic Targets; A Hallmark for Prospective Restoration of Pancreatic Insulin Production and Cancer dysregulation", Research Square Platform LLC, 2022

Publication

6

[www.pulsus.com](http://www.pulsus.com)

Internet Source

1 %

7

[docplayer.net](http://docplayer.net)

Internet Source

1 %

8

Greta Fowlie, Nicholas Cohen, Xue Ming. "The Perturbance of Microbiome and Gut-Brain Axis in Autism Spectrum Disorders", International Journal of Molecular Sciences, 2018

Publication

1 %

9

Hedieh Piraneh, Mahdia Gholami, Katayoun Sargeran, Ahmad Reza Shamshiri. "Oral health and dental caries experience among students aged 7–15years old with autism spectrum disorders in Tehran, Iran", BMC Pediatrics, 2022

Publication

1 %

10

Muneerah I. AL tamimi, Hanan A. Alfawaz, Ramesa Shafi Bhat, Shaista Arzoo et al.

"Comparative Study on the Exacerbating

Effects of Casein-Rich vs. Gluten-Rich Diets on

1 %

# Biochemical-Induced Features in Rodent Model of Autism", Journal of Molecular Neuroscience, 2022

Publication

11

[ijcrims.com](http://ijcrims.com)

Internet Source

1 %

12

[www.researchgate.net](http://www.researchgate.net)

Internet Source

1 %

13

Camron Davies, Dibyanshi Mishra, Rebecca S. Eshraghi, Jeenu Mittal et al. "Altering the gut microbiome to potentially modulate behavioral manifestations in autism spectrum disorders: A systematic review", Neuroscience & Biobehavioral Reviews, 2021

Publication

<1 %

14

[doaj.org](http://doaj.org)

Internet Source

<1 %

15

[www.ehu.eus](http://www.ehu.eus)

Internet Source

<1 %

16

[www.gacetasanitaria.org](http://www.gacetasanitaria.org)

Internet Source

<1 %

17

[revistas.ucr.ac.cr](http://revistas.ucr.ac.cr)

Internet Source

<1 %

18

K. K. Adithya, Riya Rajeev, Joseph Selvin, George Seghal Kiran. "Dietary Influence on the Dynamics of the Human Gut Microbiome:

<1 %

Prospective Implications in Interventional Therapies", ACS Food Science & Technology, 2021

Publication

19

Chang Lu, Jiaqi Rong, Changxing Fu, Wenshi Wang, Jing Xu, Xing-Da Ju. "Overall Rebalancing of Gut Microbiota Is Key to Autism Intervention", Frontiers in Psychology, 2022

Publication

<1 %

20

Margaret E. Coleman, Rodney R. Dietert, D. Warner North, Michele M. Stephenson. "Enhancing Human Superorganism Ecosystem Resilience by Holistically 'Managing Our Microbes'", Applied Microbiology, 2021

Publication

<1 %

21

[hal.archives-ouvertes.fr](http://hal.archives-ouvertes.fr)

Internet Source

<1 %

22

[srinivasuniverstrg.blob.core.windows.net](http://srinivasuniverstrg.blob.core.windows.net)

Internet Source

<1 %

23

Jianhao Liang, Ting Li, Jiajia Zhao, Cheng Wang, Haitao Sun. "Current understanding of the human microbiome in glioma", Frontiers in Oncology, 2022

Publication

<1 %

24

[etd.cput.ac.za](http://etd.cput.ac.za)

Internet Source

<1 %

25 [link.springer.com](https://link.springer.com) Internet Source <1 %

---

26 [wjgnet.com](https://wjgnet.com) Internet Source <1 %

---

27 Monia Kittana, Asma Ahmadani, Farah Al Marzooq, Amita Attlee. "Dietary Fat Effect on the Gut Microbiome, and Its Role in the Modulation of Gastrointestinal Disorders in Children with Autism Spectrum Disorder", *Nutrients*, 2021  
Publication <1 %

---

28 Katarína Janšáková, Klaudia Kyselicová, Daniela Ostatníková, Gabriela Repiská. "Potential of Salivary Biomarkers in Autism Research: A Systematic Review", *International Journal of Molecular Sciences*, 2021  
Publication <1 %

---

29 L. S. Kind, I. H. A. Aartman, M. C. M. van Gemert-Schriks, C. C. Bonifacio. "Parents' satisfaction on dental care of Dutch children with Autism Spectrum Disorder", *European Archives of Paediatric Dentistry*, 2020  
Publication <1 %

---

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# Can oral microbiome dysbiosis affect the behavior of children with Autism Spectrum Disorders (ASD)? : Narrative review

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---