



HASIL VALIDASI DAN PENILAIAN
KARYA ILMIAH DOSEN FAKULTAS KEDOKTERAN GIGI UNIVERSITAS AIRLANGGA
(MEDIA PUBLIKASI KARYA ILMIAH : JURNAL ILMIAH INTERNASIONAL)

Nomor : 54 /UN3.1.2.5.9 / PS /2023

A. Identitas Karya Ilmiah

Judul Jurnal Ilmiah (Artikel) : Oral Microbiome Profiles and Inflammation in Pregnant Women Who Used Orthodontic Appliances
Jumlah penulis : 7 (tujuh) orang
Status Pengusul : Penulis Ke-4
Identitas Jurnal Ilmiah : a. Nama Jurnal : Dentistry Journal
b. Nomor ISSN: 2304-6767
c. Volume, Nomor, bulan, tahun : Volume 10, bulan Juli, tahun 2022
d. Penerbit : Multidisciplinary Digital Publishing Institute (MDPI)
e. DOI artikel :
f. Alamat web Jurnal : <https://doi.org/10.3390/dj10070118>
g. Terindek di Scimagojr/Thomson Reuter ISI Knowledge atau di :
Scimago Q2 SJR 0,52 (2021)

B. Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional Bereputasi (SJR > 0,10)
(beri lipada kategori yang tepat) : Jurnal Ilmiah Internasional Bereputasi (SJR < 0,10)
 Jurnal Ilmiah International terindeks di Web of science clarivate analytics /
kelompok emerging sources citation indeks (tidak terindeks SJR)

C. Hasil Validasi Ketua Departemen

Telah diperiksa dan divalidasi dengan baik, dan sampai pernyataan ini dibuat sebagai karya ilmiah **original / plagiat**, sehingga kami turut bertanggung jawab bahwa karya ilmiah tersebut telah memenuhi syarat kaidah ilmiah, norma akademik, dan norma hukum, sesuai dengan Peraturan Menteri Pendidikan Nasional Nomor 17 Tahun 2010 tanggal 16 Agustus 2010 tentang Pencegahan dan Pananggulangan Plagiat di Perguruan Tinggi.

Namun demikian, apabila di kemudian hari ternyata terbukti bahwa karya ilmiah tersebut merupakan karya Ilmiah Plagiat, maka akan menjadi tanggung jawab mutlak penulis tersebut di atas, baik secara perdata maupun pidana.

Demikian surat pernyataan ini saya buat untuk dipergunakan sebagaimana mestinya

Surabaya, 1 Maret 2023

Ketua Departemen,

Prof. Dr. Soegeng Wahlujo, drg, MKes, SpKGA.,
Supsp KKA(K)

NIP : 1954111019810301003

* Coret salah satu

Form Penilaian Kualitas Karil dan Kesesuaian Bidang Ilmu	
Profil Sinta : Link Sinta https://sinta.kemdikbud.go.id/profile	
A	Identitas Karya Ilmiah
1	Judul : Oral Microbiome Profiles and Inflammation in Pregnant Women Who Used Orthodontic Appliances
2	Nama Penulis : 1. Fajar Kusuma Dwi Kurniawan , 2. Retno Indrawati Roestamadji, 3. Nobuhiro Takahashi, 4. Udijanto Tedjosongko, 5. Ida Bagus Narmada, 6. Meircurius Dwi Condro Surboyo, 7. Indeswati Diyatri
3	Nama Jurnal : Dentistry Journal
B	Peng-index : Scimago Q2 SJR 0,52 (2021)
C	<p>Relevansi kompetensi dosen dengan substansi karya ilmiah</p> <ol style="list-style-type: none"> 1. Artikel ini membahas mengenai Profil Microbiome Oral dan Peradangan pada Ibu Hamil Wanita yang Menggunakan Peralatan Ortodontik dan tidak menggunakan ortodontik dengan hasil normal tidak berpengaruh asal menjaga mulut rongga bersih dan peralatan mereka terpelihara dengan baik untuk menghindari masalah mulut. 2. Artikel ini sesuai dengan bidang keahlian pengusul yaitu Cariology. 3. Tidak ada keterkaitan dengan naskah Disertasi pengusul yang berjudul: Initial acquisition and transmission of Mutans Streptococci in children at day nursery.
D	<p>Kesesuaian antara lingkup / subjek area jurnal dengan karya ilmiah yang diusulkan</p> <ol style="list-style-type: none"> 1. Alamat Web Jurnal : https://doi.org/10.3390/dj10070118 https://www.mdpi.com/2304-6767/10/7/118/pdf 2. Kebenaran ISSN/ISBN : 2304-6767 3. Termasuk "Predatory" tidak (jurnal; penerbit) : tidak masuk pada predatory 4. Syarat komposisi Editor Board : 24 Negara 5. Syarat kontributor penulis artikel : Penulis keempat 6. Keberkalaan penerbitan : 12 terbitan pertahun 7. Subjek area dan katagori jurnal : Oral Microbiology
E	<p>Kepastian tidak ada pelanggaran integritas akademik</p> <ol style="list-style-type: none"> 1. Indikasi plagiasi (lihat check similarity) : 19% 2. Fabrikasi : tidak ada 3. Falsifikasi : tidak ada 4. Praktek kepalsuan : tidak ada
	Nilai pengusul (penulis pertama dan corespondensi 60%) =
	Nilai pengusul (penulis pertama / penulis corespondensi masing - masing 40%)
	Nilai lainnya sesuai PO PAK 2019 dan suplemennya

Surabaya,
Penilai Angka Kredit 1

Nama :
NIP :
Bidang Ilmu :



Article

Oral Microbiome Profiles and Inflammation in Pregnant Women Who Used Orthodontic Appliances

Fajar Kusuma Dwi Kurniawan ¹, Retno Indrawati Roestamadji ^{2,*}, Nobuhiro Takahashi ³,
Udijanto Tedjosongko ⁴, Ida Bagus Narmada ⁵, Meircurius Dwi Condro Surboyo ⁶ and Indeswati Diyatri ²

- ¹ Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia; kurifana@gmail.com
² Department of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia; indeswati-d@fkg.unair.ac.id
³ Graduate School of Dentistry, Tohoku University, Sendai 980-8575, Japan; nobuhiro.takahashi.a5@tohoku.ac.jp
⁴ Department of Pediatric Dentistry, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia; udijanto@fkg.unair.ac.id
⁵ Department of Orthodontics, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia; ida-b-n@fkg.unair.ac.id
⁶ Department of Oral Medicine, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia; meircurius@fkg.unair.ac.id
* Correspondence: retnoindrawati@fkg.unair.ac.id

Abstract: It is common for women to undergo orthodontic treatment during pregnancy, especially through the use of fixed orthodontic devices. In changing the oral microbiome profile, it is crucial to increase the immune responses of pregnant women using fixed orthodontics; however, changes in the microbiomes of pregnant women with orthodontic appliances can be adjusted. Therefore, we aimed to conduct research on the oral cavity microbiome profiles, specifically IL-6 and TNF- α , of pregnant women using fixed orthodontic appliances. We proposed an observational analysis of 30 third-trimester pregnant women. OHI-S was recorded, saliva collection was performed using the passive drool method for IL-6 and TNF- α , and analysis and mucosal swabs were used to determine the oral microbiome profile. Kruskal–Wallis and post hoc Bonferroni tests were used to identify any significant differences with values of $p < 0.05$. Of these pregnant women, those with orthodontic appliances developed 10 types of bacteria at similar levels (>80%) from the genera *Streptococcus*, *Lactobacillus*, and *Veillonella*. There was no difference between the oral microbiomes of the control group and the pregnant women with a history of orthodontic appliance use. While the level of TNF- α in the women with orthodontic appliances was higher compared with the control group who had never used orthodontic appliances ($p < 0.05$), there was no difference in the IL-6 levels. The IL-6 and microbiome profile produced normal results, so the use of orthodontic appliances during pregnancy should be allowed with conditions. Pregnant women with orthodontic appliances must keep the oral cavity clean and their appliances well-maintained to avoid oral problems.

Keywords: fixed orthodontic; microbiome; orthodontics; pregnancy; IL-6; inflammation; TNF- α



Citation: Kurniawan, F.K.D.; Roestamadji, R.I.; Takahashi, N.; Tedjosongko, U.; Narmada, I.B.; Surboyo, M.D.C.; Diyatri, I. Oral Microbiome Profiles and Inflammation in Pregnant Women Who Used Orthodontic Appliances. *Dent. J.* **2022**, *10*, 118. <https://doi.org/10.3390/dj10070118>


Academic Editor: Andrea Scribante

Received: 6 May 2022

Accepted: 27 June 2022

Published: 1 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Research on the diversity of human microbiomes was initiated by Antonie van Leeuwenhoek in the early 1680s when he compared the microbiota of the oral cavity and faeces [1]. It is estimated that 96% of the bacteria found in the microbiomes of the oral cavity and intestines are similar, including the phyla *Firmicutes*, *Bacteroidetes*, *Proteobacteria*, *Actinobacteria*, *Spirochaetes*, and *Fusobacteria* [2].

When the body undergoes physiological and even pathological changes, the microbiome of the body also changes [3]. During pregnancy, women experience changes in their intestinal microbiome profile [4]. These changes can be affected by nutrition and may alter foetal health [5]. However, this does not only occur in the intestines; microbiome changes

also occur in the oral cavity. In the first and third trimesters of pregnancy, it was found that the numbers of *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans* increased significantly compared to non-pregnant women [6,7]. In addition to these two bacteria, *Candida albicans* also increased during the second and third trimesters of pregnancy. *Streptococcus mutans* showed a similar increase but was not quite as dramatic in the first trimester and remained constant until the third trimester [8].

Oral hygiene is necessary to prevent caries and periodontal disease during and after a fixed orthodontic treatment. Instruction in oral hygiene is essential to all cases of orthodontic treatment and must be reinforced [9]; pregnant women are no exception. During three months of fixed orthodontic bonding, there was a significant increase in the number of *Porphyromonas gingivalis*, *Tannerella forsythia*, *Porphyromonas intermedia*, *Porphyromonas nigrescens*, and *Fusobacterium spp.*, which were previously found in the periodontal ligament [10]. Other bacteria such as *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, and *Serratia marcescens* emerged, even in people without orthodontic devices [11].

IL-6 is a multifunctional cytokine that plays an important role in acute and chronic inflammation and autoimmunity. In relation to orthodontic treatment, IL-6 plays an important role in the movement of teeth, namely the inflammatory process that supports the resorptions process in apical areas [12,13]. Increased IL-6 levels occur not only in periodontal or alveolar tissues but also in saliva [14]. Similar to IL-6, TNF- α is also affected by orthodontic treatment and plays a role in the processes of tooth movement and bone resorption [15]. Orthodontic treatment increases TNF- α levels in periodontal tissue, especially in gingival crevicular fluid [16]. With the inclusion of various inflammatory markers such as cytokines, namely TNF- α and IL-6, it also affects the hygiene and oral composition of the microbiome [17].

This current study aimed to identify the composition of bacteria in the microbiome, derived from mucosal swabs, and the levels of TNF- α and IL-6 found in saliva. Orthodontic treatments trigger changes in the microbiome profile that may cause abnormalities in both the soft and hard tissues of the oral cavity. During orthodontic treatment and pregnancy, there is a significant increase in oral microbiome profile changes, which can lead to various diseases of the oral cavity. Pregnancy naturally results in a wide range of tissue abnormalities. This includes the dental tissues and could affect oral health.

The hypotheses were as follows:

1. That the use of orthodontic appliances in pregnant women changes the microbiome profile.
2. That the levels of TNF- α and IL-6 found in saliva are higher in pregnant women that have used orthodontic appliances.

2. Materials and Methods

2.1. Study Participant

This research was an observational analytical study. The participants in this study were all pregnant women in their third trimester who visited Kendangsari Mothers and Child Hospital in Surabaya. The inclusion criteria for this research were pregnant women in their third trimester with no caries or periodontal disease. The exclusion criteria were pregnant women who take medicine that may affect oral cavity homeostasis or the presence of systemic disease.

A sample of 32 participants was selected and divided into 3 groups. The control group was pregnant women who had never used orthodontic appliances, the second group was pregnant women who had previously used orthodontic appliances, and the third group was pregnant women with current orthodontic appliances.

2.2. Oral Health Assessment

Subjects were instructed to sit on normal chairs and clinical examinations were performed. The oral hygiene index (OHI-S) is the sum of the debris index (DI) and calculus index (CI) [18]. DI and CI examinations were carried out using a periodontal probe (Probe

UNC15, Osung, Osung MND Co LTD, Gyeonggi-do, Korea) placed on the tooth surface horizontally. To obtain the DI score, the sum of the debris assessment was divided by the number of teeth examined. The sum of the calculus scores was then divided by the number of teeth examined to obtain a CI score. These scores were divided into good OHI-S (0–1.2) and fair OHI-S (>1.3).

2.3. Saliva Collection

Saliva collection was performed using the passive drool method, which involved lowering of the head and releasing saliva collected at the base of the mouth into a 15 mL Eppendorf tube. Subjects were instructed not to eat or drink for about one hour before the salivary sample collection. They were asked to gargle with water for approximately one minute, and then wait five minutes [19]. Salivary samples were collected up to 5 mL [19], and placed in a refrigerator at a temperature of $-30\text{ }^{\circ}\text{C}$.

2.4. Mucosal Swab

Subjects were instructed not to eat or drink for one hour before the procedure. The entire surface of the oral cavity was rubbed using a sterile cotton swab, and the swab then was placed in a phosphate buffer saline solution. The mucosal swab samples were kept at $-80\text{ }^{\circ}\text{C}$ until testing was carried out.

2.5. DNA Extraction

DNA extraction of samples was carried out in accordance with the factory protocol of the kit used (QIAamp DNA Stool Mini Kit—Qiagen, Hilden, Germany). The sample was taken from the Eppendorf tube and placed in a microcentrifuge tube, and then 180 μL Buffer ATL and 20 μL proteinase K were added to the same tube. The tube was vortexed for 15 s and then incubated at $600\text{ }^{\circ}\text{C}$ for 24 h. Following this, 200 μL Buffer AL was added and the tube was vortexed for 15 s, and then incubated for 10 min at $70\text{ }^{\circ}\text{C}$. Finally, 200 μL 96% ethanol was added, vortexed for 15 s, and then placed in spin down. The mixed results were placed into the QIAamp Mini spin column and centrifuged at a speed of 8000 rpm for 1 min.

The concentration and quality of DNA was determined using a Nano-Drop 1000 spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA). At the end of the extraction, 50 μL of DNA sample was obtained. This DNA was used for the qPCR running process.

2.6. qPCR for Microbiome Analysis

PCR Master mix (Intron; iNtRON Biotechnology, Gyeonggi-do, Korea), Universal Forward Primer Macrogen (5'-GAG AGT TTG ATY CTG GCT CAG-3'), Universal Reverse Primer Macrogen (5'-GAA GGA GGT GWT CCA RCC GCA-3'), and DNA Marker 1 kbp (Thermo Fisher Scientific, Wilmington, DE, USA).

2.7. TNF- α and IL-6 Levels

The TNF- α and IL-6 levels were analysed using an ELISA kit. TNF- α is human tumour necrosis factor α (Bioassay Technology Laboratory, Shanghai, China) and IL-6 is human interleukin 6 (Bioassay Technology Laboratory, Shanghai, China).

2.8. Statistical Analysis

The levels of TNF- α and IL-6 were analysed with the Kolmogorov–Smirnov test to assess the data distribution and the Levene test for data homogeneity. Next, the Kruskal–Wallis and post hoc Bonferroni tests were used to discover any significant differences with values of $p < 0.05$. SPSS version 24 (IBM SPSS Statistic 24 for mac, New York, NY, USA) was used for the analysis.

3. Results

3.1. Subject Characteristics

The pregnant women who had never used orthodontic appliances (control) had an average age of 26.2 ± 2.4 years old, the women who had previously used orthodontic appliances were, on average, 29.4 ± 5.1 years old, and those with current orthodontic appliances were, on average, 30.6 ± 3.9 years old (Figure 1A).




Figure 1. Demographic information. Ages (A); week of pregnancy (B); and OHI-S (C).

The average stage of pregnancy for women who had never used orthodontic appliances (control) was 32.4 ± 3.2 weeks; for those who had previously used orthodontic appliances, it was 34.2 ± 4.6 weeks; and for those with current orthodontic appliances, it was 33.0 ± 3.5 weeks (Figure 1B).

3.2. OHI-S Status

For OHI-S, the pregnant women who had never used orthodontic appliances (control) had an average score of 0.5, the women who had previously used orthodontic appliances

had an average score of 0.4, and those with current orthodontic appliances had an average score of 1.6 (Figure 1C).

3.3. qPCR for Microbiome Analysis

The qPCR results showed that the microbiome in pregnant women who had never used orthodontic appliances (control) consisted of seven types of bacteria with similar levels (>80%) from the genera *Streptococcus*, *Gemella*, *Lactobacillus*, and *Abiotrophia* (Figure 2A). The oral bacteria found in the control included *Streptococcus mitis* strain 1042, *Streptococcus mutants* UA159, *Streptococcus* sp. strain C17, *Streptococcus* sp. strain D19, *Gemella* strain IR3.5, *Lactobacillus fermentum*, and *Abiotrophia defective*.




Figure 2. qPCR results of the similarities in the bacteria in the oral microbiomes of pregnant women who had never used orthodontic appliances (control) (A); pregnant women who had previously used orthodontic appliances (B); and pregnant women with current orthodontic appliances (C).

The microbiome in pregnant women who had previously used orthodontic appliances consisted of seven types of bacteria with similar levels (>80%) from the genera *Streptococcus*, *Gemella*, *Lactobacillus*, and *Abiotrophia*; no differences were observed with the microbiome of pregnant women who had never used orthodontic appliances (Figure 2B). The oral bacteria consisted of *Streptococcus mitis* strain 1042, *Streptococcus* sp. strain C17, *Streptococcus* sp. strain D19, *Streptococcus mutants* UA159, *Gemella* strain IR3.5, *Lactobacillus fermentum*, and *Abiotrophia defective*.

In the pregnant women with current orthodontic appliances, there were 10 types of bacteria with similar levels (>80%) from the genera *Streptococcus*, *Lactobacillus*, and *Veillonella* (Figure 2C). The oral bacteria were *Streptococcus tigurinus*, *Streptococcus mitis* strain 1042, *Streptococcus oralis* strain PL430, *Streptococcus* sp. 11aTha1, *Streptococcus salivarius*, *Streptococcus* sp. strain D19, *Streptococcus* sp. strain C17, *Streptococcus mutants* UA159, *Lactobacillus fermentum*, and *Vellonelia dispar*.

3.4. TNF- α and IL-6 Value

The IL-6 levels in all three sample groups are shown in Table 1 and no significant differences were observed (Figure 3). The level of TNF- α in the pregnant women with current orthodontic appliances was higher compared with pregnant women who had never used orthodontic appliances (control) ($p < 0.05$) while the level of TNF- α in pregnant women with a history of orthodontic appliances showed no significant difference from those with current orthodontic appliances (Figure 3).

Table 1. Values of salivary TNF- α and IL-6 in each group.

Groups	IL-6 Value (ng/L)			TNF- α Value (ng/L)		
	Mean \pm SD	Minimum	Maximum	Mean \pm SD	Minimum	Maximum
pregnant women who had never used orthodontic appliances (control)	22.38 \pm 7.94	13.53	40.74	182.57 \pm 69.24	84.55	294.22
pregnant women who had previously used orthodontic appliances	27.23 \pm 18.22	9.78	75.69	171.41 \pm 90.50	77.76	333.18
pregnant women with current orthodontic appliances	57.13 \pm 66.71	17.13	260.46	294.78 \pm 213.35	81.60	909.38

SD: standard deviation.




Figure 3. Values of salivary TNF- α and IL-6. ^aSignificant differences were shown using Kruskal-Wallis and post hoc Bonferroni tests with a significance value of $p < 0.05$.

4. Discussion

The analysed data showed similarity in the oral microbiomes. The control group and pregnant women who had previously used orthodontic appliances had similar oral microbiomes consisting of *Streptococcus mitis* strain 1042, *Streptococcus mutants* UA159, *Streptococcus* sp. strain C17, *Streptococcus* sp. strain D19, *Gemella* strain IR3.5, *Lactobacillus fermentum*, and *Abiotrophia defective*. The pregnant women with current orthodontic appliances had oral microbiomes with five similar bacteria but did not include the *Gemella* strain IR3.5. Other bacteria also found included *Streptococcus tigurinus*, *Streptococcus oralis* strain PL430, *Streptococcus* sp. 11aTha1, *Streptococcus salivarius*, and *Vellonelia dispar*. The results of this study show that the microbiomes of all the groups were dominated by *Streptococcus* strain. These bacteria are the dominant species of the oral microbiome [20].

In pregnant women who had previously and currently had fixed orthodontic appliances, it was found that the most dominant bacteria were *Streptococcus mitis* strain 1042, *Streptococcus mutants* UA159, *Streptococcus* sp. strain C17, *Streptococcus* sp. strain D19, and *Lactobacillus fermentum*. *Streptococcus mitis* is a type of viridians streptococcus and commensal normal flora of the oral cavity. *Streptococcus mitis* is a bacterium that initiates colonisation in the human oral cavity and oropharynx. It can move from its normal habitat and cause many complications such as endocarditis, bacteraemia, and septicaemia. This bacterium is an opportunistic pathogen [21]. The presence of *Streptococcus* sp., especially *Streptococcus mutants*, was reported to increase with the placement of an orthodontic appliance. The appliance becomes an iatrogenic factor that increases the risk of plaque and biofilm formation on the teeth and the appliance itself [22]. This accumulation can lead to disease of the teeth, gingiva, and periodontium. The current study confirmed that the oral microbiome is dominated by *Streptococcus* strain in pregnant women without orthodontic appliances, with a history of orthodontic appliances, and with current orthodontic appliances. Knowledge of the microbiome is very important as it has been demonstrated that the use of probiotics [23] and natural compounds [24] can modify clinical and microbiological parameters in periodontitis patients. The use of these products could also have an effect in pregnancy, and these variables should be considered in future clinical trials [25].

The dominance of the *Streptococcus* strain was observed among both pregnant women and non-pregnant women. Not only did the species change but the number of microorganisms also increased in pregnant women [26]. In non-pregnant women, the *Streptococcus* strain was dominated by *Streptococcus agalactiae* [27]. For pregnant women, the *Streptococcus mutants* increased compared with non-pregnant women [28]. In the present study, *Streptococcus mutants* was also found in pregnant women who had never used orthodontic appliances. The changes in these bacteria [29] are related to increased levels of oxidative stress such as malondialdehyde (MDA) [28], and cause a decrease in the salivary acidity and calcium content [30]. Due to this condition, the risk of dental caries and gingival disease increases. These findings show that oral cavity health in pregnant women must be considered, even down to the decision of using or not using orthodontic appliances.

Increased oxidative stress also creates an increase in the inflammatory response in the oral cavity. Various abnormalities in the soft tissues of the oral cavity of pregnant women include gingivitis and periodontitis. These diseases lead to an increase in the salivary IL-6 and TNF- α levels of pregnant women compared with pregnant women without gingivitis and periodontitis [31]. In this research, the IL-6 levels of all three sample groups showed no significant differences. IL-6 has a role in regulating paracrine to maintain foetal growth and placental hormone production. IL-6 also contributes to the development of the immune and hematopoietic system of the foetus [32,33].

The level of TNF- α in pregnant women with orthodontic appliances was higher compared with pregnant women who had never used orthodontic appliances. Physiological changes during pregnancy lead to changes in the oral mucosa. Pregnant women commonly experience diseases such as caries and periodontitis. Periodontitis causes an increase in the salivary-soluble TNF- α receptor, which leads to an increase in the TNF- α level [34]. An increase in the salivary TNF- α level in pregnant women, especially those who are obese, can affect the weight of the infant [35]. On the other hand, tooth movement caused by orthodontic force may increase salivary TNF- α [36] and IL-6 levels [14]. It can be concluded that, in the absence of pregnancy, the concentration of salivary IL-6 and TNF- α increases due to orthodontic appliance use. The risk of inflammation in pregnant women with orthodontic appliances is also caused by high OHI-S. Orthodontic appliance use can also increase plaque formation, which may impact the health of the gingiva and periodontium [37]. Oral hygiene is important to maintain and even increase, especially in pregnant women with orthodontic appliances, in order to prevent caries, gingivitis, and periodontitis. To maintain oral hygiene, several strategies are needed, including daily use of fluoride toothpaste and mouthwash. Daily use of both shows a reduced risk of caries during orthodontic treatment [38].

It should be noted that in orthodontics, over the past few years, research on the use of wire orthodontics derived from NiTi (nickel-titanium) on the adhesion of *Streptococcus mutans* has been performed. Various concentrations of Ni in the orthodontic wire influence the adhesion of *Streptococcus mutans*. Research by Pavlic et al. showed that Ni concentrations of 1000 µg/mL significantly inhibit adhesion [39]. Other research on the prevention of bacteria adhesion on orthodontic wire by applying titanium dioxide has been carried out [40,41]. The relationship with the maintenance of oral hygiene and daily use of fluoride toothpaste is also reported have an influence on the release of Ni [42,43], so it is likely that it also has an influence on the adhesion and formation of biofilms on orthodontic appliances.

5. Conclusions

The conclusion of this study is that the microbiome profile and IL-6 value showed normal results, so the use of orthodontic appliances during pregnancy should be allowed under certain conditions. Pregnant women who have orthodontic appliances must keep the oral cavity in good condition and maintain their orthodontic appliances periodically to avoid oral cavity problems. The limitation of this study is the short observation time. Future research should be conducted in a study encompassing multiple centres to strengthen this result.

Author Contributions: Conceptualization, F.K.D.K., R.I.R. and I.B.N.; methodology, F.K.D.K., R.I.R., U.T. and I.B.N.; validation, R.I.R., U.T., N.T., I.D. and I.B.N.; formal analysis, F.K.D.K.; investigation, F.K.D.K., R.I.R. and I.B.N.; data curation, F.K.D.K., R.I.R. and I.B.N.; writing—original draft preparation, F.K.D.K., R.I.R. and I.B.N.; writing—review and editing M.D.C.S., N.T. and I.D.; visualization, M.D.C.S., R.I.R. and N.T.; supervision, R.I.R., U.T., N.T., I.D. and I.B.N.; funding acquisition, R.I.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research received funding from Universitas Airlangga in the schema Hibah Riset Mandat (886/UN3/2018).

Institutional Review Board Statement: This research received a certificate of ethics from the Ethics Committee of the Faculty of Dentistry, Universitas Airlangga, registered number: 234 /HRECC.FODM/V/2019. All the participant has obtained the informed consent and agree to participated in this study. All methods were performed in accordance with relevant guidelines and regulations.

Informed Consent Statement: The participant has received the informed consent.

Data Availability Statement: The datasets used and analysed during the current study are available from the corresponding author upon reasonable request.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Giglio, J.A.; Lanni, S.M.; Laskin, D.M.; Giglio, N.W. Oral health care for the pregnant patient. *J. Can. Dent. Assoc.* **2009**, *75*, 45–48.
2. Hartnett, E.; Haber, J.; Krainovich-Miller, B.; Bella, A.; Vasilyeva, A.; Lange Kessler, J. Oral Health Pregnancy. *J. Obstet. Gynecol. Neonatal Nurs.* **2016**, *45*, 565–573. [[CrossRef](#)] [[PubMed](#)]
3. Bajinka, O.; Darboe, A.; Tan, Y.; Abdelhalim, K.A.; Cham, L.B. Gut microbiota and the human gut physiological changes. *Ann. Microbiol.* **2020**, *70*, 65. [[CrossRef](#)]
4. Berry, A.S.F.; Pierdon, M.K.; Mistic, A.M.; Sullivan, M.C.; O'Brien, K.; Chen, Y.; Murray, S.J.; Ramharack, L.A.; Baldassano, R.N.; Parsons, T.D.; et al. Remodeling of the maternal gut microbiome during pregnancy is shaped by parity. *Microbiome* **2021**, *9*, 146. [[CrossRef](#)]
5. Yang, H.; Guo, R.; Li, S.; Liang, F.; Tian, C.; Zhao, X.; Long, Y.; Liu, F.; Jiang, M.; Zhang, Y.; et al. Systematic analysis of gut microbiota in pregnant women and its correlations with individual heterogeneity. *Biofilms Microbiomes* **2020**, *6*, 32. [[CrossRef](#)]
6. Reyes, L.; Phillips, P.; Wolfe, B.; Golos, T.G.; Walkenhorst, M.; Progulsk-Fox, A.; Brown, M. *Porphyromonas gingivalis* and adverse pregnancy outcome. *J. Oral Microbiol.* **2017**, *9*, 1374153. [[CrossRef](#)]
7. Hirano, E.; Sugita, N.; Kikuchi, A.; Shimada, Y.; Sasahara, J.; Iwanaga, R.; Tanaka, K.; Yoshie, H. The association of *Aggregatibacter actinomycetemcomitans* with preeclampsia in a subset of Japanese pregnant women. *J. Clin. Periodontol.* **2012**, *39*, 229–238. [[CrossRef](#)]
8. Sukontapatipark, W. Bacterial colonization associated with fixed orthodontic appliances. A scanning electron microscopy study. *Eur. J. Orthod.* **2001**, *23*, 475–484. [[CrossRef](#)]

9. Anuwongnukroh, N.; Dechkunakorn, S.; Kanpiputana, R. Oral Hygiene Behavior during Fixed Orthodontic Treatment. *Dentistry* **2017**, *7*, 457.
10. Lucchese, A.; Bondemark, L.; Marcolina, M.; Manuelli, M. Changes in oral microbiota due to orthodontic appliances: A systematic review. *J. Oral Microbiol.* **2018**, *10*, 1476645. [[CrossRef](#)]
11. Scarpa, M.; Grillo, A.; Faggian, D.; Ruffolo, C.; Bonello, E.; D'Inca, R.; Scarpa, M.; Castagliuolo, I.; Angriman, I. Relationship between mucosa-associated microbiota and inflammatory parameters in the ileal pouch after restorative proctocolectomy for ulcerative colitis. *Surgery* **2011**, *150*, 56–67. [[CrossRef](#)] [[PubMed](#)]
12. Kunii, R.; Yamaguchi, M.; Tanimoto, Y.; Asano, M.; Yamada, K.; Goseki, T.; Kasai, K. Role of interleukin-6 in orthodontically induced inflammatory root resorption in humans. *Korean J. Orthod.* **2013**, *43*, 294. [[CrossRef](#)] [[PubMed](#)]
13. Yang, J.H.; Li, Z.C.; Kong, W.D.; Zhang, W.; Jia, Y.P.; Zhang, Y.L.; Liu, L.B.; Han, X.P. Effect of orthodontic force on inflammatory periodontal tissue remodeling and expression of IL-6 and IL-8 in rats. *Asian Pac. J. Trop. Med.* **2013**, *6*, 757–761. [[CrossRef](#)]
14. Baeshen, H.A. Assessment of salivary pro inflammatory cytokines profile level in patients treated with labial and lingual fixed orthodontic appliances. *PLoS ONE* **2021**, *16*, e0249999. [[CrossRef](#)] [[PubMed](#)]
15. Marahleh, A.; Kitaura, H.; Ohori, F.; Noguchi, T.; Nara, Y.; Pramusita, A.; Kinjo, R.; Ma, J.; Kanou, K.; Mizoguchi, I. Effect of TNF- α on osteocyte RANKL expression during orthodontic tooth movement. *J. Dent. Sci.* **2021**, *16*, 1191–1197. [[CrossRef](#)]
16. Afshar, M.K.; Safarian, F.; Torabi, M.; Farsinejad, A.; Mohammadzadeh, I. Comparison of TNF- α and IL-1 β Concentrations in Gingival Crevicular Fluid During Early Alignment Stage of Orthodontic Treatment in Adults and Adolescents. *Pesqui. Bras. Odontopediatria Odontol. Integr.* **2020**, *20*, e0004. [[CrossRef](#)]
17. Chen, Y.; Wong, W.K.; Seneviratne, J.C.; Huang, S.; McGrath, C.; Hagg, U. Associations between salivary cytokines and periodontal and microbiological parameters in orthodontic patients. *Medicine* **2021**, *100*, e24924. [[CrossRef](#)]
18. Bashirian, S.; Seyzedeh-Sabounchi, S.; Shirahmadi, S.; Soltanian, A.R.; Karimi-shahanjarini, A.; Vahdatinia, F. Socio-demographic determinants as predictors of oral hygiene status and gingivitis in schoolchildren aged 7–12 years old: A cross-sectional study. *PLoS ONE* **2018**, *13*, e0208886. [[CrossRef](#)]
19. Mahmoud, J.; Mehri, G.; Mohsen, T.; Yousefimanesh, H.; Maryam, R. Evaluation of salivary tumor necrosis factor-alpha in patients with the chronic periodontitis: A case-control study. *J. Indian Soc. Periodontol.* **2013**, *17*, 737. [[CrossRef](#)]
20. Radaic, A.; Kapila, Y.L. The oralome and its dysbiosis: New insights into oral microbiome-host interactions. *Comput. Struct. Biotechnol. J.* **2021**, *19*, 1335–1360. [[CrossRef](#)]
21. Mitchell, J. *Streptococcus mitis*: Walking the line between commensalism and pathogenesis. *Mol. Oral Microbiol.* **2011**, *26*, 89–98. [[CrossRef](#)] [[PubMed](#)]
22. Shukla, C.; Maurya, R.; Singh, V.; Tijare, M. Evaluation of changes in *Streptococcus mutans* colonies in microflora of the Indian population with fixed orthodontics appliances. *Dent. Res. J.* **2016**, *13*, 309.
23. Butera, A.; Gallo, S.; Maiorani, C.; Preda, C.; Chiesa, A.; Esposito, F.; Pascadopoli, M.; Scribante, A. Management of Gingival Bleeding in Periodontal Patients with Domiciliary Use of Toothpastes Containing Hyaluronic Acid, Lactoferrin, or Paraprobiotics: A Randomized Control. *Clin. Trial. Appl. Sci.* **2021**, *11*, 8586. [[CrossRef](#)]
24. Iskandar, B.; Lukman, A.; Syaputra, S.; Al-Abrori, U.N.H.; Surboyo, M.D.C.; Lee, C.K. Formulation, characteristics and antibacterial effects of *Euphorbia hirta* L. mouthwash. *J. Taibah Univ. Med. Sci.* **2021**, *17*, 271–282. [[CrossRef](#)] [[PubMed](#)]
25. Ghasemi, N.; Behnezhad, M.; Asgharzadeh, M.; Zeinalzadeh, E.; Kafil, H.S. Antibacterial Properties of Aloe vera on Intracanal Medicaments against *Enterococcus faecalis* Biofilm at Different Stages of Development. *Int. J. Dent.* **2020**, *2020*, 1–6. [[CrossRef](#)] [[PubMed](#)]
26. Jang, H.; Patoine, A.; Wu, T.T.; Castillo, D.A.; Xiao, J. Oral microflora and pregnancy: A systematic review and meta-analysis. *Sci. Rep.* **2021**, *11*, 16870. [[CrossRef](#)]
27. Kardos, S.; Tóthpál, A.; Laub, K.; Kristóf, K.; Ostorházi, E.; Rozgonyi, F.; Dobay, O. High prevalence of group B streptococcus ST17 hypervirulent clone among non-pregnant patients from a Hungarian venereology clinic. *BMC Infect. Dis.* **2019**, *19*, 1009. [[CrossRef](#)]
28. Wagle, M.; Basnet, P.; Vartun, Å.; Trovik, T.A.; Acharya, G. Oxidative stress levels and oral bacterial milieu in the saliva from pregnant vs. non-pregnant women. *BMC Oral Health* **2020**, *20*, 245. [[CrossRef](#)]
29. Wagle, M.; Basnet, P.; Vartun, Å.; Acharya, G. Nitric Oxide, Oxidative Stress and *Streptococcus mutans* and *Lactobacillus* Bacterial Loads in Saliva during the Different Stages of Pregnancy: A Longitudinal Study. *Int. Environ. Res. Public Health* **2021**, *18*, 9330. [[CrossRef](#)]
30. Kamate, W.I.; Vibhute, N.A.V.; Baad, R.K. Estimation of DMFT, Salivary *Streptococcus Mutans* Count, Flow Rate, Ph, and Salivary Total Calcium Content in Pregnant and Non-Pregnant Women: A Prospective Study. *J. Clin. Diagn. Res.* **2017**, *11*, 147–151. [[CrossRef](#)]
31. Machado, V.; Mesquita, M.F.; Bernardo, M.A.; Casal, E.; Proença, L.; Mendes, J.J. IL-6 and TNF- α salivary levels according to the periodontal status in Portuguese pregnant women. *PeerJ* **2018**, *6*, e4710. [[CrossRef](#)] [[PubMed](#)]
32. Vujacic, A.; Pavlovic, J.; Konic-Ristic, A. *Current Approaches in Orthodontics, The Role of Cytokines in Orthodontic Tooth Movement*; IntechOpen: London, UK, 2018; pp. 35–39.
33. Omere, C.; Richardson, L.; Saade, G.R.; Bonney, E.A.; Kechichian, T.; Menon, R. Interleukin (IL)-6: A Friend or Foe of Pregnancy and Parturition? Evidence from Functional Studies in Fetal Membrane Cells. *Front. Physiol.* **2020**, *11*, 891. [[CrossRef](#)] [[PubMed](#)]

34. Kibune, R.; Muraoka, K.; Morishita, M.; Ariyoshi, W.; Awano, S. Relationship between Dynamics of TNF- α and Its Soluble Receptors in Saliva and Periodontal Health State. *Dent. J.* **2022**, *10*, 25. [[CrossRef](#)] [[PubMed](#)]
35. Foratori-Junior, G.A.; Mosquim, V.; Buzalaf MA, R.; de Carvalho Sales-Peres, S.H. Salivary cytokines levels, maternal periodontitis and infants' weight at birth: A cohort study in pregnant women with obesity. *Placenta* **2021**, *115*, 151–157. [[CrossRef](#)]
36. Saadi, N.; Ghaib, N.H. Effect of Orthodontic Tooth Movement on Salivary Levels of Interleukin-1beta, Tumor Necrosis Factor-Alpha and Creactive Protein. *J. Baghdad Coll. Dent.* **2013**, *25*, 120–125. [[CrossRef](#)]
37. Singla, S.; Gupta, P.; Lehl, G.; Talwar, M. Effects of Reinforced Oral Hygiene Instruction Program with and without Professional Tooth Cleaning on Plaque Control and Gingival Health of Orthodontic Patients Wearing Multibracket Appliances. *J. Indian Orthod. Soc.* **2019**, *53*, 272–277. [[CrossRef](#)]
38. Hosseinzadeh Nik, T.; Hooshmand, T.; Farhadifard, H. Effect of Different Types of Toothpaste on the Frictional Resistance Between Orthodontic Stainless Steel Brackets and Wires. *J. Dent. (Tehran)* **2017**, *14*, 275–281.
39. Pavlic, A.; Begic, G.; Tota, M.; Abram, M.; Spalj, S.; Gobin, I. Bacterial Exposure to Nickel: Influence on Adhesion and Biofilm Formation on Orthodontic Archwires and Sensitivity to Antimicrobial Agents. *Materials* **2021**, *14*, 4603. [[CrossRef](#)]
40. Raji, S.H.; Shojaei, H.; Ghorani, P.S.; Rafiei, E. Bacterial colonization on coated and uncoated orthodontic wires: A prospective clinical trial. *Dent. Res. J. (Isfahan)* **2014**, *11*, 680–683.
41. Mollabashi, V.; Farmany, A.; Alikhani, M.Y.; Sattari, M.; Soltanian, A.R.; Kahvand, P.; Banisafar, Z. Effects of TiO₂-Coated Stainless Steel Orthodontic Wires on *Streptococcus mutans* Bacteria: A Clinical Study. *Int. J. Nanomed.* **2020**, *15*, 8759–8766. [[CrossRef](#)]
42. Pritam, A.; Priyadarshini, A.; Hussain, K.; Kumar, A.; Kumar, N.; Malakar, A. Assessment of nickel and chromium level in gingival crevicular fluid in patients undergoing orthodontic treatment with or without fluoridated toothpaste. *J. Pharm. Bioallied Sci.* **2021**, *13*, 1588–1590.
43. Enerbäck, H.; Lingström, P.; Möller, M.; Nylén, C.; Ödman Bresin, C.; Östman Ros, I.; Westerlund, A. Effect of a mouth rinse and a high-fluoride toothpaste on caries incidence in orthodontic patients: A randomized controlled trial. *Am. J. Orthodont. Dentofac. Orthop.* **2022**, *162*, 6–15.e3. [[CrossRef](#)] [[PubMed](#)]