

## VERIFIKASI EMAIL CORRESPONDING AUTHOR INHSP-ENFERMERIA

2 messages

**INHSP 2019** <inhsp2019@gmail.com>  
To: rini.rachmawaty@unhas.ac.id

Thu, Jul 16, 2020 at 2:30 PM

Yth Author

Untuk memastikan validitas email corresponding author, maka kami mengirimkan email ini. Kami juga melampirkan screen shoot artikel anda. Untuk itu mohon melihat (preview) screen shoot halaman pertama artikel manuscript anda, dan pastikan NAMA\_NAMA AUTHORS dan EMAIL CORRESPONDING AUTHOR sudah BENAR.



Mohon membalas email ini sebelum tanggal 18 Juli 2020 jam 12.00 WITA.

Kerjasama Bapak Ibu sekalian dalam verifikasi email ini, sangat menentukan kelancaran proses submission di Enfermeria

Terima kasih atas kerjasamanya

Hormat Kami  
Panitia INHSP 2019

Saldy Yusuf, PhD  
Ketua



drive.google.com/file/d/TK\_F01ZGyY3jx3i3dituHGWzrn72zh4/view

### Evaluation of quality of life-based on the length of use and the number of stents of post percutaneous coronary intervention (PCI) patients

Al Amin<sup>1</sup>, Rini Rachmawaty<sup>2\*</sup>, Takdir Tahir<sup>3</sup>, Musmulyono Musmulyono<sup>4</sup>

<sup>1</sup>Master of Nursing Program, Hasanuddin University, Makassar, Indonesia  
<sup>2</sup>Dr. Wahidin Sudirohusodo Hospital, Makassar, Indonesia  
<sup>3</sup>Faculty of Nursing, Hasanuddin University, Makassar, Indonesia  
<sup>4</sup>Hasanuddin University Hospital, Makassar, Indonesia

\*Corresponding author.  
Email address: rini.rachmawaty@unhas.ac.id (R. Rachmawaty)

**Abstract**  
**Objective:** Percutaneous Coronary Intervention (PCI) is intended to improve patients' quality of life (QoL). This study aims to evaluate the QoL of post PCI patients based on the length of use and the number of stents at the cardiology outpatient clinic of Indonesian Public Hospital.  
**Method:** An analytic survey method with a cross-sectional approach was used in this study. A total of 60 samples were included using a purposive sampling method. Data were analyzed with the Spearman correlation test with a confidence level of 95%.  
**Results:** The study shows that there was a correlation of the length of use ( $p=0.039$ ;  $r=0.267$ ) and the number of stents ( $p=0.001$ ;  $r=0.406$ ) with the patients' QoL.  
**Conclusion:** Patients with more prolonged PCI use and a lower number of stents will have better QoL. Hence, nurse practitioners should provide holistic care for post-PCI patients to

drive.google.com/file/d/1vLJQWECF4SKLQe2JcGAspWNd3\_Q31/view

### Cancer Patient's Experiences Towards Nurse's Caring Demeanor Based on Watson's Theory: A Qualitative Study

Indah Gita Cahyani<sup>1</sup>, Rini Rachmawaty<sup>2\*</sup>, Ilkafah Ilkafah<sup>3</sup>, Erina Erina<sup>4</sup>

<sup>1</sup>Faculty of Nursing, Universitas Hasanuddin, Makassar, Indonesia  
<sup>2</sup>Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

\*Corresponding author  
Email: rini.rachmawaty@unhas.ac.id (R. Rachmawaty)

**Abstract**  
**Objective:** To explore the cancer patient's experiences towards the nurse's caring demeanor based on Watson's theory.  
**Method:** A qualitative descriptive study was conducted in this study. Eight participants from a large hospital were recruited through a process of purposive sampling. Data were collected using in-depth interviews and analyzed using thematic analysis.  
**Results:** Four themes emerged from this study: (1) Development of a helping-trusting relationship; (2) Supportive of faith-hope; (3) Unprotective environment; and (4) Lack of interpersonal teaching.  
**Conclusion:** Most of Watson's theory-based caring demeanor was well performed by oncology nurses included in this study. Even so, there were still some shortcomings in delivering caring. It is important to provide a training program to enhance nurses' knowledge of interpersonal teaching for cancer patients.

**Keywords:** caring, neoplasms, nurse, qualitative research, Watson's theory.

drive.google.com/file/d/11UJdzVcuANwQbYKREfUjwSRzqWg/view



### Patient Satisfaction towards Healthcare Quality in Indonesian Public Hospital

Andi Rizani Catur Wulandari<sup>1</sup>, Rini Rachmawaty<sup>2\*</sup>, Ilkafah Ilkafah<sup>3</sup>, Erina Erina<sup>4</sup>

<sup>1</sup>Faculty of Nursing, Universitas Hasanuddin, Makassar, Indonesia  
<sup>2</sup>Nursing Study Program, Universitas Airlangga, Surabaya, Indonesia

\*Corresponding author.  
Email: rini.rachmawaty@unhas.ac.id (R. Rachmawaty)

**Abstract**  
**Objective:** This study aims to evaluate patient satisfaction towards healthcare quality in Indonesian public hospital.  
**Method:** The study design was quantitative with an analytic observational approach. A total of 39 respondents were included using proportional stratified random sampling. The modified Press Ganey inpatient survey was used in this study and has been translated into the Indonesian language using cross-cultural adaptation.  
**Results:** The patient satisfaction with healthcare quality was significantly different based on the type of inpatient class ( $p=0.000$ ), and the differences were found in four dimensions: effectiveness ( $p=0.009$ ), accessibility ( $p=0.001$ ), patient-centeredness ( $p=0.003$ ), and equity ( $p=0.001$ ). Meanwhile, no statistical difference was found in patient satisfaction with healthcare quality based on patients' length of stay.  
**Conclusion:** Patients' satisfaction with healthcare quality was significantly different based on the type of inpatient class, but not based on the length of stay. The findings would provide



**Rini Rachmawaty** <rini.rachmawaty@unhas.ac.id>  
To: INHSP 2019 <inhsp2019@gmail.com>

Thu, Jul 16, 2020 at 10:19 PM

Thank you so much for your email. Yes, I received your email and this is my email address for correspondence.

Best regards,  
**Rini Rachmawaty, PhD, MN, BSN**  
Vice Dean for Academic, Research, & Innovation  
Faculty of Nursing, Universitas Hasanuddin, Indonesia  
Cell No. (+62) 812-1391-6730  
Email Addresses: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id); [rini.rachmawaty80@gmail.com](mailto:rini.rachmawaty80@gmail.com)

[Quoted text hidden]

---

## Modified Manuscript\_Rini Rachmawaty et al 20210207

1 message

---

**Rini Rachmawaty** <rini.rachmawaty@unhas.ac.id>

Mon, Feb 8, 2021 at 10:44 AM

To: INHSP 2019 <inhsp2019@gmail.com>

Cc: ilkafah@vokasi.unair.ac.id, syahrulsaid@unhas.ac.id, yudihardiantomcr@gmail.com

Asslm.

Dear INHSP Committee,

Terlampir manuscript Rini Rachmawaty et al yg telah dimodify sesuai masukan editor.

Terima kasih.

Wasslm.

Rini

**Rini Rachmawaty, PhD, MN, BSN**

*Vice Dean for Academic, Research, & Innovation*

*Faculty of Nursing, Universitas Hasanuddin, Indonesia*

*Cell No. (+62) 812-1391-6730*

*Email Addresses: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id); [rini.rachmawaty80@gmail.com](mailto:rini.rachmawaty80@gmail.com)*

---

### 2 attachments



**Rini Rachmawaty et al\_TABLE OF REVISION\_20210207.docx**

14K



**Rachmawaty et al\_Abdominal Massage\_MODIFIED\_20210207.docx**

66K

## **Abdominal massage for constipation relief in stroke patients: A participatory action research**

**Authors:** Rini Rachmawaty\*, Ilkafah Ilkafah, Syahrul Syahrul, Yudi Hardianto

\*Corresponding Author: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id)

**Table of Revision**

<b>No.</b>	<b>Editor Feedback/Additional Information</b>	<b>Revised Page</b>
1.	There is no authorization and evaluation by the ethics committee (if they provide it, it could be accepted) and it is an intervention study.	2
2.	Declaration of Conflict of Interest	5

## Abdominal massage for constipation relief in stroke patients: A participatory action research

Rini Rachmawaty<sup>a,\*</sup>, Ilkafah Ilkafah<sup>b</sup>, Syahrul Syahrul<sup>c</sup>, Yudi Hardianto<sup>d</sup>

<sup>a</sup>Department of Nursing Management, Faculty of Nursing, Universitas Hasanuddin, Indonesia.

<sup>b</sup>Nursing Study Program, Universitas Airlangga, Indonesia.

<sup>c</sup>Department of Community Health Nursing, Faculty of Nursing, Universitas Hasanuddin, Indonesia.

<sup>d</sup>Department of Physiotherapy, Faculty of Nursing, Universitas Hasanuddin, Indonesia.

\*Corresponding author.

E-mail address: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id) (R. Rachmawaty)

### Abstract

**Objective:** To improve knowledge and skills of family members and community health volunteers in applying abdominal massage to stroke patients with constipation.

**Method:** A participatory action research, involving family members and community health volunteers to participate in the training and supervision sessions about abdominal massage for constipation relief in stroke patients. Knowledge enrichment was assessed using questionnaire, while the abdominal massage skill was evaluated using direct observation. Qualitative interviews were conducted to evaluate patients' experiences regarding the intervention program.

**Results:** The program showed the improvement of participants' knowledge and skills in applying abdominal massage for stroke patients with constipation. Likewise, patients confirmed their constipation relieved after the application of abdominal massage.

**Conclusion:** This study showed that training and supervision sessions about abdominal massage for constipation alleviation in stroke patients are feasible, efficient, and beneficial programs to enhance knowledge and skill of family members and community health volunteers.

**Keywords:** Abdominal massage; community engagement program; community health volunteers; constipation; stroke.

### Introduction

Stroke is one of the common causes of death and disability in Indonesia.<sup>1</sup> It is reported that stroke prevalence increased from 7% in 2013 to 10.9% in 2018, meaning that the prevalence of people with stroke in Indonesia was ten per one thousand population.<sup>2,3</sup> In 2016, number of Indonesians died due to stroke was 212,963 and the highest prevalence was reported in the South Sulawesi province (17.9%).<sup>4,5</sup>

Besides disability, one of the most common complications among people with stroke is constipation which can lead to increased disability and death.<sup>6-8</sup> Constipation in stroke patients occurred mainly due to lack of physical mobility, difficulty to swallowing fluid and fiber, dependence on others to use the toilet, and the use of laxatives.<sup>8</sup> If the stroke patient with constipation has a history of heart failure, hypertension, or myocardial infarction; straining during defecation can lead to increased burden of the heart, rupture of blood vessels, and premature death.<sup>9-11</sup> Concerning these complications, it is crucial to prevent constipation.<sup>6,9</sup>

Abdominal massage is revealed as one of several interventions to prevent and to treat constipation among elderly, people with stroke, and those with musculoskeletal disorders,

cardiovascular diseases, chronic blood disease, and post-operative ileus.<sup>6,12-15</sup> Abdominal massage has been proved as an inexpensive and safe non-pharmacologic therapy to relieve constipation.<sup>16</sup>

Likewise, a previous study has confirmed that applying abdominal massage can prevent constipation in stroke patients hospitalized in one of Indonesian public hospital and can avoid the patients taking laxatives.<sup>6,9</sup> However, studies specifically focusing on how abdominal massage performed by family members and community health volunteers to prevent constipation in stroke patients at home is rare. Hence, this study aimed to implement abdominal massage performed by family members and community health volunteers to alleviate constipation in stroke patients at their homes.

#### **Method**

This study was participatory action research (PAR) using mix-methods data collection including interviews, FGDs, and questionnaires. The participants consisted of 11 family members of stroke patients and five community health volunteers. The intervention including training and supervision participants about the application of abdominal massage to stroke patients as well as conducting evaluation of the process and outcomes. The training was delivered in the forms of modules, audiovisual media, skill demonstration. Participants performed abdominal massage to stroke patients under the supervision of the research team. Home visits was conducted to evaluate knowledge and skills of participants in applying abdominal massage independently to stroke patients.

Regarding the ethical considerations, this study was a follow up study from the initial intervention study that was conducted at one of Indonesian public hospitals and had obtained ethical approval from the Research Ethics Committee of Faculty of Medicine, Universitas Hasanuddin (No. 513/H4.8.4.5.31/PP36-KOMETIK/2017). Furthermore, prior to data collection, the researchers reported to the Ministry of Health of Makassar City to obtain permission to conduct research at the Antara Public Health Center. Finally, informed consent was obtained from all participants prior to their participation in this study.

In conducting the training session, the principal investigator (PI) initially explained the purpose of the activity, then followed by a pre-test to assess the knowledge of family and community health volunteers about constipation and abdominal massage. After that, the research team provided health education to patients' families, community health volunteers, and the staff of public health center about stroke, constipation, and abdominal massage for constipation relief using educational videos. Next, one of the co-PIs re-demonstrated the steps of abdominal massage directly to stroke patients. After the training conducted, families and community health volunteers were required to complete the post-test to evaluate the level of knowledge of family and community health volunteers about constipation and abdominal massage.

The next activity was home visit, through which the research team and the community health volunteer visited the patients and evaluated the ability of the family members and the community health volunteers to apply abdominal massage for stroke patients. During this visit, the researcher team observed and checked using questionnaire whether the participants correctly performed the steps of abdominal massage. Participants could repeat the procedures until they re-demonstrated it properly. At the end, patients were asked about their condition after receiving abdominal massage. The last activity of this PAR was inviting all participants to the final meeting at the public health center to evaluate the overall program using focused-group discussion.

Regarding data analysis, knowledge changes was analyzed using paired t-test, while participants knowledge about constipation and abdominal massage was analyzed using repeated

**Commented [RRI]:** The authorization and evaluation by the ethics committee has been explained.

measures ANCOVA. The patients' experiences after receiving abdominal massage were analyzed using qualitative analysis.

### Result

The demographic characteristics of participants in the counseling and training sessions is shown in **Table 1**. The average age of the stroke patients' families was 36.8 and the community health volunteers was 46.8. Participants were predominated by women and most of them were high school graduates. Two third of the participants were housewives and the rest were teachers and entrepreneurs.

**Table 1** Demographic characteristics of the participants in the training and supervision sessions (n=16)

Characteristics	Family (n= 11)		Community Health Volunteer (n= 5)	
	Mean (SD)	n (%)	Mean (SD)	n (%)
Age	36.82 (8.32)		46.8 (11.78)	
<b>Gender</b>				
Female		10 (90.9)		5 (100.0)
Male		1 (9.1)		
<b>Education</b>				
Junior High School		2 (18.2)		
Senior High School		6 (54.5)		5 (100.0)
Diploma III		1 (9.1)		
Bachelor		1 (9.1)		
Master		1 (9.1)		
<b>Occupation</b>				
Housewife		5 (45.5)		3 (60.0)
Private employee		2 (18.2)		
Entrepreneur		2 (18.2)		1 (20.0)
Lecturer		1 (9.1)		
Teacher		1 (9.1)		
Retired				1 (20.0)

SD, standard deviation

**Table 2** shows that after participating in the training and supervising sessions, patients' families gained better knowledge about constipation ( $p=0.001$ ) and abdominal massage ( $p=0.000$ ), while the community health volunteers attained better knowledge only for abdominal massage ( $p=0.023$ ).

**Table 2** Participants' knowledge of constipation and abdominal massage application on stroke patients before and after counseling (n=16)

Knowledge	Family (n= 11)			Community health volunteer (n= 5)		
	Pre-test	Post-test	$p^a$	Pre-test	Post-test	$p^a$
	Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)	
Constipation	6.55 (1.75)	8.36 (0.92)	<b>0.001*</b>	6.60 (2.07)	8.00 (0.71)	0.113
Abdominal Massage	6.18 (1.25)	9.00 (0.89)	<b>0.000*</b>	5.40 (3.21)	8.60 (1.14)	<b>0.023</b>

SD, standard deviation. a: using paired t-test. \* $p<0.05$

**Table 3** depicts the ability of participants to perform abdominal massage. As clearly seen in **Table 3**, not all families and community health volunteers can provide correct and complete abdominal massage. The participants complained that they had difficulty in memorizing abdominal massage procedures due to age, lack of practice, lack of reading the provided modules, and rarely



watching provided simulation videos and YouTube. The participants reported that the abdominal massage was only applied to patients when they complained about the onset of constipation.

**Table 3.** The evaluation of participants in applying abdominal massage on stroke patients (n=12)

Ability to perform Abdominal Massage	Family (n= 7)	Community health volunteer (n= 5)
	Well performed, n (%)	
Step 1: Stroke upwards 3 times	7 (100)	5 (100)
Step 2: Stroke towards the bottom of the abdomen 3 times	7 (100)	4 (80)
Step 3: Effleurage or circular stroking	4 (57.1)	4 (80)
Step 4: Palmer kneading	7 (100)	5 (100)
Step 5: As Step 4, but moving up the stomach	6 (85.7)	4 (80)
Step 6: Repeat Steps 4 and 5	4 (57.1)	3 (60)
Step 7: Stroking	7 (100)	5 (100)
Step 8: Hand vibrations over the umbilical area	7 (100)	5 (100)
Step 1 - 8	2 (28.6)	3 (60)

**Table 4** demonstrates the enhancement of participants' knowledge and skills in applying abdominal massage for stroke patients with constipation. As depicted by **Table 4**, the family members' knowledge about constipation ( $p=0.011$ ) and abdominal massage ( $p=0.003$ ) significantly rose from pre-test to post-test 2. Likewise, the knowledge of community health volunteers about constipation ( $p=0.011$ ) and abdominal massage ( $p=0.003$ ) significantly improved overtime.

**Table 4.** Participants' knowledge of constipation and abdominal massage application on stroke patients before training, after training, and after program evaluation (n=9)

Knowledge	Family (n= 4)				Community health volunteer (n= 5)			
	Pre-test	Post-test 1	Post-test 2	$p^a$	Pre-test	Post-test 1	Post-test 2	$p^a$
	Mean (SD)	Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)	Mean (SD)	
Constipation	5.37 (0.91)	7.73 (0.36)	9.09 (0.37)	<b>0.011*</b>	6.70 (0.80)	8.02 (0.32)	9.53 (0.33)	<b>0.011*</b>
Abdominal massage	5.38 (1.39)	9.20 (0.53)	9.19 (0.36)	<b>0.003*</b>	5.30 (1.23)	8.44 (0.47)	8.85 (0.32)	<b>0.003*</b>

Pre-test: Before training session, Post-test 1: After training session, Post-test 2: After program evaluation. SD, standard deviation. a: Using repeated measures ANCOVA. \* $P<0.05$ .

Based on the qualitative interviews, patients confirmed that constipation was relieved after receiving abdominal massage for three times. They also said that having abdominal massage induced them to defecate.

### Discussion

This study is among the first studies that explored the effect of abdominal massage on constipation relief in Indonesian stroke population living in the community, and among the few studies of abdominal massage conducted in stroke population. The previous study has shown that abdominal massage is feasible and effective to prevent constipation in people with neurological condition.<sup>17</sup> Through that study, the physiotherapists taught the patients and their carer about abdominal massage and allowed them to practice. Information about abdominal massage was also provided in the CDs. Like current study, the previous study found that abdominal massage was effective to alleviate constipation.<sup>18</sup>

This study is also the first study in Indonesia to involve community health volunteers to apply abdominal massage to people with stroke. Community health volunteers are quite common in Indonesia and many developing countries, especially in villages and rural areas where health professionals are scarce.<sup>19</sup> They are valuable workforce in primary and community health care that assist health professionals to provide optimal care for patients.<sup>19</sup> They are involved in many aspects of health promotion, prevention, intervention, rehabilitation, and palliative care.<sup>20-22</sup> They have been working hand in hand with health professionals in providing community care for communicable diseases such as malaria<sup>20</sup>, tuberculosis<sup>21</sup>, and HIV/AIDS.<sup>22</sup> They also provided support for new mothers in maternal and antenatal care by working under supervision of community midwives.<sup>23</sup> Involving community health volunteers in providing abdominal massage for stroke patients with constipation is an efficient way to help people with stroke manage their constipation. Patients, family members and community health professionals has been familiar with them and thus, collaboration can be ensured.

There are some limitations of this study: No control group and small sample size. During home visit, some patients were not received abdominal massage due to contraindication. Even so, this study showed that the application of abdominal massage to stroke patients with constipation is feasible by providing training and supervision sessions for family members and community health volunteers. It is considered more economically preferred as the application of abdominal massage does not require any expensive tools. Furthermore, as it is performed by family members, the adherence to the abdominal massage application is guaranteed as it only requires several minutes to apply to the patient.

#### **Conclusion**

Participants experienced improved knowledge and skills of constipation and abdominal massage after attending training and supervision sessions. Likewise, patients confirmed ease of defecation after receiving abdominal massage. This implementation study proves that abdominal massage performed by family and health community volunteers was advantageous and effective in alleviating constipation in stroke patients. Further study is required with larger sample size, multicenter, and more rigorous methods to explore more about the effect of abdominal massage on constipation alleviation in people with stroke.

#### **Acknowledgement**

This study supported by Grant from the Institute for Research and Community Service [*Lembaga Penelitian dan Pengabdian Kepada Masyarakat*], Universitas Hasanuddin, Makassar, Indonesia. The authors would like to thank to the Head of Antara Public Health Center, drg. Weri LP. Saroengoe, and all participants involved in this study.

#### **Conflict of Interest**

The authors declare no financial or personal interests that could bias the work.

**Commented [RR2]:** Declaration of Conflict of Interest has been added.

#### **REFERENCE**

1. World Health Organization. State of Health Inequality: Indonesia. Geneva: World Health Organization; 2017.
2. Kemenkes RI. Peningkatan Gaya Hidup Sehat dengan perilaku «CERDIK». 2018:10.
3. Kementerian Kesehatan Republik Indonesia. Riskesdas 2018. Jakarta; 2018.
4. Johnson CO., Nguyen M., Roth GA., Nichols E., Alam T., Abate D., et al. Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol.* 2019;18(5):439-58, doi: 10.1016/S1474-4422(19)30034-1.

5. Indonesia KKR. Riset Kesehatan Dasar 2013. Jakarta; 2013.
6. Sinclair M. The use of abdominal massage to treat chronic constipation. *J Bodyw Mov Ther.* 2011;15(4):436-45, doi: 10.1016/j.jbmt.2010.07.007.
7. Ronald L. Hickman, FAAN Celeste M. Alfes JJF. *Handbook of Clinical Nursing: Medical-Surgical Nursing.* New York: Springer Publishing Company; 2018.
8. Li J., Yuan M., Liu Y., Zhao Y., Wang J., Guo W., et al. Incidence of constipation in stroke patients: A systematic review and meta-analysis. *Med (United States).* 2017;96(25):1-6, doi: 10.1097/MD.0000000000007225.
9. Yacoline Pailungan F., Kaelan C., Rachmawaty R. Pengaruh Pemberian Massage Abdomen Terhadap Penurunan Konstipasi Pada Pasien Stroke Iskemik Di Rsup Dr. Wahidin Sudirohusodo Makassar. *Patria Artha J Nurs Sci.* 2017;1(1):25-35.
10. Engler TMN de M., Dourado CC., Amâncio TG., Farage L., de Mello PA., Padula MP. Stroke: Bowel Dysfunction in Patients Admitted for Rehabilitation. *Open Nurs J.* 2014;8(1):43-7, doi: 10.2174/1874434601408010043.
11. Sumida K., Molnar MZ., Potukuchi PK., Thomas F., Lu JL., Yamagata K., et al. Constipation and risk of death and cardiovascular events. *Atherosclerosis.* 2019;281(May 2018):114-20, doi: 10.1016/j.atherosclerosis.2018.12.021.
12. Lämås K., Graneheim UH., Jacobsson C. Experiences of abdominal massage for constipation. *J Clin Nurs.* 2012;21(5-6):757-65, doi: 10.1111/j.1365-2702.2011.03946.x.
13. Deckers B., Von Schoen-Angerer T., Voggenreiter B., Vagedes J. External nursing applications in the supportive management of prolonged postoperative ileus description of interventions and case report. *Holist Nurs Pract.* 2016;30(4):216-21, doi: 10.1097/HNP.000000000000158.
14. Çevik K., Çetinkaya A., Yiğit Gökbel K., Menekşe B., Saza S., Tikiz C. The effect of abdominal massage on constipation in the elderly residing in rest homes. *Gastroenterol Nurs.* 2018;41(5):396-402, doi: 10.1097/SGA.0000000000000343.
15. Birimoglu Okuyan C., Bilgili N. Effect of abdominal massage on constipation and quality of life in older adults: A randomized controlled trial. *Complement Ther Med.* 2019;47:102219, doi: 10.1016/j.ctim.2019.102219.
16. Emly MC. Abdominal Massage for Constipation. En: Haslam J. LJ, editor. *Therapeutic Management of Incontinence and Pelvic Pain.* London: Springer; 2008. p. 223-5.
17. McClurg D., Hagen S., Hawkins S., Lowe-Strong A. Abdominal massage for the alleviation of constipation symptoms in people with multiple sclerosis: a randomized controlled feasibility study. *Mult Scler J.* 2010;17(2):223-33, doi: 10.1177/1352458510384899.
18. McClurg D., Hagen S., Hawkins S., Lowe-Strong A. Abdominal massage for the alleviation of constipation symptoms in people with multiple sclerosis: A randomized controlled feasibility study. *Mult Scler J.* 2011;17(2):223-33, doi: 10.1177/1352458510384899.
19. Bhattacharyya K., Winch P., LeBan K., Tien M. *Community Health Worker Incentives and Disincentives: How They Affect Motivation, Retention, and Sustainability.* Arlington: the Basic Support for Institutionalizing Child Survival Project (BASICS II) for the United States Agency for International Development; 2001.
20. Soesilo D., Wahyuni CU. Improving the Village Health Volunteers' Knowledge, Malaria Diagnostic Skill and Home Visits through the Observed Training (Study in Municipality of Tidore Islands North Maluku Province Indonesia). *Indian J Public Heal Res Dev.* 2019;10(12):1671-5.
21. Lukman M., MSi KI., Yani DI., Juniarti N. Exploring Strategies to Improve the Performance of

Community Health Volunteers for Tuberculosis Care and Prevention: A Qualitative Study. *Int J community based Nurs midwifery*. 2019;7(4):270.

22. Palaia AM. Understanding the motivation, ability and attrition of HIV/AIDS community health volunteers in resource poor settings: Implications for program sustainability. The Johns Hopkins University; 2010.
23. Tumbelaka P., Limato R., Nasir S., Syafruddin D., Ormel H., Ahmed R. Analysis of Indonesia's community health volunteers (kader) as maternal health promoters in the community integrated health service (Posyandu) following health promotion training. *Int J Community Med Public Heal*. 2018;5(3):127-35.

---

## Publishing Agreement completed for your article [ENFCLI\_1924]

1 message

---

**Elsevier - Author Forms** <Article\_Status@elsevier.com>  
To: rini.rachmawaty@unhas.ac.id

Fri, Oct 1, 2021 at 11:05 PM

# ELSEVIER

Dear Dr. Rachmawaty,

Thank you for completing the Publishing Agreement Form for your article *Abdominal massage for constipation relief in stroke patients*. Please find attached a copy of the "Journal Publishing (License) Agreement" which you completed online on October 01, 2021.

If you have any questions, please do not hesitate to contact us. To help us assist you, please quote our article reference ENFCLI1924 in all correspondence.

Now that your article has been accepted, you will want to maximize the impact of your work. Elsevier facilitates and encourages authors to share their article responsibly. To learn about the many ways in which you can share your article whilst respecting copyright, visit: [www.elsevier.com/sharing-articles](http://www.elsevier.com/sharing-articles).

We are committed to publishing your article as quickly as possible.

Kind regards,  
Elsevier Researcher Support

### Have questions or need assistance?

Please do not reply to this automated message.

For further assistance, please visit our [Elsevier Support Center](#) where you search for solutions on a range of topics and find answers to frequently asked questions.

You can also talk to our researcher support team by phone 24 hours a day from Monday-Friday and 24/7 by live chat and email.

© 2021 Elsevier Ltd | [Privacy Policy](http://www.elsevier.com/privacypolicy) <http://www.elsevier.com/privacypolicy>

Elsevier Limited, The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom, Registration No. 1982084. This e-mail has been sent to you from Elsevier Ltd. To ensure delivery to your inbox (not bulk or junk folders), please add [article\\_status@elsevier.com](mailto:article_status@elsevier.com) to your address book or safe senders list.



## Publishing Agreement

Elsevier España, S.L.U.

**Abdominal massage for constipation relief in stroke patients:**

Corresponding author	Dr. Rini Rachmawaty
E-mail address	rini.rachmawaty@unhas.ac.id
Journal	Enfermería Clínica
Our reference	ENFCLI1924
PII	S1130-8621(21)00210-2

**Your Status**

- I am one author signing on behalf of all co-authors of the manuscript

**Assignment of Copyright**

I hereby assign to Elsevier España, S.L.U. the copyright in the manuscript identified above (where Crown Copyright is asserted, authors agree to grant an exclusive publishing and distribution license) and any tables, illustrations or other material submitted for publication as part of the manuscript (the "Article"). This assignment of rights means that I have granted to Elsevier España, S.L.U., the exclusive right to publish and reproduce the Article, or any part of the Article, in print, electronic and all other media (whether now known or later developed), in any form, in all languages, throughout the world, for the full term of copyright, and the right to license others to do the same, effective when the Article is accepted for publication. This includes the right to enforce the rights granted hereunder against third parties.

**Supplemental Materials**

"Supplemental Materials" shall mean materials published as a supplemental part of the Article, including but not limited to graphical, illustrative, video and audio material.

With respect to any Supplemental Materials that I submit, Elsevier España, S.L.U. shall have a perpetual worldwide, non-exclusive right and license to publish, extract, reformat, adapt, build upon, index, redistribute, link to and otherwise use all or any part of the Supplemental Materials in all forms and media (whether now known or later developed), and to permit others to do so.

**Research Data**

"Research Data" shall mean the result of observations or experimentation that validate research findings and that are published separate to the Article, which can include but are not limited to raw data, processed data, software, algorithms, protocols, and methods.

With respect to any Research Data that I wish to make accessible on a site or through a service of Elsevier España, S.L.U., Elsevier España, S.L.U. shall have a perpetual worldwide, non-exclusive right and license to publish, extract, reformat, adapt, build upon, index, redistribute, link to and otherwise use all or any part of the Research Data in all forms and media (whether now known or later developed) and to permit others to do so. Where I have selected a specific end user license under which the Research Data is to be made available on a site or through a service, the publisher shall apply that end user license to the Research Data on that site or service.

**Reversion of rights**

Articles may sometimes be accepted for publication but later rejected in the publication process, even in some cases after public posting in "Articles in Press" form, in which case all rights will revert to the author (see <https://www.elsevier.com/about/our-business/policies/article-withdrawal>).

## Revisions and Addenda

I understand that no revisions, additional terms or addenda to this Journal Publishing Agreement can be accepted without Elsevier España, S.L.U.'s express written consent. I understand that this Journal Publishing Agreement supersedes any previous agreements I have entered into with Elsevier España, S.L.U. in relation to the Article from the date hereof.

## Author Rights for Scholarly Purposes

I understand that I retain or am hereby granted (without the need to obtain further permission) the Author Rights (see description below), and that no rights in patents, trademarks or other intellectual property rights are transferred to Elsevier España, S.L.U.. The Author Rights include the right to use the [Preprint](#), [Accepted Manuscript](#) and the [Published Journal Article](#) for [Personal Use](#) and [Internal Institutional Use](#). They also include the right to use these different versions of the Article for [Scholarly Sharing](#) purposes, which include sharing:

- the Preprint on any website or repository at any time;
- the Accepted Manuscript on certain websites and usually after an embargo period;
- the Published Journal Article only privately on certain websites, unless otherwise agreed by Elsevier España, S.L.U..

In the case of the Accepted Manuscript and the Published Journal Article the Author Rights exclude Commercial Use (unless expressly agreed in writing by Elsevier España, S.L.U.), other than use by the author in a subsequent compilation of the author's works or to extend the Article to book length form or re-use by the author of portions or excerpts in other works (with full acknowledgment of the original publication of the Article).

## Author Representations / Ethics and Disclosure / Sanctions

I affirm the Author Representations noted below, and confirm that I have reviewed and complied with the relevant Instructions to Authors, Ethics in Publishing policy, Declarations of Interest disclosure and information for authors from countries affected by sanctions (Iran, Cuba, or Syria). Please note that some journals may require that all co-authors sign and submit Declarations of Interest disclosure forms. I am also aware of the publisher's policies with respect to retractions and withdrawal (<https://www.elsevier.com/about/our-business/policies/article-withdrawal>).

For further information see the publishing ethics page at <https://www.elsevier.com/about/our-business/policies/publishing-ethics> and the journal home page. For further information on sanctions, see <https://www.elsevier.com/about/our-business/policies/trade-sanctions>

### Author representations

- The Article I have submitted to the journal for review is original, has been written by the stated authors and has not been previously published.
- The Article was not submitted for review to another journal while under review by this journal and will not be submitted to any other journal.
- The Article and the Supplemental Materials do not infringe any copyright, violate any other intellectual property, privacy or other rights of any person or entity, or contain any libellous or other unlawful matter.
- I have obtained written permission from copyright owners for any excerpts from copyrighted works that are included and have credited the sources in the Article or the Supplemental Materials.
- Except as expressly set out in this Journal Publishing Agreement, the Article is not subject to any prior rights or licenses.
- If I and/or any of my co-authors reside in Iran, Cuba, or Syria, the Article has been prepared in a personal, academic or research capacity and not as an official representative or otherwise on behalf of the relevant government or institution.
- If I am using any personal details or images of patients, research subjects or other individuals, I have obtained all consents required by applicable law and complied with the publisher's policies relating to the use of such images or personal information. See <https://www.elsevier.com/about/our-business/policies/patient-consent> for further information.
- Any software contained in the Supplemental Materials is free from viruses, contaminants or worms.

- If the Article or any of the Supplemental Materials were prepared jointly with other authors, I have informed the co-author(s) of the terms of this Journal Publishing Agreement and that I am signing on their behalf as their agent, and I am authorized to do so.

## Governing Law and Jurisdiction

This Agreement will be governed by and construed in accordance with the laws of the country or state of Elsevier España, S.L.U. ("the Governing State"), without regard to conflict of law principles, and the parties irrevocably consent to the exclusive jurisdiction of the courts of the Governing State.

For information on the publisher's copyright and access policies, please see <http://www.elsevier.com/copyright>.  
For more information about the definitions relating to this agreement click [here](#).

I have read and agree to the terms of the Journal Publishing Agreement.

1 October 2021

T-copyright-v22/2017



---

## Rights and Access form completed for your article [ENFCLI\_1924]

1 message

---

**Elsevier - Author Forms** <Article\_Status@elsevier.com>  
To: rini.rachmawaty@unhas.ac.id

Fri, Oct 1, 2021 at 11:05 PM

# ELSEVIER

Dear Dr. Rachmawaty,

Thank you for completing the Rights and Access Form for your article *Abdominal massage for constipation relief in stroke patients*: on October 01, 2021.

The Order Summary is attached to this email.

If you have any questions, please do not hesitate to contact us. To help us assist you, please quote our article reference ENFCLI1924 in all correspondence.

Now that your article has been accepted, you will want to maximize the impact of your work. Elsevier facilitates and encourages authors to share their article responsibly. To learn about the many ways in which you can share your article whilst respecting copyright, visit: [www.elsevier.com/sharing-articles](http://www.elsevier.com/sharing-articles).

Kind regards,  
Elsevier Researcher Support



### Seven strategies for you to create a brand and promote your research

Learn how to give your research the visibility it deserves with these seven strategies.

> [Access module now](#)

#### Have questions or need assistance?

Please do not reply to this automated message.

For further assistance, please visit our [Elsevier Support Center](#) where you search for solutions on a range of topics and find answers to frequently asked questions.

You can also talk to our researcher support team by phone 24 hours a day from Monday-Friday and 24/7 by live chat and email.

© 2021 Elsevier Ltd | [Privacy Policy](http://www.elsevier.com/privacypolicy) <http://www.elsevier.com/privacypolicy>

Elsevier Limited, The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom, Registration No. 1982084. This e-mail has been sent to you from Elsevier Ltd. To ensure delivery to your inbox (not bulk or junk folders), please add [article\\_status@elsevier.com](mailto:article_status@elsevier.com) to your address book or safe senders list.

---



**Order Confirmation.html**

12K

## Rights and Access

**Abdominal massage for constipation relief in stroke patients:**

Corresponding author	Dr. Rini Rachmawaty
E-mail address	rini.rachmawaty@unhas.ac.id
Journal	Enfermeria Clinica
Our reference	ENFCLI1924
PII	S1130-8621(21)00210-2

**Order Confirmation**

Thank you for taking the time to complete the Rights and Access form.

<b>Order number</b>	OACSRENFCLI19240
<b>Order date</b>	1 October 2021

<b>Research Funders</b>	Universitas Hasanuddin, Institute for Research and Community Service Grant numbers: 6128/UN4.21/PL.01.02/2018
-------------------------	--

<b>Publishing Option</b>	Subscription
--------------------------	--------------

**Publishing Agreement**

- I am one author signing on behalf of all co-authors of the manuscript

I may post the accepted manuscript in my institutional repository and make this public after an embargo period. To ensure the sustainability of peer-reviewed research in journal publications, I may not share the final article publicly, for example on ResearchGate or Academia.edu.

Further details on [Elsevier Sharing Policy here](#).

Based on information provided the embargo period/end date is 12 months.

## Share your article [ENFCLI\_1924] for free

1 message

Elsevier - Status of your article <article\_status@elsevier.com>  
To: rini.rachmawaty@unhas.ac.id

Wed, Dec 22, 2021 at 7:51 PM

# ELSEVIER

## Share your article!

Dear Dr. Rachmawaty,

We are pleased to inform you that the final version of your article *Abdominal massage for constipation relief in stroke patients*: is now available online along with full bibliographical references.

To help you access and share your work, we've created a share link, a custom URL that provides **50 days of free access** to your article. Users who click on such a link before February 10, 2022 will be redirected to the final version of the article on ScienceDirect. It is not necessary to register, subscribe or pay, you just have to click on the link to read the article.



Your custom sharing link is:  
<https://authors.elsevier.com/a/1eHuY418DR0oHs>

Click the icons below to share it with your network:



We encourage you to also use this link to download a copy of the article for your own records. It's also an easy and quick way to share your work with your peers, co-authors, and friends. And if you wish, we invite you to add it to your website or profile on social networks such as Facebook, Google+ and Twitter.

Learn more about sharing links on [Elsevier.com](https://www.elsevier.com).

Did you know that, as an author, you can use your article for a wide variety of academic, non-commercial purposes and share and disseminate it online in different ways? For more information, visit [www.elsevier.com/sharing-articles](https://www.elsevier.com/sharing-articles).

Sincerely, Elsevier  
Researcher Support

Increase the impact of your article In the "[Get Noticed](#)" guide you will find a wide variety of tips and suggestions to help you maximize the visibility **of your article**

**Have a question or need help?**

Do not reply to this automated message.

For additional help, visit [the Elsevier Support Center](#), where you can search for solutions to a wide variety of topics and find answers to frequently asked questions.

You can also speak to our investigator support team 24 hours a day, Monday through Friday, and 24/7 via chat or email.

© 2021 Elsevier Ltd | **Privacy Policy** <http://www.elsevier.com/privacypolicy>

Elsevier Limited, The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom, Registered under No. 1982084. This email has been forwarded to you from Elsevier Ltd. To ensure that it arrives correctly in your inbox (instead of junk mail or SPAM folders), add [article\\_status@elsevier.com](mailto:article_status@elsevier.com) to your address book or safe senders list.

---

## We have started the production of your article [ENFCLI\_1924] for Clinical Nursing

2 messages

---

**S.Yadav1@elsevier.com** <S.Yadav1@elsevier.com>  
To: rini.rachmawaty@unhas.ac.id

Fri, Oct 1, 2021 at 8:16 PM

-----  
Our reference: ENFCLI 1924

Article reference: Article title: Abdominal massage for constipation relief in stroke patients: To be published in: Clinical Nursing -----

Dear Dr. Rachmawaty:

Thank you for choosing to post in Clinical Nursing. Please read this message carefully as it contains important information.

**FINALIZE THE PUBLICATION OF YOUR ARTICLE:**

We make every effort to publish the articles of our authors as quickly and efficiently as possible and, therefore, we have already started the process of producing your accepted article. To ensure that your article is published according to your wishes, please complete the forms you will find here:

<http://authors.elsevier.com/authorforms/ENFCLI1924/58345e168f8c9f971d1d643bd61d04fd>

If this link does not work copy the full URL into your browser. Note that the address may be multiple lines in this message. You will need to log in with your Elsevier Profile password, which you had to create at the time you submitted your article.

**PLEASE REVIEW YOUR CONTACT DETAILS:** Please review the following contact details so that we can contact you if necessary: Dr. Rini Rachmawaty

Hasanuddin University

Nursing

Makassar

Indonesia

Phone: not available Fax: not available

Email:

[rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id)

**YOUR REFERENCE NUMBER:**

Finally, so that we can offer you the best service, please take note of the reference number of your article (ENFCLI 1924) and include it in all your messages when you contact us.

Thank you very much for your cooperation. Please do not hesitate to contact us if you have any questions.

Sincerely,

Mr Sachin Sachin

Data Administrator

Elsevier

Email: [S.Yadav1@elsevier.com](mailto:S.Yadav1@elsevier.com)

-----  
**HAVE A QUESTION OR NEED HELP?**

For more help, visit our Customer Support page, where you can search for solutions to a variety of topics and find answers to frequently asked questions (English webpage with a selection of articles in Spanish). You can also contact

our Customer Support team by phone or email (in Spanish: Monday to Friday from 09:00 to 17:00 GMT; in English: Monday to Friday, 24 hours a day) and 24/7 via live chat (English only).

You can start here: <http://service.elsevier.com/app/home/supporthub/publishing>

Copyright © 2018 Elsevier B.V. | Privacy Policy <http://www.elsevier.com/privacypolicy>

Elsevier Limited, The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom, Registration No. 1982084

---

**Rini Rachmawaty** <rini.rachmawaty@unhas.ac.id>

Tue, Jan 4, 2022 at 3:33 PM

To: ilkafah@vokasi.unair.ac.id

**Rini Rachmawaty, PhD, MN, BSN**

*Vice Dean for Academic, Research, & Innovation*

*Faculty of Nursing, Universitas Hasanuddin, Indonesia*

*Cell No. (+62) 812-1391-6730*

*Email Addresses: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id); [RINI.rachmawaty80@gmail.com](mailto:RINI.rachmawaty80@gmail.com)*

----- Forwarded message

----- From: <[S.Yadav1@elsevier.com](mailto:S.Yadav1@elsevier.com)>

Date: Fri, Oct 1, 2021 at 8:16 PM

Subject: We have started production of

your article [ENFCLI\_1924] for Enfermeria Clinica To: <[rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id)>

-----  
Our reference: ENFCLI 1924

Article reference: Article title: Abdominal massage  
for constipation relief  
in stroke

patients: To be published in: Enfermeria Clinica ----- Dear Dr. Rachmawaty:

Thank you for choosing to publish in Enfermeria Clinica. Please read this message carefully as it contains important information. FINALIZE THE PUBLICATION

OF YOUR ARTICLE: We make every effort to publish the articles of our authors as quickly and efficiently as possible and, therefore, we have already started the process of producing your accepted article.

To ensure that your article is published according to your wishes, please fill out the forms you will find here:

<http://authors.elsevier.com/authorforms/ENFCLI1924/58345e168f8c9f971d1d643bd61d04fd>

If this link does not work copy the full URL into your browser. Note that the address may take up multiple lines in this message. You will have to log in with the password of your Elsevier Profile, which you should have created at the time you submitted your article.

PLEASE REVIEW YOUR CONTACT DETAILS: Please review the following contact details so that we can contact you if necessary: Dr. Rini Rachmawaty Hasanuddin University Nursing Makassar Indonesia Phone: not available Fax: not available Email: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id)

YOUR REFERENCE NUMBER:

Finally, so that we can offer you the best service, take note of the reference number of your article (ENFCLI 1924) and include it in all your messages when you contact us.

Thank you very much for your collaboration. Please feel free to contact us if you have any queries.

Sincerely, Mr Sachin Sachin Data Administrator Elsevier Email: [S.Yadav1@elsevier.com](mailto:S.Yadav1@elsevier.com) ----- DO YOU HAVE ANY QUESTIONS OR NEED HELP?

For more help, visit our Customer Support page, where you can search for solutions to a wide variety of topics and find answers to frequently asked questions (English webpage with a selection of articles in Spanish). You can also contact our Customer Support team by phone or email (in Spanish: Monday to Friday from 09:00 to 17:00 GMT; in English: Monday to Friday, 24 hours a day) and 24/7 via live chat (English only).

You can start here: <http://service.elsevier.com/app/home/supporthub/publishing> ----- Copyright © 2018 Elsevier B.V. |

[Quoted text hidden]



---

## Proofs of [ENFCLI\_1924]

3 messages

---

**corrections.eses@elsevier.thomsondigital.com** <corrections.eses@elsevier.thomsondigital.com> Wed, Oct 13, 2021 at 2:51 AM

To: rini.rachmawaty@unhas.ac.id

Cc: p.pau@elsevier.com

P-authorproof-ESES\_Eng\_v14(a)

PLEASE DO NOT ALTER THE SUBJECT LINE OF THIS E-MAIL

Dear Author

The proof of your article, to be published in ENFERMERÍA CLÍNICA, is attached to this e-mail as a PDF file. Also attached are instructions on the annotation of PDF files (notas.pdf). A 'Query Form' is also included as the front page of the proof, detailing any questions regarding your article that have arisen during the preparation of the proof.

We will do everything possible to get your article published quickly and accurately; to do this we need your cooperation. Please respond promptly (48 hours), even if you have no corrections: the sooner we hear from you, the sooner your corrected article will appear online. Please note that any delay in returning your corrections could result in a delay in publication and that any significant changes to the article as accepted for publication will only be considered at this stage with the permission of the Editor.

Please note that proof corrections can now be annotated on-screen, which allows you to mark directly in the PDF file, and return the marked file as an e-mail attachment. See the attached instructions for further information.

Alternative methods of returning proof corrections:

If you do not wish to use the PDF annotations function, you may list the corrections (including replies to the Query Form) in an e-mail and return them to us using the 'reply' button to this e-mail. Please list your corrections quoting line number.

If, for any reason, this is not possible, mark the corrections and any other comments (including replies to the Query Form) on a printout of your proof and fax this to the number given below, or scan the pages and return them by e-mail.

We prefer to receive your corrections by e-mail or fax so that we can process your article quickly and efficiently. However, if you wish to return your corrections by post then please contact us and we will provide the full postal address.

Please use this proof for checking the typesetting, editing, completeness and correctness of the text, tables and figures.

If you submitted usable colour figures with your article they will appear, at no extra charge, in colour on the web if reproduced in colour in the attached PDF proof of your article. In the printed issue, colour reproduction depends on journal policy and whether or not you agree to bear any costs (not applicable to journals which appear only online). Any 'supplementary' material to your article (i.e., not appearing in print) will be accessible after your corrected article is placed online; such material is not part of the proofing procedure and is therefore not attached here.

Before returning your proof corrections, please ensure that you have answered any questions raised on the Query Form and that you have indicated all corrections: this is the last opportunity to make corrections before the final publication of your article.

Kind regards,

Elsevier

E-mail corrections to: [corrections.eses@elsevier.thomsondigital.com](mailto:corrections.eses@elsevier.thomsondigital.com)

Fax: +34 932 091 136

For further assistance, please visit our customer support site at <http://support.elsevier.com>. Here you can search for solutions on a range of topics. You will also find our 24/7 support contact details should you need any further assistance from one of our customer support representatives.

---

**2 attachments**

 **ENFCLI\_1924.pdf**  
258K

 **p-annotatepdf.pdf**  
157K

---

**Rini Rachmawaty** <[rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id)>

Wed, Oct 13, 2021 at 10:16 PM

To: [corrections.eses@elsevier.thomsondigital.com](mailto:corrections.eses@elsevier.thomsondigital.com)

Cc: [p.pau@elsevier.com](mailto:p.pau@elsevier.com)

Bcc: [ilkafah@vokasi.unair.ac.id](mailto:ilkafah@vokasi.unair.ac.id), [syahrulsaid@unhas.ac.id](mailto:syahrulsaid@unhas.ac.id), [yudihardiantomcr@gmail.com](mailto:yudihardiantomcr@gmail.com)

Dear Sir/Madam,

Attached please kindly find our article (ENFCLI\_1924) that has been corrected towards your queries.

Many thanks.

Rini

**Rini Rachmawaty, PhD, MN, BSN**

*Vice Dean for Academic, Research, & Innovation*


*Faculty of Nursing, Universitas Hasanuddin, Indonesia*

*Cell No. (+62) 812-1391-6730*

*Email Addresses: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id); [rini.rachmawaty80@gmail.com](mailto:rini.rachmawaty80@gmail.com)*

[Quoted text hidden]

---

 **ENFCLI\_1924\_Corrected.pdf**  
312K

---

**Rini Rachmawaty** <[rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id)>

Tue, Jan 4, 2022 at 3:33 PM

To: [ilkafah@vokasi.unair.ac.id](mailto:ilkafah@vokasi.unair.ac.id)

**Rini Rachmawaty, PhD, MN, BSN**

*Vice Dean for Academic, Research, & Innovation*

*Faculty of Nursing, Universitas Hasanuddin, Indonesia*

*Cell No. (+62) 812-1391-6730*


*Email Addresses: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id); [rini.rachmawaty80@gmail.com](mailto:rini.rachmawaty80@gmail.com)*

[Quoted text hidden]


---

**2 attachments**

 **ENFCLI\_1924.pdf**  
258K

 **p-annotatepdf.pdf**  
157K

**AUTHOR QUERY FORM**

 <b>ELSEVIER</b>	<b>Journal:</b> ENFCLI	<b>Please e-mail your responses and any corrections to:</b>
	<b>Article Number:</b> 1924	<b>E-mail:</b>

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.


<b>Location in article</b>	<b>Query / Remark: <a href="#">click on the Q link to go</a> Please insert your reply or correction at the corresponding line in the proof</b>
<b>Q1</b>	Please confirm that given name and surname are correctly identified. The different colors indicate whether tagged as first or last name. Please note that proper identification is key for correct indexing of the article.
<b>Q2</b>	Correctly acknowledging the primary funders and grant IDs of your research is important to ensure compliance with funder policies. We could not find any acknowledgement of funding sources in your text. Is this correct?
<b>Q3</b>	Have we correctly interpreted the following funding source(s) and country names you cited in your article: Institute for Research and Community Service [Lembaga Penelitian dan Pengabdian Kepada Masyarakat], Universitas Hasanuddin?
	<div style="border: 1px solid black; padding: 5px; margin-top: 20px;">           Please check this box or indicate your approval if you have no corrections to make to the PDF file           <input style="float: right; margin-left: 20px;" type="checkbox"/> </div>

Thank you for your assistance.



# Enfermería Clínica

[www.elsevier.es/enfermeriaclinica](http://www.elsevier.es/enfermeriaclinica)



## ORIGINAL ARTICLE

### Abdominal massage for constipation relief in stroke patients: A participatory action research

Rini Rachmawaty<sup>a,\*</sup>, Ilkafah Ilkafah<sup>b</sup>, Syahrul Syahrul<sup>c</sup>, Yudi Hardianto<sup>d</sup>

<sup>a</sup> Department of Nursing Management, Faculty of Nursing, Universitas Hasanuddin, Indonesia

<sup>b</sup> Nursing Study Program, Universitas Airlangga, Indonesia

<sup>c</sup> Department of Community Health Nursing, Faculty of Nursing, Universitas Hasanuddin, Indonesia

<sup>d</sup> Department of Physiotherapy, Faculty of Nursing, Universitas Hasanuddin, Indonesia

Received 4 January 2021; accepted 19 July 2021

#### KEYWORDS

Abdominal massage;  
Community  
engagement  
program;  
Community health  
volunteers;  
Constipation;  
Stroke

#### Abstract

**Objective:** To improve knowledge and skills of family members and community health volunteers in applying abdominal massage to stroke patients with constipation.

**Method:** A participatory action research, involving family members and community health volunteers to participate in the training and supervision sessions about abdominal massage for constipation relief in stroke patients. Knowledge enrichment was assessed using questionnaire, while the abdominal massage skill was evaluated using direct observation. Qualitative interviews were conducted to evaluate patients' experiences regarding the intervention program.

**Results:** The program showed the improvement of participants' knowledge and skills in applying abdominal massage for stroke patients with constipation. Likewise, patients confirmed their constipation relieved after the application of abdominal massage.

**Conclusion:** This study showed that training and supervision sessions about abdominal massage for constipation alleviation in stroke patients are feasible, efficient, and beneficial programs to enhance knowledge and skill of family members and community health volunteers.

© 2021 Elsevier España, S.L.U. All rights reserved.

Peer-review process of the 3rd International Nursing & Health Sciences Students & Health Care Professionals Conference (INHSP) 2019 is under the responsibility of the scientific committee of the 3rd International Nursing & Health Sciences Students & Health Care Professionals Conference (INHSP). Full-text and the content of it is under the responsibility of authors of the article.

\* Corresponding author.

E-mail address: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id) (R. Rachmawaty).

<https://doi.org/10.1016/j.enfcli.2021.07.035>

1130-8621/© 2021 Elsevier España, S.L.U. All rights reserved.

Please cite this article as: R. Rachmawaty, I. Ilkafah, S. Syahrul et al., Abdominal massage for constipation relief in stroke patients: A participatory action research, *Enfermería Clínica*, <https://doi.org/10.1016/j.enfcli.2021.07.035>

## Introduction

Stroke is one of the common causes of death and disability in Indonesia.<sup>1</sup> It is reported that stroke prevalence increased from 7% in 2013 to 10.9% in 2018, meaning that the prevalence of people with stroke in Indonesia was ten per one thousand population.<sup>2,3</sup> In 2016, number of Indonesians died due to stroke was 212,963 and the highest prevalence was reported in the South Sulawesi province (17.9%).<sup>4,5</sup>

Besides disability, one of the most common complications among people with stroke is constipation which can lead to increased disability and death.<sup>6-8</sup> Constipation in stroke patients occurred mainly due to lack of physical mobility, difficulty to swallowing fluid and fiber, dependence on others to use the toilet, and the use of laxatives.<sup>8</sup> If the stroke patient with constipation has a history of heart failure, hypertension, or myocardial infarction; straining during defecation can lead to increased burden of the heart, rupture of blood vessels, and premature death.<sup>9-11</sup> Concerning these complications, it is crucial to prevent constipation.<sup>6,9</sup>

Abdominal massage is revealed as one of several interventions to prevent and to treat constipation among elderly, people with stroke, and those with musculoskeletal disorders, cardiovascular diseases, chronic blood disease, and post-operative ileus.<sup>6,12-15</sup> Abdominal massage has been proved as an inexpensive and safe non-pharmacologic therapy to relieve constipation.<sup>16</sup>

Likewise, a previous study has confirmed that applying abdominal massage can prevent constipation in stroke patients hospitalized in one of Indonesian public hospital and can avoid the patients taking laxatives.<sup>6,9</sup> However, studies specifically focusing on how abdominal massage performed by family members and community health volunteers to prevent constipation in stroke patients at home is rare. Hence, this study aimed to implement abdominal massage performed by family members and community health volunteers to alleviate constipation in stroke patients at their homes.

## Method

This study was participatory action research (PAR) using mix-methods data collection including interviews, FGDs, and questionnaires. The participants consisted of 11 family members of stroke patients and five community health volunteers. The intervention including training and supervision participants about the application of abdominal massage to stroke patients as well as conducting evaluation of the process and outcomes. The training was delivered in the forms of modules, audiovisual media, skill demonstration. Participants performed abdominal massage to stroke patients under the supervision of the research team. Home visits were conducted to evaluate knowledge and skills of participants in applying abdominal massage independently to stroke patients.

Regarding the ethical considerations, this study was a follow up study from the initial intervention study that was conducted at one of Indonesian public hospitals and had obtained ethical approval from the Research Ethics Committee of Faculty of Medicine, Universitas Hasanuddin (No. 513/H4.8.4.5.31/PP36-KOMETIK/2017). Furthermore, prior

to data collection, the researchers reported to the Ministry of Health of Makassar City to obtain permission to conduct research at the Antara Public Health Center. Finally, informed consent was obtained from all participants prior to their participation in this study.

In conducting the training session, the principal investigator (PI) initially explained the purpose of the activity, then followed by a pre-test to assess the knowledge of family and community health volunteers about constipation and abdominal massage. After that, the research team provided health education to patients' families, community health volunteers, and the staff of public health center about stroke, constipation, and abdominal massage for constipation relief using educational videos. Next, one of the co-PIs re-demonstrated the steps of abdominal massage directly to stroke patients. After the training conducted, families and community health volunteers were required to complete the post-test to evaluate the level of knowledge of family and community health volunteers about constipation and abdominal massage.

The next activity was home visit, through which the research team and the community health volunteer visited the patients and evaluated the ability of the family members and the community health volunteers to apply abdominal massage for stroke patients. During this visit, the researcher team observed and checked using questionnaire whether the participants correctly performed the steps of abdominal massage. Participants could repeat the procedures until they re-demonstrated it properly. At the end, patients were asked about their condition after receiving abdominal massage. The last activity of this PAR was inviting all participants to the final meeting at the public health center to evaluate the overall program using focused-group discussion.

Regarding data analysis, knowledge changes was analyzed using paired t-test, while participants knowledge about constipation and abdominal massage was analyzed using repeated measures ANCOVA. The patients' experiences after receiving abdominal massage were analyzed using qualitative analysis.

## Result

The demographic characteristics of participants in the counseling and training sessions is shown in Table 1. The average age of the stroke patients' families was 36.8 and the community health volunteers was 46.8. Participants were predominated by women and most of them were high school graduates. Two third of the participants were housewives and the rest were teachers and entrepreneurs.

Table 2 shows that after participating in the training and supervising sessions, patients' families gained better knowledge about constipation ( $p=0.001$ ) and abdominal massage ( $p=0.000$ ), while the community health volunteers attained better knowledge only for abdominal massage ( $p=0.023$ ).

Table 3 depicts the ability of participants to perform abdominal massage. As clearly seen in Table 3, not all families and community health volunteers can provide correct and complete abdominal massage. The participants complained that they had difficulty in memorizing abdominal massage procedures due to age, lack of practice, lack of

**Table 1** Demographic characteristics of the participants in the training and supervision sessions (n = 16).

Characteristics	Family (n = 11)		Community health volunteer (n = 5)	
	Mean (SD)	n (%)	Mean (SD)	n (%)
Age	36.82 (8.32)		46.8 (11.78)	
Gender				
Female		10 (90.9)		5 (100.0)
Male		1 (9.1)		
Education				
Junior high school		2 (18.2)		
Senior high school		6 (54.5)		5 (100.0)
Diploma III		1 (9.1)		
Bachelor		1 (9.1)		
Master		1 (9.1)		
Occupation				
Housewife		5 (45.5)		3 (60.0)
Private employee		2 (18.2)		
Entrepreneur		2 (18.2)		1 (20.0)
Lecturer		1 (9.1)		
Teacher		1 (9.1)		
Retired				1 (20.0)

SD, standard deviation.

**Table 2** Participants' knowledge of constipation and abdominal massage application on stroke patients before and after counseling (n = 16).

Knowledge	Family (n = 11)			Community health volunteer (n = 5)		
	Pre-test Mean (SD)	Post-test Mean (SD)	p <sup>a</sup>	Pre-test Mean (SD)	Post-test Mean (SD)	p <sup>a</sup>
Constipation	6.55 (1.75)	8.36 (0.92)	0.001*	6.60 (2.07)	8.00 (0.71)	0.113
Abdominal massage	6.18 (1.25)	9.00 (0.89)	0.000*	5.40 (3.21)	8.60 (1.14)	0.023

SD, standard deviation

<sup>a</sup> Using paired *t*-test.

\* *p* < 0.05.

**Table 3** The evaluation of participants in applying abdominal massage on stroke patients (n = 12).

Ability to perform abdominal massage	Family (n = 7)	Community health volunteer (n = 5)
	Well performed, n (%)	
Step 1: Stroke upwards 3 times	7 (100)	5 (100)
Step 2: Stroke towards the bottom of the abdomen 3 times	7 (100)	4 (80)
Step 3: Effleurage or circular stroking	4 (57.1)	4 (80)
Step 4: Palmer kneading	7 (100)	5 (100)
Step 5: As Step 4, but moving up the stomach	6 (85.7)	4 (80)
Step 6: Repeat Steps 4 and 5	4 (57.1)	3 (60)
Step 7: Stroking	7 (100)	5 (100)
Step 8: Hand vibrations over the umbilical area	7 (100)	5 (100)
Step 1-8	2 (28.6)	3 (60)

144 reading the provided modules, and rarely watching provided  
145 simulation videos and YouTube. The participants reported  
146 that the abdominal massage was only applied to patients  
147 when they complained about the onset of constipation.

Table 4 demonstrates the enhancement of participants' knowledge and skills in applying abdominal massage for stroke patients with constipation. As depicted by Table 4, the family members' knowledge about constipation

148  
149  
150  
151

**Table 4** Participants' knowledge of constipation and abdominal massage application on stroke patients before training, after training, and after program evaluation ( $n=9$ ).

Knowledge	Family ( $n=4$ )				Community health volunteer ( $n=5$ )			
	Pre-test Mean (SD)	Post-test 1 Mean (SD)	Post-test 2 Mean (SD)	$p^a$	Pre-test Mean (SD)	Post-test 1 Mean (SD)	Post-test 2 Mean (SD)	$p^a$
Constipation	5.37 (0.91)	7.73 (0.36)	9.09 (0.37)	0.011*	6.70 (0.80)	8.02 (0.32)	9.53 (0.33)	0.011*
Abdominal massage	5.38 (1.39)	9.20 (0.53)	9.19 (0.36)	0.003*	5.30 (1.23)	8.44 (0.47)	8.85 (0.32)	0.003*

Pre-test: before training session, Post-test 1: after training session, Post-test 2: after program evaluation. SD, standard deviation.

<sup>a</sup> Using repeated measures ANCOVA.

\*  $p < 0.05$ .

( $p=0.011$ ) and abdominal massage ( $p=0.003$ ) significantly rose from pre-test to post-test 2. Likewise, the knowledge of community health volunteers about constipation ( $p=0.011$ ) and abdominal massage ( $p=0.003$ ) significantly improved overtime.

Based on the qualitative interviews, patients confirmed that constipation was relieved after receiving abdominal massage for three times. They also said that having abdominal massage induced them to defecate.

## Discussion

This study is among the first studies that explored the effect of abdominal massage on constipation relief in Indonesian stroke population living in the community, and among the few studies of abdominal massage conducted in stroke population. The previous study has shown that abdominal massage is feasible and effective to prevent constipation in people with neurological condition.<sup>17</sup> Through that study, the physiotherapists taught the patients and their carer about abdominal massage and allowed them to practice. Information about abdominal massage was also provided in the CDs. Like current study, the previous study found that abdominal massage was effective to alleviate constipation.<sup>18</sup>

This study is also the first study in Indonesia to involve community health volunteers to apply abdominal massage to people with stroke. Community health volunteers are quite common in Indonesia and many developing countries, especially in villages and rural areas where health professionals are scarce.<sup>19</sup> They are valuable workforce in primary and community health care that assist health professionals to provide optimal care for patients.<sup>19</sup> They are involved in many aspects of health promotion, prevention, intervention, rehabilitation, and palliative care.<sup>20-22</sup> They have been working hand in hand with health professionals in providing community care for communicable diseases such as malaria,<sup>20</sup> tuberculosis,<sup>21</sup> and HIV/AIDS.<sup>22</sup> They also provided support for new mothers in maternal and antenatal care by working under supervision of community midwives.<sup>23</sup> Involving community health volunteers in providing abdominal massage for stroke patients with constipation is an efficient way to help people with stroke manage their constipation. Patients, family members and community health professionals has been familiar with them and thus, collaboration can be ensured.

There are some limitations of this study: No control group and small sample size. During home visit, some patients were not received abdominal massage due to contraindication. Even so, this study showed that the application of abdominal massage to stroke patients with constipation is feasible by providing training and supervision sessions for family members and community health volunteers. It is considered more economically preferred as the application of abdominal massage does not require any expensive tools. Furthermore, as it is performed by family members, the adherence to the abdominal massage application is guaranteed as it only requires several minutes to apply to the patient.

## Conclusion

Participants experienced improved knowledge and skills of constipation and abdominal massage after attending training and supervision sessions. Likewise, patients confirmed ease of defecation after receiving abdominal massage. This implementation study proves that abdominal massage performed by family and health community volunteers was advantageous and effective in alleviating constipation in stroke patients. Further study is required with larger sample size, multicenter, and more rigorous methods to explore more about the effect of abdominal massage on constipation alleviation in people with stroke.

## Conflict of interest

The authors declare no financial or personal interests that could bias the work.

## Acknowledgement

This study supported by grant from the Institute for Research and Community Service [Lembaga Penelitian dan Pengabdian Kepada Masyarakat], Universitas Hasanuddin, Makassar, Indonesia. The authors would like to thank to the Head of Antara Public Health Center, drg. Weri LP. Saroengoe, and all participants involved in this study.


## References

1. World Health Organization. State of Health Inequality: Indonesia. Geneva: World Health Organization; 2017.

- 234 2. Kemenkes RI. Peningkatan Gaya Hidup Sehat dengan perilaku.  
235 CERDIK; 2018. p. 10.
- 236 3. Kementerian Kesehatan Republik Indonesia. Riskesdas 2018.  
237 Jakarta; 2018.
- 238 4. Johnson CO, Nguyen M, Roth GA, Nichols E, Alam T,  
239 Abate D, et al. Global, regional, and national burden of  
240 stroke, 1990–2016: a systematic analysis for the Global Bur-  
241 den of Disease Study 2016. *Lancet Neurol.* 2019;18:439–58,  
242 [http://dx.doi.org/10.1016/S1474-4422\(19\)30034-1](http://dx.doi.org/10.1016/S1474-4422(19)30034-1).
- 243 5. Indonesia KKR. Riset Kesehatan Dasar 2013. Jakarta; 2013.
- 244 6. Sinclair M. The use of abdominal massage to treat  
245 chronic constipation. *J Bodyw Mov Ther.* 2011;15:436–45,  
246 <http://dx.doi.org/10.1016/j.jbmt.2010.07.007>.
- 247 7. Ronald L, Hickman FAAN, Celeste M, Alfes JF. Handbook of  
248 clinical nursing: medical–surgical nursing. New York: Springer  
249 Publishing Company; 2018.
- 250 8. Li J, Yuan M, Liu Y, Zhao Y, Wang J, Guo W, et al.  
251 Incidence of constipation in stroke patients: a systematic  
252 review and meta-analysis. *Med (United States).* 2017;96:1–6,  
253 <http://dx.doi.org/10.1097/MD.00000000000007225>.
- 254 9. Yacoline Pailungan F, Kaelan C, Rachmawaty R. Pengaruh Pem-  
255 berian Massage Abdomen Terhadap Penurunan Konstipasi Pada  
256 Pasien Stroke Iskemik Di Rsup Dr. Wahidin Sudirohusodo Makas-  
257 sar. *Patria Artha J Nurs Sci.* 2017;1:25–35.
- 258 10. de Engler TMNM, Dourado CC, Amâncio TG, Farage L, de  
259 Mello PA, Padula MP. Stroke: bowel dysfunction in patients  
260 admitted for rehabilitation. *Open Nurs J.* 2014;8:43–7,  
261 <http://dx.doi.org/10.2174/1874434601408010043>.
- 262 11. Sumida K, Molnar MZ, Potukuchi PK, Thomas F, Lu JL,  
263 Yamagata K, et al. Constipation and risk of death and car-  
264 diovascular events. *Atherosclerosis.* 2019;281(May):114–20,  
265 <http://dx.doi.org/10.1016/j.atherosclerosis.2018.12.021>.
- 266 12. Lämås K, Graneheim UH, Jacobsson C. Experiences of abdom-  
267 inal massage for constipation. *J Clin Nurs.* 2012;21:757–65,  
268 <http://dx.doi.org/10.1111/j.1365-2702.2011.03946.x>.
- 269 13. Deckers B, Von Schoen-Angerer T, Voggenreiter B, Vagedes J.  
270 External nursing applications in the supportive management  
271 of prolonged postoperative ileus description of interven-  
272 tions and case report. *Holist Nurs Pract.* 2016;30:216–21,  
273 <http://dx.doi.org/10.1097/HNP.000000000000158>.
- 274 14. Çevik K, Çetinkaya A, Yiğit Gökbel K, Menekşe B, Saza S, Tikiz C.  
The effect of abdominal massage on constipation in the elderly  
residing in rest homes. *Gastroenterol Nurs.* 2018;41:396–402,  
<http://dx.doi.org/10.1097/SGA.0000000000000343>.
- 275 15. Birimoglu Okuyan C, Bilgili N. Effect of abdominal massage  
276 on constipation and quality of life in older adults: a random-  
277 ized controlled trial. *Complement Ther Med.* 2019;47:102219,  
278 <http://dx.doi.org/10.1016/j.ctim.2019.102219>.
- 279 16. Emly MC. Abdominal massage for constipation. In: Haslam JLJ,  
280 editor. Therapeutic management of incontinence and pelvic  
281 pain. London: Springer; 2008. p. 223–5.
- 282 17. McClurg D, Hagen S, Hawkins S, Lowe-Strong A. Abdomi-  
283 nal massage for the alleviation of constipation symptoms  
284 in people with multiple sclerosis: a randomized con-  
285 trolled feasibility study. *Mult Scler J.* 2010;17:223–33,  
286 <http://dx.doi.org/10.1177/1352458510384899>.
- 287 18. McClurg D, Hagen S, Hawkins S, Lowe-Strong A. Abdomi-  
288 nal massage for the alleviation of constipation symptoms  
289 in people with multiple sclerosis: a randomized con-  
290 trolled feasibility study. *Mult Scler J.* 2011;17:223–33,  
291 <http://dx.doi.org/10.1177/1352458510384899>.
- 292 19. Bhattacharyya K, Winch P, LeBan K, Tien M. Community health  
293 worker incentives and disincentives: how they affect motiva-  
294 tion, retention, and sustainability, Arlington: the Basic Support  
295 for Institutionalizing Child Survival Project (BASICS II) for the  
296 United States Agency for International Development; 2001.
- 297 20. Soesilo D, Wahyuni CU. Improving the village health volunteers’  
298 knowledge, malaria diagnostic skill and home visits through  
299 the observed training (study in Municipality of Tidore Islands  
300 North Maluku Province Indonesia). *Indian J Public Heal Res Dev.*  
301 2019;10:1671–5.
- 302 21. Lukman M, MSi KI, Yani DI, Juniarti N. Exploring strategies to  
303 improve the performance of community health volunteers for  
304 tuberculosis care and prevention: a qualitative study. *Int J*  
305 *Commun Based Nurs Midwif.* 2019;7:270.
- 306 22. Palaia AM. Understanding the motivation, ability and attrition  
307 of HIV/AIDS community health volunteers in resource poor set-  
308 tings: Implications for program sustainability. Johns Hopkins  
309 University; 2010.
- 310 23. Tumbelaka P, Limato R, Nasir S, Syafruddin D, Ormel H, Ahmed  
311 R. Analysis of Indonesia’s community health volunteers (kader)  
312 as maternal health promoters in the community integrated  
313 health service (Posyandu) following health promotion training.  
314 *Int J Community Med Public Heal.* 2018;5:127–35.
- 315 316



**AUTHOR QUERY FORM**

 <b>ELSEVIER</b>	<b>Journal:</b> ENFCLI	<b>Please e-mail your responses and any corrections to:</b>
	<b>Article Number:</b> 1924	<b>E-mail:</b>

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.


<b>Location in article</b>	<b>Query / Remark: <a href="#">click on the Q link to go</a> Please insert your reply or correction at the corresponding line in the proof</b>
<a href="#">Q1</a>	Please confirm that given name and surname are correctly identified. The different colors indicate whether tagged as first or last name. Please note that proper identification is key for correct indexing of the article. <b>Given name and surname of all authors are correct.</b>
<a href="#">Q2</a>	Correctly acknowledging the primary funders and grant IDs of your research is important to ensure compliance with funder policies. We could not find any acknowledgement of funding sources in your text. Is this correct? <b>No. The primary funder &amp; grant ID have been stated in the Acknowledgment section in line 224.</b>
<a href="#">Q3</a>	Have we correctly interpreted the following funding source(s) and country names you cited in your article: Institute for Research and Community Service [Lembaga Penelitian dan Pengabdian Kepada Masyarakat], Universitas Hasanuddin? <b>Yes, it is correct for the funding source, but the country name is Indonesia. And it has been completely stated in the Acknowledgment section in line 228.</b>
<a href="#">Line 7</a>	<b>There is an additional information for the second author's affiliation.</b>
	<div style="border: 1px solid black; padding: 10px; width: fit-content; margin: 0 auto;"> <p>Please check this box or indicate your approval if you have no corrections to make to the PDF file <input type="checkbox"/></p> </div>

Thank you for your assistance.



# Enfermería Clínica

[www.elsevier.es/enfermeriaclinica](http://www.elsevier.es/enfermeriaclinica)



## ORIGINAL ARTICLE

### Abdominal massage for constipation relief in stroke patients: A participatory action research

Q1 Rini Rachmawaty<sup>a,\*</sup>, Ilkafah Ilkafah<sup>b</sup>, Syahrul Syahrul<sup>c</sup>, Yudi Hardianto<sup>d</sup>

<sup>a</sup> Department of Nursing Management, Faculty of Nursing, Universitas Hasanuddin, Indonesia

<sup>b</sup> Nursing Study Program, Universitas Airlangga, Indonesia

<sup>c</sup> Department of Community Health Nursing, Faculty of Nursing, Universitas Hasanuddin, Indonesia

<sup>d</sup> Department of Physiotherapy, Faculty of Nursing, Universitas Hasanuddin, Indonesia

Received 4 January 2021; accepted 19 July 2021

#### KEYWORDS

Abdominal massage;  
Community engagement program;  
Community health volunteers;  
Constipation;  
Stroke

#### Abstract

**Objective:** To improve knowledge and skills of family members and community health volunteers in applying abdominal massage to stroke patients with constipation.

**Method:** A participatory action research, involving family members and community health volunteers to participate in the training and supervision sessions about abdominal massage for constipation relief in stroke patients. Knowledge enrichment was assessed using questionnaire, while the abdominal massage skill was evaluated using direct observation. Qualitative interviews were conducted to evaluate patients' experiences regarding the intervention program.

**Results:** The program showed the improvement of participants' knowledge and skills in applying abdominal massage for stroke patients with constipation. Likewise, patients confirmed their constipation relieved after the application of abdominal massage.

**Conclusion:** This study showed that training and supervision sessions about abdominal massage for constipation alleviation in stroke patients are feasible, efficient, and beneficial programs to enhance knowledge and skill of family members and community health volunteers.

© 2021 Elsevier España, S.L.U. All rights reserved.

Peer-review process of the 3rd International Nursing & Health Sciences Students & Health Care Professionals Conference (INHSP) 2019 is under the responsibility of the scientific committee of the 3rd International Nursing & Health Sciences Students & Health Care Professionals Conference (INHSP). Full-text and the content of it is under the responsibility of authors of the article.

\* Corresponding author.

E-mail address: [rini.rachmawaty@unhas.ac.id](mailto:rini.rachmawaty@unhas.ac.id) (R. Rachmawaty).

<https://doi.org/10.1016/j.enfcli.2021.07.035>

1130-8621/© 2021 Elsevier España, S.L.U. All rights reserved.

Please cite this article as: R. Rachmawaty, I. Ilkafah, S. Syahrul et al., Abdominal massage for constipation relief in stroke patients: A participatory action research, *Enfermería Clínica*, <https://doi.org/10.1016/j.enfcli.2021.07.035>

## Introduction

Stroke is one of the common causes of death and disability in Indonesia.<sup>1</sup> It is reported that stroke prevalence increased from 7% in 2013 to 10.9% in 2018, meaning that the prevalence of people with stroke in Indonesia was ten per one thousand population.<sup>2,3</sup> In 2016, number of Indonesians died due to stroke was 212,963 and the highest prevalence was reported in the South Sulawesi province (17.9%).<sup>4,5</sup>

Besides disability, one of the most common complications among people with stroke is constipation which can lead to increased disability and death.<sup>6-8</sup> Constipation in stroke patients occurred mainly due to lack of physical mobility, difficulty to swallowing fluid and fiber, dependence on others to use the toilet, and the use of laxatives.<sup>8</sup> If the stroke patient with constipation has a history of heart failure, hypertension, or myocardial infarction; straining during defecation can lead to increased burden of the heart, rupture of blood vessels, and premature death.<sup>9-11</sup> Concerning these complications, it is crucial to prevent constipation.<sup>6,9</sup>

Abdominal massage is revealed as one of several interventions to prevent and to treat constipation among elderly, people with stroke, and those with musculoskeletal disorders, cardiovascular diseases, chronic blood disease, and post-operative ileus.<sup>6,12-15</sup> Abdominal massage has been proved as an inexpensive and safe non-pharmacologic therapy to relieve constipation.<sup>16</sup>

Likewise, a previous study has confirmed that applying abdominal massage can prevent constipation in stroke patients hospitalized in one of Indonesian public hospital and can avoid the patients taking laxatives.<sup>6,9</sup> However, studies specifically focusing on how abdominal massage performed by family members and community health volunteers to prevent constipation in stroke patients at home is rare. Hence, this study aimed to implement abdominal massage performed by family members and community health volunteers to alleviate constipation in stroke patients at their homes.

## Method

This study was participatory action research (PAR) using mix-methods data collection including interviews, FGDs, and questionnaires. The participants consisted of 11 family members of stroke patients and five community health volunteers. The intervention including training and supervision participants about the application of abdominal massage to stroke patients as well as conducting evaluation of the process and outcomes. The training was delivered in the forms of modules, audiovisual media, skill demonstration. Participants performed abdominal massage to stroke patients under the supervision of the research team. Home visits were conducted to evaluate knowledge and skills of participants in applying abdominal massage independently to stroke patients.

Regarding the ethical considerations, this study was a follow up study from the initial intervention study that was conducted at one of Indonesian public hospitals and had obtained ethical approval from the Research Ethics Committee of Faculty of Medicine, Universitas Hasanuddin (No. 513/H4.8.4.5.31/PP36-KOMETIK/2017). Furthermore, prior

to data collection, the researchers reported to the Ministry of Health of Makassar City to obtain permission to conduct research at the Antara Public Health Center. Finally, informed consent was obtained from all participants prior to their participation in this study.

In conducting the training session, the principal investigator (PI) initially explained the purpose of the activity, then followed by a pre-test to assess the knowledge of family and community health volunteers about constipation and abdominal massage. After that, the research team provided health education to patients' families, community health volunteers, and the staff of public health center about stroke, constipation, and abdominal massage for constipation relief using educational videos. Next, one of the co-PIs re-demonstrated the steps of abdominal massage directly to stroke patients. After the training conducted, families and community health volunteers were required to complete the post-test to evaluate the level of knowledge of family and community health volunteers about constipation and abdominal massage.

The next activity was home visit, through which the research team and the community health volunteer visited the patients and evaluated the ability of the family members and the community health volunteers to apply abdominal massage for stroke patients. During this visit, the researcher team observed and checked using questionnaire whether the participants correctly performed the steps of abdominal massage. Participants could repeat the procedures until they re-demonstrated it properly. At the end, patients were asked about their condition after receiving abdominal massage. The last activity of this PAR was inviting all participants to the final meeting at the public health center to evaluate the overall program using focused-group discussion.

Regarding data analysis, knowledge changes was analyzed using paired t-test, while participants knowledge about constipation and abdominal massage was analyzed using repeated measures ANCOVA. The patients' experiences after receiving abdominal massage were analyzed using qualitative analysis.

## Result

The demographic characteristics of participants in the counseling and training sessions is shown in Table 1. The average age of the stroke patients' families was 36.8 and the community health volunteers was 46.8. Participants were predominated by women and most of them were high school graduates. Two third of the participants were housewives and the rest were teachers and entrepreneurs.

Table 2 shows that after participating in the training and supervising sessions, patients' families gained better knowledge about constipation ( $p=0.001$ ) and abdominal massage ( $p=0.000$ ), while the community health volunteers attained better knowledge only for abdominal massage ( $p=0.023$ ).

Table 3 depicts the ability of participants to perform abdominal massage. As clearly seen in Table 3, not all families and community health volunteers can provide correct and complete abdominal massage. The participants complained that they had difficulty in memorizing abdominal massage procedures due to age, lack of practice, lack of

**Table 1** Demographic characteristics of the participants in the training and supervision sessions (n = 16).

Characteristics	Family (n = 11)		Community health volunteer (n = 5)	
	Mean (SD)	n (%)	Mean (SD)	n (%)
Age	36.82 (8.32)		46.8 (11.78)	
Gender				
Female		10 (90.9)		5 (100.0)
Male		1 (9.1)		
Education				
Junior high school		2 (18.2)		
Senior high school		6 (54.5)		5 (100.0)
Diploma III		1 (9.1)		
Bachelor		1 (9.1)		
Master		1 (9.1)		
Occupation				
Housewife		5 (45.5)		3 (60.0)
Private employee		2 (18.2)		
Entrepreneur		2 (18.2)		1 (20.0)
Lecturer		1 (9.1)		
Teacher		1 (9.1)		
Retired				1 (20.0)

SD, standard deviation.

**Table 2** Participants' knowledge of constipation and abdominal massage application on stroke patients before and after counseling (n = 16).

Knowledge	Family (n = 11)			Community health volunteer (n = 5)		
	Pre-test Mean (SD)	Post-test Mean (SD)	p <sup>a</sup>	Pre-test Mean (SD)	Post-test Mean (SD)	p <sup>a</sup>
Constipation	6.55 (1.75)	8.36 (0.92)	0.001*	6.60 (2.07)	8.00 (0.71)	0.113
Abdominal massage	6.18 (1.25)	9.00 (0.89)	0.000*	5.40 (3.21)	8.60 (1.14)	0.023

SD, standard deviation

<sup>a</sup> Using paired *t*-test.

\* *p* < 0.05.

**Table 3** The evaluation of participants in applying abdominal massage on stroke patients (n = 12).

Ability to perform abdominal massage	Family (n = 7)	Community health volunteer (n = 5)
	Well performed, n (%)	
Step 1: Stroke upwards 3 times	7 (100)	5 (100)
Step 2: Stroke towards the bottom of the abdomen 3 times	7 (100)	4 (80)
Step 3: Effleurage or circular stroking	4 (57.1)	4 (80)
Step 4: Palmer kneading	7 (100)	5 (100)
Step 5: As Step 4, but moving up the stomach	6 (85.7)	4 (80)
Step 6: Repeat Steps 4 and 5	4 (57.1)	3 (60)
Step 7: Stroking	7 (100)	5 (100)
Step 8: Hand vibrations over the umbilical area	7 (100)	5 (100)
Step 1-8	2 (28.6)	3 (60)

144 reading the provided modules, and rarely watching provided  
145 simulation videos and YouTube. The participants reported  
146 that the abdominal massage was only applied to patients  
147 when they complained about the onset of constipation.

Table 4 demonstrates the enhancement of participants' knowledge and skills in applying abdominal massage for stroke patients with constipation. As depicted by Table 4, the family members' knowledge about constipation

148  
149  
150  
151

**Table 4** Participants' knowledge of constipation and abdominal massage application on stroke patients before training, after training, and after program evaluation ( $n=9$ ).

Knowledge	Family ( $n=4$ )				Community health volunteer ( $n=5$ )			
	Pre-test Mean (SD)	Post-test 1 Mean (SD)	Post-test 2 Mean (SD)	$p^a$	Pre-test Mean (SD)	Post-test 1 Mean (SD)	Post-test 2 Mean (SD)	$p^a$
Constipation	5.37 (0.91)	7.73 (0.36)	9.09 (0.37)	0.011*	6.70 (0.80)	8.02 (0.32)	9.53 (0.33)	0.011*
Abdominal massage	5.38 (1.39)	9.20 (0.53)	9.19 (0.36)	0.003*	5.30 (1.23)	8.44 (0.47)	8.85 (0.32)	0.003*

Pre-test: before training session, Post-test 1: after training session, Post-test 2: after program evaluation. SD, standard deviation.

<sup>a</sup> Using repeated measures ANCOVA.

\*  $p < 0.05$ .

( $p=0.011$ ) and abdominal massage ( $p=0.003$ ) significantly rose from pre-test to post-test 2. Likewise, the knowledge of community health volunteers about constipation ( $p=0.011$ ) and abdominal massage ( $p=0.003$ ) significantly improved overtime.

Based on the qualitative interviews, patients confirmed that constipation was relieved after receiving abdominal massage for three times. They also said that having abdominal massage induced them to defecate.

## Discussion

This study is among the first studies that explored the effect of abdominal massage on constipation relief in Indonesian stroke population living in the community, and among the few studies of abdominal massage conducted in stroke population. The previous study has shown that abdominal massage is feasible and effective to prevent constipation in people with neurological condition.<sup>17</sup> Through that study, the physiotherapists taught the patients and their carer about abdominal massage and allowed them to practice. Information about abdominal massage was also provided in the CDs. Like current study, the previous study found that abdominal massage was effective to alleviate constipation.<sup>18</sup>

This study is also the first study in Indonesia to involve community health volunteers to apply abdominal massage to people with stroke. Community health volunteers are quite common in Indonesia and many developing countries, especially in villages and rural areas where health professionals are scarce.<sup>19</sup> They are valuable workforce in primary and community health care that assist health professionals to provide optimal care for patients.<sup>19</sup> They are involved in many aspects of health promotion, prevention, intervention, rehabilitation, and palliative care.<sup>20-22</sup> They have been working hand in hand with health professionals in providing community care for communicable diseases such as malaria,<sup>20</sup> tuberculosis,<sup>21</sup> and HIV/AIDS.<sup>22</sup> They also provided support for new mothers in maternal and antenatal care by working under supervision of community midwives.<sup>23</sup> Involving community health volunteers in providing abdominal massage for stroke patients with constipation is an efficient way to help people with stroke manage their constipation. Patients, family members and community health professionals has been familiar with them and thus, collaboration can be ensured.

There are some limitations of this study: No control group and small sample size. During home visit, some patients were not received abdominal massage due to contraindication. Even so, this study showed that the application of abdominal massage to stroke patients with constipation is feasible by providing training and supervision sessions for family members and community health volunteers. It is considered more economically preferred as the application of abdominal massage does not require any expensive tools. Furthermore, as it is performed by family members, the adherence to the abdominal massage application is guaranteed as it only requires several minutes to apply to the patient.

## Conclusion

Participants experienced improved knowledge and skills of constipation and abdominal massage after attending training and supervision sessions. Likewise, patients confirmed ease of defecation after receiving abdominal massage. This implementation study proves that abdominal massage performed by family and health community volunteers was advantageous and effective in alleviating constipation in stroke patients. Further study is required with larger sample size, multicenter, and more rigorous methods to explore more about the effect of abdominal massage on constipation alleviation in people with stroke.

## Conflict of interest

The authors declare no financial or personal interests that could bias the work.

## Acknowledgement

This study supported by grant from the Institute for Research and Community Service [Lembaga Penelitian dan Pengabdian Kepada Masyarakat], Universitas Hasanuddin, Makassar, Indonesia. The authors would like to thank to the Head of Antara Public Health Center, drg. Weri LP. Saroengoe, and all participants involved in this study.

## References

1. World Health Organization. State of Health Inequality: Indonesia. Geneva: World Health Organization; 2017.

- 234 2. Kemenkes RI. Peningkatan Gaya Hidup Sehat dengan perilaku.  
235 CERDIK; 2018. p. 10.
- 236 3. Kementerian Kesehatan Republik Indonesia. Riskesdas 2018.  
237 Jakarta; 2018.
- 238 4. Johnson CO, Nguyen M, Roth GA, Nichols E, Alam T,  
239 Abate D, et al. Global, regional, and national burden of  
240 stroke, 1990–2016: a systematic analysis for the Global Bur-  
241 den of Disease Study 2016. *Lancet Neurol.* 2019;18:439–58,  
242 [http://dx.doi.org/10.1016/S1474-4422\(19\)30034-1](http://dx.doi.org/10.1016/S1474-4422(19)30034-1).
- 243 5. Indonesia KKR. Riset Kesehatan Dasar 2013. Jakarta; 2013.
- 244 6. Sinclair M. The use of abdominal massage to treat  
245 chronic constipation. *J Bodyw Mov Ther.* 2011;15:436–45,  
246 <http://dx.doi.org/10.1016/j.jbmt.2010.07.007>.
- 247 7. Ronald L, Hickman FAAN, Celeste M, Alfes JF. Handbook of  
248 clinical nursing: medical–surgical nursing. New York: Springer  
249 Publishing Company; 2018.
- 250 8. Li J, Yuan M, Liu Y, Zhao Y, Wang J, Guo W, et al.  
251 Incidence of constipation in stroke patients: a systematic  
252 review and meta-analysis. *Med (United States).* 2017;96:1–6,  
253 <http://dx.doi.org/10.1097/MD.00000000000007225>.
- 254 9. Yacoline Pailungan F, Kaelan C, Rachmawaty R. Pengaruh Pem-  
255 berian Massage Abdomen Terhadap Penurunan Konstipasi Pada  
256 Pasien Stroke Iskemik Di Rsup Dr. Wahidin Sudirohusodo Makas-  
257 sar. *Patria Artha J Nurs Sci.* 2017;1:25–35.
- 258 10. de Engler TMNM, Dourado CC, Amâncio TG, Farage L, de  
259 Mello PA, Padula MP. Stroke: bowel dysfunction in patients  
260 admitted for rehabilitation. *Open Nurs J.* 2014;8:43–7,  
261 <http://dx.doi.org/10.2174/1874434601408010043>.
- 262 11. Sumida K, Molnar MZ, Potukuchi PK, Thomas F, Lu JL,  
263 Yamagata K, et al. Constipation and risk of death and car-  
264 diovascular events. *Atherosclerosis.* 2019;281(May):114–20,  
265 <http://dx.doi.org/10.1016/j.atherosclerosis.2018.12.021>.
- 266 12. Lämås K, Graneheim UH, Jacobsson C. Experiences of abdom-  
267 inal massage for constipation. *J Clin Nurs.* 2012;21:757–65,  
268 <http://dx.doi.org/10.1111/j.1365-2702.2011.03946.x>.
- 269 13. Deckers B, Von Schoen-Angerer T, Voggenreiter B, Vagedes J.  
270 External nursing applications in the supportive management  
271 of prolonged postoperative ileus description of interven-  
272 tions and case report. *Holist Nurs Pract.* 2016;30:216–21,  
273 <http://dx.doi.org/10.1097/HNP.000000000000158>.
- 274 14. Çevik K, Çetinkaya A, Yiğit Gökbel K, Menekşe B, Saza S, Tikiz C.  
The effect of abdominal massage on constipation in the elderly  
residing in rest homes. *Gastroenterol Nurs.* 2018;41:396–402,  
<http://dx.doi.org/10.1097/SGA.0000000000000343>.
- 275 15. Birimoglu Okuyan C, Bilgili N. Effect of abdominal massage  
276 on constipation and quality of life in older adults: a random-  
277 ized controlled trial. *Complement Ther Med.* 2019;47:102219,  
278 <http://dx.doi.org/10.1016/j.ctim.2019.102219>.
- 279 16. Emly MC. Abdominal massage for constipation. In: Haslam JLJ,  
280 editor. Therapeutic management of incontinence and pelvic  
281 pain. London: Springer; 2008. p. 223–5.
- 282 17. McClurg D, Hagen S, Hawkins S, Lowe-Strong A. Abdomi-  
283 nal massage for the alleviation of constipation symptoms  
284 in people with multiple sclerosis: a randomized con-  
285 trolled feasibility study. *Mult Scler J.* 2010;17:223–33,  
286 <http://dx.doi.org/10.1177/1352458510384899>.
- 287 18. McClurg D, Hagen S, Hawkins S, Lowe-Strong A. Abdomi-  
288 nal massage for the alleviation of constipation symptoms  
289 in people with multiple sclerosis: a randomized con-  
290 trolled feasibility study. *Mult Scler J.* 2011;17:223–33,  
291 <http://dx.doi.org/10.1177/1352458510384899>.
- 292 19. Bhattacharyya K, Winch P, LeBan K, Tien M. Community health  
293 worker incentives and disincentives: how they affect motiva-  
294 tion, retention, and sustainability, Arlington: the Basic Support  
295 for Institutionalizing Child Survival Project (BASICS II) for the  
296 United States Agency for International Development; 2001.
- 297 20. Soesilo D, Wahyuni CU. Improving the village health volunteers’  
298 knowledge, malaria diagnostic skill and home visits through  
299 the observed training (study in Municipality of Tidore Islands  
300 North Maluku Province Indonesia). *Indian J Public Heal Res Dev.*  
301 2019;10:1671–5.
- 302 21. Lukman M, MSi KI, Yani DI, Juniarti N. Exploring strategies to  
303 improve the performance of community health volunteers for  
304 tuberculosis care and prevention: a qualitative study. *Int J*  
305 *Commun Based Nurs Midwif.* 2019;7:270.
- 306 22. Palaia AM. Understanding the motivation, ability and attrition  
307 of HIV/AIDS community health volunteers in resource poor set-  
308 tings: Implications for program sustainability. Johns Hopkins  
309 University; 2010.
- 310 23. Tumbelaka P, Limato R, Nasir S, Syafruddin D, Ormel H, Ahmed  
311 R. Analysis of Indonesia’s community health volunteers (kader)  
312 as maternal health promoters in the community integrated  
313 health service (Posyandu) following health promotion training.  
314 *Int J Community Med Public Heal.* 2018;5:127–35.
- 315 316