

Intercountry prevalences and practices of betel-quid use in south, southeast and eastern Asia regions and associated oral preneoplastic disorders: an international collaborative study by Asian betel-q
by Budi Utomo .

Submission date: 02-Mar-2022 10:41AM (UTC+0700)

Submission ID: 1774369854

File name: Artikel_1._Ganjil_10-11._Intercountry_prevalences.pdf (366.82K)

Word count: 8246

Character count: 37646

6

Intercountry prevalences and practices of betel-quin use in south, southeast and eastern Asia regions and associated oral preneoplastic disorders: an international collaborative study by Asian betel-quin consortium of south and east Asia

Chien-Hung Lee^{1,2}, Albert Min-Shan Ko¹, Saman Warnakulasuriya³, Bang-Liang Yin⁴, Sunarjo⁵, Rosnah Binti Zain⁶, Salah Osman Ibrahim⁷, Zhi-Wen Liu⁸, Wen-Hui Li⁸, Shan-Shan Zhang⁸, Kuntoro⁵, Budi Utomo⁵, Sandage Sunethra Rajapakse⁹, Supun Amila Warusavithana⁹, Ishak Abdul Razak¹⁰, Nortida Abdullah¹¹, Prashanta Shrestha¹², Aij-Lie Kwan¹³, Tien-Yu Shieh¹⁴, Mu-Kuan Chen¹⁵ and Ying-Chin Ko¹

¹ Center of Excellence for Environmental Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

² Department of Public Health, Kaohsiung Medical University, Kaohsiung, Taiwan

³ Department of Oral Medicine, King's College London, and WHO Collaborating Centre for Oral Cancer, London, United Kingdom

⁴ Department of Surgery, The Second Xiangya Hospital, Central South University, Mainland China, People's Republic of China

⁵ Department of Public Health & Preventive Medicine, Airlangga University, Surabaya, Indonesia

⁶ Faculty of Dentistry, Oral Cancer Research & Coordinating Centre, University of Malaya, Kuala Lumpur, Malaysia

⁷ Department of Biomedicine, University of Bergen, Bergen, Norway

⁸ Department of Stomatology, The Second Xiangya Hospital, Central South University, Mainland China, People's Republic of China

⁹ Faculty of Dental Sciences, University of Peradeniya, Peradeniya, Sri Lanka

¹⁰ Faculty of Dentistry, University of Malaya, Kuala Lumpur, Malaysia

¹¹ Oral Health Division, Ministry of Health, Malaysia

¹² B and B Hospital, Kathmandu University Teaching Hospital, Kathmandu, Nepal

¹³ Department of Neurosurgery, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

¹⁴ Department of Oral Hygiene, Kaohsiung Medical University, Kaohsiung, Taiwan

¹⁵ Oral Cancer Center, Changhua Christian Hospital, Changhua, Taiwan

Health risks stemming from betel-quin (BQ) chewing are frequently overlooked by people. Updated epidemiological data on the increased BQ use among Asian populations using comparable data collection methods have not been widely available. To investigate the prevalence, patterns of practice and associated types of oral preneoplastic disorders, an intercountry Asian Betel-quin Consortium study (the ABC study) was conducted for Taiwan, Mainland China, Malaysia, Indonesia, Nepal and Sri Lanka. A random sample of 8,922 subjects was recruited, and the data were analyzed using survey-data modules adapted for the complex survey design. Chewing rates among men (10.7–43.6%) were significantly higher than women (1.8–34.9%) in Taiwan, Mainland China, Nepal and Sri Lanka, while women's rates (29.5–46.8%) were higher than that for men (9.8–12.0%) in Malaysia and Indonesia. An emerging, higher proportion of new-users were identified for Hunan in Mainland China (11.1–24.7%), where Hunan chewers have the unique practice of using the dried husk of areca fruit rather than the solid nut universally used by others. Men in the Eastern and South Asian study communities were deemed likely to combine BQ chewing with smoking and drinking (5.6–13.6%). Indonesian women who chewed BQ exhibited the highest prevalence of oral lichen planus, oral submucous fibrosis and oral leukoplakia (9.1–17.3%). Lower schooling, alcohol drinking and tobacco smoking were identified as being associated with BQ chewing. In conclusion, the ABC study reveals the significant cultural and demographic differences contributing to practice patterns of BQ usage and the great health risks that such practices pose in the Asian region.

Key words: Asia, areca nut, betel-quin, epidemiology, oral disorders, substance abuse

Abbreviations: AN: areca nut; BQ: betel-quin; OL: oral leukoplakia; OLP: oral lichen planus; OSF: oral submucous fibrosis; TA-BQ: tobacco-added betel-quin; TF-BQ: tobacco-free betel-quin

Grant sponsor: Centre of Excellence for Environment Medicine (CEEM), **Grant sponsor:** Kaohsiung Medical University; **Grant number:**

U-EM-99-1-1

DOI: 10.1002/ijc.25809

History: Received 17 Aug 2010; Accepted 4 Nov 2010; Online 2 Dec 2010

Correspondence to: Ying-Chin Ko, The Center of Excellence for Environmental Medicine, Kaohsiung Medical University, No. 100 Shih-Chuan 1st Road, Kaohsiung 807, Taiwan, Tel.: +886-7-311-4418, Fax: +886-7-316-2725, E-mail: ycko@kmu.edu.tw, Fax: +886-7-316-2725, E-mail: ycko@kmu.edu.tw

20 Betel-quid (BQ), a mixture of areca nut (AN) in combination with betel leaf and slaked lime, 13 packed with flavoring ingredients like condiments and sweetening agents, is a chewing (masticatory) substance that is extensively used by diverse Asian populations. Although it is prepared with regional 17 variations, AN is the most important component of BQ. There are an estimated 600 million BQ users in the world, predominantly on the Indo-Pakistan subcontinent, 41 countries in South and Southeast Asia, on islands around the Pacific Rim and in migrant populations in Africa, Europe and North America.¹

While in many parts of the Indian subcontinent, 13 habitual chewers often add tobacco to the BQ, in Taiwan and Papua New Guinea, for example, tobacco is never added. Due to long-standing 59 cultural perspectives, its use has been viewed as publicly acceptable among all strata of society, including women and children.^{2,3} Epidemiological 32 surveys have reported that in the past 2 to 3 decades, 20–40% of the population in India, Nepal and Pakistan have customarily used BQ.⁴ Although a decreased trend in the consumption of BQ has been observed in certain countries or regions, 12 as in Thailand, an alarmingly high chewing prevalence has been found among the Palauans of the West Pacific (a 72–80% use of BQ, with 80% found to be consumers of tobacco-added mixtures).^{4,5} To our knowledge, population data in regard to BQ usage stemming from comparable data collection tools have 67 not been available.

There is sufficient evidence that both tobacco-added (TA-BQ) and tobacco-free (TF-BQ) BQ are carcinogenic 10 to humans.⁶ Furthermore, the mastication of AN is associated with the development and onset 46 of several premalignant lesions in the oral cavity, such as oral submucous fibrosis, which have a high potential 40 for malignant transformation.^{6,7} Evaluations performed by the International Agency for Research on Cancer 53 (IARC) have shown that BQ consumption is causally linked to cancer of the oral 16 cavity, pharynx and esophagus and, with restricted evidence, to liver cancer.⁸ Recent epidemiological studies have shown that BQ use is associated with a 1.8–2.1-, 1.6–2.0-, 1.2–1.9-, 1.3- and 2.6-fold higher 16 risk of suffering from obesity,⁹ metabolic syndromes,¹⁰ cardiovascular disease,¹¹ type 2 diabetes¹² and chronic kidney disease,¹³ a 2.4–3.7-fold risk of delivering low birth weight infants^{14,15} and a 3.6-fold risk of developing cirrhosis of the liver.¹⁶ It is increasingly clear that there are multidimensional health consequences to the use of BQ.

In recent years, the marketing of industrially 50 manufactured BQ products in Asia that contain a variety of areca mixtures with and without tobacco has made the use of this substance common and popular, especially to children and adolescents.⁴ Comprehensive data on the prevalence, usage patterns and conditions of associated oral neoplasia had previously been limited in numerous Asian populations. Such information is warranted to recognize contemporary problems so as to better facilitate the strategic development of prevention and the control of AN-related health consequences in this region.

In 2008, in conjunction with the 60 World Health Organization Collaborating Centre for Oral Cancer in the United Kingdom, the Center of Excellence for Environmental Medicine at Kaohsiung Medical University (KMU) in Taiwan initiated an international collaborative project to mobilize outreach activities in disease prevention focusing 26 on the health effects of BQ in Asian communities (the Asian Betel-quid Consortium study, termed ABC study). Six large study centers 3 participated in this international research consortium (East Asia: Taiwan and Mainland China; Southeast Asia: Malaysia and Indonesia; South Asia: Nepal and Sri Lanka), where the goals were to update the characteristics of usage prevalence and the patterns of use with other substances within the Asian region as well as the potential identification of genetic susceptibilities to oral malignancies. Our study represents phase I among a wide range of planned activities by this consortium. The specific aims herein were to outline the current prevalence of BQ use, the characteristics among the variations of practice, the lifestyle factors inherent in BQ usage and the types of associated oral preneoplastic disorders across the countries that were studied.

Material and Methods

Recruitment in the phase I study by this consortium commenced in January 2009 and ended in February 2010. The same research protocol was applied among the countries in the study, with the purpose of offering a comparative framework. This proposal was approved independently by the Research and Ethical Review Committees at the KMU (Kaohsiung) in Taiwan, the Central South University (Changsha) in Mainland China, the University of Malaya (Kuala Lumpur) in Malaysia, the Airlangga University (Surabaya) in 3 Indonesia, the Kathmandu University (Kavrepalanchok) in Nepal and the University of Peradeniya (Peradeniya) in Sri Lanka. Written informed consent was obtained from all of the participants prior to the collection of data.

Study population

The study populations comprised of residents from southern Taiwan; the Hunan Province of Mainland China; the Selangor, Sabah and Sarawak States of Malaysia; the North Sumatran, East Javan, Balese, West Nusa Tenggara, South Sulawesi and Papua Provinces of Indonesia; as well as the Middle of Nepal and the Central Province of Sri Lanka, as shown in Figure 1.

Using demographic data from census records and the household registry of individual participating countries, a multistage sampling approach was employed to select civilian, noninstitutionalized representative samples of the general population of the ages of 15 and older. In stage one, a number of cities and rural areas were independently selected to favorably make up and represent the geographic and economic status within each population to be studied. The resulting chosen regions were Kaohsiung and Pingtung of southern Taiwan; Changsha, Liuyang, Changde, Yongzhou,




Figure 1. The geographic regions showing lifetime prevalence rate of betel-quid chewing by gender distribution in the Asian countries studied in the ABC study. The international mobilizing outreach activities in prevention of potentially malignant oral disorders. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

LouDi and Xiangxi prefecture, Hunan of Mainland China; Pulau Carey, Simpang Morib and Klang (Selangor), Kampung Sembirai and Kota Belud (Sabah), Kampung, Tebedu/Mongkos and Serian (Sarawak) of Malaysia; Deli Serdang (North Sumatra), Pacitan and Banyuwangi (East Java), Jembrana (Bali), Mataram (West Nusa Tenggara), Tana Toraja (South Sulawesi) and Wamena (Papua) of Indonesia; Kathmandu, Chitwan, Nawalparasi and Pokhara of middle Nepal and Gangawata Korale, Udunuwara, Yatinuwara of Sri Lanka.

In the second stage, projected numbers for administrative districts or townships were randomly selected. In stage three, researchers listed households within each district/township, and a sample of households was randomly drawn. In the final stage, subjects were chosen to participate in our study from a list of people residing in the selected households. On average, 2.2–2.6 people were selected per household in Taiwan, Malaysia, Indonesia and Sri Lanka. Due to logistic concerns, only one participant was randomly selected from each

household in the populations of Mainland China and Nepal. With this exception, a similar sampling scheme was implemented in all of the areas that were studied.

54 Data collection

After the study samples were drawn, the research teams conducted door-to-door home visits to the participants' dwellings to conduct the interviews. Each survey team consisted of a principle investigator (PI), four to five trained interviewers and data-recording clerks. The purposes and procedures involved in the study were explained to all of the participants. The resulting total of those surveyed who completed the interviews was: 1,548 subjects in Taiwan, 2,356 in Mainland China, 1,003 in Malaysia, 1,941 in Indonesia, 1,002 in Nepal and 1,072 in Sri Lanka. The response rates were 68% for Taiwan, 79% for Mainland China, 71% for Malaysia, 100% for Indonesia, 99% for Nepal and 99% for Sri Lanka, respectively.

Data were collected by trained staff members who had recently successfully completed the necessary training

program as a prerequisite to working in each study center under the supervision of the country's PI. A standardized questionnaire was developed which was adapted from WHO surveys and sources such as national prevalence surveys. The original questionnaires were written in English and translated into the appropriate language or dialect for each country. They were later retranslated into English to verify their validity. The questions have been written in community-specific languages, and in the few situations where the participants were unable to understand the questions, study items were translated into local dialects by investigators with the assistance of local, native people. Pictograms were used to clarify constituents used to make the BQ. The interviewer-administered structured questionnaire gathered data on the socio-demographics and habits of lifetime BQ usage, alcohol intake and tobacco use (including tobacco smoking and smokeless tobacco use). In addition, information concerning the age at which subjects started using these substances, the daily consumption, duration of usage, as well as the types and frequency of substances consumed were collected. For the BQ users, further information was collected in regard to the materials added with AN. Chewers were also asked to report whether they had habitually swallowed fluids from such chewing. Moreover, in preparation for their work in the detection of oral lesions, the dentists or dental hygienists/medical officers received 1 month of standardized training in regard to performing oral cavity examination, and this was done under the supervision of the survey team's PI. The training programs included differential diagnosis for oral disorders where clinical pictures were used and pretest courses conducted in a clinical or field setting.

Our study defined BQ chewers as subjects who had consumed at least one quid per day of any type of betel/AN product for a minimum of 6 months. Alcohol drinkers and tobacco smokers were defined separately as individuals who had drunk any type of alcoholic beverage per week and/or had smoked/chewed one or more tobacco products per day for at least 6 months. Among them, current-users were those who had any of these habitual practices within the year prior to the interview; past-users were defined as those who had stopped any of the habits for at least a year before the interviews took place.

Statistical analysis

The study data was analyzed using survey-data modules implemented in Stata v11, a technique that adjusts for the complex survey design. All analyses, including means, percentages, standard errors, point estimates for prevalence and regression modeling were conducted on weighted data that took age and gender structure into account. For continuous and binary/multinomial outcomes, we respectively used multiple linear and logistic/polytomous regression models to assess across country differences in regard to the characteristics of BQ usage. A multivariate logistic regression model was employed to investigate the influence of age, gender, educa-

tional level, alcohol consumption and tobacco smoking on country-specific behaviors in BQ use. Because observed gender heterogeneity in the chewing prevalence was identified in most countries, the results were evaluated according to gender.

Results

The demographic characteristics with respect to gender, age, educational level and the status of the respective drinking and smoking are presented in Table 1. Significant differences in these factors were detected across the areas of study (all $p < 0.05$). The prevalence of tobacco smoking was found to be higher than that of alcohol drinking in every sampled country, with the highest rates occurring in Nepal for both smoking (55.2%) and drinking (29.9%).

Table 2 compares the prevalence and BQ usage among the communities that were studied according to gender. Remarkable country-dependent differences in lifetime prevalence were observed for both genders for difference (< 0.005). As shown in Figure 1, the lifetime chewing rates for men (15.6–43.6%) were significantly higher than that for women (2.3–34.9%) in areas that were sampled in Taiwan, Mainland China, Nepal and Sri Lanka, while conversely the prevalence for women (32.1–47.8%) were notably higher than men (10.3–12.4%) in the Malaysian and Indonesian regions of investigation. In regard to the BQ ever-users, a high percentage of quitters (15.3–31.1%), with a >14 years of abstinence level was found among the Taiwanese. All of the BQ users in Nepal were current-chewers, and no quitters were identified. The Hunan (Mainland Chinese) BQ users had a shorter duration of usage (8.8–9.7 years), but a higher proportion of new consumers (usage of ≤ 2 years) among current-chewers (11.1–24.7%). A reported $>49\%$ chewing rate occurred in older Malaysian and Indonesian women (>40 years). In contrast, a higher prevalence of chewing (32.0%) was seen among younger Hunan men in Mainland China (<41 years).

Significant differences in the ingredients of BQ were found in the investigated communities in our study. To assess the possible carcinogenetic impact of BQ consumption, we categorized the BQ preparation according to the addition of tobacco or its being tobacco-free.⁶ Table 3 shows that 100% of Nepalese chewers added tobacco to their BQ, and that, in Taiwan and the Hunan of Mainland China, tobacco is never added as an ingredient to BQ. An overwhelming majority of chewers in Taiwan consumed AN with a piece of betel (84.4–93.3%), and all of the chewers in Hunan consumed the dried husk of the areca fruit (not the solid AN) marinated with flavored additives, such as lime, sweeteners, oil and bitter. The highest daily quantity of BQ use by both men and women was found in Taiwan (16.2–19.5 quid). In addition, TA-BQ users chewed BQ more frequently (5.7–6.9 days per week) as compared to TF-BQ users, the latter of which reporting their chewing frequency at 4.3–5.9 days per week. AN chewers in Taiwan, Malaysia, Indonesia and Sri

Table 1. Characteristics for the study participants, the ABC study

Characteristics	Taiwan	Mainland China	Malaysia	Indonesia	Nepal	Sri Lanka
Study sample, No.	1,548	2,356	1,003	1,941	1,002	1,072
Gender (%)						
Male	47.6	52.0	38.2	49.7	66.3	35.9
Female	52.5	48.0	61.8	50.3	33.7	64.1
Age [years (%)]						
≤30	19.0	31.6	20.2	17.1	46.7	24.4
31–40	19.5	22.8	15.9	19.1	23.9	19.7
41–50	17.9	16.9	21.3	17.2	15.9	15.6
51–60	20.9	14.8	21.7	19.2	10.5	17.0
≥61	22.7	13.9	20.8	27.4	3.1	23.3
Education [years (%)]						
≤6	24.2	20.2	61.2	75.2	66.2	22.8
7–12	44.1	58.0	36.7	24.5	20.1	67.0
≥13	31.7	21.8	2.1	0.3	13.8	10.3
Alcohol drinker (%)						
No	87.1	78.3	84.7	77.1	70.1	88.0
Yes	12.9	21.7	15.4	22.9	29.9	12.0
Tobacco smoker (%)						
No	76.7	68.0	72.8	58.8	44.8	82.8
Yes	23.3	32.0	27.2	41.2	55.2	17.2

Lanka were found to be inclined to spit out the chewing quid juice; this is in stark contrast to those from Mainland China and Nepal, who swallowed it. Furthermore, in Eastern and South Asian study communities, male BQ users were found to have the highest prevalence for chewing that was combined with smoking and drinking (5.6–13.6%). With the exception of Nepal, only a handful of female chewers were found to concomitantly use cigarettes and alcohol. Among Malaysian and Indonesian women, a high proportion of substance users were observed to be exclusively BQ users (a prevalence of 27.8–45.7%).

The prevalence of oral mucosal lesions associated with BQ chewing is presented in Table 4. Indonesian women had the highest prevalences of oral lichen planus (OLP), oral submucous fibrosis (OSF) and oral leukoplakia (OL) (9.6–17.3%). Compared to nonchewers, Taiwanese male chewers had a remarkably higher likelihood of suffering from these oral pre-neoplastic disorders [an adjusted odds ratio of (aOR) = 13.5–137.4]. OSF appeared to be the most common oral pre-malignancy in Hunan, as indicated by the 3.5–6.6% among BQ users there.

The association between the practice of BQ chewing with educational levels and the habits of drinking and smoking are given in Figure 2. Compared to those who had only finished primary school (<7 years of schooling), subjects who had achieved a tertiary level of education (>12 years of education) were found to have a 0.3-fold less likelihood of being

a chewer among men in Taiwan. Women with 7 to 12 years of schooling also had a reduced chance of becoming a chewer in Malaysia and Sri Lanka (aOR = 0.2–0.3, $p < 0.05$). With the exception of Indonesia, alcohol drinkers almost always had a higher tendency of being chewers, with a 2.1–8.9-fold increased risk observed in men, and a 5.2–21.2-fold elevated risk in women in Malaysia and Nepal. In Taiwan and Hunan, male smokers had an elevated propensity of becoming a BQ chewer. However, smokers in Malaysia and Indonesia were found to have a less likelihood of being a chewer (aOR = 0.1–0.4 in men, 0.2–0.4 in women).

Discussion

Our study offers the most up-to-date and comprehensive information on the prevalence, practice patterns and related factors in regard to BQ usage for six Asian populations. The findings further reveal the most affected populations and types of potentially malignant oral disorders among the studied regions where BQ use is widely practiced.

In this investigation, a remarkable discrepancy in gender prevalence for BQ use was recognized in the countries studied. Malaysian and Indonesian women had a particularly high chewing rate (29.5–46.8%), with female chewers in these two countries having reported an average history of >23 years of BQ usage, and such women tending to have a lower educational level (Fig. 2). Further, women aged >40 years in Malaysia and >30 years in Indonesia showed a 35.9–95.9%

Table 2. Prevalence and characteristics of betel-quinid chewing, stratified by gender, the ABC study

Prevalence/category	Men										Women									
	Taiwan n = 736	Mainland China n = 1225	Malaysia n = 383	Indonesia n = 965	Nepal n = 664	Sri Lanka n = 385	Taiwan n = 812	Mainland China n = 1131	Malaysia n = 620	Indonesia n = 976	Nepal n = 338	Sri Lanka n = 687	Taiwan n = 812	Mainland China n = 1131	Malaysia n = 620	Indonesia n = 976	Nepal n = 338	Sri Lanka n = 687	p for diff. ¹	
Nonchewer (%)	84.4	71.0	89.7	79.5	61.8	78.8	97.0	97.7	67.9	32.1	63.3	83.1	97.0	97.7	67.9	32.1	63.3	83.1		
Chewer in lifetime (%)	15.6	29.0	10.3	12.4	43.6	21.2	3.0	2.3	32.1	47.8	34.9	14.5	3.0	2.3	32.1	47.8	34.9	14.5	**	
Chewing status																				
Past-chewer (%)	4.8	5.1	0.5	0.4	0.0	3.2	0.5	0.5	2.6	1.0	0.0	0.9	0.5	0.5	2.6	1.0	0.0	0.9		
Proportion of quitter in ever-chewers (%)	31.1	17.5	5.2	3.5	0.0	15.2	15.3	21.4	8.1	2.1	0.0	6.5	15.3	21.4	8.1	2.1	0.0	6.5	**	
Years of quitting ²	14.8 ± 1.9	3.6 ± 0.6	4.0 ± 3.2	9.0 ± 2.2	NA	13.8 ± 4.2	26.0 ± 2.1	2.4 ± 0.5	13.1 ± 2.7	17.4 ± 2.7	NA	10.6 ± 3.1	26.0 ± 2.1	2.4 ± 0.5	13.1 ± 2.7	17.4 ± 2.7	NA	10.6 ± 3.1	**	
Current-chewer (%)	10.7	23.9	9.8	12.0	43.6	18.0	2.5	1.8	29.5	46.8	34.9	13.5	2.5	1.8	29.5	46.8	34.9	13.5		
Age at starting use ²	20.7 ± 0.8	23.7 ± 0.6	20.9 ± 1.5	33.3 ± 0.9	18.9 ± 0.7	26.2 ± 1.5	25.4 ± 3.5	29.6 ± 2.6	23.1 ± 0.8	25.5 ± 0.6	18.7 ± 0.6	31.6 ± 1.5	25.4 ± 3.5	29.6 ± 2.6	23.1 ± 0.8	25.5 ± 0.6	18.7 ± 0.6	31.6 ± 1.5	**	
Years of chewing ²	21.1 ± 2.0	8.8 ± 0.4	30.1 ± 2.0	15.3 ± 1.2	15.4 ± 1.9	21.7 ± 2.0	33.8 ± 5.3	9.7 ± 1.7	25.2 ± 1.3	23.8 ± 1.6	16.0 ± 2.0	23.9 ± 2.1	33.8 ± 5.3	9.7 ± 1.7	25.2 ± 1.3	23.8 ± 1.6	16.0 ± 2.0	23.9 ± 2.1	**	
Proportion of new user in chewers (%) ³	7.7	11.1	0.0	14.7	13.1	15.5	0.0	24.7	10.7	7.7	18.2	20.3	0.0	24.7	10.7	7.7	18.2	20.3	**	
Age-specific prevalence (%)																				
≤30	6.7	32.0	1.8	1.1	46.1	4.3	0.0	1.3	4.7	18.9	29.1	3.4	0.0	1.3	4.7	18.9	29.1	3.4	**	
31–40	16.0	32.0	3.0	14.4	40.4	24.2	0.0	2.5	28.8	35.9	34.3	5.2	0.0	2.5	28.8	35.9	34.3	5.2		
41–50	13.0	20.6	16.2	17.4	41.5	35.8	3.8	2.6	49.1	61.2	46.5	17.4	3.8	2.6	49.1	61.2	46.5	17.4		
51–60	13.2	8.6	32.7	22.3	34.3	23.2	4.0	1.0	68.7	81.5	57.0	22.8	4.0	1.0	68.7	81.5	57.0	22.8		
≥61	4.9	3.9	20.0	20.0	53.0	19.7	7.7	1.4	61.3	95.9	56.2	36.5	7.7	1.4	61.3	95.9	56.2	36.5	**	

* p < 0.05. ** p < 0.005. ¹ p for the difference in characteristics of betel-quinid chewing across study areas was obtained adjusted for age. ² Mean ± SE. ³ New-users, defined as those who regularly used betel-quinid for ≤ 2 years. Abbreviations: NA: nonappreciable; NS: nonsignificant.

Table 3. Types of betel-quinid (BQ) chewed and habits practiced among current-chewers, stratified by gender, the ABC study

Betel-quinid chewing/category	Men							Women							p for diff. ¹				
	Taiwan			Mainland China			Sri Lanka			Taiwan			Mainland China			Sri Lanka			
	n = 731	n = 1225	n = 383	n = 965	n = 664	n = 385	n = 812	n = 1131	n = 620	n = 976	n = 338	n = 687							
Type of betel-quinid²																			
Tobacco-free BQ																			
Prevalence (%)	10.7	23.9	3.6	1.6	0.0	11.6	**	2.5	1.8	17.5	15.0	0.0	10.4	**					
% in all types	100.0	100.0	36.4	29.7	0.0	64.6		100.0	100.0	59.4	31.3	0.0	76.7						
Nut + leaf	84.4	0.0	36.4	2.3	0.0	0.0		93.3	0.0	54.6	16.5	0.0	6.7						
Nut + inflorescence	8.0	0.0	0.0	6.2	0.0	1.0		0.0	0.0	0.4	11.7	0.0	0.9						
Nut + stem	3.3	0.0	0.0	0.6	0.0	0.0		0.0	0.0	0.3	0.0	0.0	0.0						
Nut husk + favorites	0.0	100.0	0.0	0.0	0.0	0.0		0.0	100.0	0.0	0.0	0.0	0.0						
Nut + local favorites	4.3	0.0	0.0	0.6	0.0	63.6		6.7	0.0	4.1	3.1	0.0	69.1						
Daily amount of chewing ³	19.5 ± 2.7	10.0 ± 0.6	3.6 ± 1.0	3.9 ± 0.5	NA	2.1 ± 0.2	**	16.2 ± 2.2	4.9 ± 0.8	4.6 ± 0.4	3.5 ± 0.3	NA	2.0 ± 0.2	**					
Usage day per week ³	5.7 ± 0.3	5.8 ± 0.1	4.4 ± 0.9	4.3 ± 0.4	NA	5.0 ± 0.4	**	6.1 ± 0.5	5.7 ± 0.5	5.9 ± 0.3	5.7 ± 0.4	NA	4.4 ± 0.3	**					
Tobacco-added BQ																			
Prevalence (%)	0.0	0.0	6.2	10.4	43.6	6.4	**	0.0	0.0	12.0	31.7	34.9	3.2	**					
% in all types	0.0	0.0	63.6	70.4	100.0	35.5		0.0	0.0	40.6	68.7	100.0	23.3						
Daily amount of chewing ³	NA	NA	5.7 ± 0.6	6.5 ± 0.4	9.0 ± 0.3	6.9 ± 1.1	**	NA	NA	6.8 ± 0.4	6.1 ± 0.3	9.0 ± 0.9	6.2 ± 0.9	NS					
Usage day per week ³	NA	NA	6.4 ± 0.4	6.8 ± 0.1	6.0 ± 0.1	6.8 ± 0.1	**	NA	NA	6.8 ± 0.1	5.7 ± 0.2	5.8 ± 0.1	6.9 ± 0.1	**					
Habit in chewers (%)																			
Spit out the juice	41.2	15.2	82.3	72.5	0.0	76.9		64.4	16.7	89.1	71.8	0.0	65.6						
Swallow the juice	43.5	78.7	4.8	13.2	95.0	9.7		25.8	77.2	3.3	7.6	94.4	19.5						
Spit/swallow alternately	15.3	6.1	12.9	14.3	5.0	13.3		9.8	6.1	7.6	20.6	5.6	14.9						
Prevalence of combined habits (%)																			
Chewing only	1.4	2.9	1.7	10.1	13.7	3.7	**	2.4	1.5	27.8	45.7	8.1	13.4	**					
Chewing + smoking	0.8	2.5	3.8	0.1	10.4	2.7		0.0	0.1	1.2	0.0	7.7	0.2						
Chewing + drinking	2.9	8.8	0.2	1.0	6.0	4.3		0.0	0.1	0.4	1.0	6.6	0.0						
Chewing + smoking + drinking	5.6	9.8	4.1	0.8	13.6	7.3		0.1	0.1	0.1	0.1	12.6	0.0						

* p < 0.05, ** p < 0.005, ¹p for the difference in characteristics of betel-quinid chewing across study areas was obtained adjusted for age, ²There were missing data in the type of betel-quinid chewed, ³Mean ± SE.

15
Table 4. Prevalence rate (PR) and adjusted odds ratio (aOR) of oral lichen planus (OLP), 34
oral submucous fibrosis (OSF) and oral leukoplakia
(OL) associated with betel-quinid (BQ) chewing, stratified by gender, the ABC study

Gender/country	Study number		PR of OLP (%)				PR of OSF (%)				PR of OL (%)			
	Nonchewer	Chewer	Nonchewer	Chewer	aOR ¹	p	Nonchewer	Chewer	aOR ¹	p	Nonchewer	Chewer	aOR ¹	p
Men														
Taiwan	585	91	0.1	5.5	57.5	**	0.0	7.3	13.5	** ²	0.1	6.4	137.4	*
Mainland China	886	339	0.1	0.3	2.2	NS	0.0	6.6	91.8	** ²	0.1	0.6	8.1	NS
Malaysia	337	45	0.0	0.0	NA	—	0.0	0.0	NA	—	0.0	0.0	NA	—
Indonesia	718	196	6.5	9.6	5.7	**	4.8	5.5	2.7	*	9.6	6.6	1.7	NS ³
Nepal	410	254	0.0	0.0	NA	—	0.0	0.0	NA	—	0.3	1.6	4.7	NS
Sri Lanka	283	99	0.0	0.0	NA	—	0.0	0.0	NA	—	0.0	0.8	NA	—
Women														
Taiwan	715	15	0.0	0.0	NA	—	0.0	0.0	NA	—	0.0	0.0	NA	—
Mainland China	1105	26	0.1	0.0	NA	—	0.1	3.5	26.7	NS	0.0	0.0	NA	—
Malaysia	345	275	0.0	0.0	NA	—	0.0	0.0	NA	—	0.0	0.0	NA	—
Indonesia	277	660	0.8	17.3	28.4	**	1.2	9.6	6.8	*	0.2	9.1	212.8	**
Nepal	214	124	0.0	0.0	NA	—	0.2	0.0	NA	—	0.0	0.0	NA	—
Sri Lanka	571	114	0.0	0.0	NA	—	0.1	0.0	NA	—	0.0	0.0	NA	—

* $p < 0.05$. ** $p < 0.005$. ¹ p for OR of oral mucosal lesions was obtained adjusted for age, tobacco smoking and alcohol drinking. ²The odds ratio was calculated using the median unbiased estimates with the aid of exact logistic regression. ³Smoking conveys a negative confounding effect on the aOR of OL for BQ use.

Abbreviations: NA: nonappreciable; NS: nonsignificant.

high BQ chewing rate. As noted and released by the IRAC,⁶ the length of BQ usage time is an important predictor in the development of oral precancerous and neoplastic lesions. Moreover, the health risk stemming from BQ chewing was generally less awareness among those women with a lesser educational level.¹⁷ Consequently, Malaysian and Indonesian women from the study regions appear to be a high-risk group who require an increased amount of concern and educational assistance in regard to BQ-related health and disease prevention.

Although the raw areca used in Hunan is imported from Hainan in Mainland China, as well as from Thailand, the vast majority of BQ products in China, including commercial forms, are manufactured there. The Xiangtan city in Hunan province, where BQ chewing was reported to be very common (prevalence, 64.5–82.7%), is also where most BQ production factories and workshops are located.¹⁸ In our study, we found that men (11.1%) and women (24.7%) from Hunan have a high percentage of BQ new-users. In contrast to other regions, the prevalence of BQ chewing in Hunan men was higher in the younger age groups, suggesting that Hunan is an emerging region of BQ practice. The improving economy and easy access to BQ products there, along with some recently begun BQ advertising campaigns in Hunan, could be the factors for widespread use of this substance, particularly among young people.

There is great diversity in how people chew BQ due to quite a bit of habitual variation, the majority of such practices being associated with higher risks for tumor development. In 1985, TA-BQ has been deemed by the IARC to be

carcinogenic to humans.¹⁹ Moreover, AN-derived carcinogenic nitrosamines have been unequivocally identified in the saliva of TF-BQ chewers.²⁰ Currently, there is sufficient evidence that TF-BQ causes oral and esophageal cancers.^{6,8} Recent epidemiological evidence has revealed that the use of TF-BQ with betel inflorescent (containing safrole, a Group 2B carcinogen) conferred a higher carcinoma risk in the oral cavity, pharynx and esophagus than any other accompanying ingredients.^{21–24} BQ chewers who also have the habits of alcohol drinking and tobacco smoking were found to have a synergistic risk of contracting the three aforementioned cancers of the upper gastrointestinal tract (OR = 8.1–41.2).^{21–23} For users who habitually swallowed the juice from BQ chewing, a greater cancer risk was also observed.^{21,22}

The Nepalese subjects who resided in the investigative regions displayed a 34.9–43.6% high rate of BQ chewing, and all were observed to have consumed TA-BQ. The daily amount of TA-BQ used by both genders was found to be the most (nine quids) in our study, and the overwhelming majority of chewers (94.4–95.0%) typically swallowed the BQ juice. Further, 12.6–13.6% of these people practiced chewing, drinking and smoking concomitantly. This is consistent with the age-standardized incidence of oral cancer in Nepal, which is ranked at a relatively high level according to GLOBOCAN 2002 report (12.8 and 8.4 per 100,000 men and women, respectively).²⁵ Hence, BQ-associated health problems should be carefully observed and counteracted in regard to the Nepalese population.

In Taiwan, the most important campaign to prevent BQ usage was initiated in 1998 where the government named




Figure 2. Adjusted odds ratio (aOR) showing the impact of education (schooling year), alcohol drinking and tobacco smoking upon the practice of BQ chewing, the ABC study. Note: aOR was adjusted for age and covariates in this figure. NA: nonappreciable, due to null BQ chewers in the group >12 years of schooling. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

December 3 as “Betel Quid Prevention Day” due to finding that BQ chewers who concomitantly have the habits of alcohol drinking and cigarette smoking experienced a 123-fold increased oral cancer risk.²¹ Although a reduced quantity of AN consumption has been observed in Taiwan,²⁶ health promotion campaigns have been continued. The high rate of BQ quitting that was noticed in our Taiwanese male samples (31.1%) is probably a result of such activities.

Compared to the results found in a population survey of the greater Kaohsiung area that was conducted in 1994,⁷ the consumption of TF-BQ with betel inflorescence among chewers in Taiwan has substantially declined (from 51% in 1994 to 0–8% in our study). This indicates that people in

Taiwan have noticed and considered the warnings about the higher cancer risks detected as a result of the intake of safrole-contained betel inflorescence.^{21–23} On the other hand, usage pattern for BQ chewers in Hunan differs greatly from that of other regions where BQ chewing is prevalent. All of the Hunan users chewed the dried husk of areca fruit,²⁷ and the majority (77.2–78.7%) of such users habitually swallowed the chewing quid juice as well. While current data has been limited in regard to the appraisal of the carcinogenetic consequences of consuming the dried areca husk,¹⁸ a significant relationship between the consumption of this substance and OSF development was identified in our study for the male group from Hunan.

The IARC working group in 2004 reported that the prevalence of OLP, OSF and OL among BQ chewers in selected Asian populations was 0–3.7%, 0–3.0% and 0–40.1%, respectively.⁶ The annual rate of malignant transformation was estimated to be <0.1%, 0.5% and 1% for the corresponding oral disorders.²⁸ Our study revealed that male Taiwanese TF-BQ chewers had a notably greater prevalence of exhibiting these potentially malignant oral disorders than nonchewers. In Hunan, an appreciably elevated OSF prevalence was detected in male chewers, despite the fact that they consumed tobacco-free areca husk. Approximately 6–17% of Indonesian BQ chewers suffered the above oral disorders, and a 7–213-fold risk was found for female chewers. The high prevalence of oral preneoplastic lesions and conditions in Indonesia reflect the current disease loadings and oral public health problems. With no reliable marker to predict malignant transformation²⁹ as yet, these data strengthen the great importance of disease prevention in Indonesia.

In with several studies,^{30,31} subjects with higher schooling in Taiwan, Malaysia, Indonesia and Sri Lanka were found to be correlated with a lesser likelihood to be BQ chewers. These results suggest that BQ-related health education programs should target less educated people in these study communities. In contrast, Hunan subjects with higher educational levels had a greater likelihood to use BQ (OR = 1.3–1.5, though nonsignificant), regardless of gender. As previously noted, the usage of this substance is more readily available in Hunan. This implies that the significance of extensive health education in this region is particularly important.

Large-scale epidemiological studies conducted in Taiwan have shown that drinking and smoking are significantly associated with an earlier onset age for the chewing of BQ.³² Findings from a nationwide investigation further indicated that chewing practices are appreciably reduced among smokers who successfully quit smoking.³³ In this ABC study, alcohol drinking was the most important habit concomitant with BQ use. Male drinkers had a 2.1–8.9-fold likelihood of being a chewer in most countries, and female drinkers had a 5.2–21.2-fold probability of being a BQ user in Malaysia and Nepal. Interestingly, in the two Chinese populations, where chewers have been mostly men, male smokers showed a greater propensity to practice BQ chewing (aOR = 3.6–

4.3). As the habit of chewing is in sync with that of drinking and/or smoking, preventive policies should focus on BQ chewing and its link to individual alcohol and tobacco habits.³³

In summary, we have outlined the significant country and gender differences contributing to BQ use and the great health risks that such practices pose in the Asian region. This international collaborative effort reveals the important BQ chewing-related demographic factors and regional-dependent factors, such as the emerging number of users in Hunan, and the overwhelming number of female chewers in Malaysia and Indonesia. It has also shed light on the importance of the inclusion of diverse ingredients like tobacco and nontobacco-based BQ and on the types of oral premalignant conditions associated with

BQ chewing. We believe that the ABC study will greatly help to unify the awareness of the dangers of such diverse BQ practices. We also believe that it can provide an important framework to the development of more effective public health programs to help combat such serious practices

Acknowledgements

We would like to thank Jennifer Ko, the staff of the CEEM, for her great assistance in helping to organize the various centers' principal investigators. We would also like to express our appreciation to the study staff members, including Sanjeevani Jayshankar, Dipak Sapkota, Norain Abu Talib, Helen Ng Lee Ching, Tah Geok Mooi, Zaiton Tahir, Nurshaline Pauline Hj Kipli, Tian-You Ling, Ekamaya, Aris Istiaq, SKM, Lutviana, Sriana Herman, Nuril, Taufiq, Herman and Arifin, for their diligent work and excellent endeavors in this international cooperative study.

References

- Boucher BJ, Mannan N. Metabolic effects of the consumption of Areca catechu. *Addict Biol* 2002;7:103–10.
- Gupta PC, Warnakulasuriya S. Global epidemiology of areca nut usage. *Addict Biol* 2002;7:77–83.
- Yang MS, Su IH, Wen JK, Ko YC. Prevalence and related risk factors of betel quid chewing by adolescent students in southern Taiwan. *J Oral Pathol Med* 1996; 25:69–71.
- Gupta PC, Ray CS. Epidemiology of betel quid usage. *Ann Acad Med Singapore* 2004; 33:31–6.
- Tovossia S, Chen PH, Ko AM, Tu HP, Tsai PC, Ko YC. Prevalence and associated factors of betel quid use in the Solomon Islands: a hyperendemic area for oral and pharyngeal cancer. *Am J Trop Med Hyg* 2007;77:586–90.
- IARC. Betel-quid and areca-nut chewing and some areca-nut derived nitrosamines. *IARC Monogr Eval Carcinog Risks Hum* 2004;85:1–334.
- Lee CH, Ko YC, Huang HL, Chao YY, Tsai CC, Shieh TY, Lin LM. The precancer risk of betel quid chewing, tobacco use and alcohol consumption in oral leukoplakia and oral submucous fibrosis in southern Taiwan. *Br J Cancer* 2003;88:366–72.
- Secretan B, Straif K, Baan R, Grosse Y, El Ghissassi F, Bouvard V, Benbrahim-Tallaa L, Guha N, Freeman C, Galichet L, Cogliano V. A review of human carcinogens. Part E. Tobacco, areca nut, alcohol, coal smoke, and salted fish. *Lancet Oncol* 2009;10:1033–4.
- Lin WY, Pi-Sunyer FX, Liu CS, Li TC, Li CI, Huang CY, Lin CC. Betel nut chewing is strongly associated with general and central obesity in Chinese male middle-aged adults. *Obesity (Silver Spring)* 2009;17: 1247–54.
- Yen AM, Chiu YH, Chen LS, Wu HM, Huang CC, Boucher BJ, Chen TH. A population-based study of the association between betel-quid chewing and the metabolic syndrome in men. *Am J Clin Nutr* 2006;83:1153–60.
- Lin WY, Chiu TY, Lee LT, Lin CC, Huang CY, Huang KC. Betel nut chewing is associated with increased risk of cardiovascular disease and all-cause mortality in Taiwanese men. *Am J Clin Nutr* 2008;87:1204–11.
- Tung TH, Chiu YH, Chen LS, Wu HM, Boucher BJ, Chen TH. A population-based study of the association between areca nut chewing and type 2 diabetes mellitus in men (Keelung Community-based Integrated Screening programme No. 2). *Diabetologia* 2004;47:1776–81.
- Lin WY, Cheng SY, Liu JH, Cheng WC, Kang IM, Tseng YH, Shih CM, Chen W. Association between betel-nut chewing and chronic kidney disease in men. *Public Health Nutr* 2009;12:723–727.
- Yang MS, Lee CH, Chang SJ, Chung TC, Tsai EM, Ko AM, Ko YC. The effect of maternal betel quid exposure during pregnancy on adverse birth outcomes among aborigines in Taiwan. *Drug Alcohol Depend* 2008;95:134–9.
- Yang MS, Chang FT, Chen SS, Lee CH, Ko YC. Betel quid chewing and risk of adverse pregnancy outcomes among aborigines in southern Taiwan. *Public Health* 1999;113: 189–92.
- JF, Jeng JE, Chuang LY, Ho MS, Ko Lin ZY, Hsieh MY, Chen SC, Chuang WL, Wang LY, Yu ML, Lin CY, et al. Habitual betel quid chewing as a risk factor for cirrhosis: a case-control study. *Medicine (Baltimore)* 2003;82:365–72.
- Gunaseelan R, Sankaralingam S, Ramesh S, Datta M. Areca nut use among rural residents of Sriperambudur Taluk: a qualitative study. *Indian J Dent Res* 2007; 18:11–4.
- Yang X, Reichart PA. A review of betel quid chewing, oral cancer and precancer in Mainland China. *Oral Oncol* 2007;13: 424–30.
- IARC. Tobacco habits other than smoking: betel-quid and areca-nut chewing; and some related nitrosamines. IARC Working Group. Lyon, 23–30 October 1984. *IARC Monogr Eval Carcinog Risk Chem Hum* 1985;37:1–268.
- Nair J, Nair UJ, Ohshima H, Bhide SV, Kawanishi S, H. Endogenous nitrosation in the oral cavity of chewers while chewing betel quid with or without tobacco. *IARC Sci Publ* 1987;84:465–9.
- Ko YC, Huang YL, Lee CH, Chen MJ, Lin LM, Tsai CC. Betel quid chewing, cigarette smoking and alcohol consumption related to oral cancer in Taiwan. *J Oral Pathol Med* 1995;24:450–3.
- Lee KW, Kuo WR, Tsai SM, Wu DC, Wang L, Fang FM, Chiang FY, Ho KY, Wang LF, Tai CF, Kao EL, Chou SH, et al. Different impact from betel quid, alcohol and cigarette: risk factors for pharyngeal and laryngeal cancer. *Int J Cancer* 2005;117:831–6.
- Lee CH, Lee JM, Wu DC, Hsu HK, Kao EL, Huang HL, Wang TN, Huang MC, Lin MT. Independent and combined effects of alcohol intake, tobacco smoking and betel quid chewing on the risk of esophageal cancer in Taiwan. *Int J Cancer* 2005;113:475–82.
- Jiang SL, Chen PH, Lee CH, Ko AM, Lee Lin YC, Ho PS, Tu HP, Wu DC, Shieh TY, Ko YC. Up-regulation of inflammatory signalings by areca nut extract and role of cyclooxygenase-2 -1195G>A polymorphism reveal risk of oral cancer. *Cancer Res* 2008;68: 8489–98.
- Ferlay J, Bray F, Pisani P, Parkin DM. Globocan 2002, cancer incidence, mortality and prevalence worldwide, IARC, 2003. Lyon, France.

26. COA, Council of Agriculture, Executive Yuan, Republic Of China. Yearly report of Taiwan's agriculture: 1979–2007, Taipei, Council of Agriculture, Executive Yuan, Taiwan, ROC, 2009.
27. Zhang X, Li C, Liao Q, Reichart PA. Areca chewing in Xiangtan, Hunan province, China: interviews with chewers. *J Oral Pathol Med* 2008; 37:423–9.
28. [18](#) der Waal I. Potentially malignant disorders of the oral and oropharyngeal mucosa; terminology, classification and present concepts of management. *Oral Oncol* 2009;45:317–323.
29. [18](#) age G, Tilakaratne WM, Tavassoli M, [33](#) nakulasuriya S. Molecular markers in oral epithelial dysplasia: review. *J Oral Pathol Med* 2009;38:737–52.
30. [11](#) CF, Wang JD, Chen PH, Chang SJ, Yang YH, Ko YC. Predictors of betel quid chewing behavior and cessation patterns in Taiwan aborigines. *BMC Public Health* [11](#):6:271.
31. Ko YC, Chiang TA, Chang SJ, Hsieh SF. Prevalence of betel quid chewing habit in Taiwan and related sociodemographic factors. *J Oral Pathol Med* 1992;21: [68](#)–4.
32. Yap SF, Ho PS, Kuo HC, Yang YH. [35](#) Comparing factors affecting commencement and cessation of betel quid chewing behavior in Taiwanese adults. *BMC Public Health* 2008;8:199.
33. [13](#) CP, Tsai SP, Cheng TY, Chen CJ, Levy DT, Yang HJ, Eriksen MP. [28](#) overing the relation between betel quid chewing and cigarette smoking in Taiwan. *Tob Control* 2005;14 (Suppl 1):i16–22.

Intercountry prevalences and practices of betel-quid use in south, southeast and eastern Asia regions and associated oral preneoplastic disorders: an international collaborative study by Asian betel-q

ORIGINALITY REPORT

18%

SIMILARITY INDEX

10%

INTERNET SOURCES

16%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Chien-Hung Lee, Deng-Chyang Wu, Jang-Ming Lee, I-Chen Wu et al. "Anatomical subsite discrepancy in relation to the impact of the consumption of alcohol, tobacco and betel quid on esophageal cancer", International Journal of Cancer, 2007
Publication 1%
- 2 www.labome.org
Internet Source 1%
- 3 jamanetwork.com
Internet Source 1%
- 4 Ping-Ho Chen, Qaisar Mahmood, Gian Luigi Mariottini, Tai-An Chiang, Ka-Wo Lee. "Adverse Health Effects of Betel Quid and the Risk of Oral and Pharyngeal Cancers", BioMed Research International, 2017
Publication 1%
- 5 Submitted to Flinders University

Student Paper

1 %

6

www.aminer.org

Internet Source

1 %

7

ph.kmu.edu.tw

Internet Source

1 %

8

www.gcancer.org

Internet Source

1 %

9

Submitted to The University of Manchester

Student Paper

<1 %

10

hdl.handle.net

Internet Source

<1 %

11

smedph.kmu.edu.tw

Internet Source

<1 %

12

Nisha Thakur, Ravi Mehrotra. "Chapter 84-1
Carcinogenic Alkaloids Present in Areca Nut",
Springer Science and Business Media LLC,
2022

Publication

<1 %

13

bmcpublichealth.biomedcentral.com

Internet Source

<1 %

14

Albert Min-Shan Ko, Chien-Hung Lee, Allen
Min-Jen Ko, Ying-Chin Ko. "Betel quid
dependence mechanism and potential
cessation therapy", Progress in Neuro-

<1 %

Psychopharmacology and Biological Psychiatry, 2020

Publication

15 www.mdpi.com <1 %
Internet Source

16 www.searo.who.int <1 %
Internet Source

17 Ping-Ho Chen. "Monoamine oxidase A variants are associated with heavy betel quid use : MAO-A variations with betel", *Addiction Biology*, 05/2011 <1 %
Publication

18 open.library.ubc.ca <1 %
Internet Source

19 "Development of Oral Cancer", Springer Science and Business Media LLC, 2017 <1 %
Publication

20 atm.amegroups.com <1 %
Internet Source

21 gyan.iitg.ac.in <1 %
Internet Source

22 medios.economiayfinanzas.gob.bo <1 %
Internet Source

23 Andrew J. Castiglioni. "Expression and vesicular localization of mouse Trpm13 in stria vascularis, hair cells, and vomeronasal and <1 %

olfactory receptor neurons", The Journal of Comparative Neurology, 04/15/2011

Publication

24

Irene Tamí-Maury, Wei-Fen Ma, Mi-Ting Lin, Cheng-Chieh Lin et al. "A qualitative study of attitudes to and perceptions of betel quid consumption and its oral health implications in Taiwan", Community Dentistry and Oral Epidemiology, 2018

Publication

25

Lingyun Wei. "Cross-signaling among phosphoinositide-3 kinase, mitogen-activated protein kinase and sonic hedgehog pathways exists in esophageal cancer", International Journal of Cancer, 07/15/2011

Publication

26

Yogesh Dokhe, Sivakumar Vidhydharan, Krishnakumar Thankappan, Subramania Iyer. "Chapter 1 Epidemiology", Springer Science and Business Media LLC, 2021

Publication

27

docshare.tips

Internet Source

28

www.nature.com

Internet Source

29

"Areca nut and betel-quid chewing bibliography", Addiction Biology, 1/2002

Publication

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

30 Chien-Hung Lee. "The use of tobacco-free betel-quid in conjunction with alcohol/tobacco impacts early-onset age and carcinoma distribution for upper aerodigestive tract cancer : UADT cancer onset and substance use", Journal of Oral Pathology and Medicine, 10/2011 <1 %

Publication

31 Chin-Feng Lin, Jung-Der Wang, Ping-Ho Chen, Shun-Jen Chang, Yi-Hsin Yang, Ying-Chin Ko. "Predictors of betel quid chewing behavior and cessation patterns in Taiwan aborigines", BMC Public Health, 2006 <1 %

Publication

32 Shelly Arora, Christopher Squier. "Areca nut trade, globalisation and its health impact: perspectives from India and South-east Asia", Perspectives in Public Health, 2018 <1 %

Publication

33 Tien-En Chiang, Yu-Chun Lin, Chi-Tsung Wu, Sheng-Tang Wu, Yuan-Wu Chen. "Association Between Socioeconomic Status and Severity of Oral Epithelial Dysplasia Using a Taiwanese Nationwide Oral Mucosal Screening Program: A Retrospective Analysis", Research Square Platform LLC, 2021 <1 %

Publication

Submitted to University of Glasgow

<1 %

35

Wan MN Ghani, Ishak A Razak, Yi-Hsin Yang, Norain A Talib et al. "Factors affecting commencement and cessation of betel quid chewing behaviour in Malaysian adults", BMC Public Health, 2011

Publication

<1 %

36

mafiadoc.com

Internet Source

<1 %

37

"45th Annual SER Meeting Minneapolis, Minnesota June 27-30, 2012", American Journal of Epidemiology, 2012.

Publication

<1 %

38

A. Alcaide-Raya. "How well informed are our Internet savvy patients on oral premalignant disorders? : Quality of Internet patient information", Oral Surgery, 11/2010

Publication

<1 %

39

Pei-Shan Ho, Pai-Li Chen, Saman Warnakulasuriya, Tien-Yu Shieh, Yun-Kwan Chen, I-Yueh Huang. "Malignant transformation of oral potentially malignant disorders in males: a retrospective cohort study", BMC Cancer, 2009

Publication

<1 %

40

Ping-Ho Chen, Ka-Wo Lee, Chung-Ho Chen, Tien-Yu Shieh et al. "CYP26B1 is a novel candidate gene for betel quid-related oral squamous cell carcinoma", Oral Oncology, 2011

Publication

<1 %

41

Sunder Lokhande, Marewa Glover, Kyro Selket. "Chewing tobacco use among South - East Asian men in Auckland", International Journal of Migration, Health and Social Care, 2013

Publication

<1 %

42

fedetd.mis.nsysu.edu.tw

Internet Source

<1 %

43

Alameddine, I.. "Brine discharge from desalination plants: a modeling approach to an optimized outfall design", Desalination, 20070815

Publication

<1 %

44

Bmcpublikealth.Biomedcentral.Com

Internet Source

<1 %

45

Chien-Hung Lee, Jang-Ming Lee, Deng-Chyang Wu, Hon-Ki Hsu et al. "Independent and combined effects of alcohol intake, tobacco smoking and betel quid chewing on the risk of esophageal cancer in Taiwan", International Journal of Cancer, 2005

Publication

<1 %

46 Kanako Yoshikawa, Hiroyuki Yamada, Midori Matsushima. "Betel quid chewing and its associated factors in Myanmar: A quantitative study", International Journal of Drug Policy, 2020
Publication <1 %

47 Liu, C.C.. "The Impact of Cigarette Smoking, Alcohol Drinking and Betel Quid Chewing on the Risk of Calcium Urolithiasis", Annals of Epidemiology, 200908
Publication <1 %

48 Namrah Anwar, Shahid Pervez, Qurratulain Chundriger, Sohail Awan, Tariq Moatter, Tazeen Saeed Ali. "Oral cancer: Clinicopathological features and associated risk factors in a high risk population presenting to a major tertiary care center in Pakistan", PLOS ONE, 2020
Publication <1 %

49 www.oncotarget.com
Internet Source <1 %

50 A. Winstock. "Areca nut-abuse liability, dependence and public health", Addiction Biology, 1/2002
Publication <1 %

51 Chiang, C.P.. "Hamsters chewing betel quid or areca nut directly show a decrease in body <1 %

weight and survival rates with concomitant epithelial hyperplasia of cheek pouch", Oral Oncology, 200408

Publication

52

Ching-Sung Ho. "Factors related to betel chewing among junior high school students in Taiwan", Community Dentistry And Oral Epidemiology, 4/2000

Publication

<1 %

53

Guha, Neela, Saman Warnakulasuriya, Jelle Vlaanderen, and Kurt Straif. "Betel quid chewing and the risk of oral and oropharyngeal cancers: a meta-analysis with implications for cancer control", International Journal of Cancer, 2013.

Publication

<1 %

54

Hsin-Yi Yen, Ping- Ho Chen, Ying-Chin Ko, Shih-Kuang Chiang, Yevvon Yi-Chi Chang, Yung-Jong Shiah. "Betel Quid Chewing, Personality and Mood: Betel Quid Chewing Associated with Low Extraversion and Negative Mood", Substance Use & Misuse, 2018

Publication

<1 %

55

Pei-Shian Ho. "The incidence of oropharyngeal cancer in Taiwan: an endemic betel quid chewing area", Journal of Oral Pathology and Medicine, 4/2002

Publication

<1 %

56 Ping-Ho Chen, Ka-Wo Lee, Cheng-Chieh Hsu, Jeff Yi-Fu Chen et al. "Expression of a Splice Variant of CYP26B1 in Betel Quid-Related Oral Cancer", The Scientific World Journal, 2014

Publication

<1 %

57 R Botell. "Neurological rehabilitation of stroke", Journal of Neurology Neurosurgery & Psychiatry, 10/1/2005

Publication

<1 %

58 Zhang, X.. "A review of betel quid chewing, oral cancer and precancer in Mainland China", Oral Oncology, 200705

Publication

<1 %

59 annals.edu.sg

Internet Source

<1 %

60 archive.org

Internet Source

<1 %

61 escholarship.org

Internet Source

<1 %

62 www.rrdental.com.au

Internet Source

<1 %

63 C P Wen. "Uncovering the relation between betel quid chewing and cigarette smoking in Taiwan", Tobacco Control, 2005

Publication

<1 %

64

Che-Yi Lin, Tien-Szu Pan, Chun-Chan Ting, Shih-Shin Liang et al. "Cytochrome P450 Metabolism of Betel Quid-Derived Compounds: Implications for the Development of Prevention Strategies for Oral and Pharyngeal Cancers", *The Scientific World Journal*, 2013

Publication

<1 %

65

Chien-Hung Lee, Chiung-Yu Peng, Ruei-Nian Li, Yu-Chieh Chen et al. "Risk evaluation for the development of cervical intraepithelial neoplasia: Development and validation of risk-scoring schemes", *International Journal of Cancer*, 2015

Publication

<1 %

66

Hsiao-Ling Huang. "School-level contextual factors associated with betel quid chewing among schoolchildren in Taiwan", *Community Dentistry And Oral Epidemiology*, 02/2009

Publication

<1 %

67

U. Nair. "Alert for an epidemic of oral cancer due to use of the betel quid substitutes gutkha and pan masala: a review of agents and causative mechanisms", *Mutagenesis*, 07/01/2004

Publication

<1 %

68

biomedicine.cmu.edu.tw

Internet Source

<1 %

69

s3.amazonaws.com

Internet Source

<1 %

70

www.biomedcentral.com

Internet Source

<1 %

71

www.cellmolbiol.org

Internet Source

<1 %

72

Bharath Kumar Velmurugan, Kun-Tu Yeh, Chien-Hung Lee, Shu-Hui Lin et al. "Acidic leucine-rich nuclear phosphoprotein-32A (ANP32A) association with lymph node metastasis predicts poor survival in oral squamous cell carcinoma patients", *Oncotarget*, 2016

Publication

<1 %

73

Chi-Pin Lee, Shang-Lun Chiang, Chien-Hung Lee, Yi-Shan Tsai et al. "AURKA Phe31Ile polymorphism interacted with use of alcohol, betel quid, and cigarettes at multiplicative risk of oral cancer occurrence", *Clinical Oral Investigations*, 2015

Publication

<1 %

74

Kim, Hye Yeon, SiHyun Cho, Young Sik Choi, Hyo In Yang, Kyung Eun Lee, Seok Kyo Seo, and Byung Seok Lee. "Cyclooxygenase-2 (COX-2) Gene-765G/C Polymorphism and Advanced-Stage Endometriosis in Korean

<1 %

Women", American Journal Of Reproductive Immunology, 2012.

Publication

75

Wei-Chung Tsai, Ming-Tsang Wu, Guei-Jane Wang, Kun-Tai Lee et al. "Chewing areca nut increases the risk of coronary artery disease in taiwanese men: a case-control study", BMC Public Health, 2012

Publication

<1 %

76

"Abstracts of the 3rd North American Congress of Epidemiology, June 21-24, 2011 Montreal, Canada", American Journal of Epidemiology, 2011.

Publication

<1 %

77

Chien-Hung Lee, Albert Min-Shan Ko, Frances M. Yang, Chung-Chieh Hung et al. " Association of Betel-Quid Use Disorder With Oral Potentially Malignant Disorder in 6 Betel-Quid Endemic Asian Populations ", JAMA Psychiatry, 2018

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On