


Journal of the Ecological Society of America

ECOLOGY

ENVIRONMENT & CONSERVATION

ISSN 0014-1801



JOHN WILEY & SONS

Ecology, Environment and Conservation \Editorial Advisory Board

Chief Editor

Prof. (Dr.) R.K. Trivedy, Pune, India

Email id- rktrivedy@gmail.com

Tel: 91-20-46745119

Mobile No. -9975703363

EDITORIAL ADVISORY BOARD

1. Dr. Prof. Moses Inbaraj, Professor, Department of Botany (Retired), Madras Christian College, Chennai, India	Indiainbarajmoses2004@yahoo.com
2. Dr. Prof. D.J. Lee, Professor, Department of Chemical Engineering, National Taiwan University, Taiwan	djlee@ntu.edu.tw
3. Dr. T. Bahorun, Professor, Department of Biosciences, Mauritius University, Mauritius	tbahorun@uom.ac.mu
4. Dr. Prof. A.R. Ghosh, Professor, Department of Environmental Sciences, University of Burdwan, Burdwan, India	apurbaghosh2010@gmail.com
5. Dr. Prof. M. Zaman, Professor Department of Botany, Dhaka University, Dhaka, Bangladesh	zashim@du.ac.bd
6. Dr. Marcantonio Bragadin, Professor Department of Chemistry, University of Venice, Venice, Italy	bragadin@unive.it
7. Dr. Z. Fuat Topark, Professor Dicle University, Diyarbakir, Turkey	toprakzey@itu.edu.tr
8. Dr. Wilson S. Tisera, Associate Professor Christian University, Kupang, Indonesia	wilson_tisera@yahoo.com
9. Dr. Prof. Philip C. Reid, Scientist Marine Biological Association, U.K	pcre@mba.ac.uk
10. Dr. Prof. Mohd. Yusuf, Professor Deptt. of Environmental Engineering, University Sans Malaysia, Malaysia	danielmy@iium.edu.my
11. Dr. Prof. Azni Idris, Deputy Vice-Chancellor, Research and Innov., Univ. of Putra Malaysia, Malaysia	azni@upm.edu.my
12. Dr. Vikas Sharma, Associate Professor, RARS of SKUAST, Jammu, J&K, India	vsagron@gmail.com
13. Dr. A.C. Pandey, Senior Scientist, Birsa Agricultural University, Ranchi, India	azni@upm.edu.my
14. Dr. Prof. Agoes Soegianto, Professor, Dept. Biology, Airlangga Univ. Kampus C, Surabaya, Indonesia	agoes_soegianto@fst.unair.ac.id
15. Dr. Ahmed L. Mahmoudi, Professor, Water Studies Centre, King Faisal University, Saudi Arabia	mahmoudi@kfu.edu.sa
16. Dr. Siroop Chaudhuri, Associate Professor Center for Environment, Sustainability and Human	schaudhuri@jgu.edu.in

Development (CESH) O.P. Jindal Global University, Sonipat, India	
17. Dr. Amin L. Setyo, Professor, Graduate School of International Crop, Hiroshima University, Hiroshima, Japan	amin28@ub.ac.id
18. Dr. Francis Gbogbo, Associate Professor, Department of Animal Biology, University of Ghana, Ghana	fgbogbo@ug.edu.gh
19. Dr. Prof. Deenbandhu Sahoo, Professor Department of Botany, university of Delhi, Delhi, India	dbsahoo@yahoo.com
20. Dr. Prof. J.K. Ladha, Professor Emeritus University of California, Davis, California, U.S.A.	jkladha@ucdavis.edu
21. Dr. Prof. M.Z.M. Nomani, Professor Faculty of Law, Aligarh Muslim University, Aligarh, India	zafarnomani@rediffmail.com
22. Dr. G.R. Pathade, Principal (Retired) H.V. Desai College, Pune, India	girishpathade@yahoo.co.in
23. Dr. Alfred Maroyi, Associate Professor Department of Botany, University of Fort Hare, Alice 5700, South Africa	amaroyi@ufh.ac.za
24. Dr. Prof. A.V.L.N.S.H. Hariharan, Professor Dept. of Chemistry, College of Science, GITAM University, Visakhapatnam, India	ahharan@gmail.com
25. Dr. Zeinab El-Sayed Zayed, Associate Professor Department of Biotechnology, The Central Lab of Date Palm Researches and Development Agriculture Research Center, Cairo, Egypt	zemmz2005@yahoo.com
26. Dr. Raj Singh, Professor Department of Botany, MMD University, Mullana, Ambala, Haryana, India	dr.rajsingh09@gmail.com
27. Dr. Prof. Mohammed Latif Khan, Professor, Department of Botany Dr. Hari Singh Gour Vishwavidyalaya Sagar, India	khanml61@gmail.com
28. Dr. R. Dinesh Principal Scientist, ICAR-Indian Institute of Spices Research, Marikunnu, P.O. Kozhikode, India	rdinesh2005@gmail.com
29. Dr. Suresha Gududappa, Senior Environmental Scientist, Saudi Arabia	suresh@ebe.com.sa
30. Dr. Azma Hanim Ismail, School of Biological Sciences, Universiti Sains Malaysia, 11800 USM, Pulau Pinang, Malaysia.	

ECOLOGY, ENVIRONMENT AND CONSERVATION

VOL. 28 (3) : 2022

CONTENTS

- 1087–1093 Comparison of Occlusal Molar Tooth Width of the Etawa Crossbreed Goat
—*Soeharsono, Hana Elijani, Widjiati, Viski Fitri Hendrawan and Epy Muhammad Luqman*
- 1094–1097 A Survey on Fish Species at Tram Chim National Park, Vietnam
—*D. H. Truong, T. M. Duong and V. L. Loi*
- 1098–1102 Effect of soursop leaf extract solution on the death of armyworm (*Spodoptera litura* F.) on mustard plants
—*Akhmad Rizali, Noor Laili Aziza and Yusriadi Marsuni*
- 1103–1107 Teixobactin: A powerful antibiotic from natural resource
—*Parichat Phumkhachorn and Pongsak Rattanachaiakumsopon*
- 1108–1118 Impact of heavy metals and biochemical parameters on specific leaf area of roadside trees in Kathmandu, Nepal
—*Jaya Prakash Hamal and Mukesh Kumar Chettri*
- 1119–1124 Survey on the Diversity of Species Composition of Phytoplankton at Bung Binh Thien
—*D. H. Truong, T. M. Duong and V. L. Loi*
- 1125–1129 Effect of Dosage on Combination Ethylen Glycol and Dimethyl Sulfoxide as Cryoprotectant using Vitrification on Viability of Mice (*Mus musculus*) Blastocyst Post Warming
—*Zakiah Saumi, Viski Fitri Hendrawan, Widjiati and Epy Muhammad Luqman*
- 1130–1134 Domestic Waste Pollution of River Settlements, Banjarmasin City, Indonesia
—*Sidharta Adyatma and Muhammad Muhaimin*
- 1135–1138 Different Concentration Organic Carbon Waste Cassava in Mixotrophic Cultivation on Protein and Beta Carotene Content of *Dumaliella* sp.
—*Prive Widya Antika, Happy Nursyam and Arning Wilujeng Ekawati*
- 1139–1143 Aquatic Plants at an Agricultural Station, Vietnam
—*D. H. Truong, T. M. Duong and V. L. Loi*
- 1144–1152 The Effects of Formalin (Formyl Tetrahydrofolate) on CDK1 Protein on the Oocyte Maturation Process: Quantitative Structure Toxicity Relationships (QSAR) In Silico Analysis
—*Lutvia Krismayanti, Widjiati, Bambang Purwanto and Dwi Krihariyani*
- 1153–1156 Freshwater Fish Survey by Land Use Type in Cantho City, Vietnam
—*D. H. Truong, L. V. Ly, T. M. Duong and H. T. Phung*
- 1157–1170 Structuring the *Bruguiera gymnorhiza* Mangrove Area for Food Security in Paciran Village, Paciran District and Labuhan Village, Brondong District, Lamongan Regency, East Java
—*Rudianto and Dwi Setijawati*
- 1171–1177 Effects of Temperature and Different Feed Uptake on Mercury (Hg) and Cadmium (Cd) Accumulation in Nile Tilapia
—*Sunardi, Kabul Fadilah, Alfa Pratista Rakhmatul Fittri, Tiara Berliani and Keukeu Kaniawati Rosada*
- 1178–1181 Water Quality in Rice-shrimp Model
—*Lien, B.T.B., Truyen, D.M., Dan, T.H., Loi, L.V. and Nga, L.K.*
- 1182–1197 African Elephant (*Loxodonta africana*) Impact on Selected Tree Species and Assessment of Piosphere Gradients in Venetia-limpopo Nature Reserve
—*S.E. Nkosi, E. Adam, A.S. Barrett and L.R. Brown*
- 1198–1206 First record of heavy metal concentrations in the natural habitat of silkworms, Riau Province, Indonesia
—*B. Budijono, Muhammad Hasbi and Yudho Harjoyudanto*
- 1207–1215 Evaluation soil changes under agricultural management and irrigation in an arid region: the case of Ghardaïa (Zelfana)
—*Faredj Chikhi, Mohamed Benslama, Fouzi Benbrahim, Abderraouf Benslama, Sabrine Darem, Imane Hamel and Jose Navarro-Pedreño*
- 1216–1225 Sublethal Effects of Monocrotophos on Histological Architecture of Liver and Kidney of a Freshwater Fish, *Anabas testudineus*
—*B.T. Sulekha and T.V. Anna Mercy*

- 1226–1231 Experimental evaluation of the stability of chitosan based biocomposites under various environmental conditions
—*Wondalem Misganaw Golie and M.A. Azaz Ahmed*
- 1232–1245 A Modelling and Mitigation Strategy for the Impact of the Tsunami Disaster at Tamban Beach, Malang Regency, using a Geographic Information System
—*Rudianto, Maheno Sri Widodo, Guntur, Fahreza Okta Setyawan and Seftiawan Samsu Rijal*
- 1246–1251 Effect of Exposure to E-Cigarette Vapor and Cigarette Smoke on Seminiferous Tubules Diameter and Spermatozoa Quality of Mice (*Mus musculus*)
—*Muhammad Alif Zulfikar, Eka Pramytha Hestianah, Herry Agoes Hermadi, Suzanita Utama, Tatik Hernawati, Suryo Kuncorojakti and Epy Muhammad Luqman*
- 1252–1261 Radial growth dynamics in Scots pine forests of the Yalutorovsky forest district of Tyumen region
—*Epishkov A.A., Lipatkin V.A., Frolova V.A., Sidorenkov V. M., Stonozhenko L.V., Vorobyeva N.S. and Rummyantsev D.E.*
- 1262–1274 Rush fibers reinforced Adobe for green buildings
—*Wassef Ounaies*
- 1275–1280 Habitat Characteristics and Abundance of Goonch (*Bagarius yarelli*) in Perjaya Dam, Komerling River, South Sumatra, Indonesia
—*Sofian, Syaeful Anwar, Naim Pramono, Alfi Sahrin and Diah Ayu Satyari Utami*
- 1281–1287 A lizard diversity snapshot in the poorly studied Mila Region of north-eastern Algeria
—*Madaci Brahim, Derradj Lotfi and Benmerzouk Bilel*
- 1288–1292 An Introduction of Autoinducer Biosensor for *Vibrio parahemolyticus*, Pathogenic Bacteria in White Shrimp (*Litopenaeus vannamei*) Intensive Culture
—*Mohamad Fadjar, Sri Rahayu, Gumpita Windy Distiyanti and Jamilah Z. Janah*
- 1293–1300 The application of a Combination of Coconut Water and a Growth Regulator as (Effective Microorganisms) for Boosting Arabica Coffee Seedlings' Growth
—*Evy Latifah and Hanik Anggraeni Dewi*
- 1301–1308 Effect of Biological Pollution on Human Health in North East Algeria "Case of Seraidi-Annaba Region"
—*Imene Layachi and Mohamed Benslama*
- 1309–1315 The Effect of Temperature on *Goniastrea aspera* Polyp Degradation and the Post-bleaching Resistance
—*Pujiono Wahyu Purnomo, Max Rudolf Muskananfolo and Sigit Febrianto*
- 1316–1323 The Waste Management Paradigm Shift from Linear Economy to Circular Economy in Campuses and Urban Communities
—*Z. Ariani, N. Septyanun, R. R. Harun, Syawaluddin, F. Khairunnisa, L. M. I. Hadini, D. Suryati*
- 1324–1331 Biochar from waste verbena foumanat as an adsorbent to remove phenol from aqueous solution
—*Houari Bella, Hayet Bendaikha and Yasmina Bendaikha*
- 1332–1336 Impacts of Fertilizer and Pesticide consumptions on Natural resources of India
—*Avinash Sharma, Chowlani Manpoong, Hemant Sharma, Himanshu Pandey, Gaytri Soni, Supriya Sharma, Nirupa Kumari and Megha Raghavan*
- 1337–1344 Effect of Air Velocity, Type and Size of Biomass on Performance of Downdraft Gasifier
—*H. Manjunath, Vijayakumar Palled, M. Veerangouda, Devanand Maski and Ramachandra, C. T.*
- 1345–1356 Assessment of the effect of COVID-19 Lockdown on Water Quality of Baldi River Sahastradhara Sulphur Spring, Dehradun, Uttarakhand, India
—*Shail Kulshrestha*
- 1357–1366 The Relationship between water quality and Macrophyte biomass of Kongba River, Manipur, India
—*Beeteswari Khangembam and Asha Gupta*
- 1367–1375 Environmental Impact of Pesticide Usage in Cotton
—*Almaszabeen Badekhan and K. Uma Devi*
- 1376–1378 Oxygen Generation by DBH Trees - A Study inside RV College of Engineering (RVCE), Bengaluru, India
—*Vinod A.R., Prasanth Acharya and Poornima*
- 1379–1383 Effect of bio-formulations on growth and yield of Pea (*Pisum sativum* L.)
—*Michi Yarang, Chowlani Manpoong, Avinash Sharma, Megha Raghavan and V.S. Devadas*

- 1384–1389 Land Use and Land Cover Change Mapping: A Spatio Temporal and Correlational Analysis of Ramganjmandi Tehsil, Kota, Rajasthan, India
—*Arushi Rana*
- 1390–1394 Effect of herbicides on weed infestation, productivity of groundnut and their residual effect on sunflower
—*Shivanand N. Honnali and D.G. Satihal*
- 1395–1403 A Study on Water Quality and Macrophyte Diversity in Three Wetlands of Sambalpur District, Odisha from Ecosystem Management Perspective
—*Rajalaxmi Badra, Jijnasa Panda and Sunanda Sahoo*
- 1404–1410 Effect of Different Levels of Sulphur and Nitrogen on Growth Yield and Quality of Sesame (*Sesamum indicum* L.) under Bael (*Aegle marmelos* L.) based Agri-horti system
—*Shailesh Kushawaha, Alok Kumar Singh, Abhinav Singh, Bipin Kumar Singh and Srishti Dipriya Minz*
- 1411–1419 The role of tillage practices, nitrogen management by inorganic and organic manure and fertilizers and foliar applied of zinc on wheat (*Triticum aestivum* L.) crop performance and physico-chemical properties of soil under rainfed conditions
—*Manoj Kumar, Vikram Singh, Km. Nikam Kumari, Sagar Kumar, Thomas Abraham, Lipi Rina and J. J. Patel*
- 1420–1426 Study of Vitamin D₂ accumulation potential of two edible mushrooms from North-East India
—*Marium Begum, Subhankar Ghosh and Siddhartha Proteem Saikia*
- 1427–1430 Studies on Phytoplankton diversity of Barehalla tank, Shivamogga, Karnataka, India
—*H.A. Sayeswara and H.M. Ashashree*
- 1431–1445 A review on Immunostimulatory and antioxidant potential of herbs, *Curcuma longa* L., *Camellia sinensis* L. *Zingiber officinale* and *Allium sativum* Linn. in fish health: a sustainable approach for a healthy aquaculture
—*Moirangthem Kameshwar Singh, Dhriti Borah, Manash Pratim Dutta, Simi Gogoi, Chandopal Saikia, Susmita Sonowal and Saimoun Ken Manhai*
- 1446–1451 Fixed Bed Modelling Column Adsorption Studies by Activated Carbon for Removal of Fluoride
—*Keerthi B. Gurani and S.R. Mise*
- 1452–1456 Prediction of Smartphone Users in India using Trend Analysis Looking for Better Planning of E-waste
—*Shiv Kumar Singh Pundhir and Sandeep Kumar*
- 1457–1467 Toxicity of phosphamidon on the gills of a freshwater fish, *Anabas testudineus*
—*B.T. Sulekha and T.V. Anna Mercy*
- 1468–1472 Influence of bio-fertilizers and growth regulators for enhancing yield and yield attributes of Indian mustard (*Brassica juncea* L.) grown under rainfed condition
—*Lipi Rina, Joy Dawson, Manoj Kumar, Gautam Ghosh, Liyir Rina and Bhura Singh*
- 1473–1477 Biocompatible Herbal Polymeric Nano-formulation of [6]-Gingerol: Development, Optimisation, and Characterization
—*Virendra Singh, Himanshu Pandey, Vatsala Misra and Devendra Singh*
- 1478–1484 Development and Quality Assessment of Probiotic Ice-cream using Stevia as Sweetener
—*G. Nikhitha, P. Abhiram, Md. Haneef Azad and B. Amala*
- 1485–1488 Some of the Immediate Damages and Preliminary Observations on the 22 June, 2020 Champhai Earthquake, India
—*Malsawmtluanga, P.C. Zoramthara, F. Zodinthara and J.L. Pachau*
- 1489–1493 Various Causes of Urban Heat Islands, effects and their Mitigation Measures for Urban Ecology – A Review
—*Aneesa S. and Reshma J.K.*
- 1494–1497 Stage Specific food consumption and utilization by the Mexican beetle, *Zygodontia bicolorata* Pallister (Coleoptera: Chrysomelidae) on *Parthenium hysterophorus* Linnaeus
—*Amrit Mohapatra, Satya Narayan Satapathy, Dibyajyoti Swain, Soumya Sree Panda, Shrivasini Parida, Priyanka Priyadarshini and Subhashree Subhasmita Paikaray*
- 1498–1506 Efficacy appraisal of Cobalt oxide electrode module as a patron of sustainable energy harvesting super capacitor
—*Shubhangi Gavande, Shivani Gavande and B. R. Karche*

- 1507–1515 Effect of potassium nitrate on germination physiology of magnesium nitrate primed and non-primed rice seeds var. Swarna
—S. Mondal and B. Bose
- 1516–1525 Evaluation of Heavy Metals in Vegetables and their Associated Health Risk through the consumption of Vegetables in the National Capital Region, India
—Jyoti Rani, Tripti Agarwal and Sudesh Chaudhary
- 1526–1535 Tree Diversity and Timber Resources Richness of Reserve Forests under Pasighat Forest Division, Arunachal Pradesh, India
—K. Ch. Momin, N. Lyngdoh, K. Upadhyaya and B. D. Mongjam
- 1536–1538 Comparative Study of DO, BOD and COD of Indrayani River Water, Pune (Maharashtra)
—S.D. Jadhav and M.S. Jadhav
- 1539–1543 Key Information Sources and their Utility for Upliftment of Members of Tribal Farmer Producer Groups
—Katiki Srikar and M. Asokhan
- 1544–1560 Kinetic Indices of Enzyme Activity used as Sensitive Biomarkers to Monitor Mine Spoil Genesis in Chronosequence Coal Mine Overburden Spoil
—Payal Agrawal, Jitesh Kumar Maharana and Amiya Kumar Patel
- 1561–1565 Effect of Rice Establishment Methods and Nutrient Management Practices on Growth, Yield and Economics of Rice (*Oryza sativa* L.)
—Jaya Bharti, Ashok Kumar Singh, C. S. Singh, Arvind Kumar Singh and Ranveer Kumar
- 1566–1570 Influence of renowned traditional liquid organic manure in conjunction with bacterial consortia on growth and yield of foxtail millet
—Veerendra Manduri, Padmaja Bhimireddy, Malla Reddy Maduri and Triveni Sodimalla
- 1571–1575 Investigation on Mechanized Harvesting Requirements of Pokkali Paddy for Optimizing Harvester Design
—Rathinavel S., Sindhu Bhaskar, Manoj Mathew and Jayan P.R.
- 1576–1583 Water Quality Assessment of River Kuwano, Basti (U.P.), using WQI and Pollution Indices
—Gopal Ji Kushwaha, Shivendra Mohan Pandey and Pradip Kumar
- 1584–1594 Eco-biological interaction of Polyphenols among *Acanthus ilicifolius*, detritus, and *Uca lactea annulipes* in estuarine-ecosystem
—Munmun Ghosh, Santi M. Mandal and Susanta K. Chakraborty
- 1595–1610 Municipal Solid Waste Model Development through Binary Coding
—Das Lalit, Das Adyasha and Mishra Sitikantha
- 1611–1618 Biobriquette: A Mixture of Palm Kernel Shell and Coconut Shell, An Indonesian Study Case
—Efri Mardawati, Aldita Khoiril R amadhan, Agus Kusnayat, Irfan Ardiansah and Hana Nur Fitriana
- 1619–1623 Impact of differential substitution of nutrients through organics on yield, nutrient uptake and economics of Basmati rice in irrigated Shivalik foothills of J&K, India
—Monika Banotra, B.C. Sharma, Rakesh Kumar and Rohit Sharma
- 1624–1627 Spatial Pattern Analysis on Landslide Incidents in Kuala Lumpur, Malaysia
—Nuriah Abd Majid, Syaidatul Azwani Zulkafli, Sharifah Zarina Syed Zakaria, Muhammad Rizal Razman and Minhaz Farid Ahmed
- 1628–1635 Endocrine Disruption and Physiological Alteration in Fish Exposed to Hexavalent Chromium
—Rashmita Behera, Bidut Prava Mohanty, Malaya Ranjan Mahananda and Samikshya Mishra
- 1636–1643 The Ganga River Water Pollution Status in India Characterize with River Gomti
—Deepika Singh, A.K. Shukla, Shraddha Yadav, Govind Pandey and V. Dutta
- 1644–1646 Diversity and seasonality of rotifers in a lower Shiwalik lake, India
—Rajesh Kumar, Romana Akthar and Supreet Kour
- 1647–1659 Fish Gills as a Tool for the Assessment of Sublethal Effects of Monocrotophos on a Freshwater Fish, *Etroplus maculatus*
—B.T. Sulekha and T.V. Anna Mercy
- 1660–1669 Hazard Study of Earthquake and Liquefaction in Correctional Institution Palu, Central Sulawesi
—Dina Ruslanjari, Rahmat Hidayat Made, Muhammad Ulin Nuha and Inayah Bastin Al Hakim
- 1670–1672 Impact of *Aloe vera* gel, Ashwagandha, Brahmi, and *Cassia tora* plant extract on lifespan and fecundity in the model organism *Drosophila melanogaster*
—Usha Rani, Manvender Singh and Krishan Kuamr Selwal

Effect of Dosage on Combination Ethylen Glycol and Dimethyl Sulfoxide as Cryoprotectant using Vitrification on Viability of Mice (*Mus musculus*) Blastocyst Post Warming

Zakiah Saumi¹, Viski Fitri Hendrawan², Widjiati¹, Epy Muhammad Luqman^{1*}

¹Departement of Veterinary Science Faculty of Veterinary Medicine Universitas Airlangga Kampus C Unair Jalan Mulyorejo, Surabaya 60115 Indonesia

²Department of Animal Reproduction Faculty of Veterinary Medicine, Universitas Brawijaya, Malang 65151, Indonesia

(Received 2 September, 2021; Accepted 30 October, 2021)

ABSTRACT

Cryopreservation is a method of reproductive technology that is useful for embryo and preservation creatures that are almost extinct. Cryoprotectants are often used for vitrification are PROH, DMSO and EG because they have good and efficient results. The study objective was to know does the viability of mice blastocyst increase with the combination of intracellular cryoprotectant ethylene glycol (EG) and dimethyl sulfoxide (DMSO) in vitrification. This study used mice oocytes then carried out in vitro fertilization and cultured until reaching the blastocyst stage. The mice were superovulated using 5 IU PMSG and 5 IU hCG after 48 hours, then monomating using infertile males. Egg cells are taken by tearing the fertilization bag and washing using MEM. Embryo is cultured so that it reaches the blastocyst phase and is then divided into 4 groups: vitrified using 30% EG, 30% DMSO, 20% combination EG + 10% DMSO and 10% P4 combination EG and 20% DMSO. The four groups were inserted into the hemi straw which has been modified with 0.25 berurn and then put back into the hemi straw with a size of 0.5. Hemi Straw which already contained embryos was inserted into a liquid N₂ container with a temperature of -196. Warming was done after the embryo was vitrified for a week. After being removed from the container, heating is done at room temperature. The embryo hemi straw was inserted into the sucrose level alternately, i.e. 0.25 M, 0.5 M and 1M respectively for 2 minutes. Warming process to remove cryoprotectants that are in the embryo. Examination of the viability of blastocyst was done using an inverted microscope. The results of this study that there were significant differences between groups using combination cryoprotectant EG 20% and DMSO 10% ($p < 0.05$) with 84% viability. The combination of EG and DMSO could increase the viability of mice blastocyst. The ethylene glycol 20% and dimethyl sulfoxide 10% was the best concentration combination based on this research.

Keywords : Food production, Vitrification, Ethylene glycol, Dimethyl sulfoxide, Blastocyst.

Introduction

Mice are one of the experimental animals that are

widely used for experiments because the gene structure is similar to humans and is easy to obtain. Cryopreservation is a method of reproductive tech-

nology that is useful for embryo and preservation creatures that are almost extinct (Marcao, 2010). Cryopreservation has good results in various studies and one of the most widely used is vitrification. Vitrification is a fast freezing embryo at -196°C without forming ice crystals (FAO, 2012). The advantage of vitrification is that it does not form ice crystals which can be deadly to cells (Gao and Critser, 2000). In vitrification method of cell or tissue incubated using cryoprotectant that has high concentration (Fay and Wowk, 2015). There are many studies using cryoprotectant combinations, but still showing low results, decreased embryo viability often occurs due to the warming process, the temperature changes that cause damage to cells (Pamungkas, 2010) but also because cryoprotectants cannot reduce the osmotic shock during vitrification (Pamungkas, 2010).

Cryoprotectants are often used for vitrification are PROH, DMSO and EG because they have good and efficient results (nateghi, 2017 HP). Dimethyl Sulfoxide has a molecular weight 78.13 g/mol, EG has 70 g/mol and PROH has 76.06 g/mol. Cryoprotectant size and density have different effects on various species (19 categories). Cryoprotectants are toxic, therefore their use must be done properly, according to Kartberg, many studies using a combination of EG and DMSO showing good results. Research publication show that vitrification with different cryoprotectant solutions is possible for human and mammal embryo, such as DMSO combination with ethylene glycol (EG), or EG alone (Rama *et al.*, 2005; Takahashi *et al.*, 2005). The combination of DMSO and EG has been used successfully by Desai *et al.* (2007). The ultrarapid freezing and vitrification most commonly employed for mouse cryopreservation. The ultrarapid method provides dehydration similar to the obtained with the conventional curve, even without cooling, allowing embryos to be plunged directly into membrane plasma, In previous studies (Mezzalira, 1991) freezing of mice morulae showing good developmental rates (83.6 to 92.9% of blasto-

cysts) were obtained with the ultra-rapid, using 30% glycerol. Another research showing good result using morulae stage with viability 80% (Cseh *et al.*, 1997). This research is to know the viability of mice blastosis were vitrified using a combination of cryoprotectant EG and DMSO.

Materials and Methods

This study use mouse oocytes then carried out in vitro fertilization and cultured until reach the blastocyst stage. Female mice were superovulated using 5 IU PMSG and 5 IU hCG after 48 hours, then monomating using infertile males. The next day a vaginal plug is checked to ensure marriage. Female mice were then sacrificed with cervical os dislocation. Egg cells are taken by tearing the fertilization bag and washed using MEM. Embryo is cultured using MEM under mineral oil and stored in a 5% CO_2 incubator. Embryo is cultured so that it reaches the blastocyst phase and is then divided into 4 groups. The first group was vitrified using 30% EG, 30% DMSO, 20% combination EG + 10% DMSO and 10% combination EG and 20% DMSO. The four groups were inserted into the hemi straw which has been modified with 0.25 berurn and then put back into the hemi straw with a size of 0.5. Hemi Straw which already contained embryos was inserted into liquid N_2 container with a temperature of -196 .

Warming is done after the embryo is vitrified for a week. After being removed from the container, heating is done at room temperature. The embryo hemi straw was inserted into the sucrose level alternately, i.e. 0.25 M, 0.5 M and 1M respectively for 2 minutes. Warming process to remove cryoprotectants that are in the embryo. Blastosis that has gone through a warming process is observed under an inverted microscope.

Results and Discussion

The results obtained after observing blastosis under an inverted microscope all groups survived with

Table 1. The mean and standard deviation of the amount of blastocyst between treatment groups.

Treatment	Percentage of Viability
Vitrified using 30% EG	60.00 ^{ab} ± 13.693
Vitrified using 30% DMSO	67.00 ^{ab} ± 9.747
Vitrified 20% combination EG + 10% DMSO and 10%	84.00 ^a ± 4.00
Vitrified combination EG and 20% DMSO	42.00 ^b ± 2.00

different viability. In groups with the highest viability, group of Vitrified 20% combination EG + 10% DMSO and 10% (84%) and significantly different from group of Vitrified using 30% EG (60%) $p < 0.05$. the lowest result was in group of Vitrified combination EG and 20% DMSO with only 42% viability and significantly different from group of Vitrified using 30% DMSO (67%) $p < 0.05$. from the results obtained group of Vitrified 20% combination EG + 10% DMSO and 10% has the best results and group of Vitrified combination EG and 20% DMSO has the worst results.

Vitrification has different successes in each study, in the process of successful vitrification by optimizing each stage of the vitrification process, namely at the stage of oocyte collection, sperm retrieval, in vitro fertilization, clotting and warming processes. Vitrification uses cryoprotectant as a supporting material that has high toxicity, so the use and time of cryoprotectant exposure are important. Blastocyst viability can be determined by microscopic morphological examination that is not found in the pelucid degeneration zone, the blastosol is large and the shape of the blastocyst is compact. In accordance with the study of Lieberman who used 20% EG and 20% DMSO and produced a viability of 84.5%.

Lieberman's results are not much different from the group of Vitrified 20% combination EG + 10% DMSO and 10%. Recent publication shows most successful vitrification method with combination cryoprotectant using EG and DMSO each 15% concentration (Matsui *et al.*, 1995; Selman *et al.*, 2006; Desai *et al.*, 2007).

Ethylene glycol is a cryoprotectant which has a low molecular weight and high penetration power, as well as a low level of toxicity (nateghi). Dimethyl sulfoxide has a molecular weight that is not much different from EG, besides that DMSO is relatively polar and has a compact and stable structure so that to enter the cell does not cause significant damage (Kartberg, 2008). The cryoprotectant combination of EG and DMSO eliminates the deficiencies of each cryoprotectant, according to Kartberg (2008) who said the use of cryoprotectant combinants EG and DMSO causes low toxicity, because EG reduces toxicity from DMSO as well as DMSO makes penetration of this combination more stable due to more DMSO structures compact compared to EG.

According to Mukaida *et al.*, (2003) the process of vitrification and warming occurs prior to adjusting the culture medium, the effect of cryoprotectant toxin, the occurrence of physical damage due to




Fig. 1. Blastocyst cells. 1: blastocyst cells before vitrification (magnification 100x); 2: blastocyst cells after vitrification (100x magnification) there are dead blastocysts (signs); 3: blastocyst cells after vitrification (400x magnification), clearly visible pelucid zones (black arrows), large blastosols (red arrows); 4: blastocyst cells die (400x magnification), the pelucid zone looks degenerated (black arrow) and irregular shape 2: blastocysts expand

crystalline ice formation during clotting, osmotic stress during cryoprotectant release of cells during warming, and lack of embryo quality good. Therefore cell death in vitrification or warming process is determined by many influencing factors. In the group of vitrified combination EG and 20% DMSO, the results of blastocyst cell viability after warming were 33%, this was very significant ($P < 0.05$) with group of Vitrified 20% combination EG + 10% DMSO and 10%. The P4 group was a combination of 10% cryoprotectant EG and 20% DMSO. The combination of group of Vitrified combination EG and 20% DMSO showed a decrease in post-warming blastocyst cell viability, this happened because the DMSO concentration was 20% so that cryoprotectant had a high toxicity effect, 10% EG had a less stable structure when penetrating, so in the group of EG 10 combination % and 20% DMSO can reduce post-warming viability.

The results obtained in this study with 30% EG cryoprotectants had good results, 60% of these results were very significantly different ($P < 0.05$) with group of vitrified using 30% DMSO with 66% viability. Cryoprotectant use in the (Vitrified using 30% EG) and (Vitrified using 30% DMSO) groups gave viability presentations above 50%, this proves that the use of a single cryoprotectant is able to provide good results, blastocyst cell death can be caused due to the nature of each cryoprotectant.

Conclusion

The conclusion of this research is combination cryoprotectant DMSO and EG can increase the viability of mice blastocyst. Because the bad characteristic each cryoprotectant, the other concentration of combination EG and DMSO decrease the viability of mice blastocyst and the most suitable and show good result is combination cryoprotectant EG 20% + DMSO 10%.

Acknowledgments

The authors are grateful to the authorities Faculty of Veterinary Medicine Universitas Airlangga, Surabaya, East Java, Indonesia.

Authors' contributions

Research concept and design: Zakiah Saumi, Collection and/or assembly of data: Epy Muhammad Luqman, Data analysis and interpretation: Viski Fitri Hendrawan, Writing the article: Zakiah Saumi,

Widjiati; Critical revision of the article: Epy Muhammad Luqman; Final approval of the article: Zakiah Saumi, Epy Muhammad Luqman.

Competing interests

The authors declare that there is no conflict of interest.

References

- Cseh, S., Corselli, J., Nehlsen-Cannarella, S. L., Bailey, L. L. and Szalay, A. A. 1997. The effect of quick-freezing in ethylene glycol on morphological survival and *in vitro* development of mouse embryos frozen at different pre-implantation stages. *Theriogenology*. 48(1): 43-50.
- Desai, N., Blackmon, H. and Szeptycki, J. 2007. Cryoloopvitrification of human day 3 cleavage-stage embryos: post-vitrification development, pregnancy outcomes and live births. *Reprod Bio. Med. Online*. 14 : 208-213.
- Fahy, G. M. and Wowk, B. 2015. Principles of cryopreservation by vitrification. *Methods Mol Biol*. 1257: 21-82.
- Food and Agriculture Organization. 2012. Cryoconservation of animal genetic resource. FAO Animal Production and Health Guidelines 12.
- Gao, D. and Critser, J. K. 2000. Mechanisms of cryoinjury in living cells. *ILAR Journal*. 41 : 187-196.
- Kartberg, A. J., Hambiliki, F., Arvidsson, T., Stavreus, A. and Svalander, P. 2008. Vitrification with DMSO protects embryo membrane integrity better than solution without DMSO. *Reprod Biomed*. 17(3) : 378-384.
- Liebermann, J. and Tucker, M. J. 2002. Effect of carrier system on the yield of human oocytes and embryo as assessed by survival and development potential after vitrification. *Reproduction*. 124: 483-489.
- Liebermann, J., Nawroth, F., Isachenko, V., Isachenko, E., Rahimi, G., Tucker, M. J. 2002. Potential Importance of vitrification in reproductive medicine. *Biol Reprod*. 67 : 1 71-1680.
- Matsui, M., Takahashi, Y. and Hishinuma, M. 1995. Insulin and insulin-like growth factor-I (IGF-I) stimulate the development of bovine embryos fertilized in vitro. *J Vet Med Sci*. 57 : 1109-1111.
- Mezzalana, A. and Rubin, M. I. B. 1991. Effects of Pre Treatment Ano Lactose Concentration on Ultra Rapid Freezing Mouse Morulae. *Cienc Rural*. 21(2): 243-248.
- Mukaida, T., Nakamura, S., Tomiyama, T., Wada, S., Oka, C., Kasai, M. and Takahashi, K. 2003. Vitrification of human blastocysts using cryoloops: clinical outcome of 223 cycles. *Hum Reprod* 18: 384-391.
- Pamungkas, F. A. 2010. Utilization of Vitrification Method for Cryopreservation of Mammal Oocytes. *WARTAZOA*. 20(3).

- Rama, R. G. A., Haranath, G. B. and Krishna, K. M. 2005. Vitrification of human 8-cell embryos, a modified protocol for better pregnancy rates. *Reprod BioMed. Online*. 11: 434–437.
- Selman, H., Angelini, A. and Barnocchi, N. 2006. Ongoing pregnancies after vitrification of human oocytes using a combined solution of ethylene glycol and dimethyl sulfoxide. *Fertil Steril*. 86 : 997–1000.
- Takahashi, K., Mukaida, T. and Goto, T. 2005. Perinatal outcome of blastocyst transfer with vitrification using cryoloop: a 4-year follow-up study. *Fertil Steril* 84: 88–92.