

# Brain Cells Death on Infant Mice (Mus musculus) Caused by Carbofuran Exposure During the Lactation Period

*by* Epy Muhammad Luqman

---

**Submission date:** 20-Dec-2022 11:35AM (UTC+0800)

**Submission ID:** 1984834235

**File name:** C-15\_Brain\_Cells\_Death\_on\_Infant\_Mice\_Mus\_musculus.pdf (1.19M)

**Word count:** 6441

**Character count:** 34908

## Brain Cells Death on Infant Mice (*Mus musculus*) Caused by Carbofuran Exposure During the Lactation Period

Epy Muhammad LUQMAN <sup>1</sup> ✍️ Widjiati WIDJIATI <sup>1</sup> Lita Rakhma YUSTINASARI <sup>1</sup>

<sup>1</sup>Veterinary Anatomy Department, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, INDONESIA

Article Code: KVFD-2018-20045 Received: 03.05.2018 Accepted: 25.09.2018 Published Online: 26.09.2018

### How to Cite This Article

Luqman EM, Widjiati W, Yustinasari LR: Brain cells death on infant mice (*Mus musculus*) caused by carbofuran exposure during the lactation period. *Kafkas Univ Vet Fak Derg*, 24 (6): 845-852, 2018. DOI: 10.9775/kvfd.2018.20045

### Abstract

An exposure to insecticide carbofuran has been reported able to generate a reactive oxygen species (ROS) in mice brains. This study was undertaken to evaluate the oxidative damage, biochemical and histopathological alterations by respectively examining malondialdehyde (MDA), cholinesterase (ChE) levels, necrosis and apoptosis in the suckling mice whose mothers were exposed to the carbofuran. The carbofuran was exposed via an oral route at the doses of 0.0208 mg/kg and 0.0417 mg/kg BW from the first until the fourth postnatal day after the delivery (n=27). The six-day-old pups were examined for its brain's MDA and ChE levels as well as the necrotic and apoptotic Purkinje cell were counted using the Tunel assay and hematoxylin-eosin (HE) staining. The mothers' exposure to carbofuran caused an increase in MDA levels, necrosis of Purkinje cells and a decrease in ChE, but there was no significant apoptosis in lactating pups. Carbofuran altered the level of the marker parameters related to the MDA, ChE and necrosis of Purkinje cells. Consistent changes were found in MDA, ChE and necrosis of Purkinje cells of the subjected pups, especially between the control and treatment groups, and there was no change between treatments. In conclusion, the transfer of carbofuran intoxication through the mother's milk resulted in the oxidative stress, biochemical and histopathological alterations in the suckling pups.

**Keywords:** Carbofuran, Purkinje cells, Infant mice brain, Lactation

## Laktasyon Döneminde Karbofurana Maruz Kalan Yavru Farelerde (*Mus musculus*) Beyin Hücreleri Ölümü

### Öz

İnsektisit karbofurana maruz kalmanın farelerin beyinlerinde reaktif oksijen türleri (ROS) ürettiği bildirilmiştir. Bu çalışma, anneleri karbofurana maruz bırakılan ve anne sütü emen farelerde malondialdehid (MDA), kolinesteraz (ChE), nekroz ve apoptozis düzeyleri incelenerek oluşan oksidatif hasar ile biyokimyasal ve histopatolojik değişiklikleri değerlendirmek amacıyla yapılmıştır. Fareler (n=27) karbofurana, oral yoldan, doğumdan sonraki 1. günden 4. güne kadar 0.0208 mg/kg ve 0.0417 mg/kg BW dozlarında maruz bırakıldı. Altı günlük yavruların beyin MDA ve ChE düzeyleri yanı sıra, Tunel testi ve hematoksilin-eozin (HE) boyaması kullanılarak nekrotik ve apoptotik Purkinje hücre sayısı incelendi. Annelerin karbofurana maruz kalması, MDA düzeylerinde artışa, Purkinje hücrelerinin nekrozuna ve ChE'de azalmaya neden oldu, ancak emen yavrularda belirgin bir apoptoz gözlenmedi. Özellikle kontrol ve tedaviye tabi tutulan yavruların MDA, ChE ve Purkinje hücrelerinin nekrozunda tutarlı değişiklikler bulunmuş, tedavi grupları arasında ise herhangi bir değişiklik olmadığı tespit edilmiştir. Sonuç olarak, anne sütü yoluyla karbofuran zehirlenmesi, emen yavrularda oksidatif stres, biyokimyasal ve histopatolojik değişiklikler ile sonuçlanmıştır.

**Anahtar sözcükler:** Karbofuran, Purkinje hücreleri, Yavru fare beyni, Laktasyon

## INTRODUCTION

Insecticide carbofuran residues in food may be harmful to organisms which are actually not the target of insecticide itself <sup>[1]</sup>. In a flower plantation which was contaminated by carbofuran in Ecuador in 2001, there were several cases of babies born with abnormalities, such as declining reflexes and motoric skill. At the child stage, there were

some brain function developmental abnormalities, such as the degeneration of memorizing and concentrating abilities <sup>[2]</sup>. In tested animals, carbofuran contamination causes an oxidative stress and weakens the motoric, memory, and cognitive functions <sup>[3]</sup>. Like organophosphate, carbofuran <sup>[38]</sup> inductions result in a significant oxidative damage in the cerebral cortex, cerebellum, and brain-stem <sup>[4]</sup>. Carbofuran inductions in the cerebral cortex



### İletişim (Correspondence)



+62 31 5992785 Fax: +62 31 5993015



epy-m-l@fkh.unair.ac.id

strongly correlate to the decline of cognitive and motoric functions [3].

An oral administration of carbofuran has been proven to strongly stimulate the reactive oxygen species (ROS) in mice brains and increase the levels of malondialdehyde (MDA) [3]. Intraperitoneal sub-acute administrations of carbofuran increase the brain's oxidative stress as the dose increases; and thus, improve MDA levels significant [22]. The increase of oxidative stress induces the activity of antioxidant enzymes, such as superoxide dismutase (SOD) and catalase in the brain [5]. The presence of ROS could trigger the formation of hydroxyl radicals (OH\*) which breaks the DNA chains or changes the composition of nucleotides in DNA creating mutations and apoptosis [6]. Hydroxyl radicals (OH\*) as a result of an oxidative stress could also damage all membrane systems in the cell, such as creating leaks in the lysosome membrane which may cause the cell death (necrosis).

Uncontrolled increase in ROS causes the injury and death of neuron cells [7]. Cerebrum consists of 80% neuron cells and of 20% glial cells which are responsible for transmitting information to the spinal cord to control motoric functions [8]. During the embryonal stage of brain development, neuron cells develop earlier and reach their developmental peak at mid-pregnancy. Glial cells develop at mid-pregnancy until a few days before the fetus is born and reach the peak of development at the end of pregnancy [9]. The development of the cerebellum starts in the late period of pregnancy and develops at maximum (peak) at the beginning of birth (the beginning of lactation). The death of neuron cells, cerebral glial cells and Purkinje cells due to carbofuran exposure potentially degenerates reflexes and motoric functions.

A deeper understanding of the brain's cell death mechanism in mice pups due to carbofuran exposure during the lactation period is necessary to acquire the basic treatment and prevention during the lactation period. Besides, it is important to understand the brain's cell death mechanism in order to identify the most sensitive period as well as the type of cells affected by carbofuran exposure during the lactation period. If the mechanism and the type of dead cells are recognized, actions can be taken to prevent the degeneration of reflexes and motoric skills in infant mice.

The aim of this research was to determine the brain's cell death mechanism in mice pups whose mother were exposed to carbofuran during the lactation period by measuring the MDA levels as an indicator of ROS production, ChE levels as the indicator of a neural function response to carbofuran exposure, apoptotic and necrotic cerebral Purkinje cells. This study contributed in disclosing the prevention mechanism of brain's cell death in mice pups whose mother were exposed to carbofuran during the lactation period. In addition, this study also provided a scientific information of insecticide carbofuran exposure,

especially during the lactation period related to the attempts of inhibiting brain's development disorders.

## 10 MATERIAL and METHODS

### Ethics Approval

The study was approved by the Faculty of Veterinary Medicine's Animal Ethics Committee. All variables had been considered in accordance with the Ethics Committee related to the animal handling to ensure no discomfort or pain caused to the animals during sampling (2011/111-KE).

### 40 Laboratory Animals

The animals used in this study were 27 female mice (*Mus musculus*), 10 weeks old, with the weight range of 25-30 grams, and 12-weeks-old male mice. Environmental adaptation was done to female mice (*Mus musculus*) for 7 days. On the 8<sup>th</sup> day, pregnant mare serum gonadotropin (PMSG) with a dosage of 5 IU/mouse was injected into the female mice and followed by Human Chorionic Gonadotrophin (HCG) injections with a dosage of 5 IU/mouse which was performed on the 10<sup>th</sup> day. Afterwards, the female mice were mated with the 12-week-old male mice. On the 11<sup>th</sup> day, a gestation examination was carried out. The gestation of female mice was indicated by the visible mating plug covering the female mice vulva; and then the day was considered as the first day of gestation [10].

### Carbofuran Exposure

Carbofuran exposure was targeted at the suckling mice's brain and this study examined the Purkinje cells in the cerebellum. The female mice were exposed to carbofuran with a dosage of 0.0208 mg/kg (1/24 LD<sub>50</sub>) and 0.0417 mg/kg (1/12 LD<sub>50</sub>) [10] on the 1<sup>st</sup> to 4<sup>th</sup> day of the lactation period given orally using a sterile disposable syringe. The six-day-old mice pups were then tested and measured for MDA and Cholinesterase (ChE) levels and histopathologic preparations were made. A microscopic examination was conducted to estimate the number of cells experiencing necrosis and apoptosis by using HE staining and an Apoptag Apoptosis Detection Kit.

### Measuring Cholinesterase (ChE) Levels

Cholinesterase (ChE) was determined according to the manufacturer's instructions of Cholinesterase FS (DiaSys Diagnostic Systems, 11401) [11]. To measure the ChE levels on the six-day-old mice's cerebrospinal fluid, the following materials were used to create a substrate: S-Butyrylthiocholine iodide Phosphate buffer pH 7.7, 5.5 dithiobis-2-nitrobenzoate. The principle of ChE level measurement was that the process of S-Butyrylthiocholine iodide + H<sub>2</sub>O hydrolysis and with the help of ChE sample was converted to Thiocholine iodide + butyrate. Thiocholine iodide + 5,5-dithiobis-2-nitro benzoate would transform into 5-Mercapto-2-nitro benzoate - 5 Mercaptothiocholine. The

reaction solution is from Diluent 100 mL, 49 nL reagent mix (for 30 samples) with concentrations of Phosphatase buffer pH 7.7 50 mmol/L, S-Butyrylthiocholine iodide 6 mmol/L, 5,5-dithiobis-2-nitrobenzoic 0.25 mmol/L. The standard wavelength of reaction solution was 405 nm and the length of translucent light was 1 cm measured at 25, 35 and 37°C. The 10 mL of sampled fluid was collected and mixed with 100 mL of reaction solution. The mixture was examined every minute for three minutes. To determine the sampled concentration, the enzyme activity of sample (U/L) =  $\Delta A/\text{min} \times 68500$  was measured.

#### Measuring Malondialdehyde (MDA) Level

Malondialdehyde (MDA) was determined by the method of Conti et al.<sup>[12]</sup>. The measurement of malondialdehyde level on the six-day-old mice pups' brains was performed using the MDA/Thiobarbituric Acid Reactive Substance (TBARS). This method was performed by weighing 1 gram of the infant mice brain sample then putting it into a reaction tube, mixing it with 9 cold PBS and then crushing it with a spatula. The liquid was then centrifuged at 3000 rpm for 48 min. 4 mL of supernatant was collected and added to 1 mL of trichloroacetic acid (TCA) 15% solution. Then, 1 mL of 0.37% Thiobarbituric acid (TBA) solution was added into HCl 0.25 N and heated in water bowl at 80°C for 15 min. After cooling the solution at room temperature for 60 min, the solution was centrifuged at 3000 rpm for 15 min. Finally, the value of the absorbance was read against the red lines which were formed using a spectrophotometer at  $\lambda = 532$  nm.

#### Necrosis Examination Using HE Staining

Necrosis examination was performed using HE staining. The six-day-old mice's brains were fixated with 10% formalin buffer and brain tissues were processed in routine processing until they formed paraffin blocks. The paraffin blocks were cut using a microtome with a thickness of 5  $\mu\text{m}$  in a series and then glued to the object glass. The observed area was the cerebellum which was obtained through the coronal section at the position 11/1.56 mm from the edge of the posterior lobe<sup>[13]</sup>. After 24 h' fixation, the mice brain was washed in a 70% alcohol solution three times and stained using Haematoxylin Eosin (HE). Purkinje cells which experienced necrosis were characterized by cells undergoing pyknosis and karyorrhexis.

#### Apoptosis Examination Using Tunel Assay

The examination of apoptotic cells was performed by Tunel assay. The paraffin blocks were cut using a microtome with a thickness of 5  $\mu\text{m}$  in a series and then glued to the object glass using polylysine. For counting the apoptotic cells, pieces of the tissue were processed with S7101 Apoptag Plus Peroxidase. Apoptotic cells were identified by the color absorbent (dark brown).

#### Data Analysis

Data analysis and evaluation of statistical significance among different determined values was performed using one-way Analysis of variance (ANOVA) with post hoc analysis (Duncan test)<sup>[14]</sup>. The values were expressed as mean  $\pm$  SD and considered significant at  $P < 0.05$ . The statistical analysis was performed using Statistical Product and Service Solutions (SPSS) version 17.0.

## RESULTS

#### Malondialdehyde (MDA) Level

This study showed that there was an increase in MDA levels, as the results of the ANOVA test indicated a sign value of 0.000 between the control and treatment group, which was less than the significance value  $\alpha = 0.05$ . It can be interpreted that there was a difference between the control and treatment groups. Thus, carbofuran administration in doses of either 0.0208 mg/kg or 0.0417 mg/kg might potentially produce the free radicals. The Duncan test indicated a sign value of 0.115 between 0.0208 mg/kg and 0.0417 mg/kg, which was more than the significance value  $\alpha = 0.05$ . There was no significant difference in pups whose mothers were exposed to carbofuran 0.0208 mg/kg and those with 0.0417 mg/kg. Although there was no significant difference, the increase in MDA levels was very high at 58.95% and 247.47% compared to that the control group, and the increase in MDA levels reached 118% between 0.0208 mg/kg and 0.0417 mg/kg (Table 1).

#### Cholinesterase (ChE) Level

This study presented the decreasing ChE levels and the ANOVA test result indicated a sign value of 0.001 between

**Table 1.** Effect of carbofuran exposure in subacute doses for 4 days on the MDA, ChE levels, Purkinje necrotic cell and Purkinje apoptotic cell in mice pups' brains (n=27)

Parameters	Control Group (mean $\pm$ SD)	0.0208 mg/kg BW Carbofuran (mean $\pm$ SD)	0.0417 mg/kg BW Carbofuran (mean $\pm$ SD)
Malondialdehyde (nmol/mg)	25.53 $\pm$ 3.02	40.58 $\pm$ 5.77 <sup>a</sup>	88.7 $\pm$ 3.02 <sup>a</sup>
ChE levels (U/L)	801.75 $\pm$ 129.73	671.50 $\pm$ 50.53 <sup>a</sup>	606.75 $\pm$ 28.45 <sup>a</sup>
Purkinje necrotic cell	1.98 $\pm$ 1.92	6.24 $\pm$ 0.73 <sup>y</sup>	7.68 $\pm$ 1.01 <sup>y</sup>
Purkinje apoptotic cell	11.00 $\pm$ 1.92	13.00 $\pm$ 0.73	14.13 $\pm$ 1.01

Statistical difference from the control: <sup>a,y</sup> Significant at  $P \leq 0.05$



the control and treatment groups, which was less than 17 significance value  $\alpha=0.05$ . It can be concluded that there was a difference between control and treatment groups. Carbofuran exposure in the doses of 0.0208 mg/kg and 0.0417 mg/kg potentially lowered the ChE levels. The Duncan test analysis revealed a sign value of 0.707 between 0.0208 mg/kg and 0.0417 mg/kg, which was more than the significance value  $\alpha=0.05$ . There was no significant difference in pups whose mothers were exposed to carbofuran of 0.0208 mg/kg and 0.0417 mg/kg. Although there was no significant difference 32 decrease in ChE levels reached 16.25% and 24.32% compared to the control group, and the decrease in ChE levels reached 9.64% between 0.0208 mg/kg and 0.0417 mg/kg (Table 1).





#### Necrotic Cell


18 In light microscopic examinations, histopathological changes were observed in mice pups' brains of all exposed groups compared to the control groups. This study found an increasing number of Purkinje necrotic cell in the brains of six-day-old mice pups whose mothers were exposed to carbofuran on the first and the fourth day of the lactation

period. Carbofuran exposure in doses of 0.0208 mg/kg and 0.0417 mg/kg had enhanced Purkinje necrotic cell. There was no significant difference in pups whose mothers were exposed to 0.0208 mg/kg and 0.0417 mg/kg of carbofuran. Nevertheless, the increase in Purkinje necrotic cell was very high at 215.15% for a dosage of 42 208 mg/kg and 287.87% for a dosage of 0.0417 mg/kg compared to the control group. Furthermore, the increase in Purkinje necrotic cell reached 23.07% between the dose administrations of 0.0208 mg/kg and 0.0417 mg/kg. An overview of staining results using HE showed necrosis of the Purkinje cells between the control and treatment groups (Fig. 1,2,3; Table 1).

#### Apoptotic Cell

This study expressed that there was an increasing number of Purkinje apoptotic cell in the brains of six-day-old mice pups from mother mice which were exposed to the insecticide carbofuran from the 1<sup>st</sup> and 4<sup>th</sup> day of the lactation period. The results of the Kruskal-Wallis test indicated a significance level of  $0.549 > 0.05$ . There was no significant difference in Purkinje apoptotic cell in lactating





**Fig 3.** Histopathology of the cerebellum of mice pups (*Mus musculus*), yellow arrow indicates the normal Purkinje cells, red arrow indicates the Purkinje necrotic cells. H.E Staining, 1000x

pups whose mothers received 0.0208 mg/kg and 0.0417 mg/kg of carbofuran compared to the control groups (Table 1). The estimated increase of Purkinje apoptotic cell in the carbofuran groups was 18.18% and 28.45%, while the increase in Purkinje apoptotic cell reached 7.99% between the dose administrations of 0.0208 mg/kg and 0.0417 mg/kg (Table 1).

## DISCUSSION

The metabolism of carbofuran has been well studied in rats, mice, and lactating cows. Carbofuran is rapidly absorbed, metabolized, and eliminated, primarily via urine, in the species investigated<sup>[15]</sup>. Carbofuran was altered by oxidation of the number 3 carbon and of the N-methyl group, hydrolysis of the ester linkage, and conjugation of metabolites containing a hydroxyl group. Carbofuran metabolites in the milk were the 3-hydroxycarbofuran, 3-keto-carbofuran, and 3-hydroxy-N-hydroxy methyl derivatives of carbofuran<sup>[2]</sup> which were found both in free and conjugated forms. Conjugated 2,3-dihydro-2,2-dimethyl-3-keto-7-hydroxybenzofuran was the major hydrolytic product of carbofuran in the milk. The same metabolites were also found in the urine and feces. 3-Hydroxycarbofuran was one of the most rapidly formed metabolites and 3-ketocarbofuran phenol was one of the end products<sup>[16]</sup>.

3-hydroxycarbofuran and nitrosocarbofuran could induce the micronucleus formation, while 3-ketocarbofuran could not but caused a significant DNA migration in SCGE test. Moreover, 3-ketocarbofuran caused an obvious increase in damaged cells accompanied with a great decrease in undamaged cells which displayed a higher degree of cell damage than other three compounds, especially seriously damaged cells increased in number and suggested a more serious DNA damaging effects. There has been no report on the mechanisms of DNA damaging effects induced by carbofuran and its metabolites. But in the oxidation and the hydrolyze of its transformation, free radicals are a potential

outcome, which is widely known to reduce DNA damage. The presence of metabolic active system in organisms could be degraded to less toxic phenols directly, while 3-hydroxycarbofuran was metabolized to 3-ketocarbofuran with a high direct toxicity to cells, thus, 3-hydroxycarbofuran revealed a stronger *in vivo* toxicity<sup>[17]</sup>.

Purkinje cell is a cell-specific marker of the cerebellar Purkinje cell and a suitable indicator for observing the postnatal development of the cerebellum after birth. The Purkinje cell levels in the mice's cerebellum during the critical postnatal (4 days after birth) is to determine the effect of external exposure on cerebellar growth in the offspring during lactation<sup>[18]</sup>.

Many pathophysiological conditions may have oxidative stress. In normal conditions, it is balanced by the antioxidant system and this balance is disturbed due to the increased oxidative stress. MDA is one of the fairly reactive metabolic products created from the effect of free oxygen radicals on tissues and from a series of reactions during lipid peroxidation. Malondialdehyde (MDA) is the best and a sensitive indicator of lipid peroxidation and so of oxidative stress<sup>[19,20]</sup>.

The final result of lipid peroxidation is MDA and a high MDA level indicates the damaged oxidation processes or cell membrane due to free radicals<sup>[21,22]</sup>. The results of this study showed that carbofuran exposure at subacute doses (0.0208 mg/kg and 0.0407 mg/kg) during a lactation period caused a significant enhancement in MDA levels of mice pups' brains. The data suggested that the significantly elevated MDA levels in the brain in turn produced a reactive oxygen species (ROS) which caused an oxidative stress in this organ. The increase of MDA levels results in cell death which is triggered by toxicants, depending on the dose. In this study, carbofuran dose increased MDA levels but there was no significant difference in pups whose mothers were exposed to carbofuran of 0.0208 mg/kg and 0.0417 mg/kg. Although there was no significant difference, the increase



in MDA levels reached 118% between 0.0208 mg/kg and 0.0417 mg/kg (Table 1). The increased doses could increase the significant MDA levels if exposed postnatally on the 1<sup>st</sup> until 20<sup>th</sup> day [23]. In another study, the increase in MDA levels due to an acute exposure of carbofuran [34] reach up to 175.04% for the dose administrations of 0.2 mg/kg which are compared to 0.4 mg/kg in mice [5]. Furthermore, oral sub-acute exposure of carbofuran for 28 days in male mice increases the MDA levels by 65% in other study [24].

This indicates that the increasing MDA levels due to carbofuran exposure depends highly on the dose, duration of exposure, and the type of affected organ. Vulnerability of brain development depends on the agent or the active metabolite which can be achieved during the development of nervous system and is associated with the exposure period. Exposure before [33] after the organ is fully developed makes the organ less vulnerable to inhibitions than if the exposure occurs [44] during the organ development [25]. An adult mouse brain is protected by the blood-brain barrier (BBB) to chemicals, while such protection does not exist in mice fetus and 6 months old mice [26].

[23] Neuronal membranes which are rich in polyunsaturated fatty acids are the source of lipid peroxidation reaction [27,28]. Lipid peroxidation causes destruction and damage to cell membranes and also changes the fluidity/membrane permeability [28,29]. Thus, the increase in MDA is caused by pesticides induction by forming RO [37]. Acute intraperitoneal carbofuran exposure may cause a significant increase in MDA levels of the brains and livers of adult mice. The increase in MDA levels is in line with the doses of carbofuran exposed. Carbofuran is proven to be efficiently absorbed and rapidly distributed to various organs of an organism. Lipophilic nature of carbamate also causes it to be able to interact with lipid serums and tissues [30]. The forming process of lipid peroxidation starts from hydrogen ions on the side chain of polyunsaturated fatty acids (PUFA), which construct the cell membranes by free radicals, form carbon radicals. Carbon radicals are oxidized to form peroxy radicals. Furthermore, peroxy radicals draw H<sup>+</sup> ion into the side chain of adjacent PUFA and form lipid peroxidation. This process is a chain reaction because the lipid peroxidation attracts more H<sup>+</sup> ions into the side chain of adjacent PUFA until the PUFA chain is finally split into other compounds, such as MDA, 9-hydroxy-nonanal, ethane and pentane [21,22].

Cholinesterase level measurements are often conducted to determine the exposure effect to insecticides. ChE used in this measurement was collected from tissues, plasma and red blood cells [31] and ChE collected from the brain was the best sample to be used as an indicator of the exposure to insecticides [32]. In this study, the insecticide carbofuran [26] decreased the ChE levels between the control and the treatment groups. However, there was no significant difference in the ChE levels among lactating

pups whose mothers received carbofuran of 0.0208 mg/kg and the ones with 0.0417 mg/kg (Table 1). The increased doses could decrease the significant ChE levels if exposed postnatally on the 1<sup>st</sup> until the 20<sup>th</sup> day [23]. Almost all insecticide exposure resulted in the decrease of ChE levels, either during embryonic period, growth period, or adulthood. Generally, the response of the decreased ChE level is in accordance with the exposed dose. However, the decreased ChE levels due to the insecticide exposure in several phases of individual growth may induce varied responses.

There were correlations [4] between the accumulation of acetylcholine and the extent of MDA. An increased oxidative stress by carbofuran might be a result of cholinergic hyperactivity or might be due to [3] direct effect on the production of reactive oxygen [33]. The peroxidation does not only alter lipid milieu, the structural and functional integrity of the cell membrane, but also affects the activities of various membrane-bound enzymes, including acetylcholinesterase (AChE) and different ATPases. The inhibition of ATPase activities may be a causative factor of neuronal/cellular dysfunction, due to an alteration in cationic transport across the membrane and a disturbance in uptake as well as a release of certain neurotransmitters [34].

The role of ChE is activated before the synaptogenesis during the formation of neural tube. The formation of ChE is in line with the axon growth [35]. The cholinergic system in early development acts as a regulatory growth and has morphogenetic functions [36] by controlling cell proliferation, motility [47], cell differentiation and genetic expression [37]. Thus, the cholinergic system has a very important role in the cell development and brain formation [38]. Although the brains of infant mice are extremely sensitive to carbofuran exposure during the lactation period and both treatments showed a decrease in ChE levels, all infant mice were still alive with symptoms of mild poisoning. The decrease in ChE levels indicates the response of brain or adult nervous system [35].

In this study, the insecticide carbofuran increased the necrotic death of Purkinje cells both in control and treatment group. However, there was no significant difference in necrotic cells among lactating pups whose mothers received carbofuran of 0.0208 mg/kg and the ones with 0.0417 mg/kg. In this study, we found an association between [8] the increased MDA levels and the necrotic cells. The increased lipid peroxidation and lipid peroxidation products, such as MDA levels, contribute to neuronal loss in conditions associated with oxidative stress [39]. Increased MDA levels indicates a membrane damage and leads to a cellular necrosis. The attack of free radicals on a cell membrane makes it devoid of integrity and viability causing the cells to undergo necrosis [40]. Cell death caused by the swelling of cytoplasm, nucleus karyolysis and lysis are classified as necrosis [41,42].

The number of Purkinje necrotic cells was not as many as the number of cells which experienced apoptosis due to the carbofuran exposure. This was because during the neurogenesis period, the Purkinje cells had experienced more apoptosis biologically through homeostasis efforts. However, when the number of Purkinje necrotic cells was compared to the control group, there was an increase in the number of Purkinje necrotic cells in the treatment group which was significantly higher than the number of Purkinje apoptotic cells (apoptosis increases 28.45% and necrosis increases up to 287.87%) (Table 1).

In this study, the insecticide carbofuran could increase the apoptosis of Purkinje cells between control and treatment groups. However, there was no significant difference in the number of Purkinje apoptotic cells among lactating pups whose mothers received carbofuran of 0.0208 mg/kg and the ones with 0.0417 mg/kg. Such different results from those of a study by Luqman<sup>[10]</sup> could be because the brain development phases during the embryonal period and lactation period have different critical time. In addition, the duration of carbofuran exposure during embryonal period was longer (10 days), while during the lactation period the exposure lasted only for 4 days. Although cerebellum is the most sensitive organ to oxidative-stress causing neurotoxins, longer exposure time is needed for the agent to reach the target of nuclear DNA and mitochondria to induce apoptosis<sup>[43]</sup>. The exposure of carbofuran insecticide can increase activity on cerebral ROS during embryonal period and the expression of p53, caspase 3 and apoptosis. The increasing expression of p53, caspase 3 and apoptosis indicated that the insecticide carbofuran caused an apoptosis through an intrinsic pathway<sup>[10]</sup>.

In conclusion, this study revealed that the carbofuran had been distributed in pups' tissues through the milk of lactating mothers and had caused an oxidative damage of pups' brains. Carbofuran exposure indicated that mice pups' brains were particularly more vulnerable to oxidative stress, which may eventually lead to neurobehavioral disorders. In this study, we also found that the insecticide carbofuran dose in lactating mice of 0.0208 mg/kg BW had been able to increase ROS activity and Purkinje cell death. with the same dose, if converted to humans, according to dose conversion by Laurence and Bacharach (1964), it will be equal to 0.115 mg/kg BW. This result of dose conversion can be applied as a carbofuran potential standard in increasing the ROS activity and Purkinje cell death since the residual level found in cows' meat and milk is around 0.17 mg/kg BW and 0.349 mg/kg BW<sup>[44]</sup>.

The insecticide carbofuran exposure to the lactating mice made the mean of Purkinje apoptotic cell higher than the necrotic ones in all treatment doses. A high increase of apoptosis in Purkinje cells allowed an opportunity to prevent and manage the strategy to overcome Purkinje cell death due to carbofuran insecticide exposure during the lactation period, such as by providing antioxidant

variations. Some efforts to prevent the formation of ROS can be done to inhibit and countermeasure the neuronal development cell death due to the exposure to carbofuran. The administration of antioxidants, such as vitamin C, curcumin, and allopurinol during lactation is possible to reduce the oxidative stress through the inhibitor xanthine oxidase and scavenger effects of free radicals. Therefore, there are still opportunities to improve the environment of neonate's nerve by increasing the growth of axons, dendrites and synaptogenesis and myelination of axons<sup>[45,46]</sup>.

## 6 ACKNOWLEDGMENTS

The authors express sincere thanks to the Ministry of Research, Technology and Higher Education of the Republic of Indonesia for funding the research, and the Dean of the Faculty of Veterinary Medicine for providing all necessary facilities and funds in conducting this research work.

## REFERENCES

1. Eskenazi B, Rosas LG, Marks AR, Bradman A, Harley K, Holland N, Johnson C, Fenster L, Barr DB: Pesticide toxicity and the developing brain. *Basic Clin Pharmacol Toxicol*, 102, 228-236, 2008. DOI: 10.1111/j.1742-7843.2007.00171.x
2. Handal AJ, Lozoff B, Breih J, Harlow SD: Effect of community of residence on neurobehavioral development infant and young children in a flower-growing region of Ecuador. *Environ Health Perspect*, 115, 128-133, 2007. DOI: 10.1289/ehp.9261
3. Kamboj SS, Kumar V, Kamboj A, Sandhir R: Mitochondrial oxidative stress and dysfunction in rat brain induced by carbofuran exposure. *Cell Mol Neurobiol*, 28, 961-969, 2008. DOI: 10.1007/s10571-008-9270-5
4. Masoud A, Sandhir R: Increased oxidative stress is associated with the development of organophosphate-induced delayed neuropathy. *Hum Exp Toxicol*, 31, 1214-1227, 2012. DOI: 10.1177/0960327112446842
5. Rai DK, Sharma B: Carbofuran induced oxidative stress in mamalian brain. *Mol Biotechnol*, 37, 66-71, 2007. DOI: 10.1007/s12033-007-0046-9
6. Redza-Dutordoir M, Averill-Bates DA: Activation of apoptosis signalling pathways by reactive oxygen species. *Biochim Biophys Acta*, 1863, 2977-2992, 2016. DOI: 10.1016/j.bbamcr.2016.09.012
7. Gupta RC, Milatovic S, Dettbarn WD, Aschner M, Milatovic D: Neuronal oxidative injury and dendritic damage induced by carbofuran: protection by memantine. *Toxicol Appl Pharmacol*, 219, 97-105, 2007. DOI: 10.1016/j.taap.2006.10.028
8. Ideguchi M, Palmer TD, Recht LD, Weimann JM: Murine embryonic stem cell-derived pyramidal neurons integrate into the cerebral cortex and appropriately project axons to subcortical targets. *J Neurosci*, 30, 894-904, 2010. DOI: 10.1523/JNEUROSCI.4318-09.2010
9. Stiles J, Jernigan TL: The basics of brain development. *Neuropsychol Rev*, 20, 327-348, 2010. DOI: 10.1007/s11065-010-9148-4
10. Luqman EM: Mekanisme Aktivitas ROS, Ekspresi p53 dan Caspase 3 serta Kematian Sel Neuron Korteks Serebrum Embrional Mencit (*Mus musculus*) Akibat Pajanan Insektisida Karbofuran: Penelitian Eksperimental Laboratorik. [Disertation]. Universitas Airlangga. Surabaya. Indonesia. 2013. <http://repository.unair.ac.id/21561/> Accessed: 15 January 2013.
11. Anonymus: Cholinesterase FS\*Diagnostic reagent for quantitative *in vitro* determination of cholinesterase (ChE) in serum or plasma on photometric systems. DiaSys Diagnostic Systems GmbH. 2018.
12. Conti M, Morand PC, Levillain P, Lemonnier A: Improved fluorometric determination of malonaldehyde. *Clin Chem*, 37, 1273-1275. 1991.
13. Paxinos G, Halliday G, Watson C, Koutcherov Y, Wang HQ: Atlas of the developing mouse brain at E17.5, P0 and P6. 1<sup>st</sup> ed., 48, Elsevier



London UK, 2007.

- 14. Supranto J:** Statistik: Teori dan Aplikasi. Edisi 8, 146, Erlangga Jakarta, 2016.
- 15. Sharma RK, Sharma B:** *In-vitro* carbofuran induced genotoxicity in human lymphocytes and its mitigation by vitamins C and E. *Dis Markers*, 32 (3):153-163, 2012. DOI: 10.3233/DMA-2011-0870
- 16. Gupta RC:** Carbofuran toxicity. *J Toxicol Environ Health*, 43, 383-418, 1994. DOI: 10.1080/15287399409531931
- 17. Pei Z, Baofeng L, Yitong L:** DNA damaging effects of carbofuran and its main metabolites on mice by micronucleus test and single cell gel electrophoresis. *Sci China C Life Sci*, 48 (Suppl 1), 40-47, 2005. DOI: 10.1007/BF02889800
- 18. Darmanto W:** Abnormal struktur histologis korteks cerebellar tikus dengan normal foliasi akibat iradiasi sinar X masa postnatal. *Berk Pene Hayati* 11, 13-18, 2005.
- 19. Jain S, Nair A, Shrivastava C:** Evaluation of oxidative stress marker malondialdehyde level in the cord blood of newborn infants. *IJJS*, 3 (6): 73-76, 2015.
- 20. Gülbayzar S, Arica V, Hatipoğlu S, Kaya A, Arica S, Karatekin G:** Malondialdehyde level in the cord blood of newborn infants. *Iran J Pediatr*. 21 (3), 313-319, 2011.
- 21. Haggag MEYE, Elsanhoty RM, Ramadan MF:** Impact of dietary oils and fats on lipid peroxidation in liver and blood of albino rats. *Asian Pac J Trop Biomed*, 4, 52-58, 2014. DOI: 10.1016/S2221-1691(14)60208-2
- 22. Jaiswal SK, Siddiqi NJ, Sharma B:** Carbofuran induced oxidative stress in rat heart: Ameliorative effect of vitamin C. *ISRN Oxidative Med*, 2013:824102, 2013. DOI: 10.1155/2013/824102
- 23. Mansour SA, Mossa AH:** Adverse effects of lactational exposure to chlorpyrifos in suckling rats. *Hum Exp Toxicol*, 29 (2):77-92, 2010. DOI: 10.1177/0960327109357276
- 24. Kamboj A, Kiran R, Sandhir R:** N-acetylcysteine ameliorates carbofuran induced alterations in lipid composition and activity of membrane bound enzymes. *Mol Cell Biochem*, 286, 107-114, 2006. DOI: 10.1007/s11010-005-9100-8
- 25. Rice D, Barone S:** Critical periods of vulnerability for the developing nervous system: Evidence from humans and animal models. *Environ Health Perspect*, 108 (Suppl 3): 511-33, 2000. DOI: 10.1289/ehp.00108s3511
- 26. Saunders NR, Liddelow SA, Dziegielewska KM:** Barrier mechanisms in the developing brain. *Front Pharmacol*, 3:46, 2012. DOI: 10.3389/fphar.2012.00046
- 27. Sultana R, Perluigi M, Butterfield DA:** Lipid peroxidation triggers neurodegeneration: A redox proteomics view into the Alzheimer disease brain. *Free Radic Biol Med*, 62, 157-169, 2013. DOI: 10.1016/j.freeradbiomed.2012.09.027
- 28. Van der Paal J, Neyts EC, Verlaack CCW, Bogaerts A:** Effect of lipid peroxidation on membrane permeability of cancer and normal cells subjected to oxidative stress. *Chem Sci*, 7, 489-498, 2016. DOI: 10.1039/c5sc02311d
- 29. Yusupov M, Wende K, Kupsch S, Neyts EC, Reuter S, Bogaerts A:** Effect of head group and lipid tail oxidation in the cell membrane revealed through integrated simulations and experiments. *Sci Rep*, 7, 5761, 2017. DOI: 10.1038/s41598-017-06412-8
- 30. Čolović MB, Krstić DZ, Lazarević-Pašti TD, Bondžić AM, Vasić VM:** Acetylcholinesterase inhibitors: Pharmacology and toxicology. *Curr Neuropharmacol*, 11, 315-335, 2013. DOI: 10.2174/1570159X11311030006
- 31. Linhares AG, Assis CRD, Siqueira MT, Bezerra RS, Carvalho LB:** Development of a method for extraction and assay of human erythrocyte acetylcholinesterase and pesticide inhibition. *Hum Exp Toxicol*, 32, 837-845, 2013. DOI: 10.1177/0960327112468906
- 32. Santos CSA, Monteiro MS, Soares AMVM, Loureiro S:** Brain cholinesterase reactivation as a marker of exposure to anticholinesterase pesticides: A case study in a population of yellow-legged gull *Larus michahellis* (Naumann, 1840) along the northern coast of Portugal. *Environ Sci Pollut Res Int*, 23,266-272, 2016. DOI: 10.1007/s11356-015-5730-x
- 33. Yang ZP, Dettbarn WD:** Diisopropylphosphorofluoridate-induced cholinergic hyperactivity and lipid peroxidation. *Toxicol Appl Pharmacol*, 138, 48-53, 1996. DOI: 10.1006/taap.1996.0096
- 34. Mehta A, Verma RS, Srivastava N:** Chlorpyrifos-induced alterations in rat brain acetylcholinesterase, lipid peroxidation and ATPases. *Indian J Biochem Biophys*, 42 (1): 54-58, 2005.
- 35. Tang J, Carr RL, Chambers JE:** Changes in rat brain cholinesterase activity and muscarinic receptor density during and after repeated oral exposure to chlorpyrifos in early postnatal development. *Toxicol Sci*, 51, 265-272, 1999. DOI: 10.1093/toxsci/51.2.265
- 36. Launder JM, Schambra UB:** Morphogenetic roles of acetylcholine. *Environ Health Perspect*, 107 (Suppl.-1): 65-69, 1999. DOI: 10.1289/ehp.99107s165
- 37. Torrao AS, Britto LRG:** Neurotransmitter regulation of neural development: Acetylcholine and nicotinic receptors. *An Acad Bras Ciênc*, 74, 453-461, 2002. DOI: 10.1590/S0001-37652002000300008
- 38. Resende RR, Adhikari A:** Cholinergic receptor pathways involved in apoptosis, cell proliferation and neuronal differentiation. *Cell Commun Signal*, 7:20, 2009. DOI: 10.1186/1478-811X-7-20
- 39. Keller JN, Mattson MP:** Roles of lipid peroxidation in modulation of cellular signaling pathways, cell dysfunction, and death in the nervous system. *Rev Neurosci*, 9 (2): 105-116, 1998. DOI: 10.1515/REVNEURO.1998.9.2.105
- 40. Milatovic D, Gupta RC, Dekundy A, Montine TJ, and Dettbarn WD:** Carbofuran-induced oxidative stress in slow and fast skeletal muscles: Prevention by memantine and atropine. *Toxicology*, 208, 13-24, 2005. DOI: 10.1016/j.tox.2004.11.004
- 41. Trump BF, Berezsky IK, Chang SH, Phelps PC:** The pathways of cell death: Oncosis, apoptosis, and necrosis. *Toxicol Pathol*, 25, 82-88, 1997. DOI: 10.1177/019262339702500116
- 42. Martin LJ:** Mitochondrial and cell death mechanisms in neurodegenerative diseases. *Pharmaceuticals (Basel)*, 3, 839-915, 2010. DOI: 10.3390/ph3040839
- 43. Yunus J, Sari DCR:** Efek neuroprotektif vitamin D terhadap jumlah sel Purkinje cerebellum yang diinduksi etanol. *Mutiara Medika*, 12, 63-71, 2012.
- 44. Indraningsih:** Pengaruh penggunaan insektisida karbamat terhadap kesehatan ternak dan produknya. *Wartazoa*, 18, 101-114, 2008.
- 45. Ikonomidou C, Kaindl AM:** Neuronal death and oxidative stress in the developing brain. *Antioxid Redox Signal*, 14, 1535-1550, 2011. DOI: 10.1089/ars.2010.3581
- 46. Yawno T, Castillo-Melendez M, Jenkin G, Wallace EM, Walker DW, Miller SL:** Mechanisms of melatonin-induced protection in the brain of late gestation fetal sheep in response to hypoxia. *Dev Neurosci*, 34, 543-551, 2012. DOI: 10.1159/000346323

# Brain Cells Death on Infant Mice (Mus musculus) Caused by Carbofuran Exposure During the Lactation Period

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** Pei Zhou. "DNA damaging effects of carbofuran and its main metabolites on mice by micronucleus test and single cell gel electrophoresis", Science in China Series C Life Sciences, 01/2005  
Publication 2%
- 2** docplayer.net  
Internet Source 2%
- 3** Sankhadeep Pal, Manisha Ahir, Parames C. Sil. " Doxorubicin-induced neurotoxicity is attenuated by a 43-kD protein from the leaves of L. via NF-κB and mitochondria dependent pathways ", Free Radical Research, 2012  
Publication 1%
- 4** link.springer.com  
Internet Source 1%
- 5** www.sid.ir  
Internet Source 1%
- 6** impactfactor.org  
Internet Source 1%

7	Uma Nath Tripathi, Deepak Chandra. "Hepatoprotective potential of Trigonella foenum graecum in deltamethrin induced albino rats", Indian Journal of Pharmaceutical and Biological Research, 2014 Publication	1 %
8	<a href="http://nopr.niscpr.res.in">nopr.niscpr.res.in</a> Internet Source	1 %
9	<a href="http://hdl.handle.net">hdl.handle.net</a> Internet Source	1 %
10	<a href="http://vetdergi.kafkas.edu.tr">vetdergi.kafkas.edu.tr</a> Internet Source	1 %
11	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	1 %
12	<a href="http://pubag.nal.usda.gov">pubag.nal.usda.gov</a> Internet Source	1 %
13	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	<1 %
14	Lu Zhang, Yi Na Sun, Yong Mei Li, Lai Xiang Lin, Yan Ye, Yu Qin Yan, Zu Pei Chen. "Effect of Different Iodine Nutrition on Cerebellum Pcp-2 in Rat Offspring During Lactation", Biological Trace Element Research, 2011 Publication	<1 %
15	<a href="http://www.ijss-sn.com">www.ijss-sn.com</a> Internet Source	<1 %



---

16 sphinxsai.com <1 %  
Internet Source

---

17 Mansour, S. A.-K., and L. Gamet-Payrastre. <1 %  
"Ameliorative effect of vitamin E to mouse dams and their pups following exposure of mothers to chlorpyrifos during gestation and lactation periods", Toxicology and Industrial Health, 2014.  
Publication

---

18 apjtb.com <1 %  
Internet Source

---

19 Devendra K. Rai, Prashant K. Rai, Syed I. Rizvi, <1 %  
Geeta Watal, Bechan Sharma. "Carbofuran-induced toxicity in rats: Protective role of vitamin C", Experimental and Toxicologic Pathology, 2009  
Publication

---

20 moam.info <1 %  
Internet Source

---

21 Meenakshi Kanwar Chauhan, Pankaj Kumar Sharma. <1 %  
"Optimization and characterization of rivastigmine nanolipid carrier loaded transdermal patches for the treatment of dementia", Chemistry and Physics of Lipids, 2019  
Publication

---

22

[pdfs.semanticscholar.org](https://pdfs.semanticscholar.org)

Internet Source

&lt;1 %

23

[www.biomed.cas.cz](http://www.biomed.cas.cz)

Internet Source

&lt;1 %

24

Fereidoun Azizi, Parvin Mirmiran, Amir Abbas Momenan, Farzad Hadaegh et al. "The Effect of Community-Based Education for Lifestyle Intervention on The Prevalence of Metabolic Syndrome and Its Components: Tehran Lipid and Glucose Study", International Journal of Endocrinology and Metabolism, 2013

Publication

&lt;1 %

25

Mansour, S. A., and A.-T. H. Mossa. "Adverse effects of exposure to low doses of chlorpyrifos in lactating rats", Toxicology and Industrial Health, 2011.

Publication

&lt;1 %

26

Reem Siraj Alsulaimani. "Immediate and Delayed Repair of 2 Sizes of Furcal Perforations in Dogs' Teeth Using Mineral Trioxide Aggregate Cement", Journal of Endodontics, 2018

Publication

&lt;1 %

27

Ines Bini Dhouib, Mohamed Montassar Lasram, Maha Abdeladhim, Najoua Gharbi, Melika Ben Ahmed, Saloua El-Fazaa. "Immunosuppression and oxidative stress

&lt;1 %

induced by subchronic exposure to carbosulfan in rat spleen: immunomodulatory and antioxidant role of N-acetylcysteine", *Toxicology Mechanisms and Methods*, 2014

Publication

28

[epdf.tips](http://epdf.tips)

Internet Source

<1 %

29

Bhupindervir Kaur, Alka Khera, Rajat Sandhir. "Attenuation of cellular antioxidant defense mechanisms in kidney of rats intoxicated with carbofuran", *Journal of Biochemical and Molecular Toxicology*, 2012

Publication

<1 %

30

Ratnesh Kumar Sharma, Bechan Sharma. "Carbofuran Induced Genotoxicity in Human Lymphocytes and Its Mitigation by Vitamins C and E ", *Disease Markers*, 2012

Publication

<1 %

31

Tamara Yawno, Mawin Mahen, Jingang Li, Michael C. Fahey, Graham Jenkin, Suzanne L. Miller. "The Beneficial Effects of Melatonin Administration Following Hypoxia-Ischemia in Preterm Fetal Sheep", *Frontiers in Cellular Neuroscience*, 2017

Publication

<1 %

32

[bu.edu.eg](http://bu.edu.eg)

Internet Source

<1 %



- |    |  |      |
|----|--|------|
| 33 | <a href="http://www.aipro.info">www.aipro.info</a><br>Internet Source  | <1 % |
| 34 | <a href="http://www.algologia.gr">www.algologia.gr</a><br>Internet Source  | <1 % |
| 35 | <a href="http://www.animbiosci.org">www.animbiosci.org</a><br>Internet Source  | <1 % |
| 36 | <a href="http://www.degruyter.com">www.degruyter.com</a><br>Internet Source  | <1 % |
| 37 | "Free Radical Biology and Environmental Toxicity", Springer Science and Business Media LLC, 2021<br>Publication  | <1 % |
| 38 | Amit Kamboj, Nagarajan Kumaresan, Ravi Kiran, Rajat Sandhir. " Neuroprotective potential of -acetylcysteine in carbofuran neurotoxicity: A biochemical and histopathological study ", Toxicological & Environmental Chemistry, 2006<br>Publication | <1 % |
| 39 | Amit Kamboj, Ravi Kiran, Rajat Sandhir. "N-Acetylcysteine Ameliorates Carbofuraninduced Alterations in Lipid Composition and Activity of Membrane Bound Enzymes", Molecular and Cellular Biochemistry, 2006<br>Publication                         | <1 % |

40 Asmaa Abd El-Aleem Abd El-Latif, Amany Ahmed Sayed, Amel Mahmoud Soliman, Sohair Ramadan Fahmy. "Exploration of the therapeutic potential effect of Sepia officinalis in animal model of sepsis induced by cecal ligation and puncture", Injury, 2016  
Publication <1 %

---

41 Janet K. Kern, Anne M. Jones. "Evidence of Toxicity, Oxidative Stress, and Neuronal Insult in Autism", Journal of Toxicology and Environmental Health, Part B, 2006  
Publication <1 %

---

42 Musa Kar. "Determination of the expression level of stress-related genes in Cicer arietinum root cell under Cd stress and the relationship to H2O2 concentrations", Ecotoxicology, 2018  
Publication <1 %

---

43 cyberleninka.org  
Internet Source <1 %

---

44 dl.icdst.org  
Internet Source <1 %

---

45 neurosciencesweb.wordpress.com  
Internet Source <1 %

---

46 www.cellmolbiol.org  
Internet Source <1 %

---

47

"Targeting Cellular Signalling Pathways in Lung Diseases", Springer Science and Business Media LLC, 2021

Publication

<1 %

48

Mandeep Kaur. "Oxidative Stress Response in Liver, Kidney and Gills of Ctenopharyngodon Idellus(Cuvier & Valenciennes) Exposed To Chlorpyrifos", MOJ Biology and Medicine, 2017

Publication

<1 %

49

Sevgi Başalan Över, Celal Güven, Eylem Taskin, Yusuf Sevgiler. " Oxidative and apoptotic effects of fluoxetine and its metabolite norfluoxetine in ", Archives of Industrial Hygiene and Toxicology, 2020

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On



# Brain Cells Death on Infant Mice (Mus musculus) Caused by Carbofuran Exposure During the Lactation Period

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---