

Paediatrica Indonesiana

(The Indonesian Journal of Pediatrics and Perinatal Medicine)

In this issue

- Glycated hemoglobin HbA1c, waist circumference, and waist-to-height ratio in overweight and obese adolescents
- Risk factors for miliary tuberculosis in children
- Maternal 1,25-dihydroxyvitamin D and neonatal birth weight
- A hematologic scoring system and C-reactive protein for diagnosing bacterial neonatal sepsis
- Hepatitis B seroprotection in children aged 10-15 years
- Blood pressure-to-height ratio for diagnosing hypertension
- Effect of oral administration of probiotics on intestinal colonization in preterm infants
- Duration of watching TV and child language development
- Duration and dose of antiepileptic drugs and serum calcium levels in children
- Delayed diagnosis of congenital hypothyroidism results in avoidable complications: a case report



CURRENT

ARCHIVES

ABOUT ▾

EDITORIAL BOARD

PUBLICATION ETHICS & MALPRACTICES

INSTRUCTIONS TO AUTHOR

REGISTER

LOGIN

Search

Editorial Board

Editor-in-Chief

Partini P. Trihono, Pediatric Nephrology, Universitas Indonesia, Jakarta ([Scopus:5](#); [Google Scholar:9](#); ORCID: <https://orcid.org/0000-0003-2893-6049>)

Vice Editor

Amanda Soebadi, Pediatric Neurology, Universitas Indonesia, Jakarta ([Scopus:3](#); [Google Scholar:4](#); ORCID: <https://orcid.org/0000-0001-9424-0029>)

Associate Vice Editor

Fitri Primacakti, Pediatric Hemato-Oncology, Universitas Indonesia, Jakarta ([Scopus:2](#); [Google Scholar:3](#))

Editorial Board

Agustini Utari, Pediatric Endocrinology, Universitas Diponegoro, Semarang ([Scopus:9](#); [Google Scholar:12](#); ORCID: <https://orcid.org/0000-0001-5965-1981>)

Ahmad Suryawan, Developmental Behavioral & Community Pediatrics, Universitas Airlangga ([Scopus:3](#); [Google Scholar:9](#); ORCID: <https://orcid.org/0000-0002-3377-0381>)

Ariani Dewi Widodo, Pediatric Gastrohepatology, Universitas Indonesia, Jakarta ([Scopus:5](#))

Cuno S. P. M. Uiterwaal, University Medical Centre, Utrecht ([Scopus:69](#))

Dominicus Husada, Infection & Tropical Pediatrics, Universitas Airlangga, Surabaya ([Scopus:7](#); [Google Scholar:10](#); ORCID: <https://orcid.org/0000-0001-9191-9546>)

Elisabeth Siti Herini, Pediatric Neurology, Universitas Gadjah Mada, Yogyakarta ([Scopus:6](#); [Google Scholar:10](#))

Finny Fitry Yani, Pediatric Respiriology, Universitas Andalas, Padang ([Scopus:6](#); [Google Scholar:10](#); ORCID: <https://orcid.org/0000-0002-4515-6530>)

Haryanti Fauzia Wulandari, Pediatric Imaging, Universitas Indonesia, Jakarta ([Google Scholar:1](#))

Hasri Samion, Institut Jantung Negara, Kuala Lumpur ([Scopus:12](#))

I Dewa Gede Ugrasena, Pediatric Hemato-Oncology, Universitas Airlangga, Surabaya ([Scopus:3](#); [Google Scholar:7](#))

I Gusti Ayu Trisna Windiani, Developmental Behavioral & Community Pediatrics, Universitas Udayana, Denpasar ([Scopus:7](#); [Google Scholar:8](#))

I Gusti Lanang Sidiartha, Pediatric Nutrition & Metabolic Diseases, Universitas Udayana, Denpasar ([Google Scholar:11](#))

I Gusti Made Reza Gunadi Ranuh, Pediatric Gastrohepatology, Universitas Airlangga, Surabaya ([Google Scholar:12](#))

Indah Kartika Murni, Pediatric Cardiology, Universitas Gadjah Mada, Yogyakarta ([Scopus:9](#); [Google Scholar:12](#))

Inke Nadia Diniyanti Lubis, Infection & Tropical Pediatrics, Universitas Sumatera Utara ([Scopus:9](#); [Scopus:10](#); ORCID: <https://orcid.org/0000-0001-7075-9491>)

Irene Yuniar, Emergency & Pediatric Intensive Care, Universitas Indonesia, Jakarta ([Scopus:1](#); [Google Scholar:4](#))

Jose Batubara, Pediatric Endocrinology, Universitas Indonesia, Jakarta ([Scopus:5](#); [Google Scholar:12](#))

Nadira Rasyid Ridha, Pediatric Hemato-Oncology, Universitas Hasanuddin, Makassar ([Google Scholar:4](#))

Ni Putu Veny Kartika Yantie, Pediatric Cardiology, Universitas Udayana, Denpasar ([Scopus:1](#); [Google Scholar:2](#))

Paul Glasziou, University of Sydney Faculty of Medicine, Queensland ([Scopus:1013](#); [Google Scholar:141](#); ORCID: <https://orcid.org/0000-0001-7564-073X>)

Raden Muhammad Indra, Pediatric Neurology, Universitas Sriwijaya, Palembang ([Scopus:1](#); [Google Scholar:2](#))

Ratno J. M. Sidauruk, Pediatric Imaging, Universitas Indonesia, Jakarta

Reni Ghrahani, Pediatric Allergy Immunology, Universitas Padjadjaran, Bandung ([Scopus:3](#); [Google Scholar:6](#); ORCID: <https://orcid.org/0000-0003-4476-2958>)

Retno Palupi-Baroto, Pediatric Nephrology, Universitas Gadjah Mada, Yogyakarta ([Scopus:3](#); [Google Scholar:3](#); ORCID: <https://orcid.org/0000-0001-8447-8682>)

Rina Triasih, Pediatric Respiriology, Universitas Gadjah Mada, Yogyakarta ([Scopus:15](#), [Google Scholar:20](#); ORCID: <https://orcid.org/0000-0002-6907-3422>)

Ririe Fachrina Malisie, Emergency & Pediatric Intensive Care, Universitas Sumatera Utara, Medan ([Scopus:1](#); [Google Scholar:2](#))

Risky Vitria Prasetyo, Pediatric Nephrology, Universitas Airlangga, Surabaya ([Scopus:3](#); [Google Scholar:4](#); ORCID: <https://orcid.org/0000-0002-1708-5384>)

Rosalina Dewi Roeslani, Neonatology, Universitas Indonesia, Jakarta ([Scopus:3](#); [Google Scholar:5](#))

Stephen M. Graham, Melbourne ([Scopus:52](#); [Google Scholar:78](#))

Sudigdo Sastroasmoro, Pediatric Cardiology, Universitas Indonesia, Jakarta ([Scopus:11](#))

Sumadiono, Pediatric Allergy Immunology, Universitas Gadjah Mada, Yogyakarta ([Scopus:2](#); [Google Scholar:6](#))

Tetty Yuniarti, Neonatology, Universitas Padjadjaran, Bandung

Yoga Devaera, Pediatric Nutrition & Metabolic Diseases, Universitas Indonesia, Jakarta ([Scopus:3](#);

Google [Scholar:7](#); ORCID: <https://orcid.org/0000-0002-5427-9823>)

Yvan Vandenplas, Universitair Ziekenhuis, Brussels ([Scopus:68](#))



[MAKE A SUBMISSION](#)



[Author Form](#)

Our journal is indexed by:

Scopus[®]



DOAJ DIRECTORY OF
OPEN ACCESS
JOURNALS

Risk factors for miliary tuberculosis in children

Clarissa Cita Magdalena¹, Budi Utomo², Retno Asih Setyoningrum³

Abstract

Background Miliary tuberculosis (TB) is a fatal form of tuberculosis with severe clinical symptoms and complications. The mortality rate from this disease remains high, therefore, it is important to identify the risk factors for miliary TB for early detection and treatment.

Objective To identify risk factors for miliary tuberculosis in children.

Methods A case-control study of children aged 0-14 years with miliary TB was conducted in Dr. Soetomo Hospital from 2010 to 2015. Data were taken from medical records. Case subjects were children with miliary TB, and control subjects were children with pulmonary TB. Patients with incomplete medical records were excluded. Case subjects were identified from the total patient population; control subjects were included by purposive sampling, with case:control ratio of 1:1. Potential risk factors were age, nutritional status, BCG immunization status, and history of contact with TB patients. Statistical analyses were done with Chi-square and logistic regression tests. P values < 0.05 were considered to be statistically significant.

Results A total of 72 children were analyzed, with 36 case and 36 control subjects. Nutritional status had a significant association with miliary TB in children (OR 3.182; 95%CI 1.206 to 8.398; P=0.018) in both bivariate and multivariate analyses. The probability of a child with moderate or severe undernutrition developing miliary TB was 76.09%. Other factors were not significantly associated with miliary TB.

Conclusion Nutritional status is significantly associated with miliary TB in children, and moderate or severe undernutrition increases the risk for developing miliary TB. [Paediatr Indones. 2017;57:63-6. doi: <http://dx.doi.org/10.14238/pi57.2.2017.63-6>].

Keywords: miliary tuberculosis; children; risk factors

Miliary tuberculosis (TB) is caused by hematogenous and lymphatogenous dissemination of *Mycobacterium tuberculosis* bacteria in the body, infecting multiple organs. It accounts for 3–7% of all TB cases.¹ Although there have been few reports on the prevalence of miliary TB in Indonesian children, the Indonesian Ministry of Health reported 1,168 cases of pediatric pulmonary acid-fast bacilli (AFB) positive TB in 2014.² The mortality rate from miliary TB is usually around 25%, but may reach 100% if left untreated.³

Complications from the disease include respiratory distress syndrome, renal failure, pericarditis, shock, disseminated intravascular coagulation, and acute respiratory failure.^{3,4} Miliary TB in children has been closely linked to the pathogenesis of TB meningitis (TBM), the most fatal form of TB. The proportion of children with miliary TB who suffer TBM is larger than that of adults with miliary TB.^{4,5}

Some proposed risk factors for developing miliary TB in children have been younger age, malnutrition, lack of BCG immunization, and history of contact with tuberculosis patients.^{4,6-8} Since pediatric TB and the severe complications of miliary TB are a health

From the Medical Student¹, Department of Public Health and Preventive Medicine², and Department of Child Health³, Airlangga University Medical School/Dr. Soetomo Hospital Surabaya, East Java, Indonesia.

Reprint requests to: Clarissa Cita Magdalena, Airlangga University Medical School Surabaya, Jl. Mayjen Prof. Dr. Moestopo 47, Surabaya, East Java, Indonesia. Tel. +62-31-5020251; E-mail: citamagdala@gmail.com

problem in Indonesia, we aimed to identify risk factors for miliary TB in children, in order to facilitate early prevention and intervention.

Methods

We conducted a hospital-based, case-control, retrospective study using secondary data from medical records of pediatric patients admitted to the Division of Respiriology, Department of Child Health, Airlangga University Medical School, Dr. Soetomo Hospital, Surabaya from 2010 to 2015. This study was approved by the Medical Ethics Committee of Dr. Soetomo Hospital, Surabaya, East Java, Indonesia.

Subjects were divided into case and control groups. Children under 14 years of age and diagnosed with miliary TB were selected as case subjects. Children under 14 years of age and diagnosed with pulmonary TB using Indonesian Pediatric Tuberculosis Scoring System¹ with a diagnostic score ≥ 6 were selected as control subjects. The scoring system was used only for the control subjects. Children with incomplete medical records were excluded. Cases were taken from the total population of those with miliary TB, while controls were included by purposive sampling, with a case: control ratio of 1:1.

Potential risk factors analyzed were age, nutritional status, BCG immunization status, and history of contact with TB patients. Age was categorized as ≤ 2 years or > 2 years. Nutritional status was determined using the weight/height WHO curves for subjects under 5 years or the CDC curves for those over 5 years,¹ according to subject's sex. All data were taken from the medical

records. We performed bivariate Chi-square and multivariate logistic regression analyses using SPSS version 20 software. Results with P values < 0.05 were considered to be statistically significant, with 95% confidence intervals.

Results

Out of 1,184 TB patients admitted during the study period, 46 had miliary TB. Ten patients were excluded, leaving 36 subjects in the case group. Thirty-six pulmonary TB patients were included as the control group. The characteristics of the subjects are presented in **Table 1**.

Table 1. Characteristics of subjects

Characteristics	N=72
Age, n(%)	
< 2 years	25 (34.7)
≥ 2 years	47 (65.3)
Nutritional status, n(%)	
Moderate or severe undernutrition	40 (55.6)
Normal	32 (44.4)
BCG immunization status, n(%)	
No	12 (16.7)
Yes	60 (83.3)
History of contact with TB patients, n(%)	
Yes	53 (73.6)
No	19 (26.4)

Bivariate analysis of the possible risk factors showed that only nutritional status was significantly associated with miliary TB in children (OR 3.182; 95%CI 1.206 to 8.398; P=0.018). Age, BCG

Table 2. Bivariate analysis of miliary tuberculosis risk factors in children

Variables	Case (n=36)	Control (n=36)	OR	95%CI	P value
Age, n					
< 2 years	16	9	2.400	0.882 to 6.528	0.083
≥ 2 years	20	27			
Nutritional status, n					
Moderate or severe undernutrition	25	15	3.182	1.206 to 8.398	0.018
Normal	11	21			
BCG immunization status, n					
No	9	3	3.667	0.902 to 14.901	0.058
Yes	27	33			
History of contact with TB patients, n					
Yes	25	28	0.649	0.225 to 1.871	0.422
No	11	8			

immunization status, and history of contact with associated with miliary TB in children (OR 3.182; 95%CI 1.206 to 8.398; P=0.018). Age, BCG immunization status, and history of contact with TB patients were not significantly associated with miliary TB (Table 2).

Multivariate analysis revealed similar results, with nutritional status as the only significant risk factor of miliary TB in children (Table 3). The probability of a child with moderate or severe undernutrition developing miliary TB was found to be 76.09%.

from active TB disease and TB infection.

We found no statistically significant associations between age, BCG immunization status, or history of contact with TB patients and the incidence of miliary TB in children. However, we suggest that younger age and negative BCG immunization status may also increase the risk of miliary TB, as reported by previous studies.^{6,7}

The difference between our findings and those of previous studies may be due to several reasons. First, age in our study may have been affected by the

Table 3. Multivariate analysis of miliary risk factors in children

Variables	Coefficient (B)	OR	95%CI	P value
Age	0.702	2.017	0.653 to 6.235	0.223
Nutritional status	1.165	3.204	1.166 to 8.805	0.024
BCG immunization status	0.891	2.437	0.527 to 11.263	0.254

Discussion

We identified moderate/severe malnutrition as a significant risk factor for miliary TB in children. Similarly, a study in India found that malnutrition was a risk factor for miliary tuberculosis.⁴ Also, another study in India found a significant association between nutritional status and the incidence of pulmonary tuberculosis infections.⁹ A previous study in Peru also found a significant association between malnutrition and mortality in children with TB.¹⁰ Furthermore, a systematic review of studies in developed countries like the United States, Hong Kong, Finland, and Norway, found a consistent relationship between the incidence of TB and the body mass index (BMI) of the patients, with a 14% increased risk of TB for a decrease of one BMI unit.¹¹

The relationship between nutritional status in children and miliary tuberculosis incidence may be explained by Jaganath *et al.* who suggested that nutrient deprivation may have a detrimental effect on Th1 cells, which act as an important component in cell-mediated immune system defense against miliary TB.¹² A previous study reported that cell-mediated immunity is a key factor in host defense mechanisms against the progression of TB infection to active TB disease.¹³ Therefore, the compromised cellular immune system in children with undernutrition possibly increases the risk of developing miliary TB

higher prevalence of older pediatric TB patients. A previous study noted that even though an age under 2 years was a risk factor of miliary TB, most children with TB infection in endemic areas were older than 2 years, so there was a higher chance of more children over 2 years suffering from miliary TB.⁶ Another study in England and Wales also found more miliary TB in older children, which indicated the possible reactivation of latent disease.¹⁴ Thus, we suggest that miliary TB should be suspected in children of any age. Second, BCG immunization coverage was high in our study. According to a systematic review by Trunz *et al.*, the estimated efficacy of BCG prevention of miliary TB reached 77%, but in Asian countries, there might be an overestimation of the number, due to the inclusion of studies of countries with higher immunization coverage than Asian countries but with lower risk of infection, while Asian countries have higher rate of infection and reinfection despite the high immunization coverage.¹⁵ Moreover, Fine found that the protective effect of BCG was influenced by geographic location, as the protective effect declined in regions closer to the equator.¹⁶ BCG immunization itself should continue to be administered as regulated, but we suggest continuing future study for a better vaccine. Third, the majority of our subjects reported a positive history of contact with TB patients, in contrast to previous studies with more subjects who had no history of contact with TB patients.^{8,17}

Therefore, contact tracing is an important task, but the precise relationship between contact and military TB needs further study.

A limitation of this study was the use of secondary data which could create a bias in the information we received. We propose future studies with better methodologies, preferably with a prospective design.

In conclusion, moderate/severe malnutrition is a significant risk factor for military TB in children. Further study is needed to elucidate the pathogenic mechanism between undernutrition and military TB.

Conflict of interest

None declared.

References

1. Kementerian Kesehatan RI. Petunjuk teknis manajemen TB anak. Jakarta: Kementerian Kesehatan Indonesia; 2013. p.13,18, 85-96.
2. Kementerian Kesehatan RI. Profil kesehatan Indonesia 2014. Jakarta: Kementerian Kesehatan Indonesia; 2015. p.110.
3. Hagan G, Nathani N. Clinical review: tuberculosis on the intensive care unit. *Crit Care*. 2013;17:240.
4. Sharma SK, Mohan A, Sharma A. Challenges in the diagnosis & treatment of military tuberculosis. *Indian J Med Res*. 2012;135:703-30.
5. Donald PR, Schaaf HS, Schoeman JF. Tuberculous meningitis and military tuberculosis: the Rich focus revisited. *J Infect*. 2005;50:193-5.
6. Marais BJ, Gie RP, Schaaf HS, Hesselning AC, Obihara CC, Starke JJ, et al. The natural history of childhood intra-thoracic tuberculosis: a critical review of literature from the pre-chemotherapy era. *Int J Tuberc Lung Dis*. 2004;8:392-402.
7. Newton SM, Brent AJ, Anderson S, Whittaker E, Kampmann B. Paediatric tuberculosis. *Lancet Infect Dis*. 2008;8:498-510.
8. Gomes VF, Andersen A, Wejse C, Oliveira I, Vieira FJ, Joaquim LC, et al. Impact of tuberculosis exposure at home on mortality in children under 5 years of age in Guinea-Bissau. *Thorax*. 2011;66:163-7.
9. Singh M, Mynak ML, Kumar L, Mathew JL, Jindal SK. Prevalence and risk factors for transmission of infection among children in household contact with adults having pulmonary tuberculosis. *Arch Dis Child*. 2005;90:624-8.
10. Drobac PC, Shin SS, Huamani P, Atwood S, Furin J, Franke MF, et al. Risk factors for in-hospital mortality among children with tuberculosis: the 25-year experience in Peru. *Pediatrics*. 2012;130:e373-9.
11. Lönnroth K, Williams BG, Cegielski P, Dye C. A consistent log-linear relationship between tuberculosis incidence and body mass index. *Int J Epidemiol*. 2010;39:149-155.
12. Jaganath D, Mupere E. Childhood tuberculosis and malnutrition. *J Infect Dis*. 2012;206:1809-15.
13. Cegielski JP, McMurray DN. The relationship between malnutrition and tuberculosis: evidence from studies in humans and experimental animals. *Int J Tuberc Lung Dis*. 2004;8:286-98.
14. Kruijshaar M, Abubakar I. Increase in extrapulmonary tuberculosis in England and Wales 1999-2006. *Thorax*. 2009; 64:1090-5.
15. Trunz BB, Fine P, Dye C. Effect of BCG vaccination on childhood tuberculous meningitis and military tuberculosis worldwide: a meta-analysis and assessment of cost-effectiveness. *Lancet*. 2006;367:1173-80.
16. Fine PE. Variation in protection by BCG: implications of and for heterologous immunity. *Lancet*. 1995;346:1339-45.
17. Rakhmawati W, Fatimah S, Nurhidayah I. Hubungan status gizi, imunisasi & riwayat kontak dengan kejadian tuberculosis pada anak di wilayah kerja puskesmas Ciawi kabupaten Tasikmalaya [final research report]. Bandung: Universitas Padjajaran; 2008. [cited 2016 November 25]. Available from: <http://repository.unpad.ac.id/4895/>.