

Editorial Board

Editor-in-Chief

Nancy Epstein, MD

Professor of Clinical Neurosurgery,

School of Medicine, State University of New York at Stony Brook; New York, NY

Chief of Neurosurgical Spine and Education

NYU Winthrop Hospital; Mineola, NY

Emeritus Editor-in-Chief

James I. Ausman, MD, PhD

UCLA Department of Neurosurgery, Los Angeles, CA

Founding Editors

James Ausman – UCLA Department of Neurosurgery, Los Angeles, CA

Atos de Sousa (2010–2013) – Gerais Santa Casa Hospital and Lifecenter Hospital; Belo Horizonte, Brazil

Robert "Bob" Goodkin (2010–2012) – University of Washington; Seattle, WA, USA

Pat Kelly – NYU School of Medicine; New York, NY, USA

Neil Martin – Geisinger Health System; Danville, PA

Managing Editor

Jim Cook

31855 Date Palm Drive, Suite 3-400

Cathedral City, CA 92234

(760) 349-6600

Supplements

SNI: Computational

Associate Editor-in-Chief:

Pieter L. Kubben, MD, PhD

Maastricht University Medical Center; Maastricht, The Netherlands

Editors:

Alkinoos Athanasiou – Aristotle University of Thessaloniki; Thessaloniki, Greece

Philippe Bijlenga – Hôpitaux Universitaires de Genève; Geneva, Switzerland

Guillaume Charvet – Clnatec; Grenoble, France

Daniel John DiLorenzo – Loma Linda University Medical Center; Loma Linda, CA, USA
Francesco DiMeco – Istituto Nazionale Neurologico "C.Besta"; Milan, Italy
Erik Gommer – Maastricht University Medical Center; Maastricht, The Netherlands
Michael Murek – Inselspital, Universitätsspital Bern; Bern, Switzerland
Felix Scholtes – Centre Hospitalier Universitaire de Liege; Liège, Belgium
Phillippe Schucht – Inselspital, Universitätsspital Bern; Bern, Switzerland
Peter Willems – Leiden University Medical Center; Leiden, The Netherlands

SNI: Digital Medicine

Associate Editor-in-Chief:

James I. Ausman, MD, PhD

UCLA Department of Neurosurgery, Los Angeles, CA;

Editors:

Abdul Hadi Al-Khalili – Baghdad University; Baghdad, Iraq

Zahraa Al-Sharshahi – Neurosurgery Teaching Hospital; Baghdad, Iraq

Tarun Arora – University of California at San Francisco; San Francisco, CA, USA

Iype Cherian – Bharatpur College of Medical Sciences; Chitwan, Nepal

Chandrashekhara Deopujari – Hinduja Hospital; Mahim, Mumbai, India

Abdeslam El Khamlichi – Mohammed V University of Rabat; Rabat, Morocco

S. Ather Enam – Aga Khan University; Karachi, Sindh, Pakistan

Nancy Epstein – State University of New York at Stony Brook; New York, NY, USA

Moneer Faraj – University of Baghdad; Baghdad, Iraq

Marco Gonzales-Portillo – Instituto Neurociencias de Lima; Lima, Peru

Marwan Hariz – Umeå University; Umeå, Sweden

Kazuhiro Hongo – Shinshu University; Matsumoto, Japan

Samer Hoz – University of Cincinnati; Cincinnati, OH, USA

Seyed Ali Khonsary – University of California at Los Angeles; Los Angeles, CA, USA

Jorge Lazareff – University of California Los Angeles; Los Angeles, CA, USA

Jay Morgan – Sierra Neurosurgery Group; Reno, NV, USA

Hiba Abdulmir Nasir – Dr. Saad Alwitry Neurosciences Hospital; Baghdad, Iraq

Eric Nussbaum – National Brain Aneurysm & Tumor Center, Twin Cities, MN, USA

Frank Van Calenbergh – University Hospitals; Leuven, Belgium

SNI: Epilepsy

Associate Editor-in-Chief:

Warren Boling, MD

Loma Linda University Medical Center; Loma Linda, CA, USA

SNI: General Neurosurgery

Associate Editor-in-Chief:

Eric Nussbaum, MD

National Brain Aneurysm & Tumor Center, Twin Cities, MN, USA

SNI: Global Health and Education

Associate Editor-in-Chief:

Jorge Lazareff MD, FAANS

University of California Los Angeles; Los Angeles, CA, USA

Editors:

Seyed Ali Khonsary – University of California at Los Angeles; Los Angeles, CA, USA

SNI: Head and Spinal Cord Transplantation

Associate Editors-in-Chief:

Sergio Canavero, MD

XiaoPing Ren, MD

HEAVEN/GEMINI International Collaborative Group, Turin, Italy

Editors:

Farid Amirouche – University of Illinois at Chicago

SNI: History of Medicine

Associate Editor-in-Chief:

Miguel Faria, MD

Founder and Editor Emeritus, Medical Sentinel; www.haciendapublishing.com; Macon, GA, USA

Editor:

Russell Blaylock – Advanced Nutritional Concepts and Theoretical Neurosciences; Ridgeland, MS, USA

SNI: Infection

Associate Editor-in-Chief:

Ali Akhaddar, MD, IFAANS

Avicenne Military Hospital; Marrakech, Morocco

Editors:

Mohamed Benzagmout – University Hospital of Fez; Fez, Morocco

Walter Hall – State University of New York; Syracuse, NY, USA

Julio Sotelo – National Institute of Neurology and Neurosurgery; Mexico City, Mexico

Mehmet Turgut – Adnan Menderes University School of Medicine; Aydin, Turkey

SNI: Medical Legal Public Health and Policy

Associate Editor-in-Chief:

Clark Watts, MD

University of Texas School of Law; Austin, TX, USA

SNI: Neuroanatomy/Neurophysiology

Associate Editor-in-Chief:

Seyed Ali Khonsary, MD

University of California at Los Angeles; Los Angeles, CA, USA

SNI: Neuroanesthesia and Critical Care

Associate Editor-in-Chief:

Ramsis F. Ghaly, MD

University of Illinois at Chicago; Chicago, IL, USA

SNI: Neuroendoscopy

Associate Editor-in-Chief:

J. André Grotenhuis, MD

Radboud University Medical Center; Nijmegen, The Netherlands

Editor:

Chandrashekar Deopujari - Hinduja Hospital; Mahim, Mumbai, India

SNI: Neuro-Inflammation

Associate Editor-in-Chief:

Russell Blaylock, MD

Advanced Nutritional Concepts and Theoretical Neurosciences; Ridgeland, MS, USA

Editors:

F. Gómez-Pinilla - University of California at Los Angeles; Los Angeles, CA, USA

SNI: Neuro-oncology and Skull Base

Associate Editor in Chief:

Mitsutoshi Nakada, MD, PhD

Kanazawa University; Ishikawa, Japan

Editors:

Chandrashekar Deopujari - Hinduja Hospital; Mahim, Mumbai, India

Yasuhiko Hayashi - Kanazawa Medical University; Ishikawa, Japan

Lauren Hertan - Beth Israel Deaconess Medical Center; Boston, MA, USA

Imad N. Kanaan - The King Faisal Specialist Hospital and Research Center; Riyadh, Saudi Arabia

Kazuhiko Kurozumi - Okayama University; Okayama, Japan

Anand Mahadevan – Beth Israel Deaconess Medical Center; Boston, MA, USA

Aldo A. Spallone – NCL – Institute of Neurosciences; Rome, Italy

Lee E. Tessler – Neurological Surgery, P.C.; Long Island, NY, USA

SNI: Neuropathology

Associate Editor-in-Chief:

Istvan Bodi

King's College Hospital NHS Foundation Trust; London, United Kingdom

Editors:

Johannes A. Hainfellner, MD – Medical University of Vienna; Vienna, Austria

Tibor Hortobágyi, MD – Faculty of Medicine, University of Debrecen; Debrecen, Hungary

SNI: Neuropsychiatry

Associate Editors-in-Chief:

Miguel Faria, MD

Founder and Editor Emeritus, Medical Sentinel; www.haciendapublishing.com; Macon, GA, USA

Russell Blaylock, MD

Advanced Nutritional Concepts and Theoretical Neurosciences; Ridgeland, MS, USA

Editor:

C. Pritchard – Bournemouth University; Poole, England

Robert B. Young – University of Rochester School of Medicine; Rochester, NY, USA

Oswaldo Vilela-Filho – Federal University of Goiás; Goiânia, Brazil

SNI: Neurosurgery Concepts: Perspectives on Key Articles in Neurosurgery

Associate Editor-in-Chief:

Isaac Yang, MD

UCLA Medical Center; Los Angeles, CA, USA

Editors:

A.M. Bohnen – Northwestern University; Chicago, IL, USA

J.M. Cho – Ajou University School of Medicine; Suwon, South Korea

C.B. Colen – Beaumont Hospital; Grosse Pointe, MI, USA

N.S. Dahdaleh – Northwestern University; Chicago, IL, USA

E. Duckworth – Baylor College of Medicine; Houston, TX, USA

A.V. Germanwala – Loyola University School of Medicine; Maywood, IL, USA

P. Kan – Baylor College of Medicine; Houston, TX, USA

A. Khalessi – University of California at San Diego; San Diego, CA, USA

C.Y. Kim – Seoul National University Bundang Hospital; Seongnam-si, Gyeonggi-do, South Korea

S. Lam – Baylor College of Medicine; Houston, TX, USA
G. Li – Stanford University; Stanford, CA, USA
M. Lim – Johns Hopkins University; Baltimore, MD, USA
J.H. Sherman – George Washington University; Washington, DC, USA
V.Y. Wang – University of California at San Francisco; San Francisco, CA, USA
G. Zada – Keck School of Medicine at U.S.C.; Los Angeles, CA, USA

SNI: Neurovascular

Associate Editor-in-Chief:

Kazuhiro Hongo, MD
Shinshu University; Matsumoto, Japan

Editors:

Vladimir Beneš – Charles University; Prague, Czech Republic
G.W. Britz – The Methodist Hospital; Houston, TX, USA
J.A. Dye – University of California at Los Angeles; Los Angeles, CA, USA
J. Frösen – Helsinki University Central Hospital; Helsinki, Finland
Atul Goel, MD – King Edward Memorial Hospital, Mumbai, India
G. Grasso – University of Palermo; Palermo, Italy
Tetsuyoshi Horiuchi, MD, PhD – Shinshu University School of Medicine, Asahi, Japan
Kiyohiro Houkin, MD, PhD – Hokkaido University Hospital, Sapporo, Japan
Vijendra K. Jain – Max Superspecialty Hospital; New Delhi, India
Yoko Kato – Fujita Health University Hospital, Toyoake, Japan
M.J. Korja – Macquarie University Hospital; Sydney, Australia
Satoshi Kuroda, MD, PhD – University of Toyama, Toyama, Japan
G. Lanzino – Mayo Clinic; Rochester, MN, USA
Yasushi Matsumoto, MD – Kohnan Hospital, Sendai, Japan
R. Loch Macdonald – University of Toronto; Toronto, Ontario, Canada
Yuichi Murayama, MD – Jikei University School of Medicine, Tokyo, Japan
E.S. Nussbaum – St. Joseph's Hospital; Twin Cities, MN, USA
Amitendu Sekhar – Manipal Hospital; Salem, India
T. Shibuya – Nihon University School of Medicine; Tokyo, Japan
Yuichiro Tanaka, MD – St. Marianna University School of Medicine, Kawasaki, Japan
P. Taussky – University of Utah; Salt Lake City, UT, USA
K. Van Overbeeke – Maastricht University Medical Centre; Maastricht, The Netherlands
D.A. Vega-Moreno – National Autonomous University of Mexico; Mexico City, Mexico
Shinichi Yoshimura – Hyogo College of Medicine, Hyogo, Japan
J. Zabramski – Barrow Neurosurgical Associates; Phoenix, AZ, USA

SNI: Nursing

Associate Editor-in-Chief:

Nancy Epstein, MD
NYU Winthrop Hospital; Mineola NY, USA

SNI: Pain

Associate Editor-in-Chief:

Jeff Brown
Neurological Surgery, P.C.; Great Neck, NY, USA

SNI: Pediatric Neurosurgery

Associate Editor-in-Chief:

Frank Van Calenbergh, MD
University Hospitals; Leuven, Belgium

Editors:

Chandrashekhara Deopujari – Hinduja Hospital; Mahim, Mumbai, India
G. Grant – University Medical Center; Durham, NC, USA
J. Lazareff – University of California at Los Angeles; Los Angeles, CA, USA
M.Z. Tahir – Great Ormond Street Children's Hospital, NHS Foundation Trust; London, UK

SNI: Review of Randomized Controlled Trials

Associate Editor-in-Chief:

Benjamin W. Y. Lo, MD
Montreal Neurological Institute & Hospital, McGill University; Quebec, Canada

Editors:

Mark Angle – Montreal Neurological Institute & Hospital, McGill University; Quebec, Canada
Hitoshi Fukuda – Kochi University Hospital; Kochi, Japan
Masaomi Koyanagi – National Hospital Organization Himeji Medical Center (Kyoto University); Hyogo, Japan
Matthew Wai-Man Lui – Queen Mary Hospital, University of Hong Kong; Pokfulam, Hong Kong
Satoru Miyawaki – The University of Tokyo; Tokyo, Japan
Jeanne Teitelbaum – Montreal Neurological Institute & Hospital, McGill University; Quebec, Canada
Anderson C.O. Tsang – The University of Hong Kong; Hong Kong
George Kwok Chu Wong – Prince of Wales Hospital, Chinese University of Hong Kong; Shatin, Hong Kong

Ad Hoc Members:

Mitchell A. H. Levine – McMaster University; Hamilton, Ontario, Canada
Abbas Sadikot – Montreal Neurological Institute & Hospital, McGill University

SNI: Revista Argentina de Neurocirugía

Associate Editors-in-Chief:

Álvaro Campero -- Hospital Padilla, S. M. Tucumán

Matteo Baccanelli -- Hospital Italiano de Buenos Aires; Buenos Aires, Argentina

Secretario de Redacción: Rubén Mormandi – FLENI, CABA

Editor Fundador: León Turjanski – Hospital Cosme Argerich, CABA

Comité de Redacción

Matteo Baccanelli – Hospital Italiano de Buenos Aires

Martín Guevara – Hospital Fernández, CABA

Martín Saez – Hospital Gandulfo, Lomas de Zamora

Fidel Sosa – Hospital El Cruce, Florencio Varela

Pablo Rubino – Hospital El Cruce, Florencio Varela

Pablo Ajler – Hospital Italiano, CABA

Sergio Pampín – Hospital Posadas, Ramos Mejía

Tomás Funes – Sanatorio Otamendi, CABA

Jorge Bustamante – Hospital El Cruce, Florencio Varela

Federico Landriel – Hospital Italiano, CABA

Leopoldo Luque – Hospital El Cruce, Florencio Varela

Comité Asesor y de Revisión

Coordinador: Marcelo Platas

Miembros:

Jaime Rimoldi

Jorge Oviedo

Rafael Torino

Juan Mezzadri

Graciela Zúccaro

Luis Lemme Plaghos

Jorge Salvat

Julio César Suarez

Julio Antico

Horacio Fontana

Oscar Stella

Antonio Carrizo

Silvia Berner

Jorge Lambre

Francisco Papalini

León Turjanski

Alejandra Rabadán

Javier Gardella

Comité de Trabajo Supplementum Revista Argentina de Neurocirugía – Surgical Neurology International

Matteo Baccanelli – Hospital Italiano de Buenos Aires

Pablo Ajler – Hospital Italiano, CABA

Federico Landriel – Hospital Italiano, CABA,

SNI: Socioeconomics, Politics, Medicine and World Affairs

Associate Editors-in-Chief:

James I Ausman, MD, PhD

UCLA Department of Neurosurgery, Los Angeles, CA

Russell Blaylock, MD

Advanced Nutritional Concepts and Theoretical Neurosciences; Ridgeland, MS, USA

Miguel Faria, MD

Founder and Editor Emeritus Medical Sentinel; www.haciendapublishing.com

Walter Johnson, MD

World Health Organization; Geneva, Switzerland

Editors:

Youssef Fares, MD, PhD, EMBA – Lebanese University; Beirut, Lebanon

SNI: Stereotactic

Associate Editor-in-Chief:

Konstantin V. Slavin

University of Illinois at Chicago; Chicago, IL, USA

Editors:

Antonio A.F. De Salles – University of California; Los Angeles, CA, USA

Moneer Faraj – University of Baghdad; Baghdad, Iraq

Alessandra Gorgulho – University of California; Los Angeles, CA, USA

D. Kondziolka – University of Pittsburgh; Pittsburgh, PA, USA

K. Lizarraga – University of Rochester; Rochester, NY, USA

T. Mindermann – Klinik Im Park; Zurich, Switzerland

Hiba Abdulmir Nasir – Dr. Saad Alwitry Neurosciences Hospital; Baghdad, Iraq

N. Pouratian – University of California; Los Angeles, CA, USA

G. Sinclair – North Middlesex University Hospital NHS Trust, London, UK

SNI: Spine

Associate Editor-in-Chief:

Nancy Epstein, MD

NYU Winthrop Hospital; Mineola NY, USA

Editors:

K. Abumi – Hokkaido University Health; Sapporo, Japan
P. Arnold – University of Kansas; Kansas City, KS, USA
J. Baisden – University of Wisconsin, Milwaukee, WI, USA
P.G. Benton – Atlanta Medical Center, Atlanta, GA, USA
B.R. Cohen – NYU Winthrop Hospital; Mineola, NY, USA
F. Cohen – Palm Beach Gardens Medical Center; Palm Beach Gardens, FL, USA
B. Cunha – NYU Winthrop Hospital; Mineola, NY
R. A. Deyo – Oregon Health and Science University; Portland, OR, USA
D. J. DiLorenzo – Loma Linda University Medical Center; Loma Linda, CA, USA
T. B. Ducker – Retired, Johns Hopkins University; Baltimore, MD, USA
J.H. Dunn – Southern Arizona VA Health Center; Tucson, AZ, USA
C. L. Forte – Blume, Goldfaden, Berkowitz, Donnelly, Fried, & Forte; Jersey City, NJ, USA
F. W. Gamache – Cornell Medical Center; NY, NY, USA
E. N. Hanley – Carolinas Health Care; Charlotte, NC, USA
D. L. Hilton – Neurosurgical Associates of San Antonio; San Antonio, TX, USA
L. T. Holly – UCLA Medical Center; Los Angeles, CA, USA
R. Hollingsworth – NYU Winthrop Hospital; Mineola, NY, USA
G. B. Jacobs – Hackensack University Medical Center; South Hackensack, NJ, USA
P. M. Lippe – Vista de Almaden; San Jose, CA, USA
B. W. Lo – Montreal Neurological Institute & Hospital, McGill University; Quebec, Canada
J. Loeser – University of Washington; Seattle, WA, USA
D. J. Maiman – University of Wisconsin; Milwaukee, WI, USA
R. A. McGuire – University of Mississippi Medical Center; Jackson, MS, USA
H. Morgan – University of Texas Southwestern Medical Center; Dallas, TX, USA
H. Moses – Greater Baltimore Medical Center; Baltimore, MD, USA
H. Nakagawa – University of Nagoya; Nagoya, Japan
A. Okawa – Tokyo Medical and Dental University, Graduate School of Medicine; Tokyo, Japan
A. A. Patel – Loyola University Health System; Maywood, IL, USA
R. Pawl – Loyola University Hospital; Lake Forest, IL, USA
G. R. Rechtine – Florida Orthopedic Institute; Tampa, FL, USA
K. Shinomiya – Tokyo Medical and Dental University; Tokyo, Japan
M. Stecker – University of California at San Francisco; Fresno, CA, USA
P. J. Tortolani – Union Memorial Hospital; Baltimore, MD, USA
V. C. Traynelis – Hackensack University Medical Center; South Hackensack, NJ, USA
M. Visocchi – Catholic University of Rome; Rome, Italy
J. Wang – UCLA School of Medicine; Santa Monica, CA, USA
C. Watts – University of Texas School of Law; Austin, TX, USA
K. Yonenobu – Osaka–Minami Medical Center; Kawachinagano, Osaka, Japan
S. Zeidman – The Uniformed Services University of the Health Sciences; Bethesda, MD, USA

SNI: Trauma

Associate Editors-in-Chief:

Naveed Ashraf

University of Health Sciences; Lahore, Pakistan

Iype Cherian

Krishna institute of Medical Sciences; Malkapur, India

Corrado Iaccarino

University of Modena and Reggio Emilia; Modena, Italy

Jutty Parthiban

Kovai Medical Center and Hospital; Tamil Nadu, India

Editors:

P.D. Le Roux – The Brain and Spine Center; Wynnewood, PA, USA

Eric Thelin – Karolinska University Hospital; Stockholm, Sweden

SNI: Unique Case Observations

Associate Editor-in-Chief:

S. Ather Enam, MD, PhD

Aga Khan University; Karachi, Sindh, Pakistan

Editors:

Bilal Khan Afridi – Lady Reading Hospital; Peshawar, Pakistan

Mehboob Alam – POF Hospital; Wah Cantt, Islamabad, Pakistan

Muhammad Alam – The Aga Khan University Hospital; Karachi, Pakistan

Altaf Alilaghari – The Aga Khan University Hospital; Karachi, Pakistan

Hafiza Fatima Aziz – The Aga Khan University Hospital; Karachi, Pakistan

Waqas Baqai – The Aga Khan University Hospital; Karachi, Pakistan

Saad Bin Anis – Shaukat Khanum Hospital; Lahore, Pakistan

Ehsan Bari – The Aga Khan University Hospital; Karachi, Pakistan

Ali Bydon – Johns Hopkins Hospital; Baltimore, MD, United States

Jaskaran Singh Gosal – All India Institute of Medical Sciences; Jodhpur, Rajasthan, India

Waeel Hamouda – Cairo University; Cairo, Egypt

Muhammad Imran – Dow University of Health Sciences; Karachi, Pakistan

Gohar Javed – Aga Khan University, Karachi, Pakistan

Saad Akhter Khan – The Aga Khan University Hospital; Karachi, Pakistan

Sara Khan – The Aga Khan University Hospital; Karachi, Pakistan

Rajesh Kumar – Dubai Medical College; Dubai, United Arab Emirates

Ian Lee – Henry Ford Hospital; Detroit, MI, United States

Farhan Mirza – Montreal Neurological Institute; Montreal, Quebec, Canada

Fatima Mubarak – The Aga Khan University Hospital; Karachi, Pakistan

Moshiur Rahman – Holy Family Red Crescent Medical College, Dhaka, Bangladesh

Bharath Raju – Rutgers University; New Brunswick, NJ, United States

Yaseen Rauf – Patel Hospital; Karachi, Pakistan

Shahzad Shamim – The Aga Khan University Hospital; Karachi, Pakistan

Zaeem Sultan – Queen Elizabeth Hospital; Birmingham, United Kingdom

Morteza Taheri – Iran University of Medical Sciences; Tehran, Iran

Zubair Tahir – Great Ormond Street Childrens Hospital; London, England

Badar Ujjan – Memom Medical Institute; Karachi, Pakistan

Naveed Zaman – Rehman Medical Institute; Peshawar, Pakistan

Contributing Editors

Ethics: L.F. Post, MD

Genetics: R. Rosenberg, MD

SNI Publications

Owners:

James I. And Carolyn R. Ausman Educational Foundation, a 501c3, Tax Deductible Private Operating Foundation

2 Comments



Carol L. Harte

Posted December 2, 2022, 10:55 am

I had occasion to read “COVID UPDATE: What is the Truth?” written by a Russell L. Blaylock, who serves on your editorial board in several non-scientific capacities (history, socioeconomics, etc.). Although I agree with many of his assertions, I am mystified that his excessive hyperbole, personal attacks and over-the-top rhetoric would remotely qualify this article for inclusion in the SNI (or any scientific) database. Perhaps your mission statement provides some justification in its reference to peer reviews and fact-based content while, at the same time, acknowledging that readers may interpret facts differently...but that’s a real stretch here. I am apolitical but instinctively question the credibility of any medical practitioner who refers (in this instance) to a Biden or Fauci in such disparaging terms. You may want to revisit your criteria for publication and also for membership on your editorial board.

Reply



Marvin Salgado Pérez

Posted January 20, 2023, 10:18 pm


ESPERO LA OCASIÓN EN QUE MI PAÍS NICARAGUA SEA INGRESADA AL CONSEJO EDITORIAL DE SNI.

Reply



Original Article

Periosteum-induced ossification effect in skull defect through interleukin-8 and NF-κB pathway: An experimental study with *Oryctolagus cuniculus* rabbits

Tedy Apriawan¹, Widjiati Widjiati², Dwikora Novembri Utomo³, Asra Al Fauzi¹, Eko Agus Subagio¹, Budi Utomo⁴, Andi Asadul Islam⁵, Abdul Hafid Bajamal¹, I. Ketut Sudiana⁶

¹Department of Neurosurgery, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, ²Faculty of Veterinary Medicine, Universitas Airlangga, Departments of ³Orthopaedics and Traumatology, ⁴Public Health and Preventive Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, ⁵Department of Neurosurgery, Faculty of Medicine, Universitas Hasanuddin – Dr. Wahidin Sudirohusodo Central General Hospital, Makassar, South Celebes, ⁶Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, East Java, Indonesia.

E-mail: Tedy Apriawan - drtedyapri@fk.unair.ac.id; Widjiati Widjiati - widjiati@fkh.unair.ac.id; Dwikora Novembri Utomo - dwikora_utomo@yahoo.com; Asra Al Fauzi - asrafauzi@yahoo.com; Eko Agus Subagio - eko.agus@fk.unair.ac.id; Budi Utomo - budiutomo@fk.unair.ac.id; Andi Asadul Islam - andiasadul@yahoo.com; Abdul Hafid Bajamal - hfbajamal@gmail.com; *I. Ketut Sudiana - i-ketut-s@fk.unair.ac.id



*Corresponding author:

I Ketut Sudiana,
Department of Anatomical
Pathology, Faculty of Medicine,
Universitas Airlangga – Dr.
Soetomo General Academic
Hospital, Surabaya, East Java,
Indonesia.

i-ketut-s@fk.unair.ac.id

Received : 15 January 2022

Accepted : 07 March 2022

Published : 15 April 2022

DOI

10.25259/SNI_62_2022

Quick Response Code:



ABSTRACT

Background: The purpose of this study was to analyze the response of inflammatory cytokines interleukin-8 (IL-8) and NF-κB to the closure of skull defect with periosteum as a scaffolding material in bone healing used after surgery.

Methods: Thirty *Oryctolagus cuniculus* rabbits underwent a craniotomy to create a 20 mm diameter round defect in the parietal bones. The parietal bones were returned to its place and stabilized by an internal plate fixation. The defects were either left empty or implanted with periosteum. At 6 weeks, the specimens were euthanized and examined.

Results: Histological examination showed a more well-developed formation of woven bone in the periosteum group. Immunohistochemical examinations showed that the use of periosteum in the closure of skull defects reduced the NF-κB and IL-8 response which affected the ossification process.

Conclusion: The experiment showed that the use of periosteum was linked with IL-8 and NF-κB downregulation toward ossification effects at any point throughout the trial. Periosteum usage might be beneficial as a scaffolding material in bone healing for autograft cranioplasty in animal model and could be applied to clinical practice.

Keywords: Cranioplasty, Interleukin-8, NF-κB, Periosteum flap

INTRODUCTION

Decompressive craniectomy is a surgical procedure that involves the removal of the skull bone to protect the patient from increasing intracranial pressure.^[8] If the circumstances permit, the bone will be reattached; this is referred to as cranioplasty. However, this treatment has certain risks for the patient. One of these potential problems is bone resorption. This kind of bone resorption may occur in individuals who have been fitted with bone and whose bone has been compromised in

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2022 Published by Scientific Scholar on behalf of Surgical Neurology International

terms of osteogenesis and vascularization as a result of the method involved.^[9]

Numerous cranioplasty methods have been developed to address the problems of bone resorption. Duhamel and John Hunter performed one of them utilizing the periosteum membrane, which has osteogenic characteristics. The periosteal membrane is made up of two layers: an exterior (fibrous and smooth) layer and an interior layer (cambium). The outer layer is densely packed with collagen and the reticular layer is densely packed with osteoblasts and osteoprogenitor cells, while the inner layer is densely packed with osteoblasts and osteoprogenitor cells.^[2] The mechanism of how pericranium-assisted autograph cranioplasty technique accelerates bone growth has not yet been clearly defined.

MATERIALS AND METHODS

Study design

The research protocol was authorized by the authorities of Universitas Airlangga, Department of Veterinary (Experimental Animal Studies, study number 2.KE.087.10.2020). The experimental study used 10 adults *Oryctolagus cuniculus* rabbits that were at least 5 months old and weighed between 4 and 5 kg. Animals were randomly divided into two groups (Group: with periosteum and without periosteum), each containing five rabbits. After 6 weeks, all animals were sacrificed and immunohistochemistry examination was performed.

Medication of animals

All operations were performed under general anesthesia, 65 mg kg⁻¹ ketamine and 4 mg kg⁻¹ xylazine, administered intramuscularly into the hind leg. Each animal received an intramuscular injection of 100,000 IU prophylactic benzylpenicillin. Following surgery, the animals received analgesics for 3 days, once daily, muscularly using phenylbutazone.

Craniotomy protocol

The rabbits' head was shaved between the eyes and the ears for surgical site preparation. The skin was sterilized with a solution of 10% iodine. After administering a local anesthetic (1 mL lidocaine and epinephrine solution); subcutaneously, a midline incision was created, and the skin and periosteum were reflected to reveal the skull. Round bone defects (diameter 20 mm) were bored into the inner cortex using a bone trephine with dural preservation. Following bone plate removal, the bone is returned and fixed using a long plate [Figure 1]. Periosteum sheets of each animal were applied over the bone defect in one group. Miniplates (Osteonic,




Figure 1: Autograft cranioplasty with long plate fixation following craniotomy procedure. (a) Defect placement without periosteum and (b) defect placement with periosteum.

Seoul, Korea) and surgical stitching (Silk 4.0, OneMed, Surabaya, Indonesia) for the fixation of bone and pericranial membranes are used throughout. The miniplate and thread utilized in this procedure are those used in neurosurgery. The skin was sutured properly. The animals were humanely euthanized with pentobarbital after 6 weeks of observation.

Histopathological examination

For 4 h, the specimens were preserved in cold 4% formaldehyde. The tissues were frozen in optimum cutting temperature compound (Sakura Finetek, Torrance, CA). Sections were cut to a thickness of 5 mm, attached on charged slides, and allowed to air dry for 5 min before staining. Staining with hematoxylin and eosin was carried out according to the procedure. The samples were analyzed using a light microscope (Fluorescence Microscope Olympus IX71).

Immunohistochemistry examination

Immunohistochemical responses were performed on rabbit skull tissue obtained 6 weeks after periosteum implantation. A monoclonal mouse anti-human NF- κ B antibody was used to determine the expression of the NF- κ B marker (ready-to-use antibody; 2J10D7; Novus Biologicals, CO, USA). Interleukin-8 expression was quantified using antibodies, anti-interleukin-8 (IL-8) antibody (monoclonal mouse, ready to use; MBS2025703; MybioSource, San Diego, USA). At room temperature, all sections were treated with the primary antibodies for 1 h. In addition, antigen determinants of all proteins were exposed by boiling sections for 15 min at 98°C in 10 mM citrate buffer (pH 6.5). Light microscope was used to examine the materials using a Fluorescence Microscope Olympus IX71 (Olympus, Tokyo, Japan).

RESULTS

Thirty animals were utilized in the research. All animals recovered normally and were terminated as scheduled. The two groups, rabbit skull following placement of periosteum

and without periosteum, were compared from all skull fragments. The final study comprised 30 animals, and [Figure 2] illustrates the skull after 6 weeks.

In those with periosteum, the woven bone formation could be observed on the defect walls without bridging the defects. The defect with the periosteum showed woven bone formation to the level of the original cortex [Figure 3]. In one of the sections without periosteum, a small area space reaction was observed in the soft tissue covering the defect. Otherwise, the space displayed is disjointed from the fault.

NF-κB expression/6 weeks

Statistical analysis showed a significant difference between each group at week 6 posttreatment ($P < 0.05$). Bone defects without periosteum showed the highest number of NF-B compared to cranioplasty with pericranium grafts [Figures 4 and 5].

IL-8 expression/6 weeks

Statistical analysis showed no significant difference ($P > 0.05$) between the use of periosteum and without periosteum at week 6 posttreatment. Bone defects without periosteum showed the highest IL-8 expression compared to the use of periosteum [Figures 6 and 7].




Figure 2: The defect of the rabbit skull following 6 weeks of observation. (a) Defect placement without periosteum and (b) defect placement with periosteum.




Figure 3: Hematoxylin and eosin stained sections of skull from animal after 6 weeks. (a) $\times 20$ of skull defect without periosteum and (b) $\times 20$ of skull defect after placement with periosteum.

DISCUSSION

Bone regeneration is a multistep process that begins with the mitotic expansion of progenitor cells at the sites of bone formation and continues with differentiation of these progenitor cells into functional osteoblasts. Our research sought to investigate if periosteum might be utilized as a scaffold for bone grafts, assisting in the multistep process of bone regeneration. In addition, we evaluated the osteogenic capacity of periosteum through the use of




Figure 4: Staining of NF-κB antibody at week 6 provides immunoreactivity to macrophage cells indicated by green arrows. Macrophage cells that do not exhibit immunoreactivity are indicated by red arrows. (a) With periosteum and (b) without periosteum.




Figure 5: The mean difference of NF-κB expression at 6 weeks of observation.




Figure 6: Staining of IL-8 antibody at week 6 provides immunoreactivity to macrophage cells indicated by green arrows. Macrophage cells that do not exhibit immunoreactivity are indicated by red arrows. (a) With periosteum and (b) without periosteum.




Figure 7: The difference in the mean value of interleukin-8 expression at 6 weeks of observation.

different osteoblastic phenotypic markers. Any bone graft material or scaffolding must possess the characteristics required to promote the growth of bone-forming cells in an optimum microenvironment. It was shown in the histologic examination that a more well-formed woven bone formation was observed in the periosteum group, indicating a faster bone healing compared to no periosteum flap.

The osteogenic capabilities of the osteoblasts were investigated for 6 weeks in our study. For assessing the osteogenic potential, Declercq, 2004, study and Aronow, 1990, study utilized time points ranging from 2 to 6 weeks. Investigators chose these different time points based on the component of osteoblasts they were studying. The 6-week endpoint appeared acceptable because we were evaluating the total response of the osteoblasts from all sources. Periosteum itself does not provide structural support, but it is ideal for covering faults and voids in the bone. It could be employed as a stand-alone substance to boost the osteogenic process. Periosteum can reduce the amount of artificial compound required to fill a defect, resulting in smaller donor sites and lower morbidity. The periosteum produced the least amount of IL-8 and NF- κ B in our investigation. As a result of this discovery, we can deduce that the healing process will accelerate if bone defects are covered with periosteum. Our research demonstrated that when the graft is exposed to a sufficient supply of cytokines, nutrients, and hormones, osteoprogenitor cells become activated and stain positive for osteogenic potential indicators.

NF- κ B (activated B-cell core factor) is a part of the transcription factor, which was first identified as a regulator of B-cell differentiation by its ability to bind to the B site of the kappa light chain gene in B cells. They were later shown to be involved in innate and adaptive immunity in response to pathogens and autoimmune stimuli and many aspects of normal cellular function.^[3] The role of NF- κ B in bone

was first identified in the mid-1990s when two groups of researchers produced NF- κ B1/2 double knockout (dKO) mice and found unexpectedly that they had tooth eruption failure and osteopetrosis in the absence of osteoclasts by producing both NF- κ B1 and -2 single knockout mice which shown modest immune deficiency, which translate into severe B-cell and T-cell differentiation defects and are in that respect similar to RANKL and RANK KO mice, which are produced later after RANKL.^[3]

NF- κ B itself is one of the cytokines that have been known to play an essential role in the tissue healing process.^[5] Various complex cascades related to tissue healing occur in the presence of activation of NF- κ B. Activation of NF- κ B causes a further activation process of more than 200 genes involved in inflammation, cell migration, cell proliferation, cell cycle, cell survival, and inhibition of apoptosis.^[5] Specific to bone tissue, inhibition of NF- κ B has been reported to be an effective means of inhibiting osteoclast activity and bone resorption.^[1]

IL-8 has been reported by a previous study by Edderkaoui^[4] to have an effect in bone healing conditions. In the case of human long bone fractures under 72 h, Hoff *et al.*^[6] reported an increase in IL-8 levels in the hematoma area arising from fractures and the surrounding bone marrow, with higher levels in the hematoma area itself. These data suggest an increased inflammatory reaction in the lesion area, which was a fracture, which was similar to the case we studied. In our study, the same thing happened wherein the craniotomy case on the rabbit skull, IL-8 in that area was increased.^[4,6] However, the decreased levels in the case of pericranium implantation could have several explanations, one of which is a reduction in the inflammatory reaction in the area affected by bone and/or pericranial grafts. Further studies may be needed to analyze further the inflammatory factors affected by the treatment and the healing of fractured lesions that occur in the bone tissue.^[7]

This research has several limitations, two of which are the considerably low number of samples and short period of follow-up for the new bone formation. This experiment also does not evaluate the critical bone defect size, which may play an important role in the practice of autograft osteoplasty procedure.

CONCLUSION

This study demonstrates that the periosteum has the elements required for osteogenesis. Periosteum also exhibits lower IL-8 production and NF- κ B laying down properties, according to the study. Because of its porous nature and the fact that it is autogenous, it causes no immunologic reaction, making it easy to assimilate into the defect. The periosteum is ideal for covering bone defects because it is readily available and has a layer-like consistency. Periosteum proved to be effective treatment evaluated in terms of promoting ossification and

inhibiting bone resorption inside bony defects produced in a rabbit calvarium model.

Acknowledgments

The authors would like to thank all who have contributed to the process and completion of this report, including the teaching staffs and fellow residents of the Department of Neurosurgery of Faculty of Medicine, Universitas Airlangga, Dr. Soetomo General Academic Hospital, Surabaya, Indonesia. We also want to thank the Faculty of Veterinary Medicine, the Department of Orthopaedics and Traumatology and Department of Public Health and Preventive Medicine, and the Department of Anatomical Pathology Faculty of Medicine Universitas Airlangga – Dr. Soetomo General Academic Hospital. We would also like to extend our gratitude toward the Department of Neurosurgery, Faculty of Medicine Universitas Hasanuddin – Dr. Wahidin Sudirohusodo Central General Hospital.

Declaration of patient consent

Institutional Review Board (IRB) permission obtained for the study.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Abu-Amer Y. NF- κ B signaling and bone resorption. *Osteoporos Int* 2013;24:2377-86.
2. Augustin G, Antabak A, Davila S. The periosteum. Part 1: Anatomy, histology and molecular biology. *Injury* 2007;38:1115-30.
3. Boyce BF. Advances in osteoclast biology reveal potential new drug targets and new roles for osteoclasts. *J Bone Miner Res* 2013;28:711-22.
4. Edderkaoui B. Potential role of chemokines in fracture repair. *Front Endocrinol (Lausanne)* 2017;8:1-8.
5. Heng MC. Wound healing in adult skin: Aiming for perfect regeneration. *Int J Dermatol* 2011;50:1058-66.
6. Hoff P, Gaber T, Strehl C, Schmidt-Bleek K, Lang A, Huscher D, *et al.* Immunological characterization of the early human fracture hematoma. *Immunol Res* 2016;64:1195-206.
7. Jiang WG, Sanders AJ, Ruge F, Harding KG. Influence of interleukin-8 (IL-8) and IL-8 receptors on the migration of human keratinocytes, the role of PLC- γ and potential clinical implications. *Exp Ther Med* 2012;3:231-6.
8. Joaquim AF, Mattos JP, Neto FC, Lopes A, de Oliveira E. Bone flap management in neurosurgery. *Rev Neurociencias* 2009;17:133-7.
9. Zanaty M, Chalouhi N, Starke RM, Clark SW, Bovenzi CD, Saigh M, *et al.* Complications following cranioplasty: Incidence and predictors in 348 cases. *J Neurosurg* 2015;123:182-8.

How to cite this article: Apriawan T, Widjiati W, Utomo DN, Al Fauzi A, Subagio EA, Utomo B, *et al.* Periosteum-induced ossification effect in skull defect through interleukin-8 and NF- κ B pathway: An experimental study with *Oryctolagus cuniculus* rabbits. *Surg Neurol Int* 2022;13:140.