

Combination effect of
methotrexate with Narrowband
Ultraviolet B (NB-UVB)
phototherapy in psoriasis
vulgaris patients in
dermatology and venerology
outpatient clinic Dr. Soetomo
General Academic

Submission date: 11-Oct-2022 06:39AM (UTC+0700)
by Budi Utomo .

Submission ID: 1922031993

File name: putra_022_-_3071_-_Afif_Nurul_Hidayati_-_Galley_1.pdf (295.78K)

Word count: 4459

Character count: 24559

Combination effect of methotrexate with Narrowband Ultraviolet B (NB-UVB) phototherapy in psoriasis vulgaris patients in dermatology and venereology outpatient clinic Dr. Soetomo General Academic Hospital, Surabaya, Indonesia



Emma Hidayati Sasmito¹, Iskandar Zulkarnain¹,
Muhammad Yulianto Listiawan¹,
Diah Mira Indramaya¹, Linda Astari¹, Budi Utomo², Afif Nurul Hidayati^{1,3*}

¹Department of Dermatology and Venereology, Dr. Soetomo General Academic Teaching Hospital, Surabaya

²Department of Public Health and Preventive Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya

³Department of Dermatology and Venereology, Universitas Airlangga Teaching Hospital, Surabaya

*Corresponding to:
Afif Nurul Hidayati; Department of Dermatology and Venereology, Dr. Soetomo General Academic Teaching Hospital, Surabaya, Indonesia; Department of Dermatology and Venereology, Universitas Airlangga Teaching Hospital, Surabaya;
afif_nurulhidayati@fk.unair.ac.id

Received: 2022-01-03
Accepted: 2022-03-16
Published: 2022-03-24

ABSTRACT

Background: Psoriasis is a chronic, recurrent, inflammatory skin disease that alters the quality of life. Systemic therapy and biologic agents are prescribed for severe and widespread psoriasis, but these drugs may have systemic side effects and immune suppression. Narrowband Ultraviolet B (NB-UVB) phototherapy remains one of the most effective and safe treatments for psoriasis.

Methods: This study was an observational retrospective cohort with one group pretest and post-test design. A retrospective analysis was done on 16 medical records of patients who met the inclusion criteria. The efficacy of the therapy was identified and analyzed using the SPSS version 17 for Windows.

Results: This study found that 16 psoriasis patients received NB-UVB phototherapy, 11 (68.75%) in men and 5 (31.25%) in women. The mean age was 47.31 years. There was a statistically significant difference in the reduction in PASI score before and after receiving combination methotrexate with 24 sessions of NB-UVB phototherapy ($p=0.000$); the delta PASI score was 45.02%. A statistical analysis based on the number of phototherapy sessions per week showed enhanced results when patients attended 3 phototherapy sessions per week ($p=0.001$).

Conclusion: There was a statistically significant reduction in PASI score in psoriasis vulgaris patients before and after receiving combination methotrexate with 24 sessions of NB-UVB phototherapy.

Keywords: Psoriasis Vulgaris, Phototherapy, Narrowband Ultraviolet B, Methotrexate, Human and Disease.

Cite This Article: Sasmito, E.H., Zulkarnain, I., Listiawan, M.Y., Indramaya, D.M., Astari, L., Utomo B., Hidayati, A.N. 2022. Combination effect of methotrexate with Narrowband Ultraviolet B (NB-UVB) phototherapy in psoriasis vulgaris patients in dermatology and venereology outpatient clinic Dr. Soetomo General Academic Hospital, Surabaya, Indonesia *Bali Medical Journal* 11(1): 116-121. DOI: 10.15562/bmj.v11i1.3071

INTRODUCTION

Psoriasis is a chronic inflammatory skin disease involving genetic, immune defects, hormones, and environmental factors. Psoriasis vulgaris is often characterized by thick, erythematous, well-demarcated, coarse, layered, silvery-white plaques.¹ Psoriasis vulgaris, also called plaque-type psoriasis, is the most common type, occurring in about 90% of patients.¹

Psoriasis is a chronic, recurrent, inflammatory skin disease that interferes with the quality of life, so appropriate, and effective therapy is needed.² The treatment options include topical therapy,

phototherapy, systemic, and biological agents. Systemic treatment and biologic agents are prescribed for severe and widespread psoriasis but cause systemic side effects and immune suppression, which are not tolerable for the patient. It requires alternative or combination therapy.² Narrowband Ultraviolet B (NB-UVB) phototherapy is relatively safe and effective as a treatment modality if used in erythemogenic doses with wavelengths between 311-313 nm, causing good remission in psoriasis lesions.^{1,3}

Based on those mentioned above, this study aims to evaluate the effect of combination methotrexate with 24

sessions of NB-UVB phototherapy in psoriasis vulgaris patients at Dermatology and Venereology Outpatient Department Dr. Soetomo Surabaya from March 2019 to February 2020.

METHODS

This study was an observational retrospective cohort with one group pretest and post-test design that aimed to evaluate the Psoriasis and Severity Index (PASI) score of psoriasis vulgaris patients who received a combination of methotrexate with 24 sessions of NB-UVB phototherapy using secondary data in the form of the medical record.

The inclusions criteria were all patients recorded in medical records with a diagnosis of moderate-severe psoriasis vulgaris who received combination therapy methotrexate with 24 sessions of NB-UVB phototherapy at Dermatology and Venereology Outpatient Department, Dr. Soetomo Surabaya, for the period of March 2019 - February 2020. The exclusions criteria were patients who stopped NB-UVB phototherapy treatment for more than 3 sessions consecutive weeks. The variable in this study was psoriasis vulgaris patients who received combination therapy methotrexate with 24 sessions NB-UVB phototherapy and assessed based on the Psoriasis and Severity Index (PASI) score. Those data inputted into data collection sheet to be analyzed using the Statistical Package for Social Sciences (SPSS) version 17 for Windows.

RESULTS

There were 16 psoriasis patients who received NB-UVB phototherapy at Dermatology and Venereology Outpatient Department Dr. Soetomo Surabaya for March 2019-February 2020, which met the inclusion criteria. Patients' demographic data were assessed based on gender, age, Fitzpatrick's skin phototype, and the severity degree based on the PASI scores.

The demographic distribution of psoriasis vulgaris patients received NB-UVB phototherapy (Table 1). The results showed that male patients were more female, 11 (68.75%) males and 5 (31.25%) females. The median age was 47.31 years, with the youngest being 22 years and the oldest was 64 years old. Based on the PASI scores, the severity degree was 5 patients (31.25%) for moderate and 11 patients (68.75%) for severe. Skin types based on Fitzpatrick's phototype were 2 patients (12.55%) with type III, 9 patients (56.25%)

with type IV, and 5 patients (31.25%) with type V (Table 1).

For the initial dose of NB-UVB phototherapy, 16 (100.00%) patients received an initial dose of NB-UVB phototherapy of 260 mJ (Table 1). The methotrexate dosage distribution in psoriasis vulgaris patients who received 24 sessions of NB-UVB phototherapy, as many as 1 (6.25%) patient received 2,5 mg methotrexate therapy, 3 (18.75%) patients received methotrexate 5 mg, 2 (12.5%) patients received 7,5 mg methotrexate, 5 (31.25%) patients received 10 mg

methotrexate, and 5 (31.25%) patients received 15 mg methotrexate (Table 1).

The additional systemic therapy in psoriasis vulgaris patients who received NB-UVB phototherapy, as many as 16 (100.00%) patients received folic acid and 16 (100.00%) patients received cetirizine, respectively, due to the itching (Table 1). The topical therapy in psoriasis vulgaris patients who received NB-UVB phototherapy, as many as 1 (6.25%) patient received clobetasol propionate, 15 (93.75%) patients received desoximetasone, 13 (81.25%) patients

Table 1. Psoriasis vulgaris patient distribution who received a combination of methotrexate with 24 sessions NB-UVB phototherapy

Variable	Total (n=16)
Gender, n (%)	
Male	11 (68.75)
Female	5 (31.25)
Age, Median (Minimum-Maximum)	47.31 (22-64)
Psoriasis and Severity Index (PASI) scores, n (%)	
Moderate (5-10)	5 (31.25)
Severe (>10)	11 (68.75)
Fitzpatrick's Skin Phototypes, n (%)	
I	0 (0.00)
II	0 (0.00)
III	2 (12.50)
IV	9 (56.25)
V	5 (31.25)
VI	0 (0.00)
Narrowband Ultraviolet B (NB-UVB) (260 mJ), n (%)	16 (100.00)
Methotrexate Distribution (mg), n (%)	
2.5	1 (6.25)
5	3 (18.75)
7.5	2 (12.5)
10	5 (31.25)
15	5 (31.25)
Additional Systemic Therapy, n (%)	
Folic Acid	16 (100.00)
Cetirizin	16 (100.00)
Topical Therapy Distribution, n (%)	
Clobetasol propionate	1 (6.25)
Desoximetasone	15 (93.75)
Mometasone furoate	13 (81.25)
Desonide lotion	3 (18.75)
Atopiclair	7 (43.75)
Vaseline album	7 (43.75)
Urea 10%	2 (12.50)
Ketoconazole shampoo	4 (25.00)

Table 2. Evaluation of the PASI scores of psoriasis vulgaris patients before and after receiving 24 sessions of NB-UVB phototherapy

PASI scores before phototherapy (Mean±SD)	PASI scores after 24 sessions of phototherapy (Mean±SD)	Delta PASI scores (Mean±SD)	Delta PASI scores (%)	p
12.65±4.82	7.20±4.82	5.45±3.05	45.02±21.10	0.000*

PASI: Psoriasis and Severity Index; SD: Standard Deviations; *Kruskal-Wallis Test: Statistically significant if p-value less than 0.05

Table 3. The comparison of the significant reduction of PASI scores in psoriasis vulgaris patients after receiving a combination of methotrexate with NB-UVB phototherapy

Variable	n	PASI scores before phototherapy	PASI scores after 24 sessions of phototherapy	Delta PASI scores	p
Gender					
Male	11	13.28	9.20	4.00	0.080
Female	5	13.20	6.51	6.69	
Age (Years Old)					
< 60	13	13.41	7.65	5.76	0.304
≥ 60	3	12.75	6.17	6.50	
Methotrexate (mg)					
≤ 5	4	10.80	6.30	4.25	0.323
> 5-10	7	13.78	7.97	5.81	
> 10	5	15.00	7.40	7.60	
Topical					
Steroid, emollient	12	13.87	8.03	5.79	0.458
Steroid, emollient, antifungal shampoo	4	11.25	4.92	6.33	
Phototherapy session (per week)					
3 sessions	7	13.00	4.65	8.27	0.001*
< 3 sessions	9	13.48	9.66	3.82	

PASI: Psoriasis and Severity Index; *Kruskal-Wallis Test: Statistically significant if p-value less than 0.05

received mometasone furoate, 3 (18.75%) patients received desonide lotion, 7 (43.75%) patients received atopiclair, 7 (43.75%) patients received vaseline album, 2 (12.50%) patients received urea 10%, and 4 (25.00%) scalp psoriasis patients received ketoconazole shampoo (Table 1).

The evaluation of the PASI scores in psoriasis vulgaris patients before and after receiving 24 sessions of NB-UVB phototherapy with the mean PASI scores before receiving NB-UVB phototherapy was 12.65±4.82. The mean PASI scores after receiving 24 sessions of NB-UVB phototherapy was 7.20±4.82. The mean delta PASI scores or the mean reduction of PASI scores before and after receiving 24 sessions of NB-UVB phototherapy was 5.45±3.05. The mean delta of the PASI scores (%) or the mean reduction of PASI scores (%) before and after receiving 24 sessions of NB-UVB phototherapy was 45.02±21.10. From the statistical analysis, the p-value was 0.000, indicating a statistically significant difference in reducing PASI scores before and after receiving 24 sessions of NB-UVB phototherapy (Table 2).

The comparison of the significance values for reducing PASI scores by gender, age, methotrexate dosage, combination topical therapy, and the number of phototherapy sessions per week in psoriasis

vulgaris patients receiving combination of methotrexate with 24 sessions of NB-UVB phototherapy (Table 3). The mean delta of PASI scores in males was 4.00 and in females was 6.69 (p = 0.080), which indicated that there was no statistically significant difference between male and female (p>0.05) towards the reduction of PASI scores in psoriasis vulgaris patients after receiving 24 sessions of NB-UVB phototherapy (Table 3).

The mean delta of PASI scores at < 60 years of age of 5.76 and at ≥ 60 years of 6.50 (p=0.304) which indicated that there was no statistically significant difference between ages < 60 years and ≥ 60 years of age for the reduction of PASI scores in psoriasis vulgaris patients after receiving a combination of methotrexate with 24 sessions of NB-UVB phototherapy (Table 3).

The reduction in the PASI scores is based on the methotrexate dosage using the Kruskal Wallis test. Kruskal Wallis test was a different test for 3 groups for abnormal distribution obtained the mean delta PASI scores at the methotrexate dose ≤ 5 mg was 4.25, > 5 mg - 10 mg was 5.81, > 10 mg as 7.60 (p=0.323), which indicated that there was no statistically significant difference based on the differences of the methotrexate dosage in the reduction in PASI scores in psoriasis vulgaris patients

after receiving 24 sessions of NB-UVB phototherapy (Table 3).

The PASI scores reduction based on topical combination therapy showed that the mean delta PASI scores for the combination of topical steroids and emollients was 5.79 and for the combination of topical steroids, emollients, and anti-fungal shampoos was 6.33 (p = 0.458), which indicates that there was no statistically significant difference between the combination of topical corticosteroid and emollient therapy with the combination of topical corticosteroid therapy, emollient and anti-fungal shampoo on the reduction of PASI scores in psoriasis vulgaris patients after receiving the combination of methotrexate with 24 sessions of NB-UVB phototherapy (Table 3).

The reduction in PASI scores based on the number of phototherapy sessions per week showed that the mean delta PASI scores on phototherapy with 3 sessions per week was 8.27 and < 3 sessions per week as 3.82 (p = 0.001), which showed that there was a statistically significant difference between the 3 sessions per week and < 3 sessions per week of phototherapy on the reduction of PASI scores in psoriasis vulgaris patients after receiving a combination of methotrexate with 24 sessions of NB-UVB phototherapy (Table 3).

DISCUSSION

This retrospective study found that 16 psoriasis vulgaris patients received a combination of methotrexate with 24 sessions of NB-UVB phototherapy at Dermatology and Venereology Outpatient Department Dr. Soetomo Surabaya from March 2019 to February 2020. The gender distribution of psoriasis vulgaris showed that males were more than females, 11 (68,75%) males and 5 (31,25%) females. The clinical manifestations of psoriasis in males were often more severe, so males often came to the hospital for treatment.^{1,3} Male also tend to have a higher BMI than women, and the influence of smoking and alcohol consumption, likely to be higher in males, is a risk factor³⁸ for psoriasis.^{4,5} A study in Korea found that the prevalence of psoriasis increased around 50 years old.⁶ This was due to the increasing comorbid risk factors in old age, namely obesity, metabolic disease, stress, and susceptibility to infections which could be a risk factor for psoriasis.⁷

A nonrandomized control trial on moderate-severe patients given NB-UVB phototherapy was shown to reduce the PASI scores. In general, NB-UVB phototherapy is recommended for moderate-severe¹³ psoriasis vulgaris.⁹ NB-UVB was the first-line phototherapy for moderate-severe plaque-type psoriasis vulgaris with an extensive skin involvement accompanied by thick plaques. In moderate-severe psoriasis vulgaris, there was an increase in the intensity of redness, desquamation, and extensive plaque thickness. NB-UVB phototherapy could improve redness, desquamation, and² extensive plaque thickness by inducing apoptosis of keratinocytes and T cells¹⁶ the epidermis and dermis, increasing immunosuppression by promoting migration of Langerhans cells from the epidermis and reducing degranulation of mast cells and release of histamine, and inducing changes in the cytokine of psoriasis.⁹

The characteristics of Asian skin based on Fitzpatrick's²² phototype obtained types III and IV in Chinese and Japanese and types IV and V in South Asian, Indian and Pakistani populations (10). The distribution of the initial dose of NB-UVB phototherapy was 16 (100%)

patients received an initial dose of 260 mJ. Erythema skin reactions due to burning during and after Narrowband ultraviolet B (NB-UVB) phototherapy may occur, which caused⁴⁸ dose adjustment.^{1,8} Determination of the initial dose of NB-UVB phototherapy with 2⁴⁴ mJ in all psoriasis vulgaris patients who received narrowband ultraviolet B (NB-UVB)⁵² phototherapy may be considered to avoid the risk of unwanted side effects such as erythema⁴⁶ due to burning during and after receiving NB-UVB phototherapy and the initial dose of NB-UVB phototherapy with 260 mJ was already efficient in achieving improvement.

In moderate-severe psoriasis, methotrexate could be combined with NB-UVB phototherapy.¹⁰ Methotrexate is a folic acid analog that competitively inhibits dihydrofolate reductase and other folate-related enzymes. The main effect of methotrexate is to inhibit the synthesis of thymine and purines, which results in a reduction of DNA and RNA synthesis, inhibits mitosis and cell proliferation,¹ including psoriatic keratinocytes. Inhibition of nucleic acid formation in activated T cells and keratinocytes is the main mechanism of action of methotrexate as an antiproliferative and immunomodulation in psoriasis vulgaris.¹¹

A recent meta-analysis of randomized controlled trials showed a 79% reduction in mucosal and gastrointestinal side effects when folic acid supplementation was given to the patients who received systemic methotrexate therapy for psoriasis vulgaris. Various⁴⁹ studies have been conducted to prevent the side effects of methotrexate by using folic acid supplementation while maintaining the therapeutic benefits of methotrexate.¹² The treatment for pruritus in psoriasis vulgaris is still debated because its pathogenesis is still not fully understood. Pruritus is more common in psoriasis vulgaris than in other types of psoriasis. It can interfere with the patients' quality of life. Many investigators had suggested impaired innervation and dysregulation of³ neuropeptide expression in psoriasis. As an important mediator in allergic, especially urticaria, histamine is not a causative agent in psoriasis.¹³

A study conducted topical administration of high potency

corticosteroids could improve in 75% of patients with moderate-severe psoriasis vulgaris. The mechanism⁶⁰ of action of topical corticosteroids is anti-inflammatory, antiproliferative, immunosuppressive, and vasoconstrictive. Topical corticosteroids in moderate-severe scalp psoriasis have also been effective in reducing¹⁹ plaque thickness. Combination with topical therapies such as corticosteroids and emollients while¹⁹ undergoing Narrowband Ultraviolet B (NB-UVB) phototherapy had been proven safe, effective and had the potential to increase efficacy.¹⁴

Anti-fungal ketoconazole was shown to be effective in reducing the colonization of *Pityrosporum* sp. in scalp psoriasis.¹⁵ It interacted with microbial products by causing an abnormal skin response in psoriasis. Microbial products that had a role in the pathogenesis of psoriasis were *Malassezia ovalis* (*Pityrosporum ovale*), a normal flora in the scalp, which could enhance the activation of the pathogenesis pathway, which could lead to psoriasis lesions. *Malassezia ovalis* is also believed to increase the development of scalp psoriasis.¹⁶

There was no difference between gender and age in the clearance rate of psoriasis vulgaris lesions after receiving Narrowband Ultraviolet B (NB-UVB) phototherapy. IL-17 and IL-23 levels after receiving NB-UVB phototherapy in both males and females, as well as ages, decreased significantly.⁴⁶ IL-17 and IL-23 levels are known to be associated with the pathogenesis of psoriasis as a predictive factor for the prognosis and the successful response of NB-UVB phototherapy.¹⁸ There was no difference between gender and age in reducing the PASI scores after receiving NB-UVB phototherapy. The significant difference in lowering PASI scores was the NB-UVB phototherapy.¹⁹

A study recommends the dose of methotrexate between 5-15 mg per week, 60% of patients achieved a reduction⁶² PASI scores by 75% for the initial dose of methotrexate 15 mg per week for 16 weeks, for a dose of 7,5 mg per week, a reduction in PASI scores was 40%.⁵⁷ A study on 39 patients who received a combination of narrowband ultraviolet B (NB-UVB)⁵⁵ phototherapy with methotrexate 0.4 mg/kg/week and maximum dosage of 25

mg/week showed that this combination therapy gave a rapid clinical improvement and there was a significant reduction in PASI scores in all psoriasis vulgaris patients.²¹

Although the dose of methotrexate at the set of NB-UVB phototherapy varies, the combination of methotrexate and NB-UVB phototherapy will work synergistically. Methotrexate and NB-UVB phototherapy both have antimitotic and antiproliferative functions against T lymphocytes, and both have an anti-inflammatory property. Methotrexate can help reduce the squama and infiltration of psoriasis lesions, thus allowing NB-UVB phototherapy to penetrate deeper into the dermis. It has greater effectiveness in reducing dermal T lymphocytes.²²

A study in Taiwan showed that 98,4% of psoriasis vulgaris patients received topical therapy with corticosteroids and emollients, there was a significant improvement in psoriasis vulgaris plaque lesions.²³ Other topical treatments combined with topical anti-fungal therapy were safe and effective, especially in scalp psoriasis. In psoriasis, there is an involvement of *Malassezia ovalis* (*Pityrosporum ovale*), a normal flora in the scalp. Still, in some psoriasis patients, even though *Malassezia ovalis* (*Pityrosporum ovale*) is a normal flora, it can enhance the activation of cytokines, chemokines, and PGE2. It is associated with Th1 and Th2 from mononuclear cells, leading to new scalp lesions.¹⁶

The three sessions per week of NB-UVB phototherapy caused a high cumulative dose, which increased inflammatory factor expression changes through the NF- κ B signaling pathway.²⁴ NB-UVB phototherapy with a high cumulative dose could also increase inhibition of pro-inflammatory factor expression and increase anti-inflammatory factor expression through increased induction of lymphocyte apoptosis, reduced pro-inflammatory cytokine production, down-regulation of the Th17 signaling pathway, and reduction of T cell depletion, so it could be faster in reducing or thinning plaque in psoriasis lesions which was an important mechanism for psoriasis therapy.²⁵ The limitation of this study is the relatively small group of patients and the data was based on medical records, which

caused a numerous risk of bias. Further study is needed to evaluate the PASI scores over 24 sessions of NB-UVB phototherapy to know the number of sessions that effectively reduce the PASI-75 scores in psoriasis vulgaris patients.

CONCLUSION

The conclusion is there was a statistically significant difference in the reduction in PASI scores before and after receiving combination of methotrexate with 24 sessions of NB-UVB phototherapy. Combining methotrexate with 24 sessions of NB-UVB phototherapy may be considered for patients with moderate-severe psoriasis vulgaris who failed with previous therapy. The importance of routine control is for the compliance to receive 3 sessions of narrowband ultraviolet B (NB-UVB) phototherapy per week in order to achieve a reduction in the PASI-75.

CONFLICT OF INTEREST

There is no conflict of interest regarding the manuscript

ETHICS CONSIDERATION

This research has obtained ethical approval from the Ethics Committee of Dr. Soetomo General Academic Teaching Hospital Surabaya (0142/LOE/301.4.2/X/2020).

FUNDING

None.

AUTHOR CONTRIBUTIONS

All authors equally contribute to the study from the conceptual framework, data acquisition, data analysis until reporting the results of the study through publication.

REFERENCES

1. Armstrong AW, Read C. Pathophysiology, Clinical Presentation, and Treatment of Psoriasis: A Review. *JAMA*. 2020;323(19):1945-1960.
2. Mahmood AN, Saeed MY. Extracutaneous manifestations of adult psoriatic patients attending Dermatology Teaching Center in Slemani, Iraq. *Bali Medical Journal*. 2020;9(1):121-124.

3. Dogra S, De D. Narrowband ultraviolet B in the treatment of psoriasis: the journey so far!. *Indian J Dermatol Venereol Leprol*. 2010;76(6):652-661.
4. Ito T, Takahashi H, Kawada A, Iizuka H, Nakagawa H; Japanese Society For Psoriasis Research. Epidemiological survey from 2009 to 2012 of psoriatic patients in Japanese Society for Psoriasis Research. *J Dermatol*. 2018;45(3):293-301.
5. Colombo D, Cassano N, Bellia G, Vena GA. Gender medicine and psoriasis. *World J Dermatol*. 2014;3(3):36-44.
6. Han JH, Lee JH, Han KD, Seo HM, Bang CH, Park YM, et al. Epidemiology and Medication Trends in Patients with Psoriasis: A Nationwide Population-based Cohort Study from Korea. *Acta Derm Venereol*. 2018;98(4):396-400.
7. Parisi R, Symmons DP, Griffiths CE, Ashcroft DM; Identification and Management of Psoriasis and Associated Comorbidity (IMPACT) project team. Global epidemiology of psoriasis: a systematic review of incidence and prevalence. *J Invest Dermatol*. 2013;133(2):377-385.
8. Van TN, Van TH, Minh PPT, Trong HN, Van TC, Huu ND, et al. Efficacy of Narrow - Band UVB Phototherapy versus PUVA Chemophototherapy for Psoriasis in Vietnamese Patients. *Open Access Maced J Med Sci*. 2019;7(2):227-230.
9. Nakamura M, Farahnik B, Bhutani T. Recent advances in phototherapy for psoriasis. *F1000Res*. 2016;5:F1000 Faculty Rev-1684.
10. Chan IL, Cohen S, da Cunha MG, Maluf LC. Characteristics and management of Asian skin. *Int J Dermatol*. 2019;58(2):131-143.
11. Pathirana D, Ormerod AD, Saig P, Smith C, Spuls PI, Nast A, et al. European S3-guidelines on the systemic treatment of psoriasis vulgaris. *J Eur Acad Dermatol Venereol*. 2009;23 Suppl 2:1-70.
12. Rai T, Singh SK. Role of folic acid supplementation in reducing side effects of oral methotrexate in patients of psoriasis: A study. *Indian J Drugs Dermatol*. 2016;2(1):83-86.
13. Domagala A, Szeptowski J, Reich A. Antihistamines in the treatment of pruritus in psoriasis. *Postepy Dermatol Alergol*. 2017;34(5):457-463.
14. Elmets CA, Lim HW, Stoff B, Connor C, Cordoro KM, Lebwohl M, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management and treatment of psoriasis with phototherapy. *J Am Acad Dermatol*. 2019;81(3):775-804.
15. Wang TS, Tsai TF. Managing Scalp Psoriasis: An Evidence-Based Review. *Am J Clin Dermatol*. 2017;18(1):17-43.
16. Blakely K, Gooderham M. Management of scalp psoriasis: current perspectives. *Psoriasis (Auckl)*. 2016;6:33-40.
17. Elghandour TM, Youssef Sel S, Aly DG, Abd Elhameed MS, Abdel Moneim MM. Effect of Narrow Band Ultraviolet B Therapy versus Methotrexate on Serum Levels of Interleukin-17 and Interleukin-23 in Egyptian Patients with Severe Psoriasis. *Dermatol Res Pract*. 2013;2013:618269.

18. Rácz E, Prens EP, Kurek D, Kant M, de Ridder D, Mourits S, et al. Effective treatment of psoriasis with narrow-band UVB phototherapy is linked to suppression of the IFN and Th17 pathways. *J Invest Dermatol.* 2011;131(7):1547-58.
19. Saleky S, Bulur I, Saraçoğlu ZN. Narrowband UVB treatment increases serum 25-hydroxyvitamin D levels in patients with chronic plaque psoriasis. *Cutis.* 2017;99(6):431-435.
20. Dogra S, Mahajan R. Systemic methotrexate therapy for psoriasis: past, present and future. *Clin Exp Dermatol.* 2013;38(6):573-588.
21. Khadka D, Agrawal S, Dhali T. Methotrexate plus narrow band ultraviolet B (NB-UVB) versus methotrexate alone in the treatment of moderate to severe plaque psoriasis: A randomized clinical trial. *Nepal Journal of Dermatology, Venereology & Leprology.* 2016;13(1):12-23.
22. Al-Hamamy HR, Al-Mashhadani SA, Mustafa IN. Comparative study of the effect of narrowband ultraviolet B phototherapy plus methotrexate vs. narrowband ultraviolet B alone and methotrexate alone in the treatment of plaque-type psoriasis. *Int J Dermatol.* 2014;53(12):1531-1535. d
23. Chang YT, Chen TJ, Liu PC, Chen YC, Chen YJ, Huang YL, et al. Epidemiological study of psoriasis in the national health insurance database in Taiwan. *Acta Derm Venereol.* 2009;89(3):262-6.
24. Matos TR, Ling TC, Sheth V. Ultraviolet B radiation therapy for psoriasis: Pursuing the optimal regime. *Clin Dermatol.* 2016;34(5):587-593.
25. Ye J, Huang H, Luo G, Yin L, Li B, Chen S, et al. NB-UVB irradiation attenuates inflammatory response in psoriasis. *Dermatol Ther.* 2020;33(4):e13626.



This work is licensed under a Creative Commons Attribution

Combination effect of methotrexate with Narrowband Ultraviolet B (NB-UVB) phototherapy in psoriasis vulgaris patients in dermatology and venereology outpatient clinic Dr. Soetomo General Academic

ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** D Pathirana. "European S3-Guidelines on the systemic treatment of psoriasis vulgaris", Journal of the European Academy of Dermatology and Venereology, 10/2009
Publication 1%
- 2** f1000researchdata.s3.amazonaws.com
Internet Source 1%
- 3** www.advrehab.org
Internet Source 1%
- 4** erepo.unud.ac.id
Internet Source 1%
- 5** www.nature.com
Internet Source 1%
- 6** Jianzhou Ye, Hong Huang, Guangyun Luo, Lihua Yin, Bocheng Li, Sixuan Chen, Hongying Li, Yang Yang, Xuesong Yang. " irradiation 1%

attenuates inflammatory response in psoriasis ", Dermatologic Therapy, 2020

Publication

-
- | | | |
|-----------------|--|------|
| 7 | Razvigor Darlenski, Radostina Deliyska, Lina Tabet Al - Sadek, Evgeniya Hristakieva, Joachim W. Fluhr. "Epidermal carotenoid levels in vivo of patients with plaque psoriasis: Effects of narrow band UVB phototherapy", Photodermatology, Photoimmunology & Photomedicine, 2020 | 1 % |
| Publication | | |
| 8 | "Pigmentary Skin Disorders", Springer Science and Business Media LLC, 2018 | 1 % |
| Publication | | |
| 9 | Hayashi, Mitsuha, Yoshinori Umezawa, Osamu Fukuchi, Toshihiro Ito, Hidehisa Saeki, and Hidemi Nakagawa. "Efficacy and safety of ustekinumab treatment in elderly patients with psoriasis", The Journal of Dermatology, 2014. | <1 % |
| Publication | | |
| 10 | oamjms.eu | <1 % |
| Internet Source | | |
| 11 | repository.ubaya.ac.id | <1 % |
| Internet Source | | |
| 12 | Parlak, Nehir, Nihal Kundakci, Adem Parlak, and Bengu Nisa Akay. "Narrowband | <1 % |

ultraviolet B phototherapy starting and incremental dose in patients with psoriasis: comparison of percentage dose and fixed dose protocols : Narrow band ultraviolet B regimens", Photodermatology Photoimmunology & Photomedicine, 2014.

Publication

13

Sandra Pena, Dane Hill, Steven R. Feldman. "Ultraviolet B Phototherapy", Elsevier BV, 2016

Publication

<1 %

14

abs.kafkas.edu.tr

Internet Source

<1 %

15

www.pubfacts.com

Internet Source

<1 %

16

Nilesh Rarokar, Shailendra Gurav, Dadasaheb M. Kokare, Vijay Kale, Nishikant A. Raut. "Phototherapy: A critical review", Elsevier BV, 2022

Publication

<1 %

17

moam.info

Internet Source

<1 %

18

A. Pacifico, G. Damiani, P. Iacovelli, R.R.Z. Conic et al. "Photoadaptation to ultraviolet B TL01 in psoriatic patients", Journal of the European Academy of Dermatology and Venereology, 2020

Publication

<1 %

19 Engin Sezer, Ilker Etikan. "Local narrowband UVB phototherapy vs. local PUVA in the treatment of chronic hand eczema", Photodermatology, Photoimmunology & Photomedicine, 2007
Publication <1 %

20 Submitted to Intercollege
Student Paper <1 %

21 link.springer.com
Internet Source <1 %

22 www.researchgate.net
Internet Source <1 %

23 Jacobi, Arnd, Stephan J. Rustenbach, and Matthias Augustin. "Comorbidity as a predictor for drug survival of biologic therapy in patients with psoriasis", International Journal of Dermatology, 2015.
Publication <1 %

24 Submitted to University of Ulster
Student Paper <1 %

25 www.royan.org
Internet Source <1 %

26 Asmaa M. Ele-Refaei, Fatma M. El-Esawy. "Effect of Narrow-Band Ultraviolet B Phototherapy and Methotrexate on MicroRNA

(146a) Levels in Blood of Psoriatic Patients",
Dermatology Research and Practice, 2015

Publication

27

Caren Garber, Ami Saraiya, Shiu-chung Au, Yahya Argobi, Ryan Mansfield, Natalie Smith, Alice Gottlieb. "Analysis of Gender Differences in the Efficacy of Psoriasis Treatment Modalities Resulting in Clearance", Psoriasis Forum, 2018

Publication

<1 %

28

Joanna Narbutt, Adam Reich, Zygmunt Adamski, Grażyna Chodorowska et al. "Psoriasis in children. Diagnostic and therapeutic recommendations of the Polish Dermatological Society. Part 2", Dermatology Review, 2021

Publication

<1 %

29

basicmedicalkey.com

Internet Source

<1 %

30

coek.info

Internet Source

<1 %

31

e-journal.unair.ac.id

Internet Source

<1 %

32

encyclopedia.pub

Internet Source

<1 %

33

libproxy.galencollege.edu

Internet Source

<1 %

34

"Handbook of Psoriasis", Wiley, 2004

Publication

<1 %

35

H. S. Midde, M. Priyadarssini, M. Rajappa, M. Munisamy, P. S. Mohan Raj, S. Singh, G. Priyadarshini. "Interleukin - 9 serves as a key link between systemic inflammation and angiogenesis in psoriasis", Clinical and Experimental Dermatology, 2020

Publication

<1 %

36

Jiaoquan Chen, Xin Zeng, Quan Chen, Bihua Liang, Liqian Peng, Huaping Li, Yi Tang, Shanshan Ou, Huilan Zhu. "Efficacy of NB-UVB as Add-on Therapy to Antihistamine in the Treatment of Chronic Urticaria: A Systematic Review and Meta-analysis", Dermatology and Therapy, 2021

Publication

<1 %

37

Liisa Väkevä, Sirkka Niemelä, Meri Lauha, Rafael Pasternack et al. "Narrowband ultraviolet B phototherapy improves quality of life of psoriasis and atopic dermatitis patients up to 3 months: Results from an observational multicenter study", Photodermatology, Photoimmunology & Photomedicine, 2019

Publication

<1 %

38

Yi - Hsin Ho, Hsiao - Yun Hu, Yun - Ting Chang, Chung - Pin Li, Chen - Yi Wu.

<1 %

"Psoriasis is associated with increased risk of bullous pemphigoid: A nationwide population - based cohort study in Taiwan",
The Journal of Dermatology, 2019

Publication

39

core.ac.uk

Internet Source

<1 %

40

journals.plos.org

Internet Source

<1 %

41

medwinpublishers.com

Internet Source

<1 %

42

profiles.sc-ctsi.org

Internet Source

<1 %

43

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

44

www.e-agmr.org

Internet Source

<1 %

45

www.ejmanager.com

Internet Source

<1 %

46

www.mdpi.com

Internet Source

<1 %

47

www.medwave.cl

Internet Source

<1 %

48

www.thieme-connect.com

Internet Source

<1 %

49

www.tobaccoinduceddiseases.org

Internet Source

<1 %

50

Akasaka, Emiko, Tomotaka Mabuchi, Yasuaki Manabe, Eiichiro Yahagi, Azusa Yamada-Hiruma, Hanako Yamaoka, Tomoko Kojima, Masayuki Kato, Norihiro Ikoma, Akira Ozawa, and Yasuo Haruki. "Long-term efficacy of psoriasis vulgaris treatments: Analysis of treatment with topical corticosteroid and/or vitamin D3 analog, oral cyclosporin, etretinate and phototherapy over a 35-year period, 1975-2010", *The Journal of Dermatology*, 2013.

Publication

<1 %

51

Amit G. Pandya, John E. Harris, Mark Lebwohl, Iltefat H. Hamzavi, Kathleen Butler, Fiona I. Kuo, Shaoceng Wei, David Rosmarin. "Addition of Narrow-Band UVB Phototherapy to Ruxolitinib Cream in Patients With Vitiligo", *Journal of Investigative Dermatology*, 2022

Publication

<1 %

52

European Handbook of Dermatological Treatments, 2015.

Publication

<1 %

53

I.H. Kwon. "Could colorimetric method replace the individual minimal erythema dose (MED) measurements in determining the initial dose of narrow-band UVB treatment for psoriasis

<1 %

patients with skin phototype III-V? :
Colorimetric L* value as the initial dose of
narrow-band UVB treatment", Journal of the
European Academy of Dermatology and
Venereology, 02/2012

Publication

54

Khadiga Sayed Sayed, Eman Raafat Said,
Faisal Nouredin Mohammed, Mira Algorhary,
Laila Ahmed Rashed. "Down-regulation of
tissue levels of serine protease inhibitor
(vaspin) in psoriasis vulgaris patients: a
possible mechanism of narrowband
ultraviolet B radiation", Archives of
Dermatological Research, 2019

Publication

<1 %

55

Khadka, Dhan Keshar, S Agrawal, and T K
Dhali. "Methotrexate plus narrow band
ultraviolet B (NBUVB) versus methotrexate
alone in the treatment of moderate to severe
plaque psoriasis: A randomized clinical trial",
Nepal Journal of Dermatology Venereology &
Leprology, 2016.

Publication

<1 %

56

Metwally, D., K. Sayed, R. Abdel Hay, and L.
Rashed. "Reduction in tissue plasmin: a new
mechanism of action of narrowband
ultraviolet B in psoriasis", Clinical and
Experimental Dermatology, 2014.

Publication

<1 %

57

N. Oiso. "Successful treatment of lichen amyloidosis associated with atopic dermatitis using a combination of narrowband ultraviolet B phototherapy, topical corticosteroids and an antihistamine", *Clinical and Experimental Dermatology*, 12/2009

Publication

<1 %

58

Teng - Li Lin, Chun - Ying Wu, Yun - Ting Chang, Chao - Keui Juan, Chi - Chiang Chen, Shi - Hang Yu, Yi - Ju Chen. "Risk of skin cancer in psoriasis patients receiving long - term narrowband ultraviolet phototherapy: Results from a Taiwanese population - based cohort study", *Photodermatology, Photoimmunology & Photomedicine*, 2019

Publication

<1 %

59

"RHEUMATOID ARTHRITIS ? TREATMENT", *APLAR Journal of Rheumatology*, 8/2006

Publication

<1 %

60

Amitava Mitra, Ercem Atillasoy. "Chapter 7 Topical Therapies for Psoriasis", *IntechOpen*, 2012

Publication

<1 %

61

Aybala Erek Toprak, Emin Ozlu, Tugba Kevser Uzuncakmak, Emre Yalcinkaya, Sadik Sogut, Ayşe Serap Karadag. "Neutrophil/Lymphocyte Ratio, Serum Endocan, and Nesfatin-1 Levels in Patients with Psoriasis Vulgaris Undergoing

<1 %

Phototherapy Treatment", Medical Science Monitor, 2016

Publication

62

Kelly W. Burak, Stefan J. Urbanski, Mark G. Swain. "Successful treatment of refractory type 1 autoimmune hepatitis with methotrexate", Journal of Hepatology, 1998

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On