


# Berkala Ilmu Kesehatan Kulit dan Kelamin



*Periodical of Dermatology and Venereology*

## Research / Retro

### Cat Contact as A Risk Factor for Tinea Capitis Infection

DOI : 10.20473/bikk.V34.1.2022.1-4

Evy Ervianti , Addia Salsabila , Yuri Widia , Arthur Pohan Kawilarang

1-4

 Abstract : 558

 PDF : 231

[PDF](#)

### Ceramide is More Effective than Shea Butter in Maintaining Skin Acidity

DOI : 10.20473/bikk.V34.1.2022.5-9

Zilpa Widyastuti , Suci Widhiati

5-9

 Abstract : 593

 PDF : 363

[PDF](#)

### Characteristics of Atopic Dermatitis Patients who Underwent Skin Prick Test


DOI : 10.20473/bikk.V34.1.2022.10-14

Wisnu Triadi Nugroho , Evy Ervianti , Iskandar Zulkarnain , Afif Nurul Hidayati , Damayanti Damayanti , Budi

Utomo , Cita Rosita Sigit Prakoeswa

10-14

 Abstract : 406

 PDF : 195

[PDF](#)


### Comparison of the Efficacy of Topical Clindamycin versus Niacinamide in the Treatment of Mild to Moderate Acne Vulgaris: a Systematic Review

DOI : 10.20473/bikk.V34.1.2022.15-22

Eden Leonita , Trisniartami Setyaningrum , Mohammad Fathul Qorib , Damayanti -

15-22

 Abstract : 795

 PDF : 350


[PDF](#)

### Distribution of Leprosy Patients with and without Plantaris Ulcers

DOI : 10.20473/bikk.V34.1.2022.23-28

Vida Cahlia Novita Sari , Medhi Denisa Alinda , Subur Prajitno , Cita Rosita Sigit Prakoeswa

23-28

 Abstract : 266

 PDF : 125

[PDF](#)

### Epidemiology of Leprosy in Indonesia: a Retrospective Study

DOI : 10.20473/bikk.V34.1.2022.29-35

Cita Rosita Sigit Prakoeswa , Ramona Sari Lubis , Qaira Anum , Fifa Argentina , Sri Linuwih Menaldi , Hendra Gunawan , Renni Yuniati , Nur Rachmat Muliando , Agnes Sri Siswati , Dhelya Widasmara , Luh Made Mas Rusyati , Enricco Hendra Mamuja , Vitayani Muchtar , Regitta Indira Agusni , Bagus Haryo Kusumaputra , Medhi Denisa Alinda , Muhammad Yulianto Listiawan

29-35

📄 Abstract : 1397

📄 PDF : 651

[PDF](#)

### Skin Aging Profile in Tertiary Hospital: a Descriptive Study

DOI : 10.20473/bikk.V34.1.2022.36-45

Tasya Wikassa , Damayanti Damayanti , Irfiansyah Irwadi , Trisniartami Setyaningrum , Rahmadewi Rahmadewi , Diah Mira Indramaya , Menul Ayu Umborowati

36-45

📄 Abstract : 292

📄 PDF : 151

[PDF](#)

### The Profile of Leprosy Patients in Aceh: Retrospective Study

DOI : 10.20473/bikk.V34.1.2022.46-51

Nanda Earlia , Sulamsih Sri Budini , Dea Silvia Ramadana , Mikyal Bulqiah , Aqil Yuliawan Tasrif , Randika Richard R , Muhammad Adji Pangestu , Annisa Raihani , Auni Batrisyia , Eva Cahyaningsih , Fanny Adlin Nurafika , Istiqomah Genepo , Nadira Sabrina Mufti

46-51

📄 Abstract : 317

📄 PDF : 160

[PDF](#)

### Validity and Reliability Test of Vitiligo Quality of Life Index (VitiQoL) Indonesian Version in Vitiligo Patients

DOI : 10.20473/bikk.V34.1.2022.52-58

Jeffrey Giantoro , Fajar Waskito , Sunardi Radiono

52-58

📄 Abstract : 299

📄 PDF : 124

[PDF](#)

## Literature Review

### Secukinumab Therapy in Psoriasis Management

DOI : 10.20473/bikk.V34.1.2022.59-65

Ira Yunita , Sylvia Anggraeni

59-65

📄 Abstract : 372

📄 PDF : 227

[PDF](#)

## Case Report

### A Case Report of Tinea Capitis in Children: Utility of Trichoscopy

DOI : 10.20473/bikk.V34.1.2022.66-72

Citra Dwi Harningtyas , Evy Ervianti , Linda Astari , Sylvia Anggraeni , Yuri Widia

66-72

📄 Abstract : 438

📄 PDF : 236

[PDF](#)

### **Recurrent Verruca Vulgaris Treated with Combination of 80% Trichloroacetate and Electrosurgery: a Case Report**

DOI : 10.20473/bikk.V34.1.2022.73-76

**Nyoman Yoga Maya Pramita , Ketut Kwartantaya Winaya , I Gusti Nyoman Darmaputra**

73-76


📄 Abstract : 852

📄 PDF : 483

[PDF](#)

---

---



## Editorial Team



**Prof. Dr. Cita Rosita Sigit Prakoeswa, dr., Sp.KK(K),**

**FINSDV, FAADV**

**Editor In Chief**

Faculty of Medicine, Airlangga University Surabaya, Indonesia, Indonesia

200000-0003-3232-095X

pL7tjnEAAAAJ&hl=en

**Scopus'** 57189894608



**dr. Damayanti Damayanti, Sp.KK(K), FINSDV**

**Editorial Manager**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

**Scopus'** -



**dr. Irmadita Citrashanty, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

**Scopus'** 57204810428



**dr. Maylita Sari, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

**Scopus'** -



**dr. Septiana Widyantari, Sp.KK, FINSDV**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

**Scopus'** 57210805038





**dr. Medhi Denisa Alinda, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

**Scopus**<sup>®</sup> 57202009260

-



**dr. Hasnikmah Mappamasing, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

-

**Scopus**<sup>®</sup> 57210793193

-



**dr. Sylvia Anggraeni, Sp.KK**

**Editorial Handling**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

ugPVNPQAAAAJ&hl

**Scopus**<sup>®</sup> 57210634125

-



**Dr. Afif Nurul Hidayati, dr., Sp.KK(K), FINS DV, FAADV**

**Editorial Board**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

ctqbEVQAAAAJ

**Scopus**<sup>®</sup> 57196050143

-



**dr. Evy Ervianti, Sp.KK(K), FINS DV, FAADV**

**Editorial Board**

Faculty of Medicine, Airlangga University Surabaya, Indonesia, Indonesia, Indonesia

-

Qsz4JHgAAAAJ

**Scopus**<sup>®</sup> 57201074072

-



**dr. Linda Astari, Sp.KK(K), FINS DV, FAADV**

**Editorial Board**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-

kLaGucOAAAAJ&hl

**Scopus**<sup>®</sup> 57196052464

-








**dr. Astindari Astindari, Sp.KK**

**Editorial Board**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

 -  
 -  
**Scopus'** -  
 -



**Prof. Dr. med. Isaak Effendy**

**Editorial Board**

Academic Teaching Hospital Bielefeld Klinikum Bielefeld, Germany  
Dermatology and Allergology, Germany




 -  
 -  
**Scopus'** 7005243490  
 -



**Prof. dr. Taruna Ikrar, M.Pharm, MD., Ph.D**

**Editorial Board**

Biomedical Science, The National Health University, California, United States




 -  
 xtqyDXsAAAAJ&hl  
**Scopus'** 14621460200  
 -



**dr. Sri Manovita Pateda, S.Ked., M.Kes**

**Editorial Board**

Ehime University, Japan, Japan

 -  
 -  
**Scopus'** -  
 -



**dr. Jimmy Rusdian Masjkur, M.Sc., Ph.D**

**Editorial Board**

University Hospital Carl Gustav Carus Dresden, Germany, Germany




 -  
 -  
**Scopus'** 36621238000  
 -



**Robert Tungadi, S.Si., M.Si., Apt**

**Editorial Board**

Aachen University, Germany, Germany

 -  
 \_zunygwAAAAJ&hl  
**Scopus'** 56609263400  
 -





**dr. David Sudarto Oeiria, Sp.KK, FINSDV, FAADV**

**Editorial Board**

Division of Dermatologic Surgery, Dept. of Dermatology, Huashan Hospital - Fudan University, Shanghai, China

-  
 -  
**Scopus'** -  
 -



**dr. Yuri Widia, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-  
 vgSrq4oAAAAJ&hl  
**Scopus'** 57210799099  
 -



**dr. Menul Ayu Umborowati, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-  
 -  
**Scopus'** -  
 -



**dr. Bagus Haryo Kusumaputra, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-  
 -  
**Scopus'** 57202017415  
 -



**dr. Putri Hendria Wardhani, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-  
 -  
**Scopus'** 57210789350  
 -



**dr. Regitta Indira Agusni, Sp.KK**

**Assistant Editor**

Faculty of Medicine, Airlangga University Surabaya, Indonesia

-  
 JWEo4rIAAAAJ&hl  
**Scopus'** -  
 -





# Characteristics of Atopic Dermatitis Patients who Underwent Skin Prick Test

Wisnu Triadi Nugroho<sup>1</sup>, Evy Ervianti<sup>1</sup>, Iskandar Zulkarnain<sup>1</sup>, Afif Nurul Hidayati<sup>1</sup>, Damayanti<sup>1</sup>, Budi Utomo<sup>2</sup>, Cita Rosita Sigit Prakoeswa<sup>1</sup>

<sup>1</sup> Department of Dermatology and Venereology, Faculty of Medicine, Universitas Airlangga/ Dr. Soetomo General Academic Hospital, Surabaya-Indonesia

<sup>2</sup> Department of Public Health Science, Faculty of Medicine, Universitas Airlangga, Surabaya-Indonesia

## ABSTRACT

**Background:** Skin prick test (SPT) is indicated in type I hypersensitivity reactions (rapid type), among which is Atopic Dermatitis (AD). However, the results of SPT for the tested allergens are not always accompanied by history of known allergens in AD patients. **Purpose:** To identify the characteristics of AD patients who underwent SPT. **Methods:** This was a descriptive retrospective study to determine the demographics, allergy history, SPT results, and positive and negative clinical relevance in AD patients treated at Allergy Immunology Division of Dermatology and Venereology Outpatient Clinic Dr. Soetomo General Academic Hospital Surabaya-Indonesia from January 2017 to December 2019. **Results:** A total of 284 AD patients were included. The adult age group (20-59 years) predominated the AD patient's visit every year with 58 patients (60.42%), 47 patients (46.08%), and 65 patients (75.58%) in 2017, 2018, and 2019, respectively. Only 60 AD patients underwent SPT, where most patients were allergic to dust with 30 patients (50.0%). House dust mite (HDM) allergen had the highest percentage of positive SPT result with 41 patients (63.3%). The greatest clinical relevance was found in cat fur allergen, where all patients (100.0%) had clinical relevance to their SPT result. **Conclusion:** This study shows that most AD patients were in adult age group and had a history of dust allergy. The highest percentage of positive SPT result was found in HDM allergen and the greatest clinical relevance was found in cat fur allergen.

**Keywords:** allergen, atopic dermatitis, clinical relevance, skin prick test, human and medicine.

Correspondence: Cita Rosita Sigit Prakoeswa, Department of Dermatology and Venereology, Faculty of Medicine, Universitas Airlangga/ Dr. Soetomo General Academic Hospital, Surabaya, Jl. Mayjen Prof. Dr. Moestopo No. 6-8 Surabaya 60131, Indonesia. Phone: (031) 5501609, e-mail: cita-rosita@fk.unair.ac.id

## BACKGROUND

Atopic dermatitis (AD) is a chronic, often recurrent, inflammatory skin disease and also multifactorial disease involving the interaction between host (impaired skin barrier function), agent (microbial or viral), and environment (allergen exposure). Exposure to allergens, which often causes AD, might come from vegetable and animal protein sources.<sup>1</sup> Skin prick test (SPT) is indicated in type I hypersensitivity reactions (rapid type), among which is AD. This examination could identify an individual's sensitivity to various allergens.<sup>2</sup> However, the results of SPT for the tested allergens are not always accompanied by history of known allergens in AD patients.

This study was a descriptive retrospective study that aimed to determine the characteristics of AD patients who underwent SPT. This study could provide information regarding the characteristics of AD patients who underwent SPT to better support AD diagnosis and management in the future.

## METHODS

This study was a descriptive retrospective study to determine the demographics, allergy history, SPT results, and positive and negative clinical relevance in AD patients treated at Allergy Immunology Division of the Dermatology and Venereology Outpatient Clinic of Dr. Soetomo General Academic Hospital Surabaya, Indonesia from January 2017 to December 2019. The sampling technique used was total population technique, by collecting secondary data from AD patient's medical record. The inclusion criteria were all patients recorded in the medical record with diagnosis of AD based on Hanifin and Rajka criteria and underwent SPT procedures. The data and results obtained were then entered into a data collection sheet for analysis using the Statistical Package for Social Science (SPSS) version 17. This study has received ethical approval from the Hospital Ethics Committee of Dr. Soetomo General Academic Hospital Surabaya (Ref.No: 0059/LOE/301.4.2/VII/2020).

**RESULTS**

There were total of 284 AD patients with 96, 102, and 86 patients in 2017, 2018, 2019, respectively. The adult age group (20-59 years) predominated AD

patient's visits every year with 58 patients (60.42%), 47 patients (46.08%), and 65 patients (75.58%) in 2017, 2018, and 2019, respectively. Table 1 outlines AD patient demographics by age group.

**Table 1** The age group of atopic dermatitis patients

Age Group	Visit Year					
	2017		2018		2019	
	n	Percentage (%)	n	Percentage (%)	n	Percentage (%)
Children (<10 years)	0	0.00	2	1.96	4	4.65
Adolescent (10-19 years)	28	29.17	45	44.12	7	8.14
Adult (20-59 years)	58	60.42	47	46.08	65	75.58
Elderly (≥60 years)	10	10.42	8	7.84	10	11.63
Total	96	100.00	102	100.00	86	100.00

**Table 2** History of allergy distribution on atopic dermatitis patients

Allergen	(n)	Percentage (%)
Dust	30	50.0
Broiler chicken	9	15.0
Shrimp	15	25.0
Cow's milk	1	1.7
Egg	14	23.3
Kapok	0	0.0
Beef	1	1.7
Milk fish	0	0.0
Sea fish	10	16.7
Crab	0	0.0
Peanut	3	5.0
Chocolate	0	0.0
Rice	0	0.0
Wheat flour	0	0.0
Potato	0	0.0
Banana	0	0.0
Papaya	1	1.7
Pineapple	0	0.0
Orange	0	0.0
Organic chicken	0	0.0
Chicken skin	0	0.0
Dog's fur	0	0.0
Cat's fur	0	0.0

<sup>a</sup>One patient might have more than one history of allergen.

Among all AD patients who visited the Allergy Immunology Division of the Dermatology and Venerology Department Outpatient Clinic of Dr.

Soetomo General Academic Hospital within 2017-2019, there were 60 patients who underwent SPT. Prior to SPT, history of the patient's allergy was obtained.

**Table 3** Skin prick test results on all allergens of atopic dermatitis patients

Allergen Type	Positive SPT	Positive SPT	Negative SPT	Negative SPT
	Result	Result Percentage (%)	Result	Result Percentage (%)
HDM	41/60	63.3	19/60	36,7
Broiler chicken	8/60	13.3	52/60	86,7
Shrimp	5/60	8.3	55/60	91,7
Cow's milk	3/60	5.0	57/60	95,0
White egg	2/45	4.4	43/45	95,6
Kapok	3/15	20.0	12/15	80,0
Beef	1/15	6.7	14/15	93,7
Milk fish	4/15	26.7	11/15	73,3
Tuna	0/15	0.0	15/15	100,0
Pindang fish	3/15	20.0	12/15	80,0
Crab	5/15	33.3	10/15	66,7
Peanut	1/15	6.7	14/15	93,3
Chocolate	3/15	20.0	12/15	80,0
Rice	4/15	26.7	11/15	73,3
Wheat flour	4/15	26.7	11/15	73,3
Potato	3/15	20.0	12/15	80,0
Banana	4/15	26.7	11/15	73,3
Papaya	1/15	6.7	14/15	93,3
Pineapple	2/15	13.3	13/15	86,7
Orange	2/15	13.3	13/15	86,7
Organic chicken	4/15	26.7	11/15	73,3
Chicken's skin	2/15	13.3	13/15	86,7
Dog's fur	4/9	44.4	5/9	55,6
Cat's fur	0/4	0.0	4/4	100,0

SPT = Skin Prick Test; HDM = House Dust Mite

**Table 4** Clinical relevance between allergy history and skin prick test result of atopic dermatitis patients

Allergen Type	Clinical Relevance	
	Yes	No
HDM	37 (61.7%)	23 (38.3%)
Broiler chicken	47 (78.3%)	13 (31.7%)
Shrimp	44 (73.3%)	16 (26.7%)
Cow's milk	38 (84.4%)	7 (15.6%)
White egg	56 (93.3%)	4 (6.7%)
Kapok	12 (80.0%)	3 (20.0%)
Beef	5 (55.6%)	4 (44.4%)
Milk fish	4 (100.0%)	0 (0.0%)
Tuna	14 (93.3%)	1 (6.7%)
Pindang fish	14 (93.3%)	1 (6.7%)
Crab	8 (53.3%)	7 (46.7%)
Peanut	7 (46.7%)	8 (53.3%)
Chocolate	10 (66.7%)	5 (33.3%)
Rice	12 (80.0%)	3 (20.0%)
Wheat flour	14 (93.3%)	1 (6.7%)
Potato	11 (73.3%)	4 (26.7%)
Banana	11 (73.3%)	4 (26.7%)
Papaya	12 (80.0%)	3 (20.0%)
Pineapple	12 (80.0%)	3 (20.0%)
Orange	14 (93.3%)	1 (6.7%)
Organic chicken	13 (86.7%)	2 (13.3%)
Chicken's skin	13 (86.7%)	2 (13.3%)
Dog's fur	11 (73.3%)	4 (26.7%)
Cat's fur	13 (86.7%)	2 (13.3%)

HDM = House Dust Mite

Table 2 outlines allergy distribution on AD patients. Of all 60 patients, most patients were allergic to dust, with 30 patients (50.0%). Table 3 showed SPT results on all allergens of AD patients. Among all patients who underwent SPT, house dust mite (HDM) allergen had the highest percentage of positive SPT results, with 41 patients (63.3%). House dust mite, broiler chicken, shrimp, and cow's milk allergen were tested on all patients (60 patients), egg white allergen was tested on 45 patients, dog's fur allergen was tested on 9 patients, cat's fur allergen was tested on 4 patients, and the remaining allergens were tested on 15 patients.

Table 4 describes the clinical relevance between allergy history and SPT results in AD patients. The greatest clinical relevance was found in cat's fur allergen, where all patients (100.0%) had clinical relevance to their SPT results.

## DISCUSSION

The adult (20-59 years) and adolescent (10-19 years) age group were the largest age groups who visited the outpatient clinic in 2017-2019. These results were similar to a retrospective study of AD patient's profile at Dermatology and Venereology Outpatient Clinic of Prof. Dr. R. D. Kandou Hospital, Manado conducted by Febriansyah *et al.* Among 461 AD patients, 269 patients (58.35%) belonged to the >12 years age group.<sup>3</sup> A retrospective study by Ratnaningtyas and Hutomo at Allergy Immunology Division of the Dermatology and Venereology Outpatient Clinic of Dr. Soetomo General Academic Hospital Surabaya-Indonesia reported that among 272 new AD patients, the largest age group was 15-24 years (33.3%).<sup>4</sup>

In this study, the highest prevalence of allergy history was dust allergy (50.0%). This result was consistent with a study by Wistiani and Notoatmojo who studied the correlation between allergen exposure to allergic diseases in children at Dr. Kariadi General Hospital, Semarang. From 44 children with allergy, 18 patients (34.1%) were diagnosed with AD and most patients (53.8%) had dust allergy.<sup>5</sup> Another study by Ludfi *et al* involving samples from junior high school students in Surabaya, Indonesia evaluated the factors that might affect allergic diseases in children. Using multivariate logistic regression test, they concluded that birth order ( $p=0.005$ ) and HDM exposure ( $p=0.018$ ) affected clinical manifestations of children with AD.<sup>6</sup> Aeroallergens or inhaled allergens can be divided into two types, indoor and outdoor allergens. Indonesia is a tropical country, where household allergens such as HDM are more common.<sup>7</sup>

Among the 284 AD patients who visited our outpatient clinic, only 60 patients underwent SPT. This

could be caused by the patient's condition at the time of visit who might have absolute or relative contraindications to SPT. Absolute contraindications for SPT include diffuse dermatological conditions (examination should be performed on healthy and normal skin), severe dermographism, uncooperative patients, and patients who could not stop their antihistamine or other drug use that might affect SPT results. Relative contraindications for SPT include unstable/severe persistent asthma, pregnancy (because of the mild risk of anaphylaxis with hypotension and uterine contractions), infants less than two years of age, and patients taking beta blockers and angiotensin-converting enzyme (ACE) inhibitors.<sup>8</sup>

The allergen with the highest positive SPT result was HDM allergen (63.3%). This result was in accordance with a study by Zeppa *et al* on AD patients aged 18 years and over at the Dermatology and Venereology Clinic in a hospital in Italy involving 332 AD patients, where they reported that positive SPT results for aeroallergens and food allergens were found in 227 (68,4 %) AD patients.<sup>9</sup> A retrospective study by Natallya and Barakbah at Dermatology and Venereology Outpatient Clinic of Dr. Soetomo General Academic Hospital, Surabaya from 2007 to 2012 reported that out of 19 AD patients with positive SPT results, most patients showed positive result to HDM allergen with 8 patients (42%).<sup>10</sup> A retrospective descriptive study using medical records and SPT results of patients by Garna *et al* at Allergy Clinic of Immanuel Hospital, Bandung which aimed to determine the profile of allergic patients reported a total of 206 patients with a history of allergies, where most allergic patients who came to the clinic had clinical presentation of dermatitis with 89 patients (43.20%). From all allergic patients, HDM gave the most positive SPT results with 108 patients (52.43%).<sup>11</sup>

Clinical relevance is determined if there is a match between allergy history and SPT result. In this study, the greatest clinical relevance was found in cat's fur allergen (100.0%), followed by cow's milk, beef, milkfish, and papaya (93.3%). These results were in accordance with a study by Asha'ari *et al* at Allergy Clinic of a hospital in Malaysia which compared the specific IgE test result with SPT result for allergy diagnosis and involved 99 patients with allergy history. They stated that the sensitivity and specificity of SPT for cat's fur were 90.5% and 44.9%, respectively.<sup>12</sup> Another study by Čelakovská *et al* investigated the specificity and sensitivity of SPT, specific IgE test, and Atopy Pacht Test (APT) in 301 AD patients aged over 14 years with food allergies at Dermatology Department of a hospital in Czech Republic. They found that the specificity and sensitivity of SPT for

cow's milk were 97.9% and 33.3%, respectively.<sup>13</sup> A meta-analysis examining the accuracy of SPT, specific IgE test, and APT in food allergy patients involving 24 studies with 2831 samples reported that SPT had a sensitivity of 88% (76–94%) and a specificity of 68% (56–77%) for cow's milk.<sup>14</sup> A study from Anggraeni *et al* that compared SPT results of new local and imported allergen extracts with specific IgE in 45 AD patients at Dr. Soetomo General Academic Hospital, Surabaya, Indonesia reported that cow's milk allergen had a specificity of 88.23%, while its sensitivity only reached 27.27%.<sup>15</sup> Sensitivity was defined as proportion of patients who had a history of a particular disease and reported having a positive test result, meanwhile specificity was defined as proportion of patients who did not have a history of a particular disease and reported having a negative test result.<sup>12</sup> However, only four allergens (HDM, broiler chicken, cow's milk, and shrimp) were tested in all patients in this study. This might explain the very high percentage (100.0%) of clinical relevance for cat's fur allergen.

Based on the results of this study, it can be concluded that most AD patients were in adult age group and had dust allergy history. The highest percentage of positive SPT result was found in HDM allergen. The greatest clinical relevance was found in cat's fur allergen, followed by egg white, beef, milkfish, chocolate, and papaya allergen.

## REFERENCES

1. Simpson EL, Leung DYM, Eichenfield LF, Boguniewicz M. Atopic dermatitis. Fitzpatrick's Dermatology 9<sup>th</sup> Edition. 2018. p. 363–84.
2. Percival E, Bhatia R, Preece K, McElduff P, McEvoy M, Collison A, *et al*. Reproducibility of serum IgE, Ara h2 skin prick testing and fraction of exhaled nitric oxide for predicting clinical peanut allergy in children. Allergy Asthma Clin. Immunol 2016; 12: 1–9.
3. Febriansyah JPE, Kapantow GM, Hariyanto A. Profil dermatitis atopik di poliklinik kulit dan kelamin RSUP Prof. dr. R.D Kandou Manado periode Januari 2010 - Desember 2012. J. Biomedik 2015; 7: 23–8.
4. Ratnaningtyas WD, Hutomo M. Penelitian retrospektif: pengobatan topikal pada pasien dermatitis atopik (a retrospective study: topical therapy in atopic dermatitis patient). Berkala Ilmu Kesehatan Kulit dan Kelamin 2015; 28(3): 210-6.
5. Wistiani W, Notoatmojo H. Hubungan pajanan alergen terhadap kejadian alergi pada anak. Sari Pediatri 2016; 13(3):185-90.
6. Ludfi AS, Agustina L, Fetarayani D, Baskoro A, Gatot S, Effendi C. Asosiasi penyakit alergi atopi anak dengan atopi orang tua dan faktor lingkungan. Jurnal Penyakit Dalam 2012; 13: 53-62.
7. Siregar SP. Peran alergi makanan dan alergen hirup pada dermatitis atopik. Sari Pediatri 2016; 6: 155-8.
8. Heinzerling L, Mari A, Bergmann KC, Bresciani M, Burbach G, Darsow U, *et al*. The skin prick test - European standards. Clin. Transl. Allergy 2013; 3: 1–10.
9. Zeppa L, Bellini V, Lisi P. Atopic dermatitis in adults. Dermatitis 2011; 22: 40–6.
10. Natallya FR, Barakbah J. Retrospective Study of Skin Prick Test in Atopic Dermatitis Patients at DermatoVenereology Outpatient Clinic of Dr. Soetomo General Hospital Surabaya During 2007-2012. Berkala Ilmu Kesehatan Kulit dan Kelamin 2015; 27(1): 9-16.
11. Garna DR, Lucianus J, Ivone J. Descriptive study on skin prick test in allergy clinic Immanuel hospital Bandung Indonesia. J. Med. Heal 2017; 1: 558–67.
12. Asha'ari ZA, Suhaimi Y, Yusof RA, Rushdan I, Maraina CHC. Comparison of serum specific ige with skin prick test in the diagnosis of allergy in malaysia. Med. J. Malaysia 2011; 66: 202–6.
13. Čelakovská J, Krcmova I, Bukac J, Vaneckova J. Sensitivity and specificity of specific IgE, skin prick test and atopy patch test in examination of food allergy. Food Agric. Immunol 2017; 28: 238–47.
14. Soares-Weiser K, Takwoingi Y, Panesar SS, Muraro A, Werfel T, Hoffmann-Sommergruber K, *et al*. The diagnosis of food allergy: A systematic review and meta-analysis. Allergy Eur. J. Allergy Clin. Immunol 2014; 69: 76–86.
15. Anggraeni S, Umborowati MA, Damayanti, Endaryanto A, Prakoeswa CRS. The accuracy of Indonesian new local Skin Prick Test (SPT) allergen extracts as diagnostic tool of IgE-mediated atopic dermatitis. Indian Journal of Forensic Medicine & Toxicology 2021; 15(3): 33.