

The Asian Pacific Journal of Cancer Prevention (APJCP) is a monthly electronic journal, publishing papers in a wide spectrum of cancer sciences. The journal was launched in 2000 as the official publication of the Asian Pacific Organization for Cancer Prevention (APOCP) and its first volume with 51 articles was published in 2000. Since then, the journal has published 17 volumes and close to 8,000 articles.



The APJCP with its distinguished, diverse, and Asia-wide team of editors, reviewers, and readers, ensure the highest standards of research communication within the cancer sciences community across Asia as well as globally. The APJCP accepts a manuscript on the whole spectrum of cancer sciences.

All manuscripts published in the Asia Pacific Journal of Cancer Prevention, are under the terms of the Creative Commons Attribution Licence. This permits anyone to copy, distribute, transmit and adapt the published work, provided the original work and source are appropriately cited.

New Editorial policy

Close to 10% of manuscripts submitted and accepted for publication in APJCP suffer a low quality of English writing which is a burden on the Journal's editorial system. In order to address this, the journal's editorial team has set up a quality assurance policy. Starting from July 1, 2017, all manuscripts.. Read more

Events:

Publication Information

- Publisher**
West Asia Organization for Cancer Prevention (WAACP), APOCP's West Asia Chapter
- Editor-in-Chief**
Alireza Mosavi Jarrahi, MSPH, Ph.D.
- Honorary Editor-in-Chief**
Malcolm Anthony Moore, Ph.D.
- Associate Editor**
Prof. David Roder
Dr. Abhishek Shankar
- Founder**
Yip Cheng-Har, MD, FACS
Keun-Young Yoo, MD, Ph.D.
Kazuo Tajima, MD
- International Advisory Board**
Prof. Niveen Abu-Rmeileh
Prof. Rema Gvamichava
Prof. Mohammad Esmail Akbari
Prof. Ali Montazeri
Dr. Shahab Uddin
Prof. Goura Kishor Rath
Dr. Farhana Badar



EDITORIAL BOARD APJCP



ASIAN PACIFIC JOURNAL OF CANCER PREVENTION



Official publication of the Asian Pacific Organization for Cancer Prevention (APOCP)

Home Browse Journal Info Guide for Authors Submit Manuscript Reviewers Contact Us Login Register

Editorial Board

Editor-in-Chief

Alireza Mosavi Jarrahi, MSPH, Ph.D.

Associate Prof. of Epidemiology

President, Asian Pacific Organization for Cancer Prevention
Associate Prof. of Epidemiology
Dept of Social Medicine, Medical School, Shahid Beheshti
University of Medical Sciences, Tehran, Iran
Faculty of Health Sciences, Simon Fraser University, Burnaby,
BC, Canada

rmosavi@yahoo.com

[+ More](#)

Honorary Editor-in-Chief

Malcolm Anthony Moore, Ph.D.

Toxicologist

Director-Coordination, Asia Pacific Organization for Cancer
Prevention, Korean Cancer Center, Goyang, Korea

apocpcontrol@gmail.com

Associate Editor

Prof. David Roder

Professor of Cancer Epidemiology

Chair Cancer Epidemiology and Population Health, Beat
Cancer Project, University of South Australia, Adelaide,
Australia

ca.linkedin.com/in/david-roder-54699329

Dr. Abhishek Shankar

Preventive Oncology, Radiation Oncology

Assistant Professor

Department of Preventive Oncology,
Dr. B. R. Ambedkar Institute Rotary Cancer Hospital
All India Institute of Medical Sciences,
Delhi, India

doc.abhishankar@gmail.com

[+ More](#)

Founder

Yip Cheng-Har, MD, FACS

Surgery

Prof. of Gynaecological Oncology
Faculty of Medicine, University Tunku Abdul Rahman, Kuala
Lumpur, Malaysia

chenghar.yip@gmail.com

[+ More](#)

Keun-Young Yoo, MD, Ph.D.

Prof. of Cancer Epidemiology

President, APOCP


President, Armed Forces Capital Hospital
Professor, Seoul National University College of Medicine, Seoul,
Korea

bkyyoo@gmail.com

[+ More](#)

 **Kazuo Tajima, MD**


Pathology

 Prof. of Pathology


Director, Aichi Cancer Center Research Institute, Ozaka, Japan

 ktajima@clin.medic.mie-u.ac.jp


 More

 **Prof. You Lin Qiao, M.D. Ph.D.,**

Epidemiology

 Director, Dept. of Cancer Epidemiology


Cancer Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College
Beijing, China


 qiaoy@cicams.ac.cn


International Advisory Board


 **Dr. Khuseynov Zafardzhon**

Oncologist


 Director of the Republic Oncological Scientific Center,
Ministry of public health, Republic of Tajikistan

 zafhab@gmail.ru

 +992907710535

 **Prof. Niveen Abu-Rmeileh**

Community Medicine

 Associate prof. of Community Medicine


Institute of Community and Public Health,
Birzeit University,
Birzeit - Palestine

 nrmcleh@gmail.com


 More

 **Prof. Rema Gvamichava**

Prof. of Surgery


 A. Gvamichava National Cancer Center,

Tbilisi State Medical University
Tbilisi, Georgia

 doctorrema@yahoo.com

 **Prof. Mohammad Esmail Akbari**

Prof. of Surgery

 Cancer Research Center,

Shahid Beheshti University of Medical Sciences, Tehran, Iran.

 profmeakbari@gmail.com

 **Dr. Mohsen Asadi-Lari**


Prof. of Epidemiology

 Deputy for International Relations

Ministry of Health, Tehran, Iran

Prof. of Epidemiology, School of Public Health

Iran University of Medical Sciences, Tehran, Iran

 mohsen.asadi@yahoo.com


 **Dr. Shahab Uddin**

Cancer Biology

 Hamad Medical Corporation

Academic Health System


Doha, Qatar

 suddinkhan@gmail.com

 More

 **Prof. Goura Kishor Rath**


Radiation Oncology

 Chief, DR BRAIRCH,

Professor, Department of Radiation Oncology, and


Head, National Cancer Institute,

AIIMS, Ansari Nagar, New Delhi-110029, India

 gkrath2006@gmail.com

 **Dr. Farhana Badar**


Cancer Registration and Epidemiology

 Sr. Biostatistician & Cancer Epidemiologist

Cancer Registry & Clinical Data Management

Shaukat Khanum Memorial Cancer Hospital & Research Center,

Lehore, Pakistan

 farhana@skm.org.pk

 **Prof. Kourosh Holakouie-Naieni**

Epidemiology

📍 *Professor in Epidemiology,
Department of Epidemiology, School of Health, Tehran
University of Medical Sciences;
General Secretary, Iranian Epidemiological Association, Tehran,
Iran*

✉ holakoik@hotmail.com

Editorial Board

 **Prof. Jeff Dunn**

📍 *Chief Executive Officer of Cancer Council Queensland,
School of Social Science, University of Southern Queensland,
Queensland, Australia*

✉ jeffdunn@cancerqld.org.au

[+ More](#)

 **Dr. Xinen Huang**

Prof. of Clinical Oncology

📍 *Department of Oncology, Nanjing Medical University
Affiliated Cancer Hospital of Jiangsu Province, Nanjing,
Jiangsu 210009, China*

✉ huangxinen06@163.com

 **Prof. Shahid Pervez**

Histopathology

📍 *Departments of Pathology & Laboratory Medicine
Faculty of Health Sciences, Medical College, The Aga Khan
University, Karachi, Pakistan*

✉ shahid.pervez@aku.edu

 **Prof. Salim M. Adib**

Prof. of Public Health

📍 *Professor of Public Health, American University of Beirut,
Beirut, Lebanon
WHO, East Mediterranean Regional Office
Nasr City, Cairo, Egypt*

✉ adibs@emro.who.int

 **Dr. Anton Barchuk**

Thoracic surgery

📍 *Petrov Research Institute of Oncology
Saint Petersburg, Russian Federation*

✉ barchuk.anton@gmail.com

 **Prof. Murat Gultekin**

Gynaecological Oncologist

📍 *Sağlık Bakanlığı-Turkish Ministry of Health, Hacettepe
University, European Society of Gynaecological Oncology
(ESGO)*

🌐 www.linkedin.com/in/murat-gultekin-268136100

✉ mrgultekin@yahoo.com

[+ More](#)

 **Prof. Abas Ghaderi**

Cancer Immunologist

📍 *Shiraz Institute for Cancer Research (A.G.), Medical School,
Shiraz University of Medical Sciences, Shiraz, Iran*

✉ ghaderia@gmail.com

 **DR. Elsayed I Salim**

📍 *Faculty of Science, Tanta University, Tanta, Egypt*

✉ elsalem_777@yahoo.com

 **Dr. Reza Shirkohi**

Medical Genetic

📍 *Cancer Institute, Tehran University of Medical Sciences,
Tehran, Iran*

✉ r_shirkohi@yahoo.com

 **Prof. Mehrnoosh Doroudchi**

Immunology

📍 *Associate Professor of Immunology
Director of Memory T cell Laboratory
Department of Immunology, School of Medicine
Shiraz University of Medical Sciences, Shiraz-Iran*

✉ mdoroud@sums.ac.ir



DAFTAR ISI

 **Volume & Issue: Volume 22, Issue 10, October 2021, Pages 3053-3404** 

1. COVID-19 Vaccination and Cancer, the Need for more Data

Pages 3053-3054

Seyed Alireza Javadinia; James S Welsh; Alireza Mosavi Jarrahi

View Article  PDF 201.52 K

2. Opportunities and Challenges of Tobacco Control Policy at District Level in Indonesia: A Qualitative Analysis

Pages 3055-3060

Sepri Yunarman; Aries Munandar; Abdillah Ahsan; Ali Akbarjono; Dian Kusuma

View Article  PDF 655.82 K

3. Association between Smokeless Tobacco Use and Risk of Periodontitis in Asian Countries: A Systematic Review and Meta-Analysis

Pages 3061-3074

Abhishek Mehta; Venkitachalam Ramanarayanan; Vineetha Karuveettil; Chandrashekar Janakiram

View Article  PDF 916.25 K

4. Overexpression of c-Met is Associated with Poor Prognosis in Glioblastoma Multiforme: A Systematic Review and Meta-Analyses

Pages 3075-3080

Jellyca Anton; Sudibio Sudibio; Handoko Handoko; Tiara Bunga Mayang Permata; Henry Kodrat; Endang Nuryadi; Henry Sofyan; Eka Susanto; Rahmad Mulyadi; Renindra Ananda Aman; Soehartati Gondhowiardjo

View Article  PDF 372.31 K

5. Factors Affecting Delayed Presentation and Diagnosis of Breast Cancer in Asian Developing Countries Women: A Systematic Review

Pages 3081-3092

Farida Briani Sobri; Adang Bachtiar; Sonar Soni Panigoro; Dumilah Ayuningtyas; Hardya Gustada; Patria Wardana Yuswar; Aqsha Azhary Nur; R. Cita Resti Anantia Putri; Anggindita Diah Widihiyati

View Article  PDF 511.1 K

6. Mobile Health Interventions for Improving Colorectal Cancer Screening Rates: A Systematic Review and Meta-Analysis

18. Evaluation of Silver Nanoparticles Caffeic Acid Complex Compound as New Potential Therapeutic Agent against Cancer Incidence in Mice

Pages 3189-3201

Tahany Saad Abdelwahab; Rania Ellisy Abdelhamed; Eman Noaman Ali; Nahla Ahmed Mansour; Mohga Shafik Abdalla

[View Article](#)  PDF 1.74 MB

19. Direct Relationship between Interleukin-10 Gene Polymorphism and Hepatocellular Carcinoma Complicated by Direct Acting Antiviral Treatment of Hepatitis C Virus

Pages 3203-3210

Samar Ebrahim Ghanm; Nashwa Abuel-Fetuh Shebl; Ibrahim El Tantawy El Sayed; Hamed Mohamed Abdel- Bary; Basant Farid Saad; Warda Othman Saad

[View Article](#)  PDF 401.72 K

20. Laser Therapy as a Preventive Approach for Oral Mucositis in Cancer Patients Undergoing Chemotherapy: The Potential Role of Superoxide Dismutase

Pages 3211-3217

Beatriz Coutens de Menezes; Marcela Marçal Thebit; Lucas André Silva Bonela; Karine Gadioli de Oliveira; Washington Luiz Gonçalves; Nazare Souza Bissoli; Carmem Luiza Sartorio; Sonia Alves Gouvea

[View Article](#)  PDF 344.62 K

21. p16INK4A Expression in Condyloma Acuminata Lesions Associated with High-Risk Human Papillomavirus Infection

Pages 3219-3225

Gondo Mastutik; Alphania Rahniayu; Afria Arista; Dwi Murtiastutik; Nila Kurniasari; Trisniartami Setyaningrum; Anny Setijo Rahaju; Erna Sulistyani

[View Article](#)  PDF 452.75 K

22. Lactate Dehydrogenase Levels in the Saliva of Cigarette and E-Cigarette Smokers (Vapers): A Comparative Analysis

Pages 3227-3235

Anitha Krishnan Pandarathodiyil; Anand Ramanathan; Ranjana Garg; Jennifer Geraldine Doss; Fazliny Binti Abd Rahman; Wan Maria Nabillah Ghani; Saman Warnakulasuriya

[View Article](#)  PDF 390.06 K

23. CD34+/CD38- Stem Cell Burden Could Predict Chronic Myeloid Leukemia Patients' Outcome

Pages 3237-3243

Noura Fathy El-Metwaly; Salah Aref; Mohamed Ayed; Manal Abdel Hamid; Ahmed M.A El-Sokkary

View Article  PDF 376.96 K

24. Frequency of FLT3 Internal Tandem Duplications in Adult Syrian Patients with Acute Myeloid Leukemia and Normal Karyotype

Pages 3245-3251

Ismael F Al-Arbeed; Abdulsamad Wafa; Faten Moassass; Bassel AL-Halabi; Walid Al-Achkar; Imad Abou-Khamis

View Article  PDF 356.88 K

25. Patient Satisfaction with Private Genetic Counselling for Familial Cancer in Western Australia: A Prospective Audit

Pages 3253-3259

Charmi N Perera; Sarah O'Sullivan; Nicholas Pachter; Jason Jit-Sun Tan; Paul A Cohen

View Article  PDF 300.85 K

26. Evaluation of PathTezt™, a Liquid-Based Cytology System

Pages 3261-3266

Nur Syuhada Mohd Nafis; Anani Aila Mat Zin

View Article  PDF 538.11 K

27. Assessment of the Anti-Cancer Efficiency of Silver Moringa oleifera Leaves Nano-extract against Colon Cancer Induced Chemically in Rats

Pages 3267-3286

Wael Mahmoud Aboulthana; Wafaa Ghoneim Shousha; Ehab Abdel-Raouf Essawy; Mahmoud Hosny Saleh; Alaa Hamed Salama

View Article  PDF 1.94 MB

28. Non-Invasive Saliva-based Detection of Gene Mutations in Oral Cancer Patients by Oral Rub and Rinse Technique

Pages 3287-3291

Audrey D'Cruz; Pandyanda Nanjappa Dechamma; Marina Saldanha; Sahana Maben; Pushparaja Shetty; Anirban Chakraborty

View Article  PDF 310.89 K

p16^{INK4A} Expression in Condyloma Acuminata Lesions Associated with High-Risk Human Papillomavirus Infection

Gondo Mastutik^{1*}, Alphania Rahniayu^{1,2}, Afria Arista², Dwi Murtiastutik^{1,2}, Nila Kurniasari^{1,2}, Trisniartami Setyaningrum^{1,2}, Anny Setijo Rahaju^{1,2}, Erna Sulistyani³

Abstract

Objective: The objective of this study was to discover the possible correlation between p16^{INK4A} expression and the LR/HR-HPV infection in condyloma acuminata (CA) lesions. **Materials and Method:** This cross-sectional study was conducted during January-December 2017 on 33 CA patients. The expression of p16^{INK4A} was detected by immunohistochemistry (IHC) staining. The positive interpretation was carried out by scoring which score 0 was negative, score 1 was sporadic, score 2 was focal, and score 3 was diffuses. The HPV genotypes were identified by reverse line blot, and 40 genotypes of HPV detected, including HR-HPV (HPVs 16, 18, 26, 31, 33,35, 39, 45, 51, 52, 53, 56, 58, 59, 66, 67, 68a, 68b, 69, 73, and 82) and LR-HPV (HPVs 6, 11, 40, 42, 43, 44, 54, 55, 61, 62, 64, 70, 71, 72, 81, 83, 84, 87, 89, and 90). **Results:** The expression of p16^{INK4A} was significantly correlated with HR-HPV infection. Patients infected with HR-HPV had 0.644 times higher possibility to express p16^{INK4A} gene compared to those infected with LR-HPV. LR-HPV genotypes detected in CA patients were HPVs 6, 11, 42, 61, 54, 81, 87, 89, and 90 and HR-HPV genotypes were HPVs 18, 26, 45, 51, 52, 67, 68B, 69, and 82. LR-HPV was found in 19/33 of patients and HR-HPV was in 14/33 of patients. The expression of p16^{INK4A} in CA lesions was diffuse in 15.2% of patients, was focal in 24.2% of patients, was sporadic in 39.4% of patients, and was negative in 21.2% of patients. In LR-HPV group, there was no diffuse expression, focal expression was observed in 15.8%, sporadic in 47.4%, and negative in 36.8%, while in HR-HPV group, p16^{INK4A} expression was detected in all lesions, in a way that its expression was diffuse in 35.7%, focal in 35.7%, and sporadic in 28.6%. **Conclusion:** IHC is a routine method in histopathological diagnosis, therefore the detection of p16^{INK4A} expression by IHC can be used as a biomarker for HR-HPV infection diagnosis.

Keywords: p16^{INK4A}- high-risk HPV- low-risk HPV- condyloma- sexually transmitted diseases

Asian Pac J Cancer Prev, 22 (10), 3219-3225

Introduction

Condyloma acuminata (CA) is the most frequent sexually transmitted disease worldwide (Santegoets et al., 2012). It is also known as genital warts or anogenital warts. These lesions are usually single or multiple appearing in the anogenital region and causing symptoms of itching, vaginal discharge, and bleeding. CA can be flat or lobulated and appears as pearl-like, filiform, plaque eruption, or cauliflower projection (Patel et al., 2013; Léonard et al., 2014). Its prevalence is increasing worldwide. The prevalence of CA in Italian female population during 2009-2010 was 3.8 cases per 1,000 women per year (Suligo et al., 2017), in the UK population during 2010-2012 was 3.8% in males and 4.6% in females (Sonnenberg et al., 2019), and in the US population during 2013-2014 was 2.9% (Daugherty et al.,

2018). A systematic review on the incident of anogenital warts showed that their prevalence ranged from 160 to 289 per 100,000 persons. New incident of anogenital warts ranged from 103 to 168 per 100,000 persons among males and 76 to 191 per 100,000 persons among females (Patel et al., 2013).

The most common cause of CA is Human Papillomavirus (HPV). There are two groups of HPV, including high risk (HR) HPV and low risk (LR) HPV. HR-HPV are HPV genotypes of 16, 18, 26, 31, 33, 35, 39, 45, 51, 52, 53, 56, 58, 59, 61, 73, and 82, and LR-HPV are HPV genotypes of 6, 11, 40, 42, 43, 44, 54, 61, 70, 72, and 81 (Braaten et al., 2008; Gutiérrez-Xicoténcatl et al., 2009). LR-HPV causes CA, but a study in China showed that the infection of HR-HPV was also found in CA lesions. In some cases, CA is caused by the combination of LR and HR-HPV infections (Lu et al., 2014). Persistent

¹Department of Anatomic Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. ²Department of Dermatology and Venereology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. ³Department of Oral Medicine, Faculty Dentistry, Jember University, Jember, Indonesia *For Correspondence: gondomastutik@fk.unair.ac.id

infection of LR or HR-HPV is a risk factor for the transformation of epithelial to benign hyperplasia or premalignant lesions. HR-HPV is related to the occurrence of malignancy in women, as cervical cancer (Santegoets et al., 2012). Some literature considers HR-HPV as the main causative agent responsible for the cervical cancer (Braaten et al., 2008). The detection of HPV genotype is very important to prevent, establish early diagnosis, and initiate the treatment in cervical cancer. Determination of LR or HR-HPV in anogenital warts can be used as a factor to predict the progression of lesions to benign or malignant lesions.

The E7 HPV protein plays a role in the cell transformation process. It binds to important proteins, such as the pRB and cyclin A/CDK2 complex, inhibiting the interaction between Rb and E2F. The E7 protein of HR-HPV deactivates pRB, resulting in the accumulation of p16^{INK4A} protein. The expression of p16^{INK4A} can also be considered as a marker of E7 gene activity (Izadi-Mood et al., 2012; Romagosa et al., 2011). p16^{INK4A} plays a role in cell cycle regulation and it is involved in the processes of apoptosis, angiogenesis, cell invasion. This activity may be associated with overexpression in cancer (Romagosa et al., 2011). The expression of p16^{INK4A} is a marker to determine the prognosis of a malignancy caused by HPV infection (Missaoui et al., 2010), suggesting that p16^{INK4A} can be a specific marker for HPV infection and may correlate with the type of HR-HPV or LR-HPV. The objective of this study was to analyze the correlation between p16^{INK4A} expression and the LR-HPV or HR-HPV in CA lesions. This study identified the genotype of HPV and performed immunohistochemical analysis of p16^{INK4A} staining of.

Materials and Methods

The samples collection

This cross-sectional study was conducted at Outpatient Clinic of Department Dermatology and Venereology, Dr. Soetomo General Academic Hospital, Surabaya, Indonesia, from January 2017 to December 2017. The study was ethically approved by the Medical Ethic Research at Dr. Soetomo General Academic Hospital, Surabaya (ethical code: 382/Panke.KKE/V/2016).

All patients with CA, both male and female, who were willing to participate in the study, were included. Informed consent was obtained from the participants. Menstruating and pregnant women, those who suffered from an active pelvis and / or acute cervicitis, men or women with a diagnosis of HIV and AIDS, and those who were not willing to participate in the study were excluded.

The specimens were 33 tissues of CA. Tissue from each patient was divided into 2 parts, one part for tissue processing into paraffin block preparations followed by histopathological diagnosis and immunohistochemistry (IHC) staining, while another for the examination of the HPV genotype. Histopathological diagnosis of CA and analysis of IHC staining was performed by a pathologist.

Expression of p16^{INK4A}

The expression of p16^{INK4A} was detected by

IHC staining using Anti-CDKN2A/ p16^{INK4A} Antibody (clone 1E12E10) IHC-plus™ LS-B5261 (LS Bio). The interpretation was positive if cells were stained in the nucleus or combined in the nucleus and cytoplasm. The assessment was carried out based on guidelines presented by Klaes et al. (2001) as follows:

Score 0 (negative): if cells were stained positive <1% of all cells

Score 1 (sporadic): if cells were stained positive <5% of all cells

Score 2 (focal): if cells were stained positive <25% of all cells

Score 3 (diffuses): if cells were stained positive > 25% of all cells.

Genotyping of HPV

Virus extraction was carried out from CA tissues using the QIAamp DNA Mini Kit (Qiagen) kit and according to the manufacturer's protocol. Genotyping of HPV was performed by Polymerase Chain Reaction (PCR), then followed by reverse line blot using the Ampliquality HPV type express v 3.0 kit (Ab Analitica). This kit could detect 40 genotypes of HPV, including HR-HPV were HPV 16, 18, 26, 31, 33,35, 39, 45, 51, 52, 53, 56, 58, 59, 66, 67, 68 (68a, 68b), 69, 73, and 82 and LR-HPV were HPV 6, 11, 40, 42, 43, 44, 54, 55, 61, 62, 64, 70, 71, 72, 81, 83, 84, 87, 89, and 90.

Statistical analysis

The difference in the expression of p16^{INK4A} on CA patients with HR-HPV or LR-HPV groups infection was analyzed by Mann Whitney test (significant if $p < 0.05$). The correlation between variables was analyzed by two-tailed Spearman's rho (significant if $p < 0.05$).

Results

Characteristics of patient

This study was performed on 33 patients suffering from CA. Patients' characteristic, such as sex, age, sexual partner, duration of the symptoms, medical history, presence of lesion in the partner, history of treatment, and shape, type, and location of the lesion were investigated. The patients consisted of 12 (36.4%) males and 21 (63.6%) females. The patients aged from 18 to 64 years. The highest frequency was allocated to the age group of 15-24 years old (51.5%). Regarding sexual partner, it was found that 30/33 of the patients (90.9%) were heterosexual (men having sex with women), 2/33 (6.1%) were bisexual (having sex with the same and different sex), and 1/33 (3.0%) were homosexual (having sex with the same sex). Regarding symptoms duration, the highest frequency was allocated to the duration of 1 to 3 months. Mostly, the patients experienced CA for the first time, there was no lesion in their partner, and received TCA therapy, respectively. Considering the shape, type, and location of the lesion, it was revealed that the highest frequencies were allocated to papule shape, multiple type of lesions, and gland penis for male and labia majora for female (Table 1).

HPV genotype in CA lesions




Figure 1. The Histopathological Feature and p16^{INK4A} Expression of Condyloma Acuminata. A, Condyloma acuminata from HR-HPV infected patients, in HE staining (40 x magnification); B, In the same patients, p16^{INK4A} expression showed in diffuse (score 3) (40 x magnification); C, Condyloma acuminata from LR-HPV infected patients, in HE staining (40 x magnification); D, In the same patients, p16^{INK4A} expression showed negative (score 0) (40 x magnification); E, p16^{INK4A} expression in negative (score 0) (400 x magnification); F, p16^{INK4A} expression in diffuse (score 3) (400 x magnification).

The genotype of HPV included the infection of HPV-HR and LR, both single and multiple infections. The single or multiple infections of HPV-LR was assumed as infection of LR-HPV. The multiple infection of LR-HPV with HR-HPV was assumed as infection of HR-HPV. The genotype of LR-HPV were HPV 6, 11, 42, 61, 54, 81, 87, 89, and 90, and of HR-HPV were 18, 26, 45, 51, 52, 67, 68B, 69, and 82. LR-HPV was found in 19/33 (57.6%) of the patients and HR-HPV was seen in 14/33 (42.4%) of the patients (Table 2). The most dominant HPV was HPV 11 that infected 24/56 (42.9%) times more, then followed by HPV 6 (16.1%) and HPV 18, HPV 51, and HPV 82 for 5.4%, respectively.

p16^{INK4A} expression in CA lesions

The expression of p16^{INK4A} in CA lesions showed that nucleus or the combination of nucleus and cytoplasm of cell was stained in brown color, indicating in sporadic, focal, or diffuse (Figure 1). The results showed that 7/33 (21.2%) of the lesions were negative, 13/33 (39.4%) were sporadic, 8/33 (24.2%) were focal, and 5/33 (15.2%)

were diffuse. In LR-HPV group, the results showed that negative, sporadic, and focal lesions were 7/19 (36.8%), 9/19 (47.4%), and 3/19 (15.8%), respectively. In HR-HPV group, it was found that that the frequency of sporadic lesions was 4/14 (28.6%), focal was 5/14 (35.7%), and diffuse was 5/14 (35.7%) (Table 3). There was a significant different between LR-HPV and HR-HPV groups in terms of p16^{INK4A} expression in CA lesion ($p = 0,000$). Correlation between p16^{INK4A} expression and LR-HPV and HR-HPV infections was moderate ($r = 0,644$, $p = 0,000$).

Discussion

The microscopic features of CA lesions based on haematoxylin eosin (HE) staining are parakeratosis, hyperkeratosis, hypergranulosis, basal cell hyperplasia, and koilocytic (Léonard et al., 2014). In this study, CA lesions were seen as papules, cauli flowers, and flat-tapping papules. The microscopic features found in this study were hyperkeratosis, parakeratosis, papillomatosis, hypergranulosis, and hyperplasia of basal

Table 1. Characteristics of Patients with Condyloma Acuminata

| Characteristics of Patients | N (%) | Single or multiple infection LR/LR HPV N (%) | Multiple infection of LR/HR HPV N (%) |
|------------------------------|-----------|---|--|
| Sex | | | |
| Male | 12 (36.4) | 7 (21.2) | 5 (15.1) |
| Female | 21 (63.6) | 12 (36.4) | 9 (27.3) |
| Age | | | |
| 15-24 years | 17 (51.5) | 10 (30.3) | 7 (21.2) |
| 25-34 years | 6 (18.2) | 3 (9.1) | 3 (9.1) |
| 35-44 years | 7 (21.2) | 4 (12.1) | 3 (9.1) |
| 45-54 years | 2 (6.1) | 1 (3.0) | 1 (3.0) |
| 55-64 years | 1 (3.0) | 1 (3.0) | 0 |
| Sexual Partner | | | |
| Heterosexuals | 30 (90.9) | 18 (54.5) | 12 (36.4) |
| Homosexual | 1 (3.0) | 1 (3.0) | 0 |
| Bisexual | 2 (6.1) | 0 | 2 (6.1) |
| Duration of symptom | | | |
| < 1 month | 5 (15.1) | 1 (3.0) | 4 (12.1) |
| 1- 3 months | 20 (61.6) | 14 (42.4) | 6 (18.2) |
| 4 - 6 months | 6 (18.2) | 3 (9.1) | 3 (9.1) |
| > 6 months | 2 (6.1) | 1 (3.0) | 1 (3.0) |
| History | | | |
| First lesion | 26 (78.8) | 15 (45.4) | 11 (33.3) |
| Recurrent lesion | 7 (21.2) | 4 (12.1) | 3 (9.1) |
| Lesion on the partner | | | |
| Have lesion on the partner | 4 (12.1) | 2 (6.1) | 2 (6.1) |
| No lesions on the partner | 29 (87.9) | 17 (51.5) | 12 (36.4) |
| Therapy | | | |
| TCA | 19 (57.6) | 11 (33.3) | 8 (24.2) |
| Cautery | 14 (42.4) | 8 (24.2) | 6 (18.2) |
| Shape | | | |
| Papule | 28 (84.8) | 16 (48.5) | 12 (36.4) |
| Cauli flower | 4 (12.1) | 2 (6.1) | 2 (6.1) |
| Flat tapped papule | 1 (3.0) | 1 (3.0) | 0 |
| Type | | | |
| Multiple | 31 (93.9) | 18 (54.5) | 13 (39.4) |
| Solitary | 2 (6.1) | 1 (3.0) | 1 (3.0) |
| Location | | | |
| Penile | 7 (21.2) | 4 (12.1) | 3 (9.1) |
| Anus | 4 (12.1) | 2 (6.1) | 2 (6.1) |
| Penile and anus | 1 (3.0) | 1 (3.0) | 0 |
| Labia (majora and minora) | 16 (48.5) | 10 | 6 (18.2) |
| Vulva | 3 (9.1) | 1 (3.0) | 2 (6.1) |
| Introitus vagina | 1 (3.0) | 0 | 1 (3.0) |
| Perineum | 1 (3.0) | 1 (3.0) | 0 |

cell, as well as the koilocytes that usually is associated with HPV infection.

The most common cause of CA is infection by HPV. There are more than 40 genotypes of HPV that can infect the anogenital area that are usually infected by LR-HPV including HPV 6 and HPV 11 in single infection, but most commonly co-infection with LR-HPV or HR-HPV

(Léonard et al., 2014; Hasanzadeh et al., 2019). The results of this study showed that all the tissues taken from the patients, who were clinically diagnosed with CA, were positive for LR-HPV or HR-HPV infections, in single infection or multiple infection. Most of the patients were infected by LR-HPV, while 42.4% were infected by LR-HPV and co-infected with HR-HPV. In line with this

Table 2. Distribution of HPV in Condyloma Acuminata Lesions

| Genotype of HPV | HR or LR HPV genotypes | Frequency N (%) |
|-------------------|------------------------|-----------------|
| LR HPV | | |
| HPV 6 | LR | 4 |
| HPV 11 | LR | 12 |
| HPV 6, 11 | LR | 2 |
| HPV 6,81,87,89 | LR | 1 |
| HR HPV | | |
| HPV 6,11,18,51,82 | HR | 1 |
| HPV 6,42,51,61 | HR | 1 |
| HPV 11, 18 | HR | 2 |
| HPV 11,18,45 | HR | 1 |
| HPV 11, 26 | HR | 1 |
| HPV 11,67 | HR | 1 |
| HPV 11, 51, 82 | HR | 1 |
| HPV 11, 52, 54 | HR | 1 |
| HPV 11, 52,69,90 | HR | 1 |
| HPV 11, 68B | HR | 1 |
| HPV 11, 82 | HR | 2 |
| Total | | 33 |

study, a cross-sectional study in Kuwait on 156 patients with genital warts showed that 102/156 (65.4%) of the patients were infected by LR-HPV and 54/156 (34.6%) of the patients were infected by HR-HPV. About 88.4% of the patients in the aforementioned study had single infection and 11.6% had multiple infections (Al-Awadhi et al., 2019). Another study in Spain showed that the frequency of LR-HPV infection was 63/138 (45.6%) in their patients and 71/138 (41.4%) of anogenital warts patients were infected by HR-HPV (Arroyo et al., 2016). The other study on 66 anogenital warts specimens showed that LR-HPV infected 42/66 (62.1%) which was dominated by HPV 6 (47%), and HPV-11 (13.6%), as well as HPV 18 and HPV 3 (Ozaydin-Yavuz et al., 2019). We discovered that that beside LR-HPV, CA was also co-infected with HR-HPV that could develop to malignant cancer. Therefore, identification of HPV genotype can predict the risk of developing related diseases. In addition, determination of HPV genotype affects treatment management, patients' follow-up, and prevention strategies

The findings of this study revealed that 57.6% of CA patients were infected by LR-HPV and the rest were

infected by both LR-HPV and HR-HPV infections. The most common HPV genotypes were HPV 11 and HPV 6 for LR-HPV, followed by HPV 18, HPV 51, and HPV 82 for HR-HPV. The incidence of CA and the progression of the disease can be prevented by vaccination. Recently, there are 3 commercial vaccines against HPV infection, namely Gardasil to prevent infections caused by HPV 6, 11, 16, and 18, Cervarix to prevent infections caused by HPV 16, and 18, and Gardasil 9 to prevent infections caused by HPV 6, 11, 61, 18, 31, 33, 45, 52, and 58 (Gupta et al., 2017). A previous study on anogenital benign lesions showed that HPV genotype in CA patients was dominated by HPV 11 and HPV 6 among 13 female patients (Arista et al., 2019) and 12 male patients (Murtiastutik et al., 2019). In pre-cancerous lesion and cancerous lesion of uterine cervix, those were dominated by HPV 16 (62.68%), then followed by HPV 18 (20.9%), HPV 45 (5.97%), HPV 52 (5.97%), and HPV 67 (4.48%) (Mastutik et al., 2018). Vaccination programs is expected to reduce the incidence of these diseases, but there are some genotypes of HPV that cannot be targeted by the current vaccines. Therefore, strategies to prevent the incidence of CA or the progression of malignancy still need to be developed.

Oncoprotein E6 and E7 of HR-HPV play a role in cancer development. HR-HPV E6 mediates p53 inactivation by binding to the conserved domains of E6AP (E6-linked protein) to form the E6/ E6AP/p53 complex. This complex causes degradation of p53 by ubiquitination mechanism. HR-HPV E7 targets to degrade the retinoblastoma protein (pRB). In normal cells, when cells are prevented from entering the S phase, the pRB binds to the E2F family of transcription factor, so that the cell stops at the checkpoint of G1-S phase, activating cell cycles arrest. pRb, which is phosphorylated by cyclin D1/CDK 4/6 complex, causes E2F to be released and enter the nucleus. The cell enters the S phase which then starts the activation of gene transcription (Munger et al., 2013, Pal and Kudu, 2020). Furthermore, phosphorylated pRB is a p16^{INK4A} feedback mechanism. HPV E7 induces degradation of pRB by an ubiquitin proteasome pathway that causes the loss feedback mechanism of p16^{INK4A} and then leads to accumulation of p16^{INK4A}, presenting as overexpression of p16^{INK4A} (Lassen et al., 2009; Faraji et al., 2017).

The overexpression of p16^{INK4A} in this study was sporadic (39.4%), focal (24.2%), diffuse (15.2%), and negative (21.2%). Another study on p16^{INK4A} expression in 24 CA specimens showed that overexpression of p16^{INK4A} were sporadic in 11/24 of the specimens (45.8%),

Table 3. The p16^{INK4A} Expression on Condyloma Acuminata that Infected by LR or HR-HPV

| p16 ^{INK4A} expression | HPV genotype | | Total N (%) | p-value |
|---------------------------------|-----------------|-----------------|-----------------|---------|
| | LR-HPV N (%) | LR/HR-HPV N (%) | | |
| Score 0 (Negative) | 7 (36.8) | 0 (0) | 7 (21.2) | 0.000* |
| Score 1 (Sporadic) | 9 (47.4) | 4 (28.6) | 13 (39.4) | |
| Score 2 (Focal) | 3 (15.8) | 5 (35.7) | 8 (24.2) | |
| Score 3 (Diffuse) | 0 (0) | 5 (35.7) | 5 (15.2) | |
| Total | 19 (100) | 14 (100) | 33 (100) | |

*Mann Whitney, Asymptotic Significance (2-tailed) with p = 0,000 (p < 0,05); Correlation is significant at the 0.01 level (2-tailed) with Spearman's rho (r) = 0,644, p = 0,000

were focal in 7/24 of the specimens (29.2%), and were negative in 6/24 of the specimens (25%) (Kazlouskaya et al., 2013). The expression of p16^{INK4A} in cervix tended to increase from cervical normal epithelium to invasive cervical cancer (Missaoui et al., 2010; Izadi-Mood et al., 2012) and cervical adenocarcinoma (Mastutik et al., 2021). The expression of p16^{INK4A} in HR-HPV showed that all specimens were positive ranging from sporadic to diffuse. In LR-HPV were sporadic and focal, whereas 36.8% of specimens were not expressed in p16^{INK4A} and none of specimens expressed in diffuse category. Other study on anal lesion with HR HPV infection showed that all specimens expressed p16^{INK4A} ranging from sporadic to diffuse, whereas those infected with LR-HPV were mostly patchy (Leeman et al., 2019). In oropharyngeal squamous cell carcinomas and tonsillar dysplasia that were positive for HPV 16, p16^{INK4A} expression was also diffuse. However, in benign and pre-malignant lesions were positive for HPV 6 and HPV 11 showed variations from negative to strong positive (Mooren et al., 2014). All cervical lesions infected with HR-HPV had a significant, strong, and diffuse p16^{INK4A} expression, whereas those infected with LR-HPV showed mild p16^{INK4A} expression (score 1) (Missaoui et al., 2014).

These findings highlighted that the expression of p16^{INK4A} significantly correlated with HR-HPV infection, in a way that CA lesions infected by HR-HPV had 0.644 times more chances to express p16^{INK4A} compared to CA lesions infected by LR-HPV. As previous studies showed that p16^{INK4A} expression was associated with HR-HPV infection in oropharyngeal squamous (Liu et al., 2015), mucosal squamous cell carcinomas of the head and neck (Antonsson et al., 2015), cervical squamous intraepithelial lesion (Yildiz et al., 2007), and invasive cervical and vaginal carcinomas (Missaoui et al., 2010; Missaoui et al., 2014). HR-HPV is integrated into the host cell genome, whereas LR-HPV prefers extra chromosome as episome, so that the expression of E6 and E7 oncoproteins are within the regulatory framework of E1 and E2 HPV (Boulet et al., 2007). HR-HPVs, such as HPV 18, 33, HPV 51, HPV 58, and HPV 59, were found to be integrated, while HPVs 30, 35, 39, 44, 45, 53, 56, 59, 74 and 82 did not tend to be integrated, but to be in episome form (Nkili-Meyong et al., 2019). In this study, it was found that HR-HPVs 18, 51, and 82. As reported by Nkili-Meyong et al., (2019), HPV 18 was founded to be integrated in 55% of the positive HPV 18 specimen, HPV 51 was integrated in 25% of the positive HPV 51 specimen, and HPV 82 was founded in extra chromosome. This integration was associated with partial or total deletion of E1 and E2 genes, leading to the overexpression of E6 and E7 due to loss of feedback mechanism by E2 protein (Woodman et al., 2007; Nkili-Meyong et al., 2019). In addition, the HR-HPV E7 oncoproteins have higher affinity to bind pRB than LR-HPV E7, increasing the accumulation of p16^{INK4A}. Therefore, all specimens infected by HR-HPV in this study were positive for the expression of p16^{INK4A} in sporadic, focal, and diffuse forms, while those infected by LR-HPV had sporadic and focal, but not diffuse expression.

In conclusion, this study found significant differences between CA specimens infected by LR-HPV (HPV 6, 11,

42, 61, 54, 81, 87, 89, 90) and HR-HPV (HPV 18, 26, 45, 51, 52, 67, 68B, 69, 82) regarding the expression of p16^{INK4A}. p16^{INK4A} expression was correlated with HR-HPV infection moderately. IHC is a routine method to perform the diagnostic of histopathological. Therefore, IHC of p16^{INK4A} could be used as biomarker for HR-HPV infection to predict the malignancy development of CA lesions.

Author Contribution Statement

Concepts, design, definition of intellectual content: Gondo Mastutik. Literature search: Gondo Mastutik, Afria Arista. Clinical studies: Gondo Mastutik, Dwi Murtiastutik, Trisniartami Setyaningrum. Experimental studies: Gondo Mastutik, Alphania Rahniayu, Afria Arista. Data acquisition: Nila Kurniasari, Alphania Rahniayu, Anny Setijo Rahaju. Data analysis: Gondo Mastutik, Alphania Rahniayu, Anny Setijo Rahaju. Statistical analysis: Gondo Mastutik, Nila Kurniasari, Anny Setijo Rahaju. Manuscript preparation and manuscript editing: Gondo Mastutik, Afria Arista, Anny Setijo Rahaju. Manuscript review: Gondo Mastutik, Alphania Rahniayu, Dwi Murtiastutik, Trisniartami Setyaningrum, Erna Sulistyani.

Acknowledgement

We are thankful of Government of the Republic of Indonesia, the Ministry of Research, Technology and Higher Education, and the Faculty of Medicine, Airlangga University for funding in this research with the university's operational funding for the 2017 fiscal year.

Funding statement

This study was supported by the Faculty of Medicine, Airlangga University, and the university's operational funding for the 2017 fiscal year. This manuscript have not been presented at a scientific event or submitted to any other journals.

Ethical statement

Ethical study was approved by the Medical Ethic Research at Dr. Soetomo General Academic Hospital Surabaya, Indonesia (ethical code:382/Panke. KKE/V/2016).

Statement conflict of interest

The authors declare no conflict of interest

References

- Al-Awadhi R, Al-Mutairi N, Albatineh AN, Chehadeh W (2019). Association of HPV genotypes with external anogenital warts: a cross sectional study. *BMC Infect Dis*, **19**, 375-82.
- Antonsson A, Neale RE, Boros S, et al (2015). Human papillomavirus status and p16(INK4A) expression in patients with mucosal squamous cell carcinoma of the head and neck in Queensland, Australia. *Cancer Epidemiol*, **39**, 174-81.
- Arista A, Murtiastutik D, Setyaningrum T, Mastutik G (2019). Genotyping of Human Papilloma Virus (HPV) in female condyloma acuminata patient in Dr. Soetomo Hospital, Surabaya. *Dermatol Rep*, **11**, 122-3.

- Arroyo LS, Basaras M, Arrese E, et al (2016). Distribution of genital human papillomavirus genotypes in benign clinical manifestations among men from Northern Spain. *BMC Public Health*, **16**, 81-8.
- Boulet G, Horvath C, Vanden Broeck D, Sahebali S, Bogers J (2007). Human papillomavirus: E6 and E7 oncogenes. *Int J Biochem Cell Biol*, **39**, 2006-11.
- Braaten KP, Laufer MR (2008). Human Papillomavirus (HPV), HPV-Related Disease, and the HPV Vaccine. *Rev Obstet Gynecol*, **1**, 2-10.
- Daugherty M, Byler T (2018). Genital Wart and Human Papillomavirus Prevalence in Men in the United States From Penile Swabs: Results From National Health and Nutrition Examination Surveys. *Sex Transm Dis*, **45**, 412-6.
- Faraji F, Zaidi M, Fakhry C, Gaykalova DA (2017). Gaykalova DA. Molecular mechanisms of human papillomavirus-related carcinogenesis in head and neck cancer. *Microbes Infect*, **19**, 464-75.
- Gupta G, Glueck R, Patel PR (2017). HPV vaccines: Global perspectives. *Hum Vaccin Immunother*, **13**, 1421-24.
- Gutiérrez-Xicoténcatl L, Plett-Torres T, Madrid-González CL, Madrid-Marina V (2009). Molecular diagnosis of human papillomavirus in the development of cervical cancer. *Salud Publica Mex*, **51**, S479-88.
- Hasanzadeh M, Rejali M, Mehramiz M, et al (2019). The interaction of high and low-risk human papillomavirus genotypes increases the risk of developing genital warts: A population-based cohort study. *J Cell Biochem*, **120**, 12870-4.
- Izadi-Mood N, Asadi K, Shojaei H, et al (2012). Potential diagnostic value of P16 expression in premalignant and malignant cervical lesions. *J Res Med Sci*, **17**, 428-33.
- Kazlouskaya V, Shustef E, Allam SH, Lal K, Elston D (2013). Expression of p16 protein in lesional and perilesional condyloma acuminata and bowenoid papulosis: clinical significance and diagnostic implications. *J Am Acad Dermatol*, **69**, 444-9.
- Klaes R, Friedrich T, Spitkovsky D, et al (2001). Overexpression of p16(INK4A) as a specific marker for dysplastic and neoplastic epithelial cells of the cervix uteri. *Int J Cancer*, **92**, 276-84.
- Lassen P, Eriksen JG, Hamilton-Dutoit S, et al (2009). Effect of HPV-associated p16^{INK4A} expression on response to radiotherapy and survival in squamous cell carcinoma of the head and neck. *J Clin Oncol*, **27**, 1992-8.
- Leeman A, Jenkins D, Marra E, et al (2019). Grading immunohistochemical markers p16^{INK4a} and HPV E4 identifies productive and transforming lesions caused by low- and high-risk HPV within high-grade anal squamous intraepithelial lesions. *Br J Dermatol*, **182**, 1026-33.
- Léonard B, Kridelka F, Delbecq K, et al (2014). A clinical and pathological overview of vulvar condyloma acuminatum, intraepithelial neoplasia, and squamous cell carcinoma. *Biomed Res Int*, **2014**, 1-11.
- Liu SZ, Zandberg DP, Schumaker LM, Papadimitriou JC, Cullen KJ (2015). Correlation of p16 expression and HPV type with survival in oropharyngeal squamous cell cancer. *Oral Oncol*, **51**, 862-9.
- Lu LH, Yu J, Zhang YH, Wu XF, Liu XY (2014). HPV Infection and Distribution of HPV Genotypes on Patients with Condyloma Acuminatum. *J Reproduct Contraception*, **25**, 26-31.
- Mastutik G, Alia R, Rahniayu A, et al (2018). Genotyping of human papillomavirus in cervical precancerous lesion and squamous cell carcinoma at Dr. Soetomo Hospital, Surabaya, Indonesia. *Afr J Infect Dis*, **12**, 7-12.
- Mastutik G, Rahniayu A, Kurniasari N, Rahaju AS, Alia R (2021). Correlation between the expression of E6 HPV with P53 and P16^{INK4A} at cervical adenocarcinoma and cervical normal. *Indian J Forensic Med Toxicol*, **15**, 1827-34.
- Missaoui N, Trabelsi A, Hmissa S, et al (2010). p16^{INK4A} overexpression in precancerous and cancerous lesions of the uterine cervix in Tunisian women. *Pathol Res Pract*, **206**, 550-5.
- Missaoui N, Abdelkarim SB, Mokni M, Hmissa S (2014). p16^{INK4A} expression in squamous cell carcinomas of the vagina and the vulva in Tunisian women. *Asian Pac J Cancer Prev*, **15**, 10803-8.
- Mooren JJ, Gültekin SE, Straetmans JM, et al (2013). P16(INK4A) immunostaining is a strong indicator for high-risk-HPV-associated oropharyngeal carcinomas and dysplasias, but is unreliable to predict low-risk-HPV-infection in head and neck papillomas and laryngeal dysplasias. *Int J Cancer*, **134**, 2108-17.
- Munger K, Gwin TK, McLaughlin-Drubin ME (2020). p16 in HPV-associated cancers. *Oncotarget*, **4**, 1864-5.
- Murtiastutik D, Mastutik G, Rahniayu A, Arista A, Setyaningrum T (2019). The Genotype of Human Papilloma Virus of Male Patients with Anogenital Warts. *Folia Med Indonesiana*, **55**, 100-6.
- Nkili-Meyong AA, Moussavou-Boundzanga P, Labouba I, et al (2019). Genome-wide profiling of human papillomavirus DNA integration in liquid-based cytology specimens from a Gabonese female population using HPV capture technology. *Sci Rep*, **9**, 1504-14.
- Ozaydin-Yavuz G, Bilgili SG, Guducuoglu H, et al (2019). Determinants of high-risk human papillomavirus infection in anogenital warts. *Postepy Dermatol Alergol*, **36**, 76-81.
- Pal A, Kundu R (2020). Human Papillomavirus E6 and E7: The Cervical Cancer Hallmarks and Targets for Therapy. *Front Microbiol*, **10**, 1-15.
- Patel H, Wagner M, Singhal P, Kothari S (2013). Systematic review of the incidence and prevalence of genital warts. *BMC Infect Dis*, **13**, 39-52.
- Romagosa C, Simonetti S, López-Vicente L, et al (2011). p16(INK4a) overexpression in cancer: a tumor suppressor gene associated with senescence and high-grade tumors. *Oncogene*, **30**, 2087-97.
- Santegoets LA, van Baars R, Terlou A, et al (2012). Different DNA damage and cell cycle checkpoint control in low- and high-risk human papillomavirus infections of the vulva. *Int J Cancer*, **130**, 2874-85.
- Sonnenberg P, Tanton C, Meshier D, et al (2019). Epidemiology of genital warts in the British population: implications for HPV vaccination programmes. *Sex Transm Infect*, **95**, 386-90.
- Suligoï B, Vittori G, Salfa MC, et al (2017). Prevalence and incidence of external genital warts in a sample of Italian general female population. *BMC Infect Dis*, **17**, 126-33.
- Woodman CB, Collins SI, Young LS (2007). The natural history of cervical HPV infection: unresolved issues. *Nat Rev Cancer*, **7**, 11-22.
- Yildiz IZ, Usubütün A, Firat P, Ayhan A, Küçükali T (2007). Efficiency of immunohistochemical p16 expression and HPV typing in cervical squamous intraepithelial lesion grading and review of the p16 literature. *Pathol Res Pract*, **203**, 445-9.



This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.

SCOPUS Q2

<https://www.scopus.com/sourceid/40173>

Source details

[Feedback >](#) [Compare sources >](#)

Asian Pacific Journal of Cancer Prevention

Open Access ⓘ

Scopus coverage years: from 2000 to Present

Publisher: Asian Pacific Organization for Cancer Prevention

ISSN: 1513-7368

Subject area: [Medicine: Public Health, Environmental and Occupational Health](#) [Medicine: Epidemiology](#) [Medicine: Oncology](#)

[Biochemistry, Genetics and Molecular Biology: Cancer Research](#)

Source type: Journal

[View all documents >](#) [Set document alert](#) [Save to source list](#) [Source Homepage](#)

| | | |
|----------------|-------|---|
| CiteScore 2021 | 3.1 | ⓘ |
| SJR 2021 | 0.464 | ⓘ |
| SNIP 2021 | 0.814 | ⓘ |

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology ⓘ
CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ⓘ

$$3.1 = \frac{6,446 \text{ Citations } 2018 - 2021}{2,072 \text{ Documents } 2018 - 2021}$$

Calculated on 05 May, 2022.

CiteScoreTracker 2022 ⓘ

$$2.6 = \frac{4,782 \text{ Citations to date}}{1,841 \text{ Documents to date}}$$

Last updated on 05 September, 2022 - Updated monthly


CiteScore rank 2021 ⓘ

| Category | Rank | Percentile |
|--|----------|------------|
| Medicine | | |
| Public Health, Environmental and Occupational Health | #242/562 | 57th |


SCIMAGO SJR 2021: 0.46

<https://www.scimagojr.com/journalsearch.php?q=40173&tip=sid&clean=0>

Asian Pacific Journal of Cancer Prevention

| | | | |
|--|---|--|--|
| COUNTRY Thailand  Universities and research institutions in Thailand | SUBJECT AREA AND CATEGORY Biochemistry, Genetics and Molecular Biology └ Cancer Research Medicine └ Epidemiology └ Oncology └ Public Health, Environmental and Occupational Health | PUBLISHER Asian Pacific Organization for Cancer Prevention | H-INDEX 80 |
| PUBLICATION TYPE Journals | ISSN 15137368 | COVERAGE 2000-2021 | INFORMATION Homepage How to publish in this journal rmosavi@yahoo.com |

SCIMAGO SJR 2021 0.46



The dashboard displays several key metrics and trends:

- Citation Trends:** A line chart showing citation growth from 2000 to 2021, with a significant peak around 2015. It includes three series: Cites / Doc. (4 years), Cites / Doc. (3 years), and Cites / Doc. (2 years).
- International Collaboration:** A line chart showing the percentage of international collaboration, fluctuating between approximately 9% and 27% over the period.
- Document Types:** A stacked area chart comparing Citable documents (green) and Non-citable documents (red) from 2000 to 2021.
- Cited vs. Uncited Documents:** A stacked area chart showing Cited documents (green) and Uncited documents (red). A callout indicates that there were 1108 cited documents in 2021.
- Summary Widget:** A small widget on the bottom left shows the journal's Q3 ranking in Epidemiology, its SJR 2021 score of 0.46 (marked as a best quartile), and is powered by scimagojr.com.
- SCImago Graphica:** A section on the bottom right promotes the SCImago Graphica tool, describing it as a new data visualization tool for exploring and communicating data.



**KOMITE ETIK PENELITIAN KESEHATAN
RSUD Dr. SOETOMO SURABAYA**

**KETERANGAN KELAIKAN ETIK
(" ETHICAL CLEARANCE ")**

382 / Panke.KKE / V / 2016

KOMITE ETIK RSUD Dr. SOETOMO SURABAYA TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA PENELITIAN DENGAN JUDUL :

"Eksplorasi Tipe Human Pappiloma Virus yang Menginfeksi Penderita Kondiloma Akuminata"

PENELITI UTAMA : Dr. Gondo Mastutik, drh., M.Kes

**PENELITI LAIN : 1. Dwi Murtiastutik, dr., Sp.KK(K)
2. Alphania Rahniayu, dr., Sp.PA
3. Afria Arista, dr**

UNIT / LEMBAGA / TEMPAT PENELITIAN : RSUD Dr. Soetomo Surabaya

DINYATAKAN LAIK ETIK

SURABAYA, 18 MAY 2016
KETUA

(Prof. Hari Sukanto, dr., Sp.KK (K))
NIP. 19471115 1973 03 1 001