



Source details

New Armenian Medical Journal

Scopus coverage years: from 2009 to 2022

Publisher: Yerevan State Medical University

ISSN: 1829-0825

Subject area: Medicine: General Medicine

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

CiteScore 2021

0.4



SJR 2021

0.135



SNIP 2021

0.158



[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▼

$$0.4 = \frac{85 \text{ Citations 2018 - 2021}}{221 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$0.4 = \frac{91 \text{ Citations to date}}{228 \text{ Documents to date}}$$

Last updated on 05 February, 2023 • Updated monthly

CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine		
General Medicine	#664/826	19th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.





New Armenian Medical Journal

<p>COUNTRY</p> <p>Armenia</p> <ul style="list-style-type: none"> Universities and research institutions in Armenia Media Ranking in Armenia 	<p>SUBJECT AREA AND CATEGORY</p> <p>Medicine</p> <ul style="list-style-type: none"> Medicine (miscellaneous) 	<p>PUBLISHER</p> <p>UNIPRINT</p>	<p>H-INDEX</p> <p>7</p>
<p>PUBLICATION TYPE</p> <p>Journals</p>	<p>ISSN</p> <p>18290825</p>	<p>COVERAGE</p> <p>2009-2021</p>	<p>INFORMATION</p> <p>Homepage</p> <p>namj.ysmu@gmail.com</p>

SCOPE

Information not localized






Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS

options

<p>1</p> <p>Journal of Investigative Medicine</p> <p>USA</p> <p>53%</p> <p>similarity</p>	<p>2</p> <p>Biomedical papers of the Medical Faculty of the</p> <p>CZE</p> <p>50%</p> <p>similarity</p>	<p>3</p> <p>Medicine (United States)</p> <p>USA</p> <p>50%</p> <p>similarity</p>	<p>4</p> <p>American Journal of the Medical Sciences</p> <p>USA</p> <p>50%</p> <p>similarity</p>	<p>5</p> <p>Medicina (Lithuania)</p> <p>CHE</p> <p>49%</p> <p>similarity</p>
---	---	--	--	--



New Armenian Medical Journal

← Show this widget in your own website

Q4 Medicine (miscellaneous) best quartile

SJR 2021 0.14

powered by scimagojr.com

Just copy the code below and paste within your html code:

```
<a href="https://www.scimi
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.

Metrics based on Scopus® data as of April 2022

Leave a comment

Name

Email
(will not be published)

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on [@ScimagoJR](#)

Scimago Lab. Copyright 2007-2022. Data Source: [Scopus®](#)

EST MODUS IN REBUS
Horatio (Satire 1.1, 106)

[Edit Cookie Consent](#)



The Journal is founded by
Yerevan State Medical
University after M. Heratsi.

Rector of YSMU

Armen A. Muradyan

Address for correspondence:

Yerevan State Medical University
2 Koryun Street, Yerevan 0025,
Republic of Armenia

Phones:

(+37410) 582532 YSMU

(+37410) 580840 Editor-in-Chief

Fax: (+37410) 582532

E-mail: namj.ysmu@gmail.com, ysmi@mail.ru

URL: <http://www.ysmu.am>

*Our journal is registered in the databases, EBSCO
and Thomson Reuters (in the registration process)*



SCOPUS



EBSCO



THOMSON
REUTERS

Copy editors: Tatevik R. Movsisyan

Yana K. Karapetyan

Printed in SP "Vardan Mkrtychyan Feodori"

Director: Vardan F. Mkrtychyan

Address: Armenia, 0002, Yerevan,

Khorenatci St., 4 alley, Building 69

Phone: (+374 55) 020627,

E-mail: 75vardanm@mail.ru

Editor-in-Chief

Arto V. Zilfyan (Yerevan, Armenia)

Deputy Editors

Hovhannes M. Manvelyan (Yerevan, Armenia)

Hamayak S. Sisakyan (Yerevan, Armenia)

Executive Secretary

Stepan A. Avagyan (Yerevan, Armenia)

Editorial Board

Armen A. Muradyan (Yerevan, Armenia)

Drastamat N. Khudaverdyan (Yerevan, Armenia)

Levon M. Mkrtychyan (Yerevan, Armenia)

Foreign Members of the Editorial Board

Carsten N. GUTT (Memmingen, Germany)

Muhammad MIFTAHUSSURUR (Surabaya, Indonesia)

Alexander WOODMAN (Dharhan, Saudi Arabia)

Edita MARGARYAN (MOSCOW, RUSSIA)

Coordinating Editor (for this number)

Renata L. Markosyan (Yerevan, Armenia)

Editorial Advisory Council

Ara S. Babloyan (Yerevan, Armenia)

Aram Chobanian (Boston, USA)

Luciana Dini (Lecce, Italy)

Ruben V. Fanarjyan (Yerevan, Armenia)

Gerasimos Filippatos (Athens, Greece)

Gabriele Fragasso (Milan, Italy)

Samvel G. Galstyan (Yerevan, Armenia)

Edita G. MARGARYAN (MOSCOW, RUSSIA)

Arthur A. GRIGORIAN (Macon, Georgia, USA)

Armen Dz. Hambardzumyan (Yerevan, Armenia)

Seyran P. Kocharyan (Yerevan, Armenia)

Aleksandr S. Malayan (Yerevan, Armenia)

Mikhail Z. Narimanyan (Yerevan, Armenia)

Levon N. Nazarian (Philadelphia, USA)

Linda F. Noble-Hausslein (San Francisco, USA)

Eduard S. Sekoyan (Yerevan, Armenia)

Arthur K. SHUKRYAN (Yerevan, Armenia)

Suren A. STEPANYAN (Yerevan, Armenia)

Hakob V. Topchyan (Yerevan, Armenia)

Armen A. Trchunyan (Yerevan, Armenia)

Alexander Tsiskaridze (Tbilisi, Georgia)

Konstantin B. Yenkovyan (Yerevan, Armenia)

**SARGSYAN D., CABRERA J., KOSTIS J.B., FAHIM M.,
BEAVERS T., ZINONOS S., HSU V., MEKINIAN A., KOSTIS W.J.**

A STATEWIDE STUDY OF CARDIOVASCULAR OUTCOMES IN
PATIENTS WITH ANKYLOSING SPONDYLITIS

AVAGYAN S.A., ZILFYAN A.V., MURADYAN A.A.

NEW APPROACHES RELATED TO THE USE OF POLYAMINE-FREE
AND POLYAMINE-DEFICIENT DIETS IN THE LIST OF NUTRITIONAL
PRODUCTS FOR COVID-19 PATIENTS

**WARDHANA M.P., TUMANGGER D., JUWONO H.J.,
ERNAWATI E., RIFDAH S.N., Wafa I.A., KUNTAMAN K.,
DACHLAN E.G.**

THE EXPLORATION OF INFLAMMATORY AND COAGULATION
BIOMARKERS BETWEEN PREGNANT WOMEN WITH AND WITHOUT
COVID-19

**HOVHANNISYAN A.H., ASOYAN V.A., GYULAZYAN N.M.,
MADATYAN A.A., POGHOSYAN A.H., MOHAMMADI M.,
BARSEGHYAN E.S.**

COVID-19 INFECTION AND BUERGER'S SYNDROME: A CASE
REPORT

**MAKSIMOVA E.V., KLIARITSKAIA I.L., STILIDI E.I.,
GRIGORENKO E.I., MOSHKO YU.A.**

INFLUENCE OF CHANGES IN THE INTESTINAL MICROBIOME ON
THE COURSE AND PROGRESSION OF METABOLICALLY
ASSOCIATED FATTY LIVER DISEASE

**ARTONO A., PURNAMI N., HANDOKO E., MOON I.S.,
JANITRA S.N.**

CORRELATION BETWEEN THE PERFORATION SIZE AND PATENCY
OF EUSTACHIAN TUBE AND GRAFT UPTAKE IN INTACT CANAL
WALL TYMPANOPLASTY SURGERY: A STUDY OF 32 BENIGN-TYPE
CHRONIC SUPPURATIVE OTITIS MEDIA PATIENTS

PUTRI F.R., KURNIAWATI E.M., TIRTHANINGSIH N.W.

RISK FACTORS FOR POSTPARTUM HEMORRHAGE CAUSED BY
UTERINE ATONY

MOTAMED H., MEHRABI M.

CAN SERUM AMYLASE LEVEL EVALUATION FACILITATE EARLY
DIAGNOSIS OF ACUTE APPENDICITIS, AS AN ADJUNCTIVE

ELLANNY D.D., PERDANA R.F.,

Vol. 11, No. 3, 2017 (Supplement)
The New Armenian Medical Journal

CASE REPORT OF FATAL DEEP NECK ABSCESS: A COMPLICATION OF AERODIGESTIVE FOREIGN BODIES

No. 11-12, November 2017
Future doctor

**EBRAHIMI S.M., MOTAMED H., KALANTAR H.,
A., RAHIM F.**

HOSPITAL ADMISSIONS DUE TO SHORT-TERM EXPOSURE TO AIR POLLUTION: A SCOPING REVIEW

Medicine, science and education. All
electronic periodicals

KARIMPOUR F.F., AFROUGHI S .

PREVALENCE OF WEIGHT STATUS AND ASSOCIATED FACTORS OF UNDERWEIGHT AMONG THE MEDICAL STUDENTS IN IRAN

7000

Future doctor

MARKOSYAN R. L., BABAYAN H.N.

GRAVES DISEASES WITH SEVERE PROGRESSIVE OPHTHALMOPATHY AFTER THYROIDECTOMY. CASE REPORT

contact

**KHANCHI M., MATKERIMOV A.ZH., TERGEUSSIZOV A.S.,
DEMEUOV T.N., ZHAKUBAYEV M.A., KHANCHI M.M.,
BAUBEKOV A.A., TAJIBAYEV T.K., YERKINBAYEV N.N.,
SADUAKAS A.E., MAKKAMOV R.O.**

SURGICAL TREATMENT OF VISCERAL AND RENAL ABDOMINAL ARTERY ANEURYSMS OF VARIOUS ETIOLOGY

ALARSAN S.F. PALLIATIVE CARE: A CONCEPT ANALYSIS

PALLIATIVE CARE: A CONCEPT ANALYSIS

ASHWANI K., RAGHAVENDRA R., SUJATHA B.

EFFECTIVENESS OF PLATELET INDICES IN PREDICTING TYPE 2 DIABETES MELLITUS MICROVASCULAR COMPLICATIONS



CORRELATION BETWEEN THE PERFORATION SIZE AND PATENCY OF EUSTACHIAN TUBE AND GRAFT UPTAKE IN INTACT CANAL WALL TYMPANOPLASTY SURGERY: A STUDY OF 32 BENIGN-TYPE CHRONIC SUPPURATIVE OTITIS MEDIA PATIENTS

ARTONO A.^{1*}, PURNAMI N.¹, HANDOKO E.², MOON I.S.³, JANITRA S.N.¹

¹Department of Otorhinolaryngology Head and Neck Surgery, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

²Department of Otorhinolaryngology Head and Neck Surgery, Faculty of Medicine, Universitas Brawijaya – Dr. Saiful Anwar General Hospital, Malang, Indonesia

³Department of Otorhinolaryngology, Yonsei University, South Korea

Received 15.01.2022; accepted for printing 30.05.2022

ABSTRACT

Background: Factors that contribute to the success of Intact canal wall Tympanoplasty have yet to be certainly known. Several studies show varied success rates and success criteria.

Objective: To investigate the correlation between the perforation size and patency of Eustachian Tube and the results of graft uptake on benign-type chronic suppurative otitis media patients that undergo intact canal wall tympanoplasty surgery.

Methods: This research used observational descriptive analysis. The participants were patients diagnosed with benign-type chronic suppurative otitis media who were treated with intact canal wall tympanoplasty surgery in 2018. Data such as demographic data, smoking history, clinical findings including perforation size of tympanic membrane, Eustachian Tube patency, results of pure-tone audiometry examination, results of graft uptake, and type of Tympanoplasty were collected from each participant. The statistic tests in use included contingency coefficient correlation test and Fisher exact test.

Result: Most of the participants were male (53.13%), the average age was 28.03 ± 12.32 years old, and most participants do not smoke (71.87%). The overall graft uptake success rate is 71.87% ($n = 23$), while the failure rate is 28.12% ($n = 9$). Pure Tone Average (PTA) was 37.19 dB at pre-op and 25.79 dB at post-op, which is a significant improvement ($p=0.000$) with average hearing improvement (PTA-Gain) of 15.75 dB. The patent Eustachian Tube functions give the best graft uptake results of 88.90%, followed by partial patency at 75% and non-patent at 40% ($p=0.020$). The highest average of PTA in non-patent function of the tube is 20.94 dB. Hearing threshold improvement (PTA-Gain) on patent, partial, and non-patent ETF groups differs significantly ($p=0.046$). Results of graft uptake on small and medium perforation size show the highest rate of (81.80%), followed by large perforation (50%). The highest PTA in total perforation size is (18.90 dB), then medium (15.14 dB), and small (6.22 dB). The perforation size of TM before surgery does not correlate with the success of graft uptake ($p=0.297$). The size of perforation correlates with improvement of hearing threshold ($p=0.011$).

Conclusion: Eustachian Tube's function has correlation with the success of graft uptake and hearing threshold improvement (PTA-Gain), while the perforation size of TM before surgery correlates with hearing threshold improvement (PTA-Gain).

KEYWORDS: Benign-type chronic suppurative otitis media, Eustachian tube Patency, perforation size, intact canal wall Tympanoplasty, human and disease, biomedical

CITE THIS ARTICLE AS:

Artono A, Purnami N, Handoko E, Moon IS, Janitra SN; Correlation between the perforation size and patency of eustachian tube and graft uptake in intact canal wall tympanoplasty surgery: a study of 32 benign-type chronic suppurative otitis media patients; NAMJ v.16 (2022) no.6, p. 45-50; DOI: <https://doi.org/10.56936/18290825-2022.16.2-45>

ADDRESS FOR CORRESPONDENCE:

Artono A; Department of Otorhinolaryngology Head and Neck Surgery, Faculty of Medicine, Universitas Airlangga – , Airlangga, Gubeng, Surabaya, East Java 60286, Indonesia
E-mail: artono@fk.unair.ac.id; Tel.: (+6231) 55 01 649

INTRODUCTION

Chronic suppurative otitis media is a chronic middle ear infection that lasts more than two months characterized by persistent perforation of the tympanic membrane (TM) and continuous or intermittent discharge of secretions from the ear [Acuin J, 2004]. Chronic suppurative otitis media is known to be one of the most common diseases in developing countries, one of which is Indonesia. Data from WHO (World Health Organization) shows that the prevalence of chronic suppurative otitis media in developing countries such as Malaysia, Philippines and Thailand are still relatively high, namely 2-4% compared to that of developed countries in Europe such as Australia, England, Denmark and Finland which is around 0.4% [Mahadevan M et al., 2012; Thakur SK et al., 2017]. Persistent tympanic membrane perforation and continuous discharge from the ear can cause conductive hearing loss up to 60 dB, which is considered a serious disability. If intervention is not carried out, it can cause serious complications [Morris P, 2012]. The surgical therapy approach to benign-type chronic suppurative otitis media is the intact canal wall tympanoplasty. The removal of middle ear and mastoid disease is performed by maintaining the posterior wall of the external acoustic canal. The goals of intact canal wall tympanoplasty surgery are to eradicate disease in the middle ear and mastoid and to reconstruct the hearing mechanism with or without tympanic membrane grafts [Syms III CA et al., 2010]. Factors influencing the success of intact canal wall tympanoplasty surgery are still in debate.

many factors have been investigated to determine what influences the success of graft closure and hearing improvement. various studies that have been conducted have shown varying success rates and success criteria. several studies have shown that the success factor of intact canal wall tympanoplasty surgery depends on the size and location of

the perforation, the ossicular status, the type of graft and the function of the Eustachian tube. Other influencing factors include operating technique, operator experience, previous surgical history and smoking status [Alam M et al., 2017; Dangol K, Shrivastav RP, 2017]. The purpose of this study is to learn the relationship between the size of the perforation and patency of the Eustachian tube and the results of graft uptake in benign-type chronic suppurative otitis media patients undergoing intact canal wall tympanoplasty surgery.

MATERIALS AND METHODS

Participants of this study were benign-type chronic suppurative otitis media patients who underwent surgery in 2018 at Dr. Soetomo Hospital, Surabaya, Indonesia. This study used an observational descriptive analysis. Inclusion criteria included patients diagnosed with benign-type chronic suppurative otitis media who underwent intact canal wall tympanoplasty surgery. Exclusion criteria were the presence of underlying disease such as diabetes or immunodeficiency, cholesteatoma and revised surgical cases and incomplete data. There were 136 patients diagnosed with benign-type chronic suppurative otitis media and 56 of them (41.18%) underwent intact canal wall tympanoplasty surgery. The number of samples that met the inclusion criteria was 32 patients (57.14%). Participants were identified on their demographic data, smoking history, clinical findings including size of eardrum perforation, Eustachian tube patency, pure tone audiometry (PTA) examination, graft growth results and type of tympanoplasty. Perforation size was assessed in small (less than 50%), medium (50% to 75%) and large (more than 75%). Eustachian tube patency is measured by means of ETF-P (Eustachian Tube Function-Perforated) in Impedance tympanometry. Results of the PTA before and after surgery are measured in decibels (dB). Results of postoperative PTA examination and graft uptake were evaluated after 3 months after surgery and analyzed. Data analysis was conducted using contingency coefficient correlation Test and Fisher exact test on SPSS statistical package (version 16.0; SPSS). The level of significance was $p < 0.05$.



To overcome it is possible, due to the uniting the knowledge and will of all doctors in the world

RESULTS

A total of 56 participants (41.18%) underwent intact canal wall tympanoplasty surgery, 32 of which (57.14%) met the inclusion criteria. The majority of participants were male (53.13%). The average age was 28.03 ± 12.32 years, with median age of 25.50 (13.00-68.00) years. The youngest participant was 13.00 years old and the oldest was 68.00 years old. Most of the participants do not smoke (71.87%). The distribution of patients according to graft uptake success rate is provided in Table 1. Out of the 32 participants that underwent intact canal wall tympanoplasty, the overall success rate of graft uptake rate is 71.87% (n = 23), failure rate is 28.12% (n=9), and the highest success rate of graft uptake is in ≤ 20 years (81.81%) age group. Smoking has significant correlation with graft uptake success with the rate of 82.60% (p=0.031). Tympanoplasty type does not correlate with the graft uptake success (p=0.447) (Table 1).

PTA at pre-op amounts to 37.19 ± 18.52 dB




FIGURE 1. Comparison of PACuInTA Results Before (A) and After (B) Surgery

(with a range of 23.00-86.25 dB), and 25.79 ± 15.04 dB at post-op, which statistically saw significant improvement of p=0.000 with PTA-Gain of 11.40 dB (Figure 1). The patent Eustachian Tube's function provides the best graft uptake results of 88.90%, followed by partial (75%) and non-patent (40%). Eustachian tube patency significantly correlates with the success of the graft uptake (p=0.020). The largest PTA in non-patent tube function is (20.94 dB). The results of PTA-Gain in patent, partial and non-patent ETF groups are significantly different with p=0.046. The strength of the relationship is weak (r=0.304). The graft uptake on small and medium perforations shows the same results, namely 81.80%, while for large perforations it shows 50%. PTA is greatest at total perforation size (18.90 dB) followed by medium (15.14 dB) and small (6.22 dB). The preoperative size of TM perforation does not correlate with the success of graft uptake (p=0.297), but it correlates with improved hearing threshold (PTA-Gain) (p=0.011). This relationship is weak (r=0.403) (Table 2).

DISCUSSION

This research does not show a tendency for sex differences. Out of 32 participants, the age varies from 13 to 68 years old, with an average of 28.03 years old. The highest result is shown by ≤ 20 years age group (81.81%). Several studies show similar results [Shetty S, 2012; Das A et al., 2015]. This indicates the fact that chronic suppurative otitis media is primarily a middle ear infection that tends to occur in the first decade of life [Das A et al., 2015]. This study shows overall graft uptake success rate of

TABLE 1.

Success Rate of Graft Uptake				
	Quantity	Graft Uptake		Success Rates
		Suc	Fai	
Sex				
• Male	15 (53.13%)	12	3	80.00%
• Female	17 (46.88%)	11	6	64.70%
Age				
• ≤ 20	11 (34.38%)	9	2	81.81%
• 21-30	13 (40.60%)	9	4	69.20%
• >30	8 (25.00%)	5	3	62.50%
Education				
• Elementary School	0 (0.00%)	0	0	0.00%
• Junior High School	3 (9.38%)	2	1	66.67%
• Senior High School	20 (62.50%)	14	6	70.00%
• Bachelor	9 (28.12%)	7	2	77.78%
Smoking				
• Yes	9 (28.12%)	4	5	44.44%
• No	23 (71.87%)	19	4	82.60%
p=0.031				
Type of Tympanoplasty				
• Type 1	25 (78.13%)	20	5	75.00%
• Type 2	2 (6.25%)	2	0	100.00%
• Type 3	3 (9.38%)	1	2	33.33%
• Type 4	1 (3.12%)	1	0	100.00%
• Type 5	1 (3.12%)	0	1	0.00%
(p=0.477)				

NOTES: Suc -Success, Fai- Failure

TABLE 2.

Correlation between Eustachian Tube Patency and Perforation Size and PTA Gain

	Participant No.	Graft Uptake		PTA		
		Rate	Rate (%)	PRE-OP (dB)	POST-OP (dB)	Gain (dB)
Eustachian Tube Patency						
Patent	18	16	88.90%	49.72	41.06	8.66
Partial	4	3	75.00%	68.75	47.25	21.50
Non Patent	10	4	40.00%	60.94	40.00	20.94
			p=0.020			r=0.304 p=0.046
Perforation Size						
Small	11	9	81.80%	46.67	40.44	6.22
Medium	11	9	81.80%	60.56	45.42	15.14
Total	10	5	50.00%	56.10	37.20	18.90
			p=0.297			r=0.403 p=0.011

71.87% (n=23). This is in accordance with the study conducted by Alam which reported graft uptake results of 71.4% [Alam M et al., 2017]. Several other studies show different results. Shiromany M and coauthor reported 83.6% success rate of graft uptake and Naderpour M et al., reported similar results at 93.3% [Naderpour M et al., 2016; Shiromany A, Belaldavar BP, 2016]. Results of tympanoplasty surgery among smokers and non-smokers continue to be a controversial issue.

In this study, smoking participants have significant correlation with graft uptake success with the rate of 82.60% (p=0.031). This is in line with Cantrell's study [Cantrell RW, 1970]. Several studies claimed that there are differences in the results of graft uptake and hearing improvements in smoking and non-smoking patients [Naderpour M et al., 2016; Dangol K, Shrivastav RP 2017]. Eustachian Tube dysfunction refers to disruption of middle ear ventilation function caused by abnormal opening of the Eustachian Tube. Good middle ear aeration is essential for the success of the tympanoplasty procedure [Alam M et al., 2017; Li R et al., 2020]. This study shows that patent Eustachian tube function gives the best graft uptake results (88.90%). Eustachian Tube Function (ETF) is significantly related to the success of graft uptake (p=0.020). The rates of improvement of hearing threshold (PTA-Gain) in patent, partial, and non-patent (block) ETF groups differ significantly with p=0.046, despite the weak relationship (r=0.304).

These results are in line with that of several other researchers. Alam M reported significantly different results of average PTA between patients with patent and non-patent (block) Eustachian tubes (p=0.022, t=2.63), while no significant difference in graft uptake results (p=0.629) [Alam M et al., 2017; Perdana RF et al., 2020; Wibowo AP et al., 2020]. Holmquist, Manning et al., and Merchant et al., also claimed that good Eustachian tube function will increase Tympanoplasty success rate [Holmquist J, 1968; Manning SC et al., 1987; Merchant SN et al., 2003]. Different results were reported by Li R et al., who claimed that Eustachian tube function may not have effects on the results of type I tympanoplasty for chronic suppurative otitis media [Li R et al., 2020].

Results of surgery in this study were measured based on the graft uptake success rate and post-op hearing threshold improvement. Various studies used different criteria to assess hearing improvement after Tympanoplasty, such as hearing gain method or average of Air Bone Gap in each frequency [Sarker MZ et al., 2011]. In this study, the TM perforation sizes that give the best graft uptake results are small and medium sizes (81.80%). However, TM perforation size before surgery does not correlate with graft uptake success (p=0.297). This perforation size correlates with hearing threshold improvement (PTA-Gain) (p=0.011), which this study shows 15.75 dB in average. Correlation between perforation size and graft uptake

success and improvement of hearing threshold is reported by several researchers. Al-Ghamdi SA et al., reported that perforation size is a factor in graft uptake results, although statistically there is no significant differences between small and medium perforation in the graft uptake success [Al-Ghamdi SA, 1994]. Saleh et al., in their study reported the highest myringoplasty success rate on small perforation at 93.3%, medium and large perforation at 87.5% and 71.4% respectively. A study by Pfammatter et al., claimed success of complete closure of AB Gap on 20% of the patients and AB Gap of 10 dB HL in 80% of the cases. They also reported that perforation size has the most positive impact on the outcome. Thiel et al., reported AB Gap closure up to 10 dB HL on 53% of the patients and

claimed that in cases with larger perforations, improvement of hearing is possible [Pfammatter A et al., 2013; Thiel G et al., 2013; Saleh C et al., 2020]. Different claim was made by Vartiainen and Nuutinen, in which perforation size affects neither ABG closure and hearing improvement [Vartiainen E, Nuutinen J 1993].

CONCLUSIONS

Eustachian tube patency correlates with the success rate of graft uptake and hearing threshold improvement (PTA-Gain), while TM perforation size before surgery does not correlate with graft uptake success, but does with hearing threshold improvement (PTA-Gain).

ACKNOWLEDGEMENT : We would like to thank our research assistance, Dini Fitrilia as manuscript editor and Sakinah Ramadhani for helping us in collecting data and assisting in translating our manuscripts.

Ethics approval and consent to participate: Ethical clearance was obtained from the Senate Research and Publications Committee of Dr. Soetomo General Hospital

REFERENCES

1. Acuin, J. (2004). Global burden of disease due to chronic suppurative otitis media: Disease, deafness, deaths and DALYs Chronic Suppurative Otitis Media-Burden of Illness and Management Options. Geneva: World Health Organisation , pp. 9–23.
2. Al-Ghamdi, S.A. (1994). Tympanoplasty: factors influencing surgical outcome. Annals of Saudi Medicine 14(6), pp. 483–485.
3. Alam, M., Sharma, S.C. and Hasan, S.A. (2017). Does Eustachian tube patency affect the outcome of graft uptake rate and average audiological gain in Intact canal wall and Canal wall down Mastoidectomies—A study of 50 patients. International Journal of Medical Science and Public Health 6(2), pp. 418–422.
4. Cantrell, R.W. (1970). Myringoplasty failure related to smoking: a preliminary report. Otolaryngologic clinics of North America 3(1), pp. 141–145.
5. Dangol K. and Shrivastav, R.P. (2017). Study of various prognostic factors affecting successful Myringoplasty in a tertiary care Centre. International Archives of Otorhinolaryngology 21, pp. 250–254.
6. Das A, Sen B, Ghosh D, Sengupta A. Myringoplasty: Impact of size and site of perforation on the success rate. Indian J Otolaryngol Head Neck Surg [Internet]. 2015;67(2):185–9. Available from: <http://dx.doi.org/10.1007/s12070-014-0810-7>
7. Holmquist J. (1968). The role of the eustachian tube in myringoplasty. Acta oto-laryngologica 66(1–6), pp. 289–295.
8. Li R., Wu, N., Zhang, J., Hou, Z. and Yang, S. (2020). Analysis on the correlation between Eustachian tube function and outcomes of type I tympanoplasty for chronic suppurative otitis media. Acta Oto-Laryngologica 140(8), pp. 656–659.
9. Mahadevan M, Navarro-Locsin G, Tan HKK, Yamanaka N, Sonsuwan N, Wang PC, et al (2012). A review of the burden of disease due to otitis media in the Asia-Pacific. International Journal of Pediatric Otorhinolaryngology 76(5), pp. 623–635.
10. Manning, S.C., Cantekin, E.I., Kenna, M.A. and Bluestone, C.D. (1987). Prognostic value of eustachian tube function in pediatric tympanoplasty. The Laryngoscope 97(9), pp. 1012–1016.

11. Merchant, S.N., McKenna, M.J., Mehta, R.P., Ravicz, M.E. and Rosowski, J.J. (2003). Middle ear mechanics of Type III tympanoplasty (stapes columella): II. Clinical studies. *Otology & neurotology* 24(2), pp. 186–194.
12. Morris, P. (2012). Chronic suppurative otitis media. *BMJ clinical evidence* 2012
13. Naderpour, M., Moghadam, Y.J., Ghanbarpour, E. and Shahidi, N. (2016). Evaluation of factors affecting the surgical outcome in tympanoplasty. *Iranian journal of otorhinolaryngology* 28(85), p. 99.
14. Perdana, R.F., Herawati, S., Surarso, B. and Aksono, E.B. (2020). Correlation of Aggressivity Papilloma Recurrent Respiratory Tract With Human Papillomavirus Types 6 And 11. *Medico Legal Update* 20(1), pp. 471–476.
15. Pfammatter, A., Novoa, E. and Linder, T. (2013). Can myringoplasty close the air-bone gap? *Otology & Neurotology* 34(4), pp. 705–710.
16. Saleh, C., Hayat, H., Sumartono, S. And Pratiwi, R.N. (2020). Moderating of Religiosity on Reward and Engagement: Empirical Study in Indonesia Public Service. *The Journal of Asian Finance, Economics, and Business* 7(6), pp. 287–296.
17. Sarker, M.Z., Ahmed, M., Patwary, K., Islam, R. and Joarder, A.H. (2011). Factors affecting surgical outcome of myringoplasty. *Bangladesh Journal of Otorhinolaryngology* 17(2), pp. 82–87.
18. Shetty S. Pre-Operative and Post-Operative Assessment of Hearing following Tympanoplasty. *Indian J Otolaryngol Head Neck Surg.* 2012;64(4):377–81.
19. Shiromany, A. and Belaldavar, B.P. (2016). Effect of Eustachian tube function on tympanoplasty outcome in chronic otitis media patients: Cross-sectional study. *Indian Journal of Health Sciences and Biomedical Research (KLEU)* 9(3), p. 279.
20. Syms III, C.A., Syms, M.J. and Sheehy, J.L. procedure. In: (2010). *Mastoidectomy—intact canal wall Otologic surgery.* Elsevier, pp. 195–207.
21. Thakur, S.K., Singh, S.K., Acharya, R., Anwar, A. and Ghimire, N. (2017). Sociodemographic profile and the associated factors of chronic otitis media in rural areas of eastern Nepal., *International Journal of Otorhinolaryngology and Head and Neck Surgery* 3(2), pp. 222–227.
22. Thiel, G., Mills, R.P. and Mills, N. (2013). Factors affecting hearing improvement following successful repair of the tympanic membrane. *The Journal of Laryngology & Otology* 127(4), pp. 349–353.
23. Vartiainen, E. and Nuutinen, J. (1993). Success and pitfalls in myringoplasty: follow-up study of 404 cases. *The American journal of otology* 14(3), pp. 301–305.
24. Wibowo A.P., Perdana, R.F. and Herawati, S. (2020). Case report : Management of achalasia with esophageal candidiasis and bradycardia at tertiary hospital., *6937(June 2019)*, pp. 6931–6937.



CONTENTS

- 4. SARGSYAN D., CABRERA J., KOSTIS J.B., FAHIM M., BEAVERS T., ZINONOS S., HSU V., MEKINIAN A., KOSTIS W.J.**
A STATEWIDE STUDY OF CARDIOVASCULAR OUTCOMES IN PATIENTS WITH ANKYLOSING SPONDYLITIS
- 14. AVAGYAN S.A., ZILFYAN A.V., MURADYAN A.A.**
NEW APPROACHES RELATED TO THE USE OF POLYAMINE-FREE AND POLYAMINE-DEFICIENT DIETS IN THE LIST OF NUTRITIONAL PRODUCTS FOR COVID-19 PATIENTS
- 25. WARDHANA M.P., TUMANGGER D., JUWONO H.J., ERNAWATI E., RIFDAH S.N., WAFA I.A., KUNTAMAN K., DACHLAN E.G.**
THE EXPLORATION OF INFLAMMATORY AND COAGULATION BIOMARKERS BETWEEN PREGNANT WOMEN WITH AND WITHOUT COVID-19
- 33. HOVHANNISYAN A.H., ASOYAN V.A., GYULAZYAN N.M., MADATYAN A.A., POGHOSYAN A.H., MOHAMMADI M., BARSEGHYAN E.S.**
COVID-19 INFECTION AND BUERGER'S SYNDROME: A CASE REPORT
- 38. MAKSIMOVA E.V., KLIARITSKAIA I.L., STILIDI E.I., GRIGORENKO E.I., MOSHKO YU.A.**
INFLUENCE OF CHANGES IN THE INTESTINAL MICROBIOME ON THE COURSE AND PROGRESSION OF METABOLICALLY ASSOCIATED FATTY LIVER DISEASE
- 45. ARTONO A., PURNAMI N., HANDOKO E., MOON I.S., JANITRA S.N.**
CORRELATION BETWEEN THE PERFORATION SIZE AND PATENCY OF EUSTACHIAN TUBE AND GRAFT UPTAKE IN INTACT CANAL WALL TYMPANOPLASTY SURGERY: A STUDY OF 32 BENIGN-TYPE CHRONIC SUPPURATIVE OTITIS MEDIA PATIENTS
- 51. PUTRI F.R., KURNIAWATI E.M., TIRTHANINGSIH N.W.**
RISK FACTORS FOR POSTPARTUM HEMORRHAGE CAUSED BY UTERINE ATONY
- 60. MOTAMED H., MEHRABI M.**
CAN SERUM AMYLASE LEVEL EVALUATION FACILITATE EARLY DIAGNOSIS OF ACUTE APPENDICITIS, AS AN ADJUNCTIVE BIOMARKER?
- 66. BELLANNY D.D., PERDANA R.F.,**
CASE REPORT OF FATAL DEEP NECK ABSCESS: A COMPLICATION OF AERODIGESTIVE FOREIGN BODIES
- 76. EBRAHIMI S.M., MOTAMED H., KALANTAR H., KALANTARI A., RAHIM F.**
HOSPITAL ADMISSIONS DUE TO SHORT-TERM EXPOSURE TO AIR POLLUTION: A SCOPING REVIEW
- 91. KARIMPOUR F.F., AFROUGHI S.**
PREVALENCE OF WEIGHT STATUS AND ASSOCIATED FACTORS OF UNDERWEIGHT AMONG THE MEDICAL STUDENTS IN IRAN
- 100. MARKOSYAN R. L., BABAYAN H.N.**
GRAVES DISEASES WITH SEVERE PROGRESSIVE OPHTHALMOPATHY AFTER THYROIDECTOMY. CASE REPORT.
- 104. KHANCHI M., MATKERIMOV A.ZH., TERGEUSSIZOV A.S., DEMEUOV T.N., ZHAKUBAYEV M.A., KHANCHI M.M., BAUBEKOV A.A., TAJIBAYEV T.K., YERKINBAYEV N.N., SADUAKAS A.E., MAKKAMOV R.O.**
SURGICAL TREATMENT OF VISCERAL AND RENAL ABDOMINAL ARTERY ANEURYSMS OF VARIOUS ETIOLOGY
- 113. ALARSAN S.F.**
PALLIATIVE CARE: A CONCEPT ANALYSIS
- 118. ASHWANI K., RAGHAVENDRA R., SUJATHA B.**
EFFECTIVENESS OF PLATELET INDICES IN PREDICTING TYPE 2 DIABETES MELLITUS MICROVASCULAR COMPLICATIONS



The Journal is founded by
Yerevan State Medical
University after M. Heratsi.

Rector of YSMU

Armen A. Muradyan

Address for correspondence:

Yerevan State Medical University
2 Koryun Street, Yerevan 0025,
Republic of Armenia

Phones:

(+37410) 582532 YSMU

(+37410) 580840 Editor-in-Chief

Fax: (+37410) 582532

E-mail: namj.ysmu@gmail.com, ysmiu@mail.ru

URL: <http://www.ysmu.am>

*Our journal is registered in the databases of Scopus,
EBSCO and Thomson Reuters (in the registration process)*



SCOPUS



EBSCO



THOMSON
REUTERS

Copy editor: Tatevik R. Movsisyan

Printed in "VARM" LLC
Director: Ruzanna Arakelyan
Armenia, 0018, Yerevan,
Tigran Mec 48, 43
Phone: (+374 91) 19 29 00,
E-mail: armana6@mail.ru

Editor-in-Chief

Arto V. **Zilfyan** (Yerevan, Armenia)

Deputy Editors

Hovhannes M. **Manvelyan** (Yerevan, Armenia)

Hamayak S. **Sisakyan** (Yerevan, Armenia)

Executive Secretary

Stepan A. **Avagyan** (Yerevan, Armenia)

Editorial Board

Armen A. **Muradyan** (Yerevan, Armenia)

Drastamat N. **Khudaverdyan** (Yerevan, Armenia)

Levon M. **Mkrtchyan** (Yerevan, Armenia)

Foregin Members of the Editorial Board

Carsten N. **GUTT** (Memmingen, Germany)

Muhammad **MIFTAHUSSURUR** (Indonesia)

Alexander **WOODMAN** (Dharhan, Saudi Arabia)

Hesam Adin **Atashi** (Tehran, Iran)

Coordinating Editor (for this number)

Inkar **Sagatov** (Almaty, Kazakhstan)

Editorial Advisory Council

Ara S. **Babloyan** (Yerevan, Armenia)

Aram **Chobanian** (Boston, USA)

Luciana **Dini** (Lecce, Italy)

Azat A. **Engibaryan** (Yerevan, Armenia)

Ruben V. **Fanarjyan** (Yerevan, Armenia)

Gerasimos **Filippatos** (Athens, Greece)

Gabriele **Fragasso** (Milan, Italy)

Samvel G. **Galstyan** (Yerevan, Armenia)

Arthur A. **Grigorian** (Macon, Georgia, USA)

Armen Dz. **Hambardzumyan** (Yerevan, Armenia)

Seyran P. **Kocharyan** (Yerevan, Armenia)

Aleksandr S. **Malayan** (Yerevan, Armenia)

Mikhail Z. **Narimanyan** (Yerevan, Armenia)

Levon N. **Nazarian** (Philadelphia, USA)

Yumei **Niu** (Harbin, China)

Linda F. **Noble-Haeusslein** (San Francisco, USA)

Arthur K. **Shukuryan** (Yerevan, Armenia)

Suren A. **Stepanyan** (Yerevan, Armenia)

Gevorg N. **Tamamyan** (Yerevan, Armenia)

Hakob V. **Topchyan** (Yerevan, Armenia)

Alexander **Tsiskaridze** (Tbilisi, Georgia)

Konstantin B. **Yenkoyan** (Yerevan, Armenia)

Peijun **Wang** (Harbin, China)