



# Scientifica

Hindawi Publishing Corporation  
<http://www.hindawi.com>

Volume 2017

Scientifica

Journal overview 	For authors	For reviewers	For editors	Table of Contents	Special Issues 
--	-------------	---------------	-------------	-------------------	--

Scientifica / Editorial Board


On this page
Editorial Board

# Editorial Board

## Academic Editors

- Agronomy 
- Allergy 
- Anatomy 
- Anesthesiology 
- Biochemistry 
- Bioinformatics 
- Biomaterials 
- Biophysics 
- Biotechnology 
- Botany 
- Cardiology 
- Parasitology 

- **Zbynek Bozdech**, Nanyang Technological University, Singapore
- **Fabrizio Bruschi** , Università di Pisa, Italy
- **David J. Bzik**, The Geisel School of Medicine at Dartmouth, USA
- **Lewis B. Coons**, University of Memphis, USA
- **Omar S. Harb** , University of Pennsylvania, USA
- **Jude M. Przyborski** , Philipps-Universität Marburg, Germany
- **Andrew Taylor-Robinson** , Central Queensland University, Australia
- **Alessandra Torina**, IZS, Italy
- **Carsten Wrenger** , Bernhard Nocht Institute for Tropical Medicine, Germany





**Journal metrics**

Acceptance rate	14%
Submission to final decision	101 days
Acceptance to publication	18 days
CiteScore	3.300
Journal Citation Indicator	0.590
Impact Factor	-









[See full report](#)

**APC** \$925

 **Submit**

 **Author guidelines**

## Table of Contents

2022 	
Scientifica - Volume 2022 - Article ID 1405900 - Review Article	
<b>Biofungicides for Improvement of Potato (<i>Solanum tuberosum</i> L.) Production</b>	
John Ehiobu   Emrobowansan Idamokoro   Anthony Afolayan	
03 Oct 2022	Download PDF 
Scientifica - Volume 2022 - Article ID 1116011 - Research Article	
<b>An Economic Analysis of Tourism Contribution for Urban Poverty Reduction in Major Tourist Towns of North Shewa Zone, Amhara Region, Ethiopia</b>	
Abebaw Hailu Fikire   Zemenu Bires   Girma Mulugeta Emeru	
30 Sep 2022	Download PDF 
Scientifica - Volume 2022 - Article ID 6059880 - Research Article	
<b>Optimization and Validation of ELISA for Aflatoxin B1 Detection in Fermented Forages and Feeds</b>	
Enikő Horváth   Tünde Pusztahelyi   ...   István Pócsi	
30 Sep 2022	Download PDF 
Scientifica - Volume 2022 - Article ID 3552491 - Research Article	
<b>Antimalarial Activity of Extract and Fractions of <i>Sauropus androgynus</i> (L.) Merr.</b>	
Wiwied Ekasari   Dwi Fatmawati   ...   Ram K. Sahu	
08 Sep 2022	Download PDF 
Scientifica - Volume 2022 - Article ID 9130252 - Research Article	
<b>A Potential Anticancer Mechanism of Finger Root (<i>Boesenbergia rotunda</i>) Extracts against a Breast Cancer Cell Line</b>	
Muhammad Hermawan Widyananda   Septian Tri Wicaksono   ...   Nashi Widodo	
05 Sep 2022	Download PDF 
Scientifica - Volume 2022 - Article ID 6729248 - Research Article	
<b>Food Waste Behaviour and Awareness of Malaysian</b>	
Chooi Lin Phooi   Elisa Azura Azman   ...   Evelyn Shin Rou Koay	
29 Aug 2022	Download PDF 
Scientifica - Volume 2022 - Article ID 6882147 - Research Article	
<b>Inflammation Modulating Activity of the Hydroethanol Stem Bark Extract of <i>Bombax costatum</i> in Murine Models</b>	
Meshack Antwi-Adjei   Kofi Oduro Yeboah   ...   Newman Osafo	
22 Aug 2022	Download PDF 

# Scientifica

## + Journal Menu

Scientifica - Volume 2022 - Article ID 8625345 - Research Article		
<b>Rate and Associated Factors of Suicidal Behavior among Adolescents in Bangladesh and Indonesia: Global School-Based Student Health Survey Data Analysis</b>		
M. Marthoenis   S. M. Yasir Arafat		
05 Aug 2022	<a href="#">Download PDF</a>	
Scientifica - Volume 2022 - Article ID 1068111 - Research Article		
<b>The Study of Estimation of Landfill Capacity through Dynamic System Approach</b>		
Lina Warlina   Sri Listyarini		
19 Jul 2022	<a href="#">Download PDF</a>	
Scientifica - Volume 2022 - Article ID 9675041 - Research Article		
<b>Detection of <i>Bacillus</i> Species with Arsenic Resistance and Plant Growth Promoting Efficacy from Agricultural Soils of Nepal</b>		
Lil Budha Magar   Binod Rayamajhee   ...   Pramod Poudel		
19 Jul 2022	<a href="#">Download PDF</a>	
Scientifica - Volume 2022 - Article ID 5616683 - Research Article		
<b>Salinity Stress Response of Rice (<i>Oryza sativa</i> L. cv. Luem Pua) Calli and Seedlings</b>		
Worasitikulya Taratima   Titirat Chomarsa   Pitakpong Maneerattanarungroj		
11 Jul 2022	<a href="#">Download PDF</a>	
Scientifica - Volume 2022 - Article ID 4230397 - Research Article		
<b>Antimicrobial Potential of Extract from a <i>Pseudomonas aeruginosa</i> Isolate</b>		
Francis Kwaku Dzideh Amankwah   Stephen Yao Gbedema   ...   Vivian Etsiapah Boamah		
29 Jun 2022	<a href="#">Download PDF</a>	
Scientifica - Volume 2022 - Article ID 7970435 - Research Article		
<b>Woody Species Composition, Plant Communities, and Environmental Determinants in Gennemar Dry Afromontane Forest, Southern Ethiopia</b>		
Shemsu Ahmed   Debissa Lemessa   Abera Seyum		
22 Jun 2022	<a href="#">Download PDF</a>	
Scientifica - Volume 2022 - Article ID 6309679 - Research Article		
<b>Improving the Genetics of Tuber Yield and Resistance to Mite to Avoid Mite Incident and to Increase the Productivity of Cassava (<i>Manihot esculenta</i> Crantz)</b>		
Sholihin   Sri W. Indiaty   ...   Rohmad Budiono		
16 Jun 2022	<a href="#">Download PDF</a>	

## Research Article

# Antimalarial Activity of Extract and Fractions of *Sauropus androgynus* (L.) Merr.

Wiwied Ekasari <sup>1</sup>, Dwi Fatmawati <sup>1</sup>, Siti M. Khoiriah <sup>1</sup>, Wenda A. Baquiuddin <sup>1</sup>,  
Hawi Q. Nisa <sup>1</sup>, Adinda A. S. Maharupini <sup>1</sup>, Tutik S. Wahyuni <sup>1</sup>, Rice D. Oktarina <sup>1</sup>,  
Eko Suhartono <sup>2</sup>, and Ram K. Sahu <sup>3</sup>

<sup>1</sup>Department of Pharmaceutical Sciences, Faculty of Pharmacy, Airlangga University, Surabaya 60115, East Java, Indonesia

<sup>2</sup>Department of Medical Chemistry/Biochemistry, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin 70122, South Kalimantan, Indonesia

<sup>3</sup>Department of Pharmaceutical Science, Assam University (A Central University), Silchar 788011, Assam, India

Correspondence should be addressed to Wiwied Ekasari; [wiwied-e@ff.unair.ac.id](mailto:wiwied-e@ff.unair.ac.id)

Received 19 May 2022; Accepted 18 August 2022; Published 8 September 2022

Academic Editor: Carsten Wrenger

Copyright © 2022 Wiwied Ekasari et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Sauropus androgynus* (L.) Merr., in the Indonesian local name known as “Katuk,” is a tropical shrub plant of the family Euphorbiaceae. Based on genus and chemotaxonomic approaches, as well as in vitro testing of *Plasmodium falciparum*, leaves of *S. androgynus* are presumed to have an active compound content as an antimalarial. The current study aims to investigate the antimalarial activity of 96% ethanol extract and fractions of *S. androgynus* leaves. The ethanolic extract was fractionated using the vacuum liquid chromatography (VLC) method with three solvents of different polarities (n-hexane, chloroform, and 96% ethanol). The fraction obtained was then evaluated for antimalarial activity against *P. falciparum* 3D7 strain. The ethanolic extract was evaluated for antimalarial suppressive and prophylactic activity against *P. berghei*-infected mice, as well as inhibitory activity against the heme detoxification process in vitro. Fractionation of ethanolic extract resulted in seven combined fractions, with the most active fraction being FV (50% inhibitory concentration (IC<sub>50</sub>) = 2.042 µg/mL). The ethanolic extract showed good parasitic suppressive (therapeutic) activity with a median effective dose (ED<sub>50</sub>) value of 15.35 mg/kg body weight. In a prophylactic test, ethanolic extract showed parasite growth inhibitory activity of 67.74 ± 9.21% after the administration of 400 mg/kg body weight for 4 days before infection, and 65.30 ± 10.44% after the administration of 200 mg/kg body weight for 8 consecutive days (4 days before and after infection). The ethanolic extract also showed an effect in inhibiting the formation of β-hematin of about 26.87–79.36% at a concentration of 0.1–4 mg/mL and an IC<sub>50</sub> value of 0.479 mg/mL. The *S. androgynus* leaves were shown to have antimalarial activity in vitro and in vivo, where ethanolic extract were more active compared with the fraction obtained. The antimalarial properties of the extract showed a higher suppressive activity than prophylactic activity.

## 1. Introduction

Malaria is an infectious disease caused by a protozoan parasite of the genus *Plasmodium*. The spread of *Plasmodium* into the human body is through the bite of female mosquitoes of the genus *Anopheles*. Malaria in humans is caused by five parasitic species of the genus *Plasmodium*, including *P. falciparum*, *P. vivax*, *P. ovale*, *P. malariae*, and *P. knowlesi* [1]. Malaria is one of the infectious diseases that can cause death, anemia, and decrease work productivity [2, 3].

Globally, there are an estimated 241 million malaria cases in 85 malaria-endemic countries in 2020, and there is a 12% increase in malaria-related deaths compared with 2019 [4]. Nationally, the number of malaria cases in Indonesia has decreased by more than 50% from 465,764 cases (annual parasite incidence (API) = 1.96%) in 2010 to 222,085 cases (API = 0.84%) in 2018 [5]. However, there was a stagnation in malaria cases after 2014, which indicates that interventions in elimination programs that have already been implemented are not enough to reduce cases drastically.

Efforts in controlling malaria are essential including prevention, diagnosis, and treatment. The diagnosis of malaria can be seen from the results of the examination of blood preparations, while chloroquine is used to treat malaria after a report of resistance, malaria treatment is not given as a single drug but in combination with artemisinin-based combination therapy (ACT) [6]. Efforts are needed to develop new antimalarial drugs from synthetic materials and natural materials that empirically have antimalarial activity, mainly plant-derived. One plant of the genus *Sauropus* was shown to have antimalarial activity against *P. falciparum* K1 strain from the 90% methanol fraction of *S. spatulifolius* leaves with an  $IC_{50}$  of 6.10  $\mu\text{g/mL}$  [7].

Based on genus and chemotaxonomic approaches, it is thought that the leaves of *S. androgynus* also have the content of compounds as antimalarials against *P. falciparum*. *S. androgynus* belongs to the family Euphorbiaceae known as star gooseberry which is a tropical shrub plant as a leafy vegetable and is a medicinal plant that has high antioxidant potential, can lose weight, and launch breastfeeding [8]. In addition, it can also be used for the treatment of several diseases such as antimicrobial, anti-inflammatory, antipyretic, antispasmodic, antiulcer, anti-allergic, antioxidant, antitumor, anticancer, antiulcer, immunostimulant, and lipooxygenase inhibitor [9]. *S. androgynus* contains macro-nutrient and micronutrient components. It also contains polyphenols, vitamins, saponins, tannins, alkaloids, glycosides, and essential minerals including sodium (Na), potassium (K), phosphorus (P), iron (Fe), magnesium (Mg), copper (Cu), zinc (Zn), manganese (Mn), and cobalt (Co) [8]. Scientific information on biological activity, especially in relation to malaria, of *S. androgynus* leaves is still scarce. In vitro antiplasmodial activity of *S. androgynus* against the human malaria parasite *P. falciparum* was first reported by Mahardiani et al. [10]. In the preliminary study, n-hexane, chloroform, and 96% ethanol extract from *S. androgynus* leaves showed antiplasmodial activity against *P. falciparum* 3D7 strain with  $IC_{50}$  values of 1.23, 0.85, and 1.88  $\mu\text{g/mL}$ , respectively, and were shown to be nontoxic in human liver cancer cell lines. According to Berthi et al. [11], the antiplasmodial activity of the extract is considered to be very active with  $IC_{50} < 5 \mu\text{g/mL}$ , promising with  $IC_{50}$  6–15  $\mu\text{g/mL}$ , moderate with  $IC_{50}$  16–30  $\mu\text{g/mL}$ , low with  $IC_{50}$  31–50  $\mu\text{g/mL}$ , and inactive if  $IC_{50} > 50 \mu\text{g/mL}$ . Thus, the antiplasmodial activity shown by n-hexane, chloroform, and ethanol extracts in the study was classified as very active ( $IC_{50} < 5 \mu\text{g/mL}$ ). Nevertheless, n-hexane and chloroform are among the categories of solvents that should be restricted in pharmaceutical products due to their toxicity properties [12]. Meanwhile, ethanol has been known as a good solvent for polyphenol extraction and is safe for human consumption [13]. Therefore, this study aims to identify the active fraction of 96% ethanol extract of *S. androgynus* leaves, and investigate the antimalarial activity of ethanolic extract using a mouse infection model as well as the inhibitory activity of the heme detoxification process in vitro.

## 2. Materials and Methods

**2.1. Plant Sample Collection.** *S. androgynus* leaves were obtained and determined from Purwodadi-LIPI Botanical Garden, Pasuruan, Indonesia with No. 074/68/102.7/2018.

**2.2. Preparation of Fraction of Ethanolic Crude Extracts.** The 96% ethanol extract of *S. androgynus* leaves produced by Mahardiani et al. [10] was used in this study and fractionated using VLC. The extract (2.505 g) was added with 85.015 g of silica gel 60 (0.040–0.063 mm) to dry and homogeneous, then put into a sintered glass containing two-thirds of full silica gel 60. The number of mobile phases used in each elution was 40 mL with a mixture of solvents of increased polarity, namely n-hexane, chloroform, and 96% ethanol. The mobile phase is poured into the sintered glass through filter paper and drawn off with a vacuum pump. The filtrate is housed in a vial for subsequent phytochemical screening using thin layer chromatography (TLC). Filtrates showing the same stains at phytochemical screening were combined and concentrated in a rotary evaporator at 45 rpm and 40°C, then concentrated to dryness with an oven at 36°C to obtain a dry fraction.

**2.3. In Vitro Antimalarial Screening.** *P. falciparum* 3D7 strain (chloroquine-sensitive) was used in this study to determine the in vitro antimalarial activity of *S. androgynus* leaf fraction. Parasites are routinely maintained according to the method described by Trager and Jensen [14] with slight modifications in the Laboratory of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia. *P. falciparum* cultures were maintained in human red blood cells (O+ blood type) with 5% hematocrit in Roswell Park Memorial Institute (RPMI) 1640 containing 2-(4-(2-hydroxyethyl)-1-piperazinyl)-ethanesulfonic acid (HEPES) 22.3 mM, hypoxanthine, sodium bicarbonate, gentamicin, and 10% human O+ plasma, and incubated at 37°C with 5%  $\text{CO}_2$ .

The test was performed in duplicate in a 24-well well plate with 1% initial parasitemia. The *S. androgynus* leaf fraction was dissolved with dimethyl sulfoxide (DMSO) and prepared in various concentrations, and DMSO was used as the negative control. After 48 h of incubation, thin blood strips were prepared on glass slides, fixed in methanol, and stained with Giemsa. The percentage of parasitemia was calculated microscopically, then compared with a negative control for the determination of the percentage of inhibition of parasite growth. The  $IC_{50}$  value was calculated using probit analysis.

**2.4. Experimental Animals and Parasite Inoculation.** BALB/c mice, male, aged 6–8 weeks, and weighing 25–35 g, obtained from Pharmaceutical Veterinary Center (PUS-VETMA), Surabaya, Indonesia, were used for the test. Mice were maintained in the veterinary laboratory of the Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia, under standard conditions at room temperature by exposing



them to a cycle of 12 h of light and 12 h of dark, with food and ad libitum water. Mice are handled based on internationally accepted guidelines [15], and animal testing protocols have been approved by the Research Ethics Commission of the Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia (No: 2.TO118.07.2018).

*P. berghei* ANKA strain was obtained from the Eijkman Institute for Molecular Biology, Jakarta, Indonesia and stored as frozen-infected erythrocytes stock in the parasitology laboratory of the Department of Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. Parasites were maintained by inoculating *P. berghei*-infected erythrocytes in mice, and serial passage of blood from infected mice to uninfected mice was performed. Donor mice infected with increased parasitemia by 20–30% were used to infect test mice. Donor mice were dissected and intracardially blood was collected in microtubes containing 1% ethylenediaminetetraacetic acid (EDTA) as anticoagulant to avoid variability in parasitaemia. The blood was then diluted with phosphate-buffered saline (PBS) so that each 0.2 mL of blood contained  $1 \times 10^7$  *P. berghei*-infected erythrocytes. Each mouse used in the experiment was then inoculated intraperitoneally with 0.2 mL of blood solution.

## 2.5. In Vivo Antimalarial Screening

**2.5.1. Evaluation of Suppressive Activity of Ethanolic Crude Extracts.** In vivo antimalarial activity testing of ethanolic leaf extracts from *S. androgynus* was performed against *P. berghei* according to the method described by Peters et al. [16] by randomly assigning 36 male mice to 6 groups (4 treatment groups and 2 control groups). Four treatment groups received 1, 10, 100, and 200 mg/kg body weight of extract, respectively. Both controls (negative and positive) received 0.5% sodium carbocymethyl cellulose (Na CMC) and dihydroartemisinin-piperazine (DHA-P, 20.8 + 166.4 mg/kg body weight), respectively. Na CMC, plant extracts, and DHA-P are administered once a day for 4 days orally. Treatment began after parasitemia in mice reached 1% ( $D_0$ ), using a hypodermic needle, and then continued for an additional 3 days (from  $D_1$  to  $D_3$ ). On day 5 ( $D_4$ ), thin blood strips were made from the tails of each mouse on a microscope slide. The blood was fixed with methanol, stained with 10% of Giemsa at pH 7.2 for 15 min and parasitemia was observed microscopically to determine the percent parasitaemia and the percent inhibition of parasites. Furthermore, data on the percent inhibition of parasites were used to calculate the  $ED_{50}$  value through probit analysis.

**2.5.2. Evaluation of Prophylactic Activity of Ethanolic Crude Extracts.** Prophylactic activity of ethanolic leaf extracts of *S. androgynus* and doxycycline was assessed using the method described by Peters [17] with modifications. Mice were randomly divided into eight groups, five each. Groups 1–3 were orally administered 100, 200, and 400 mg/kg body weight of extract, groups 4 and 5 were orally administered doxycycline (standard drug), and groups 6 and 7 were orally

administered 100 and 200 mg/kg body weight of extract, respectively. Group 8 was given 0.5% Na CMC as a negative control. Administration of plant extracts or standard drugs in groups 1–4 was performed for 4 consecutive days ( $D_0$ – $D_3$ ). On day 4 ( $D_3$ ), mice were inoculated with *P. berghei* (see Figure 1(a)). While administration of standard drugs, plant extracts, or Na CMC in groups 5–8 was carried out for 4 consecutive days ( $D_0$ – $D_3$ ), on the fourth day ( $D_3$ ) inoculation with *P. berghei* was carried out, then followed by administration of extracts, standard drugs, or Na CMC for 4 consecutive days ( $D_4$ – $D_7$ ; see Figure 1(b)) [18]. The percent parasitemia of each mouse from the entire group was assessed by blood elimination 72–168 h after parasite inoculation ( $D_6$ – $D_{10}$ ).

Mortality was monitored daily and the number of days from parasite inoculation to death was recorded for each mouse in the trial and control group during the experimental period. The mean survival time (MST) for each group is calculated using the equation.

$$MST = \frac{\text{Total survival time of all mice in the group (day)}}{\text{Total number of mice in the group}} \quad (1)$$

**2.6. Heme Detoxification Inhibitory Activity of Ethanolic Crude Extracts.** The heme detoxification inhibitory activity was assessed using the Basilio method [19] with slight modifications. Each 10 mg of ethanolic leaf extract of *S. androgynus* and chloroquine diphosphate (positive control) were dissolved in 1 mL as a test stock solution (100  $\mu$ L DMSO and 900  $\mu$ L aquadest), then a solution of the test material was prepared with different concentrations at 4, 2, 1, 0.5, 0.25, and 0.1 mg/mL. About 100  $\mu$ L of a 1 mM hematin solution in 0.2 M NaOH was introduced into the microtubes, then 50  $\mu$ L of the test material solution was added. The heme detoxification reaction was initiated by adding 50  $\mu$ L of glacial acetic acid (pH 2.6) and incubating at 37°C for 24 h to allow optimum detoxification. The microtubes were then centrifuged at 8000 rpm for 10 min, the filtrate was separated, and the remaining pellets were resuspended with 200  $\mu$ L of DMSO. Washing with DMSO was performed three times. The pellet obtained is dissolved in 200  $\mu$ L of 0.1 M NaOH, herein after referred to as hematin alkaline solution. About 100  $\mu$ L of hematin alkaline solution was distributed into 96-well microplates, and the absorbance was read using an ELISA reader at a wavelength of 405 nm. The preparation of the standard curve for hematin starts from the preparation of a 1 mM hematin solution in NaOH 0.2 M which is then made up of the concentrations of hematin standard solutions 500, 250, 125, 62.5, 31.25, and 15.625 mM.

**2.7. Statistical Analysis.** The results of the study were expressed as mean  $\pm$  standard error of mean (SEM). Data were analyzed using probit analysis. One-way analysis of variance (ANOVA) was used to determine statistical significance for the comparison of parasitemia and percent




FIGURE 1: Test scheme of antimalarial activity prophylaxis in vivo of ethanolic leaf extract of *S. androgynus* against *P. berghei*. (a) Administration of extracts 4 consecutive days before parasitic infection (groups 1–4). (b) Administration of the extract 4 days before the parasitic infection and continued 4 days after the parasitic infection (group 5–8).

TABLE 1: IC<sub>50</sub> value of the fraction from ethanolic leaf extract of *S. androgynus*.

Fractions	Yields (%)	IC <sub>50</sub> (μg/mL)
FI (1, 2, 3)	0.12	5044.719
FII (4, 5)	0.12	17.401
FIII (6, 7, 8, 9)	0.72	575175.293
FIV (10, 11, 12, 13)	1.08	1908.618
FV (14, 15, 16)	39.24	2.042
FVI (17, 18, 19)	51.30	2576.625
FVII (20, 21)	11.94	988.354

inhibition, among groups. The analysis was performed with a 95% confidence interval and a *p*-value of less than 0.05 was considered statistically significant.

### 3. Results

**3.1. In Vitro Antimalarial Activity of Fractions of *S. androgynus*.** The ethanolic leaf extract of *S. androgynus* was selected for activity-guided fractionation with VLC using a gradient of n-hexane-chloroform-ethanol 96% with a 10% concentration gradient. The fractionation results were then tested for in vitro antimalarial activity against *P. falciparum*. Based on the results of probit analysis, the IC<sub>50</sub> value for fraction FV was 2.042 μg/mL which is included in the active category as antimalarial. While fraction FII has an IC<sub>50</sub> value of >10 μg/mL and fractions FI, FIII, FIV, FVI, and FVII have an IC<sub>50</sub> value of >50 μg/mL, it is declared inactive antimalarial (see Table 1).

**3.2. Effect on Suppressive Activity of Ethanolic Crude Extracts of *S. androgynus*.** The extract shows a dose-dependent chemo-suppressive effect on parasitemia. This effect was statistically significant compared with the control (*p* < 0.05). The percent inhibition ranged from 33.73 to 66.11% (see Table 2). However, the effect of the extract was weak compared with the standard drug, DHA-P, with 100% inhibition (see Figure 2).

**3.3. Effect on Prophylactic Activity of Ethanolic Crude Extracts of *S. androgynus*.** The ethanolic leaf extract of *S. androgynus* showed prophylactic effects during prophylactic studies. This effect was statistically significantly different compared with the negative control (*p* < 0.05), but weak compared with the standard drug, doxycycline (see Table 3).

In addition, the percent inhibition between the administration groups 4 days before the infection (doses of 100 and 200 mg/kg body weight) resulted in a lower percentage of inhibition than the administration group continued after the infection (4 days before and after infection; doses of 100 and 200 mg/kg body weight), in the positive control group the administration before the infection and the administration continued after the infection resulted in a not too different percentage of inhibition which in the administration before the infection resulted in 69.75% inhibition (see Table 3) while in the administration group, continued after the infection resulted in 70.23% inhibition (see Table 4).

The group given the ethanolic leaf extract of *S. androgynus* survived longer than the negative control group (see Table 5).

**3.4. Heme Detoxification Inhibitory Activity of Ethanolic Crude Extracts of *S. androgynus*.** The ethanolic leaf extract of *S. androgynus* showed an effect in inhibiting the formation of β-hematin greater than 60% at a concentration of 2 mg/mL. The percentage of inhibition of β-hematin formation of ethanol extract and chloroquine diphosphate (positive control) shows that the higher the level of test material, the higher the percentage value of inhibition of β-hematin formation. The IC<sub>50</sub> values of ethanolic leaf extract of *S. androgynus* and chloroquine diphosphate were 0.479 and 0.569 mg/mL, respectively (see Table 6). According to Baelmans et al. [20], compounds that have an IC<sub>50</sub> value of inhibition of heme detoxification are smaller than the IC<sub>50</sub> value of chloroquine sulfate, which is 12 mg/mL, so the compound can be categorized as having activity in inhibiting heme detoxification. So that the ethanolic leaf extract of *S. androgynus* can be said to have heme detoxification inhibitory activity.

### 4. Discussion

*Sauropus androgynus* (L.) Merr., based on a genus and chemotaxonomic approach with *S. spatulifolius* proven to have antimalarial activity, was investigated for in vitro and in vivo antimalarial activities using standard models. The results of in vitro research revealed that *S. androgynus* leaf extract has antimalarial activity against chloroquine-sensitive strain (3D7) of *P. falciparum* with an IC<sub>50</sub> value of 96% ethanol extract of 1.88 μg/mL which is classified as very active [10]. Therefore, fractionation was performed on 96%



TABLE 2: In vivo suppressive activity of ethanolic leaf extract of *S. androgynus* (4-day suppressive test) against *P. berghei*-infected mice.

Drug/Extract	Dose (mg/kg/day)	% Parasitaemia		% Suppression	ED <sub>50</sub> (mg/kg bw)
		Day 0	Day 4		
Na CMC	—	0.92 ± 0.27	7.59 ± 0.97	—	—
Ethanolic leaf extract of <i>S. androgynus</i>	1	2.37 ± 0.64	6.79 ± 0.67	33.73	15.35
	10	1.95 ± 0.46	5.44 ± 0.38	47.67	
	100	0.76 ± 0.25	3.41 ± 0.51*	60.26	
	200	1.02 ± 0.34	3.28 ± 0.86*	66.11	
DHA-P	20.8 + 166.4	1.83 ± 0.83	0.00 ± 0.00*	100.00	—

Data are expressed as mean ± SEM;  $n = 6$ ; significance relative to the negative control, \*  $p < 0.05$ .




FIGURE 2: The effect of ethanolic leaf extract of *S. androgynus* on percent parasitaemia of *P. berghei*-infected mice on 4-day suppression test. Data are expressed as mean ± SEM;  $n = 6$ ; DHA-P = dihydroartemisinin-piperazine, EESA = ethanolic extract; numbers refer to doses in mg/kg/day.

TABLE 3: Percentages of inhibition of parasite growth in mice infected with *P. berghei* treated orally by doses of ethanolic leaf extract of *S. androgynus* for four consecutive days before infection ( $D_0$ – $D_3$ ).

Treatments	Dose (mg/kg/day)	% Parasitaemia		% Inhibition
		Day 6	Day 10	
Na CMC	—	0.26 ± 0.26	11.65 ± 1.50	—
Ethanolic leaf extract of <i>S. androgynus</i>	100	1.16 ± 0.39	6.61 ± 1.06	43.36 ± 18.38
	200	0.73 ± 0.27	5.70 ± 1.27	51.07 ± 21.77
	400	0.58 ± 0.38	3.76 ± 0.48*	67.74 ± 9.21
Doxycycline	13	0.20 ± 0.12	3.52 ± 0.44*	69.75 ± 8.51

Data are expressed as mean ± SEM;  $n = 5$ ; significance relative to the negative control, \*  $p < 0.05$ .

TABLE 4: Percentages of inhibition of parasite growth in mice infected with *P. berghei* treated orally by doses of ethanolic leaf extract of *S. androgynus* for 8 consecutive days before and after infection ( $D_0$ – $D_7$ ).

Treatments	Dose (mg/kg/day)	% Parasitaemia		% Inhibition
		Day 6	Day 10	
Na CMC	—	0.26 ± 0.26	11.65 ± 1.50	—
Ethanolic leaf extract of <i>S. androgynus</i>	100	1.00 ± 0.14	5.51 ± 0.13*	52.70 ± 1.93
	200	0.92 ± 0.15	4.04 ± 0.54*	65.30 ± 10.44
Doxycycline	13	0.00 ± 0.00	3.47 ± 0.36*	70.23 ± 6.88

Data are expressed as mean ± SEM;  $n = 5$ ; significance relative to the negative control, \*  $p < 0.05$ .

TABLE 5: The mean survival time (MST) of mice receiving various doses of ethanolic leaf extract of *S. androgynus* during prophylactic study.

Treatments	Dose (mg/kg/day)	MST	
		Day 10	Day 17
Na CMC	—	0.8	0
Ethanolic leaf extract of <i>S. androgynus</i> for 4 consecutive days ( $D_0$ – $D_3$ )	100	0.8	0.4
	200	0.8	0.4
	400	1	0.6
Doxycycline for 4 consecutive days ( $D_0$ – $D_3$ )	13	1	0.6
Ethanolic leaf extract of <i>S. androgynus</i> for 8 consecutive days ( $D_0$ – $D_7$ )	100	0.6	0.4
	200	1	0.8
Doxycycline for 8 consecutive days ( $D_0$ – $D_7$ )	13	1	0.8

TABLE 6: The  $IC_{50}$  values of ethanolic leaf extract of *S. androgynus*, chloroquine diphosphate, negative control based on the inhibition test of heme detoxification.

Drug/Extract	Concentration (mg/mL)	Concentration of hemozoin (mM)	% Inhibition	$IC_{50}$ (mg/mL)
DMSO	—	166.13 ± 11.48	—	—
Ethanolic leaf extract of <i>S. androgynus</i>	4	34.28 ± 1.90	79.36 ± 1.14	0.479
	2	50.29 ± 2.50	69.73 ± 1.50	
	1	67.59 ± 5.81	59.31 ± 3.50	
	0.5	81.57 ± 2.98	50.90 ± 1.80	
	0.25	95.31 ± 2.68	42.63 ± 1.62	
	0.1	121.48 ± 3.66	26.87 ± 2.20	
	Chloroquine diphosphate	4	44.80 ± 3.10	
2		56.42 ± 2.60	66.04 ± 1.56	
1		69.82 ± 4.23	57.97 ± 2.55	
0.5		84.69 ± 2.25	49.02 ± 1.35	
0.25		102.66 ± 3.09	38.21 ± 1.86	
0.1		118.17 ± 4.31	28.87 ± 2.60	

ethanol extract and in vitro antimalarial activity test on the results of fractionation against *P. falciparum*. The results showed that fraction FV of the seven fractions had the highest antimalarial activity ( $IC_{50} = 2.042 \mu\text{g/mL}$ ). However, the antimalarial activity of fraction FV was lower than that of 96% ethanol extract ( $IC_{50} = 1.88 \mu\text{g/mL}$ ). This result is in line with other findings by Ochieng et al. [21], where raw extracts from the aerial part of *Gardenia ternifolia* showed potent in vitro antimalarial activity against *P. falciparum* compared with its fractions and pure isolates. Crude extract of *G. ternifolia* roots also exhibited promising antimalarial activity against rodent malaria models, higher than its solvent fractions [22]. The reduction in antimalarial activity in the fraction compared with crude extract can be caused by the loss of synergistic activity between the compounds in the fraction [23]. Thus, 96% ethanol extract of *S. androgynus* leaves was selected for further evaluation in mice infected with *P. berghei* and its inhibitory activity against heme detoxification.

In vivo antimalarial activity can be classified as moderate, good, and very good if the extract displays, respectively, a percentage of parasite suppressors equal to or greater than 50% at doses of 500, 250, and 100 mg/kg body weight per day [24, 25]. Based on this classification, the ethanolic leaf extract of *S. androgynus* is considered to have shown good antimalarial activity, with dose-dependent inhibition against *P. berghei* infection in mice. In addition,

comparisons between suppressive and prophylactic models have revealed that extract at 200 mg/kg body weight is more effective in inhibiting *P. berghei* growth in mice with inhibitions of 66.11 and 51.07%, respectively. Lower inhibitory effects in prophylactic tests may arise from rapid metabolism that inactivates the active components of the extract responsible for antimalarial activity [26]. This effect can also be seen in two different prophylactic models where at 200 mg/kg body weight of extract provides greater inhibition when administered for 8 days compared with 4 days of administration. This finding is consistent with other studies in which the inhibitory effect of prophylactic tests was lower than the 4-day suppressive test [27, 28]. However, the overall ethanolic leaf extract of *S. androgynus* significantly reduced parasitemia in prophylactic and suppressive models of *P. berghei*-infected mice confirming the antimalarial potential of this extract. These findings were further supported by the results of the mean survival time of mice given extracts that were significantly prolonged compared with those from the control group, indicating a significant protective potential of the extracts.

In testing the mechanisms and targets of antimalarial active compounds in killing parasites, several methods were carried out, one of which was by testing the inhibitory activity of heme detoxification into hemozoin by *P. falciparum* which was the focus of antimalarial drug research [19]. Some properties of hemozoin that are

structurally similar to  $\beta$ -hematin make  $\beta$ -hematin an ideal product of the heme synthesis detoxification process [29]. The inhibitory effect of the heme detoxification process by ethanolic leaf extract of *S. androgynus* was studied in accordance with the results of the  $\beta$ -hematin barrier test in vitro. The test results revealed that the extract had the effect of inhibiting the formation of hemozoin which was characterized by the small value of  $\beta$ -hematin levels formed after incubating for 24 h (see Table 6). The inhibitory effect given was about 26–79% at a concentration of 0.1–4 mg/mL, and resulted in an  $IC_{50}$  value of 0.479 mg/mL. Frölich et al. [30] suggested that compounds with inhibition of hemozoin formation greater than 60% were declared to have good potential as hemozoin inhibitors. Thus, the ethanolic leaf extract of *S. androgynus* can be considered to have heme detoxification inhibitory activity. This inhibitory activity can result in death of parasites, which will minimize the pathogenesis of malaria, reduce the level of parasitemia, and reduce the accumulation of hemozoin in the liver and spleen of malaria patients which is one of the causes of hepatosplenomegaly.

Several secondary metabolites from plants such as alkaloids, flavonoids, and triterpenoids have been reported to have antimalarial activity [31–33]. Terpenoids have an important role as antimalarial agents by inhibiting the *Plasmodium* parasites' growth from ring forms to trophozoites, and can inhibit nutrient uptake by inhibiting the permeation pathway [34, 35]. Some antimalarial terpenoids isolated from other Euphorbiaceae plants have been reported, such as betulinic acid from *Uapaca nitida* Müll-Arg [36]; 8, 9-secokaurane diterpenes from *Croton kongensis* Gagnep [37]; geranylgeraniol from *Croton lobatus* L. [38]; poly-*O*-acylated jatrophone diterpenes from *Pedilanthus tithymaloides* (L.) Poit [39]; steenkrotin A from *Croton steenkampianus* Gerstner [40]; 2 $\alpha$ -hydroxyjatrophone from *Jatropha integerrima* Jacq [41]; jatrophone diterpenes from *Jatropha isabelli* Müll. Arg [42]; 6-hydroxy neomacrolactone from *Neoboutonia macrocalyx* L. [43]; samvisterin from *Uapaca paludosa* [44]; euphorbesulin G from *Euphorbia esula* L. [45]; and many more. All these compounds exhibited good antimalarial activity with  $IC_{50}$  of  $\leq 5 \mu\text{g/mL}$  against various *P. falciparum* strains. This current study did not isolate the pure compound of *S. androgynus*, however, it was reported to contain various compounds. A number of 20 compounds were identified and the major compounds in the leaves extract that were detected in the present study were L-(+)-ascorbic acid 2, 6-dihexadecanoate (27.82%). They are followed by hexadecanoic acid, ethyl ester (17.85%); ethyl 9, 12, 15-octadecatrienoate (16.32%); ethyl (9Z, 12Z)-9, 12-octadecadienoate (9.40%); 9, 12, 15-octadecatrienoic acid, (Z, Z, Z)-(7.47%); 2, 6, 10-trimethyl, 14-ethylene-14-pentadecane (4.27%); phytol, acetate (4.12%); 9, 12-octadecadienoic acid (Z, Z)-(3.76%); 3, 7, 11, 15-tetramethyl-2-hexadecen-1-ol (2.04%); 2, 4-imidazolidinedione, 1-((5-nitro-2-furanyl) methylene)amino)-(1.94%); octadecanoic acid, ethyl ester (1.39%); 7-octadecyne, 2-methyl-(0.94%); heptadecanoic acid, ethyl ester (0.59%); cyclopentasiloxane, decamethyl (0, 53%); cis-vaccenic acid (0.35%); 2, 6, 8-trimethyl-bicyclo [4.2.0]oct-2-ene-1, 8-diol (0.28%); 2-

pentadecanone, 6, 10, 14-trimethyl- (0.27%); 9-octadecenoic acid (Z)- (0.26%); cyclohexasiloxane, dodecamethyl- (0.26%); and 1, 2-benzenedicarboxylic acid, 2-ethoxy-2-oxoethyl methyl ester (0.14%) [46]. Those compounds may be the responsible for antimalarial activities. Mahardiani et al. [10] also have been reported that ethanolic leaf extract of *S. androgynus* was found in this study to contain terpenoids. The terpenoid content of *S. androgynus* leaves includes sesquiterpenoids and triterpenoids [9]. Thus, we suspect that the terpenoid content contained in this plant extract may have contributed to the antimalarial activity of this extract and therefore explains the mechanism of the antimalarial effect of the extract.

## 5. Conclusions

In general, the results of the current study indicate that 96% ethanol extracts and its fractions of *S. androgynus* leaves have potent antimalarial activity. The results also confirm that the plant has a suppressive and prophylactic effect on the growth of parasites. Ethanolic extract was also found to be active in inhibiting the heme detoxification process. This adds to the supporting evidence that the plant *S. androgynus* has the potential to be antimalarial from natural sources.

## Data Availability

The data used to support the findings of this study can be obtained from the corresponding author upon request.

## Conflicts of Interest

The authors declare that there are no conflicts of interest.

## Acknowledgments

This research was supported by the Directorate of Research and Community Service, Ministry of Research and Technology/National Research and Innovation Agency, Republic of Indonesia, through Mandate Research Grant 367/UN3.14/PT/2020.

## References

- [1] WHO, *Guidelines for the Treatment of Malaria*, WHO Press, Geneva, Switzerland, 3rd edition, 2015.
- [2] N. J. White, "Anaemia and malaria," *Malaria Journal*, vol. 17, no. 1, 2018.
- [3] A. T. Lukwa, R. Mawoyo, K. N. Zablon, A. Siya, and O. Alaba, "Effect of malaria on productivity in a workplace: the case of a banana plantation in Zimbabwe," *Malaria Journal*, vol. 18, no. 1, p. 390, 2019.
- [4] WHO, *World Malaria Report 2021*, WHO Press, Geneva, Switzerland, 2021.
- [5] Ministry of Health Republic of Indonesia, *National Action Plan for Acceleration of Malaria Elimination 2020-2024*, Ministry of Health Republic of Indonesia, Jakarta, Indonesia, 2020.
- [6] WHO, *Artemisinin Resistance and Artemisinin-Based Combination Therapy Efficacy: Status Report*, WHO Press, Geneva, Switzerland, 2018.

- [7] Y. S. Zou, K. Foubert, E. Tuentner et al., "Antiplasmodial and cytotoxic activities of *Striga asiatica* and *Sauropus spatulifolius* extracts, and their isolated constituents," *Phytochemistry Letters*, vol. 6, no. 1, pp. 53–58, 2013.
- [8] H. E. Khoo, A. Azlan, and A. Ismaila, "Sauropus androgynus leaves for health benefits: hype and the science," *The Natural Products Journal*, vol. 5, no. 2, pp. 115–123, 2015.
- [9] S. V. Selvi and B. Anusha, "Phytochemical analysis and GC-MS profiling in the leaves of *Sauropus androgynus* (L.) Merr," *International Journal of Drug Development and Research*, vol. 4, no. 1, pp. 162–167, 2012.
- [10] A. Mahardiani, S. Suciati, and W. Ekasari, "In vitro anti-malarial and cytotoxic activities of *Sauropus androgynus* leaves extracts," *Tropical Journal of Natural Product Research*, vol. 4, no. 9, pp. 558–562, 2020.
- [11] W. Berthi, A. González, A. Rios, S. Blair, A. Cogollo, and A. Pabon, "Anti-plasmodial effect of plant extracts from *Picrolemma huberi* and *Picramnia latifolia*," *Malaria Journal*, vol. 17, no. 1, p. 151, 2018.
- [12] International Council for Harmonisation, *ICH Q3C (R8) Impurities: Guideline for Residual Solvents Q3C (R8)*, International Council for Harmonisation, Geneva, Switzerland, 2021.
- [13] Q. D. Do, A. E. Angkawijaya, P. L. Tran-Nguyen et al., "Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*," *Journal of Food and Drug Analysis*, vol. 22, no. 3, pp. 296–302, 2014.
- [14] W. Trager and J. B. Jensen, "Human malaria parasites in continuous culture," *The Journal of Parasitology*, vol. 91, no. 3, pp. 484–486, 2005.
- [15] National Academy of Sciences, *Guide for the Care and Use of Laboratory Animals*, Institute for Laboratory Animal Research, Washington, DC, USA, 2011.
- [16] W. Peters, J. H. Portus, and B. L. Robinson, "The chemotherapy of rodent malaria, XXII. The value of drug-resistant strains of *P. berghei* in screening for blood schizontocidal activity," *Annals of Tropical Medicine and Parasitology*, vol. 69, no. 2, pp. 155–171, 1975.
- [17] W. Peters, "Drug resistance in *Plasmodium berghei* vincke and lips, 1948. I. Chloroquine resistance," *Experimental Parasitology*, vol. 17, no. 1, pp. 80–89, 1965.
- [18] P. N. Soh, B. Witkowski, D. Olagnier et al., "In vitro and in vivo properties of ellagic acid in malaria treatment," *Antimicrobial Agents and Chemotherapy*, vol. 53, no. 3, pp. 1100–1106, 2009.
- [19] N. Basilico, E. Pagani, and D. Monti, "A microtitre-based method for measuring the haem polymerization inhibitory activity (HPIA) of antimalarial drugs," *Journal of Antimicrobial Chemotherapy*, vol. 42, no. 1, pp. 55–60, 1998.
- [20] R. Baelmans, E. Deharo, V. Muñoz, M. Sauvain, and H. Ginsburg, "Experimental conditions for testing the inhibitory activity of chloroquine on the formation of  $\beta$ -hematin," *Experimental Parasitology*, vol. 96, no. 4, pp. 243–248, 2000.
- [21] C. O. Ochieng, J. Ogweni Mid, and P. Okinda Owu, "Antiplasmodial and larvicidal effects of surface exudates of *Gardenia ternifolia* aerial parts," *Research Journal of Pharmacology*, vol. 4, no. 2, pp. 45–50, 2010.
- [22] D. Nureye, S. Assefa, and T. Nedi, "In vivo antimalarial activity of the 80% methanolic root bark extract and solvent fractions of *Gardenia ternifolia* Schumacher and Thonn (Rubiaceae) against plasmodium Berghei," *Evidence-Based Complementary And Alternative Medicine*, vol. 2018, Article ID 9217853, 10 pages, 2018.
- [23] P. Rasoanaivo, E. Deharo, and S. Ratsimamanga-Urverg, "Guidelines for the nonclinical evaluation of the efficacy of traditional antimalarials," in *Traditional Medicinal Plants and Malaria*, CRC Press, Boca Raton, FL, USA, 2004.
- [24] V. Muñoz, M. Sauvain, G. Bourdy et al., "A search for natural bioactive compounds in Bolivia through a multidisciplinary approach," *Journal of Ethnopharmacology*, vol. 69, no. 2, pp. 127–137, 2000.
- [25] E. Deharo, G. Bourdy, C. Quenevo, V. Munoz, G. Ruiz, and M. Sauvain, "A search for natural bioactive compounds in Bolivia through a multidisciplinary approach. Part V. Evaluation of the antimalarial activity of plants used by the Tacana Indians," *Journal of Ethnopharmacology*, vol. 77, no. 1, pp. 91–98, 2001.
- [26] O. A. Salawu, A. Y. Tijani, and H. Babayi, "Anti-malarial activity of ethanolic stem bark extract of *Faidherbia albida* (Del) a. Chev (Mimosoidae) in mice," *Archives of Applied Science Research*, vol. 2, no. 5, pp. 261–268, 2010.
- [27] E. G. Unekwujo, O. James, and A. R. Olubunmi, "Suppressive, curative and prophylactic potentials of *Morinda lucida* (Benth) against erythrocytic stage of mice infective chloroquine sensitive *Plasmodium berghei* NK-65," *British Journal of Applied Science and Technology*, vol. 1, no. 3, pp. 131–140, 2011.
- [28] D. Misganaw, E. Engidawork, and T. Nedi, "Evaluation of the anti-malarial activity of crude extract and solvent fractions of the leaves of *Olea europaea* (Oleaceae) in mice," *BMC Complementary and Alternative Medicine*, vol. 19, no. 1, p. 171, 2019.
- [29] B. L. Tekwani and L. A. Walker, "Targeting the hemozoin synthesis pathway for new antimalarial drug discovery: technologies for in vitro -hematin formation assay," *Combinatorial Chemistry and High Throughput Screening*, vol. 8, no. 1, pp. 63–79, 2005.
- [30] S. Frölich, C. Schubert, U. Bienzle, and K. Jenett-Siems, "In vitro antiplasmodial activity of prenylated chalcone derivatives of hops (*Humulus lupulus*) and their interaction with haemin," *Journal of Antimicrobial Chemotherapy*, vol. 55, no. 6, pp. 883–887, 2005.
- [31] G. C. Kirby, M. J. O'Neill, J. D. Phillipson, and D. C. Warhurst, "In vitro studies on the mode of action of quassinoids with activity against chloroquine-resistant *Plasmodium falciparum*," *Biochemical Pharmacology*, vol. 38, no. 24, pp. 4367–4374, 1989.
- [32] J. D. Phillipson and C. W. Wright, "Antiprotozoal compounds from plants sources," *Planta Medica*, vol. 57, no. 7, pp. 553–559, 1991.
- [33] S. B. Christensen and A. Kharazmi, "Antimalarial natural products. Isolation, characterization and biological properties," in *Bioactive Compounds from Natural Sources: Isolation, Characterization and Biological Properties*, Taylor and Francis, London, UK, 2001.
- [34] H. B. Gabriel, R. A. C. Sussmann, and E. A. Kimura, "Terpenes as potential antimalarial drugs," in *Terpenes and Terpenoids*, IntechOpen, London, UK, 2018.
- [35] M. Muhaimin, Y. Yusnaidar, W. Syahri et al., "Antiplasmodial activity of ethanolic extract of *Macaranga gigantea* leaf and its major constituent," *Pharmacognosy Journal*, vol. 11, no. 6, pp. 1181–1188, 2019.
- [36] J. C. P. Steele, D. C. Warhurst, G. C. Kirby, and M. S. J. Simmonds, "In vitro and in vivo evaluation of betulinic

- acid as an antimalarial,” *Phytotherapy Research*, vol. 13, no. 2, pp. 115–119, 1999.
- [37] J. Thongtan, P. Kittakoop, N. Ruangrunsi, J. Saenboonrueng, and Y. Thebtaranonth, “New antimycobacterial and antimalarial 8, 9-secokaurane diterpenes from *Croton kongensis*,” *Journal of Natural Products*, vol. 66, no. 6, pp. 868–870, 2003.
- [38] B. Attioua, B. Weniger, and P. Chabert, “Antiplasmodial activity of constituents isolated from *Croton lobatus*,” *Pharmaceutical Biology*, vol. 45, no. 4, pp. 263–266, 2007.
- [39] W. Mongkolvisut and S. Sutthivaiyakit, “Antimalarial and antituberculous poly-O-acylated jatrophone diterpenoids from *Pedilanthus tithymaloides*,” *Journal of Natural Products*, vol. 70, no. 9, pp. 1434–1438, 2007.
- [40] A. M. Adelekan, E. A. Prozesky, A. A. Hussein et al., “Bioactive diterpenes and other constituents of *Croton steenkampianus*,” *Journal of Natural Products*, vol. 71, no. 11, pp. 1919–1922, 2008.
- [41] S. Sutthivaiyakit, W. Mongkolvisut, S. Prabpai, and P. Kongsaree, “Diterpenes, sesquiterpenes, and a Sesquiterpene–Coumarin conjugate from *Jatropha integerrima*,” *Journal of Natural Products*, vol. 72, no. 11, pp. 2024–2027, 2009.
- [42] V. Hadi, M. Hotard, T. Ling et al., “Evaluation of *Jatropha isabelli* natural products and their synthetic analogs as potential antimalarial therapeutic agents,” *European Journal of Medicinal Chemistry*, vol. 65, pp. 376–380, 2013.
- [43] J. Namukobe, B. T. Kiremire, R. Byamukama et al., “Cycloartane triterpenes from the leaves of *Neoboutonia macrocalyx* L.,” *Phytochemistry*, vol. 102, pp. 189–196, 2014.
- [44] J. T. Banzouzi, P. N. Soh, S. Ramos et al., “Samvisterin, a new natural antiplasmodial betulin derivative from *Uapaca paludosa* (Euphorbiaceae),” *Journal of Ethnopharmacology*, vol. 173, pp. 100–104, 2015.
- [45] B. Zhou, Y. Wu, S. Dalal, M. B. Cassera, and J. M. Yue, “Euphorbesulins A-P, structurally diverse diterpenoids from *Euphorbia esula*,” *Journal of Natural Products*, vol. 79, no. 8, pp. 1952–1961, 2016.
- [46] S. C. Mohan, R. Bose, M. S. Kumar, and A. Manivel, “Chemical constituents of *Sauropus androgynus* and evaluation of its antioxidant activity,” *Research Journal of Phytochemistry*, vol. 12, no. 1, pp. 7–13, 2018.





**KOMISI ETIK PENELITIAN  
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA  
*Animal Care and Use Committee (ACUC)***

**KETERANGAN KELAIKAN ETIK  
“ ETHICAL CLEARANCE ”**

**No : 2.KE.181.10.2018**

**KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE)  
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA,  
TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG  
DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :**

**PENELITIAN BERJUDUL** : Efikasi Ekstrak Etanol Daun *Cassia spectabilis* Sebagai Obat Antimalaria dari Bahan Alami

**PENELITI UTAMA** : Wiwied Ekasari

**UNIT/LEMBAGA/TEMPAT PENELITIAN** : Fakultas Farmasi Universitas Airlangga

**DINYATAKAN** : LAIK ETIK

Surabaya, 30 Oktober 2018

Ketua,

Dr. Nusdianto Triakoso, M.P.,Drh.  
NIP. 196805051997021001

Mengetahui,  
Dekan FKH-Unair,

Prof. Dr. Rudi Sianto, M.Kes.,Drh.  
NIP. 195601051986011001



# Bukti – Subject Area and Category, Quartile dan SJR

also developed by scimago

SCIMAGO INSTITUTIONS RANKINGS

SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name



Home Journal Rankings Country Rankings Viz Tools Help About Us

## Scientifica

### COUNTRY

Egypt



Universities and research institutions in Egypt

### SUBJECT AREA AND CATEGORY

Agricultural and Biological Sciences  
└ Agricultural and Biological Sciences (miscellaneous)  
  
Environmental Science  
└ Environmental Science (miscellaneous)

### PUBLISHER

Hindawi Limited

### H-INDEX

26

### PUBLICATION TYPE

Journals

### ISSN

2090908X

### COVERAGE


2016-2021

### INFORMATION

[Homepage](#)

[How to publish in this journal](#)

### Quartiles



### SJR

In a journal, it expresses how central to the global scientific discussion an average article of the journal is.

Year	SJR
2017	0.291
2018	0.424
2019	0.498
2020	0.474
2021	0.460

### Total Documents



# Bukti – Scopus Coverage, Publisher dan ISSN



Scopus Preview

Author Search

Sources



Create account

Sign in

## Source details

Feedback > Compare sources >

### Scientifica

Open Access ⓘ

Scopus coverage years: from 2016 to Present

Publisher: Hindawi

E-ISSN: 2090-908X

Subject area: Agricultural and Biological Sciences: General Agricultural and Biological Sciences Environmental Science: General Environmental Science

Source type: Journal

View all documents >

Set document alert

Save to source list

Source Homepage

CiteScore 2021

3.3 ⓘ

SJR 2021

0.460 ⓘ

SNIP 2021

1.193 ⓘ

CiteScore CiteScore rank & trend Scopus content coverage

#### Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021

$$3.3 = \frac{509 \text{ Citations 2018 - 2021}}{154 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$4.2 = \frac{621 \text{ Citations to date}}{147 \text{ Documents to date}}$$

Last updated on 05 August, 2022 • Updated monthly

#### CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Agricultural and Biological Sciences	#52/211	75th
General Agricultural and Biological Sciences		
Environmental Science	#89/228	61st
General Environmental Science		