

# Antioxidant, Antimicrobial, and Antiplasmodial Activities of Sonchus arvensis L. Leaf Ethyl Acetate Fractions

*by* Wiwied Ekasari

---

**Submission date:** 15-Feb-2023 04:29PM (UTC+0800)

**Submission ID:** 2014700769

**File name:** Phcog J\_naskah.pdf (897.33K)

**Word count:** 5591

**Character count:** 28593

# Antioxidant, Antimicrobial, and Antiplasmodial Activities of *Sonchus arvensis* L. Leaf Ethyl Acetate Fractions

Dwi Kusuma Wahyuni<sup>1,2\*</sup>, Anindya Nariswari<sup>1</sup>, Agus Supriyanto<sup>1</sup>, Hery Purnobasuki<sup>1</sup>, Hunsu Punnapayak<sup>2</sup>, Wichanee Bankeeree<sup>2</sup>, Sehanat Prasongsuk<sup>1,2,\*</sup>, Wiwied Ekasari<sup>3,\*</sup>

Dwi Kusuma Wahyuni<sup>1,2\*</sup>,  
Anindya Nariswari<sup>1</sup>, Agus  
Supriyanto<sup>1</sup>, Hery Purnobasuki<sup>1</sup>,  
Hunsu Punnapayak<sup>2</sup>, Wichanee  
Bankeeree<sup>2</sup>, Sehanat  
Prasongsuk<sup>1,2,\*</sup>, Wiwied Ekasari<sup>3,\*</sup>

<sup>1</sup>Department of Biology, Faculty of Science and  
Technology, Airlangga University Surabaya,  
East Java, 60115, INDONESIA.

<sup>2</sup>Plant Biomass Utilization Research Unit,  
Department of Botany, Faculty of Science,  
Chulalongkorn University, Bangkok,  
THAILAND.

<sup>3</sup>Department of Pharmaceutical Science,  
Faculty of Pharmacy, Airlangga University  
Surabaya, East Java, 60115, INDONESIA.

## Correspondence

Dwi Kusuma Wahyuni

Department of Biology, Faculty of Science  
and Technology, Airlangga University  
Surabaya, East Java, 60115, INDONESIA.

E-mail: dwi-k-w@fst.unair.ac.id

Sehanat Prasongsuk

Plant Biomass Utilization Research Unit,  
Department of Botany, Faculty of Science,  
Chulalongkorn University, Bangkok,  
THAILAND.

E-mail: sehanat.p@chula.ac.th

Wiwied Ekasari

Department of Pharmaceutical Science,  
Faculty of Pharmacy, Airlangga University  
Surabaya, East Java, 60115, INDONESIA.

E-mail: wiwied-e@ff.unair.ac.id

## History

- Submission Date: 24-10-2022;
- Review completed: 02-12-2022;
- Accepted Date: 05-12-2022.

DOI : 10.5530/pj.2022.14.202

## Article Available online

http://www.phcogj.com/v14/i6

## Copyright

© 2022 Phcogj.Com. This is an open-  
access article distributed under the terms  
of the Creative Commons Attribution 4.0  
International license.



## ABSTRACT

Infection is one of the health problems and a disease that mainly causes death. Malaria is a parasitic infection that is transmitted through the *Anopheles* sp. The female then causes infection and besides malaria, other contaminants that caused infection are bacteria such as *Escherichia coli* and *Staphylococcus aureus*. This study aims to determine the antioxidant, antimicrobial, and antiplasmodial activity of *Sonchus arvensis* L. ethyl acetate fractions. *In vitro* antiplasmodial activity was carried out by Rieckman methods against *Plasmodium falciparum* strain 3D7. *In vitro* antioxidant activity was conducted by Prieto method against (2,2-diphenyl-1-picrylhydrazyl) (DPPH). Then antimicrobial activity was performed using well diffusion method against *Escherichia coli* and *Staphylococcus aureus*. Maceration of *S. arvensis* L. dried leaves used *n*-hexane and ethyl acetate successively. Then the ethyl acetate extract was fractionated by vacuum column chromatography, using *n*-hexane and ethyl acetate as mobile phases. There are five fraction groups based on thin-layer chromatography (TLC) analysis. The IC<sub>50</sub> of antioxidant and antiplasmodial activity showed that fraction IV was the lowest value and categorized as active for antioxidant (IC<sub>50</sub>=22.56 µg/mL), for antiplasmodial (IC<sub>50</sub>=12.07 µg/mL). Fraction IV also had antimicrobial activity, with diameter of inhibition zone (DIZ) of 19.22 mm against *Escherichia coli* and 17.167 mm against *Staphylococcus aureus*.

**Key words:** Biological activities, Malaria, *Plasmodium falciparum*, *Sonchus arvensis* L., *Staphylococcus aureus*, *Escherichia coli*.

## INTRODUCTION

Infection is one of the serious global health problems.<sup>1</sup> In Indonesia, as many as 28.1% of the main causes of death in Indonesia are caused by infection.<sup>2</sup> Infections were caused by various microorganisms such as bacteria, viruses, fungi, and protozoa.<sup>3</sup> During global covid pandemic, the number of malaria cases increases by 205 million.<sup>4</sup> Besides malaria, other contaminants that cause infectious diseases are bacteria. *Escherichia coli* and *Staphylococcus aureus* are bacteria that cause diarrheal disease. Malaria and diarrhea are the ten most popular infectious diseases in Indonesia.<sup>2</sup>

Recently, the problem of antibiotic and antimalarial resistance has emerged. The World Health Organization (WHO) has issued a list of priority pathogens to increase research efforts in the search for new antibiotics and antimalarials, to overcome the problem of resistance.<sup>4</sup> One source of searching for new drugs is to use medicinal plants derived from nature. Some studies showed that the active components in medicinal plants have antimicrobial effects that differ in their mechanism of action from antibiotics that have been around so far. This shows that medicinal plants have the potential to overcome the problem of resistance.<sup>5</sup>

One of the medicinal plants that are often used in Indonesia is a plant from the Asteraceae family, a plant from the Asteraceae family which is widely known by the public as a plant that is easy to grow and has been widely used in medicine, *S. arvensis* L.<sup>6,7</sup> Based on the phytochemical screening results from the research of Wahyuni *et al* (2020b) it could be seen that *S. arvensis* L. leaves contain

various bioactive components such as flavonoids, terpenoids, and polyphenols that can act as antioxidants, hepatoprotective, diuretic, and have the potential as antimalarial agents.<sup>8,9</sup> *S. arvensis* L. also has antimicrobial activity. This is based on the chemical content of the *S. arvensis* L. flavonoids can affect the polarity of the lipids that make up bacterial DNA. Alkaloids can damage the peptidoglycan constituents of bacterial cells.<sup>10</sup>

*S. arvensis* L. has potential as an antimalarial agent based on previous research conducted by Wahyuni *et al.* (2020) that the methanol extract from *S. arvensis* L. callus treated with 1-5% sucrose was tested for antimalarial *in vitro* using the Trager and Jensen methods and tested against *Plasmodium falciparum* strain 3D7 has an IC<sub>50</sub> value of 0.343 g/ml.<sup>8</sup> According to research conducted by Tapan (2016), The (High-Performance Liquid Chromatography (HPLC) on 80% ethanol extract of *S. arvensis* L. leaves produced ascorbic acid, gallic acid, catechin, and kaempferol compounds which are believed to act as antibacterial.<sup>11</sup> Based on research by Xia *et al.* (2011), *S. arvensis* L. methanol extract had a total phenolic value of 417.3 ± 38.3 mg/g GAE dry weight and the antibacterial test had a minimum inhibitory concentration (MIC) value of 9.5-16 mg/ml.<sup>12</sup> Based on research conducted by Kanaani and Sani (2015), *S. arvensis* L. methanol extract has a Minimum Inhibitory Concentration (MIC) value of 50-100 mg/ml which were tested on *Bacillus cereus*, *Staphylococcus aureus*, *Salmonella enterica*, and *Escherichia coli*.<sup>13</sup> Another study also reported that the *n*-hexane extract of *S. arvensis* L. leaves using column chromatography and separated by several solvent fractions was able to inhibit the growth of

**Cite this article:** Wahyuni DK, Nariswari A, Supriyanto A, Purnobasuki H, Punnapayak H, Bankeeree W, et al. Antioxidant, Antimicrobial, and Antiplasmodial Activities of *Sonchus arvensis* L. Leaf Ethyl Acetate Fractions. Pharmacogn J. 2022;14(6)Suppl: 993-998.

*Staphylococcus aureus* bacteria by 9 mm, and *Escherichia coli* by 8 mm with a concentration of 1000 ppm.<sup>14</sup>

Although many studies on the bioactivity of *S. arvensis* L have been carried out, there have been no reports on the antiplasmodial activity of the ethyl acetate fraction of *S. arvensis* L. to isolate the target compound. Therefore, this study aims to determine the antioxidant, antimicrobial, and antiplasmodial activities of the ethyl acetate fraction of *S. arvensis* L.. The results of this study are expected to provide basic knowledge as a guideline for isolating compounds that have the potential as antioxidants, antimicrobials, and antiplasmodial from *S. arvensis* L..

## MATERIALS AND METHODS

### Sample preparation

*S. arvensis* L. were obtained from the Taman Husada Graha Famili (Medicinal Plant Garden), Surabaya, Indonesia. The plant was identified by the Purwodadi Botanical Gardens Herbarium, East Java, Indonesia. The voucher specimen was collected in Plant Systematic Laboratory, Department of Biology, Faculty of Science and Technology, Universitas Airlangga (Voucher No. SA.0010292021). The leaves from vegetative stage were collected, then dried and mashed to form a powder, and then weighed of 1 kg to continue the extraction process.

### Extraction

One hundred grams of *S. arvensis* L. leaf powder was extracted by maceration using 1000 mL of *n*-hexane until all the powder was submerged and stirred, then covered and stored for three days. Stirring was done approximately three times a day. Furthermore, filtering was carried out so that only the powder remains. Then the powder was soaked in 1000 mL of ethyl acetate solution for three days. Then the filtrate was concentrated with a vacuum rotary evaporator to obtain a thick extract and then weighed.

### Fractionation

Five grams of *S. arvensis* L. ethyl acetate extract was fractionated by vacuum chromatography. The 75 grams of silica gel 60G was added was used as stationary phase. The mobile phase use *n*-hexane:ethyl acetate in 16 combinations (100, 95:5, 90:10, 85:15, 80:20, 75:25, 70:30, 65:35, 60:40, 50:50, 55:45, 40:60, 30:70, 20:80, 10:90, 100). The dried extract was put into a sintered glass and filter paper was placed on top of the extract. The mobile phase was poured into a sintered glass and withdrawn with a suction pump until the liquid did not drip back. Then eluted starting from low polarity then the polarity was increased slowly and the column was sucked dry at each collection of fractions.<sup>15</sup>

### Thin Layer Chromatography analysis

GF254 silica plates were used. One hundred milligram fractions were dissolved in 100 µL ethyl acetate. Each fraction was spotted on silica plate at a distance of ± 1 cm from the bottom (± 1.5 cm) with a capillary tube, then dried and eluted with each mobile phase of the compound group. The terpenoid compound group was made into a mobile phase consisting of *n*-hexane: ethyl acetate (4:1). After the plate was eluted to the mark, taken, and allowed to dry, it was sprayed with sulfuric acid *p*-anisaldehyde reagent, then heated for 5 minutes at 105°C. The presence of terpenoids was indicated by the formation of a blue-violet or red-violet spot.<sup>9</sup>

### Antioxidant activity assay

Antioxidant activity was evaluated by a modified method of 2,2-diphenyl-1-picryl-hydrazyl-hydrate (DPPH) assay.<sup>16</sup> In brief, 100 µl of methanolic DPPH reagent (0.2 mM) was mixed with 100 µl of each sample in methanol at different concentrations (3.15, 6.25, 10.00, 12.50, 15.00, 25.00, 35.00, 50.00, 75.00, 100.00, 150.00, and 200.00 µg/

ml) or methanol as the control. The mixtures were incubated for 30 min in the dark at room temperature and the absorbance was measured at 517 nm. The inhibition of DPPH was calculated using the following equation:

$$\text{DPPH inhibition (\%)} = (A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}} \times 100\% \quad (1)$$

where  $A_{\text{sample}}$  is the absorbance of the sample and  $A_{\text{control}}$  is the absorbance of the DPPH reagent at the wavelength of 512 nm. The percentage of inhibition results at different concentrations was then plotted and regressed linearly to obtain the  $IC_{50}$  values of DPPH.

### In vitro antiplasmodial activity assay

Cultures of *Plasmodium falciparum* strain 3D7 were cultivated using the Trager and Jensen method (1976), as adapted by Ekasari et al. (2009),<sup>17</sup> in Roswell Park Memorial Institute 1640 medium (Gibco, Carlsbad, CA, USA) supplemented with human O-type red blood cells, 5% hematocrit, 22.3 mM HEPES buffer (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; Sigma-Aldrich Corporation, St. Louis, MO, USA), 50 µg/mL hypoxanthine, 2 mg/mL sodium bicarbonate, and 10% human O<sup>+</sup> plasma.<sup>15</sup> The sample was dissolved in 10% dimethyl sulfoxide (DMSO) in various concentrations (0.01, 0.1, 1, 10, and 100 µL). Chloroquine diphosphate was used as a positive control and DMSO as the negative control. An antiplasmodial assay was conducted using a 24-well microplate and incubation at 37°C for 48 h. The incubated materials were then collected, thinly smeared on a glass slide, fixed with methanol, and stained with Giemsa to assess the number of parasites under a microscope as compared with the negative control to determine the  $IC_{50}$  value to achieve inhibition of parasitic growth. The data was used to calculate the percent growth and percent inhibition using the following formulas:

$$\% \text{ Growth} = \% \text{ Parasitemia} - D0 \quad (2)$$

$$\text{Percent inhibition} = 100\% - [(Xu/Xk) \times 100\%] \quad (3)$$

Where D0 is the percentage of growth at the 0-hour, whereas Xu and Xk are the percentages of growth in the test solution and negative control, respectively. Based on the percent inhibition data, statistical analysis was carried out using Probit analysis of the SPSS version 20 program to determine the  $IC_{50}$  value or the concentration of the test material that inhibits parasitic growth by 50%.

### Antimicrobial activity assay

Well diffusion method was used in this study to determine the antimicrobial activity against *Escherichia coli* ATCC 25922 and *Staphylococcus aureus* ATCC 25923. The 10 mg of fractions were weighed and dissolved in dimethyl sulfoxide (DMSO). The extract was added with 2 mL of 20% DMSO gradually and 8 mL of sterile distilled water and then homogenized. Chloramphenicol (1000 mg/L) was used as positive control and DMSO (20%) as negative control. Next, mix 1 mL of pure culture into a test tube containing 9 mL of nutrient broth media (peptone 10 mg, yeast extract 2 g, beef extract 1 g, sodium chloride 5 g, pH 6.8 ± 0.2 at 25°C. The mixture was homogenized by vortex and the turbidity was standardized to 0.5 nm Mc Farland using a spectrophotometer with a wavelength of 625 nm. Bacteria that have been standardized were taken as much as 1 mL using a micropipette and put into a petri dish. The warm nutrient agar media (± 60°C, 25 mL) was poured into a plate containing 1 mL of bacteria. Make a well in the agar using a tool that has been sterilized with 70% alcohol and heated on fire. Putting the extract into the wells with a concentration of 25% and 50%, respectively, with positive control of chloramphenicol and negative control of DMSO. Pour plate cultures were incubated at 37°C for 24 hours. Inhibition was revealed based on the diameter of the clear zone formed around the well (diameter of inhibition zone/DIZ). The measurement used a calliper using quantitative observations and expressed in millimetres.<sup>18</sup>

The value of DIZ was used to calculate the percentage of inhibition with the formula:

$$\text{Percentage of inhibition (\%)} = (\text{DIZ sample/DIZ control}) \times 100\% \quad (4).$$

## RESULT AND DISCUSSION

### The yield of *S. arvensis* L. ethyl acetate fractions

The yield of *S. arvensis* L. *n*-hexane fraction groups were 11.14%, 12.88%, 20.18%, 24.28%, and 16.44% for fractions no I, II, III, IV, and V respectively. Yield is the ratio of the dry weight of the extract to the amount of raw material. The yield value is related to the amount of bioactive content contained. The higher yield means the higher content of substances that are attracted to raw material. The yield of extract depended on the solvent and the methods of extraction.<sup>19,21</sup> In this study, group fractions IV were the highest yield. They were isolated by the *n*-hexane: ethyl acetate (40:60, 30:70, 20:80). Some studies showed that the more polar solvent resulted in higher yield of extract.<sup>19,20</sup>

### Chromatogram profile of *S. arvensis* L. ethyl acetate fractions by Thin Layer Chromatography (TLC)

The chromatogram profile of *S. arvensis* L. ethyl acetate fractions by thin layer chromatography (TLC) showed various spots. There were bioactive compounds that can be seen from the appearance of anisaldehyde stains. The sulfuric acid-anisaldehyde solution was used to detect the presence of terpenoids, steroids, and essential oils. After spraying on the TLC plate, it was heated in an oven at 100°C for 5 -10 minutes. Fraction I (combined extract numbers 4 and 5) showed red color. Fraction II (combined extract numbers 6, 7, and 8) showed purple color. Fraction III (combined extract numbers 9, 10, and 11) showed red, purple, blue, and green colors. Fraction IV (combined extract numbers 12, 13, and 14) shows blue and green colors. And fraction V (combined extract numbers 15 and 16) showed a brown color ring. The significant spots were calculated as the retention factor (Rf). Fraction I had 2 spots (Rf. 1.00 and 0.94), fraction II had one spot (Rf. 0.69), fraction III had one spot (0.34), fraction IV had 4 spots (Rf. 0.34, 0.24, 1.90, and 0.13), and fraction V had no spot (Figure 1). TLC was used to separate plant extracts' secondary metabolites.<sup>22</sup> The Rf value indicated a significant diversity of terpenoid compounds separated from various extracts.<sup>23</sup>

### Antioxidant activity

Antioxidant activity of *S. arvensis* L. ethyl acetate fractions showed active to moderate activities. Fraction I-II had moderate activity with the IC<sub>50</sub> value were 145.07 µg/L and 110.52 µg/L respectively. Fraction III-V had active activity with the IC<sub>50</sub> value were 46.62 µg/L, 22.56 µg/L, and 22.82 µg/L respectively (Table 2).

Prieto, 2012<sup>24</sup> categorized the antioxidant activity based on the IC<sub>50</sub> value. The extracts exhibited potent antioxidant activities (IC<sub>50</sub> < 50 µg/L) and moderate antioxidant activities (101 µg/L > IC<sub>50</sub> < 250 µg/L). The IC<sub>50</sub> value (the antioxidant activity) of fraction IV was the lowest. Moreover, compared to the other studies, the ethyl acetate fraction III-V extract from *S. arvensis* L. was lower than plants and callus *Trifolium pratense* L.,<sup>25</sup> *Callisia fragrance* leave juice,<sup>26</sup> and *Centella asiatica* L. leave that have been previously reported as high antioxidant compound. The potent antioxidant activity of the *S. arvensis* L. fraction was probably due to the presence of active ingredients with antioxidant activities, such as polyphenols and flavonoids.<sup>26,27</sup>

### Antimicrobial activity

The antibacterial activity test used the well diffusion method by calculating the diameter of the inhibition zone (DIZ) against *Escherichia coli* and *Staphylococcus aureus* in the media. Based on the previous

report<sup>28</sup>, a compound acts as a very potent antimicrobial if the diameter of the inhibition zone value was > 8 mm, potent if the inhibition zone value was 5 mm > DIZ < 8 mm, and not potent if the inhibition zone value was < 5 mm. This study showed that fractions III-V acted as very potent antimicrobials, with DIZ of fraction III, 10.33±2.17 mm (25%) and 12.10±2.33 mm (50%); fraction IV, 16.78±0.70 mm (25%) and 19.22±0.07 (50%); fraction V, 9.53±5.24 mm (25%) and 13.01±5.22 mm (50%) against *E. coli*; fraction IV, 10.37±0.55 mm (25%) and 15.17±1.04 (50%); and fraction V, 0.08±3.02 mm (50%) against *S. aureus* bacteria because they had DIZ value of > 8 mm. The fraction that acted as a potent antimicrobial was fraction III, 5.09±5.31 mm (50%) against *S. aureus* because it had DIZ value of 5 mm > DIZ < 8 mm. Fractions with no antimicrobial activity were fractions I and II. because they had an inhibition zone value of < 5 mm (Table 3).


The percentage of inhibition of the ethyl acetate fractions group IV was the highest, 102.05±15.34 % (25%), 117.38±23.11% (50%) against *E. coli* and 49.35±13.30% (25%), 83.77±17.32 % (50%) against *S. aureus*. Compared to other studies, the diameter of inhibition zone value was higher than ethanol (15.9 + 0.3 mm) and water (12.2 + 0.7 mm) extract of thyme, water extract of roselle, clove, and rosemary against *E. coli*. The diameter of inhibition zone was also higher than the water extract of thyme, rosemary, and clove against *S. aureus*.<sup>29</sup> They were also higher than ethanolic extract of *Justicia flava*, *Myrianthus arboreus*, and *Momordica charantia*, however lower than aqueous extract, both of *Alchomea cordifolia* and *Psidium guajava* extract against *E. coli*.<sup>30</sup> Some compounds have been industrially applied for antimicrobial activity in doses below 1000 mg.<sup>31</sup>

### Antiplasmodial activity

Antiplasmodial activity of *S. arvensis* L. ethyl acetate fractions showed various activity with different IC<sub>50</sub> against *Plasmodium falciparum* strain 3D7. The IC<sub>50</sub> value of antiplasmodial activity fraction I-V were 494.95 µg/mL, 51.32 µg/mL, 153.66 µg/mL, 12.07 µg/mL, and 13.34 µg/mL, respectively (Table 3-4). Some studies categorized the antiplasmodial activity based on the IC<sub>50</sub> value, an extract with an IC<sub>50</sub> value < 10 µg/mL was classified as very active, an extract's antimalarial activity with a value of 10 µg/mL < IC<sub>50</sub> < 50 µg/mL was classified as active, and the antimalarial activity of an extract with an IC<sub>50</sub> value > 50 µg/mL was classified as inactive.<sup>32,33</sup> Based on the criteria, fractions I-III had inactive activity and fractions IV-V had active activity. This is because the content of bioactive compounds in each fraction is different. The content of compounds such as terpenoids, flavonoids, polyphenols, and anthraquinones has potential as antimalarials.<sup>34</sup> The compounds in *S. arvensis* L. that have been analyzed include the flavonoid and sesquiterpene compounds from the terpenoid group.<sup>7</sup> Due to the *in vitro* antiplasmodial activity of leaf extracts from *Vernonia fimbriifera* Less. (Asteraceae), a bioactivity-guided fractionation was carried out. Three sesquiterpene lactones were isolated, namely 8-(4'-hydroxymethacrylate)-dehydromelitsin, onopordopicrin, and 8α-[4'-hydroxymethacryloyloxy]-4-epi-sonchucarpolide. Their structures were elucidated by spectroscopic methods (1D and 2D NMR and MS analyses) and by comparison with published data. The isolated compounds exhibited antiplasmodial activity with IC<sub>50</sub> values ≤ 5 µg/mL.<sup>35</sup>

## CONCLUSION

This study highlighted the antioxidant, antimicrobial, and antiplasmodial activity of *S. arvensis* L. ethyl acetate fractions. Fraction IV of *S. arvensis* L. ethyl acetate extract had the lowest IC<sub>50</sub> of antioxidant activity against DPPH and antiplasmodial activity against *Plasmodium falciparum* 3D7. Fraction IV also possessed the highest diameter zone inhibition value zone (DIZ) against *Escherichia coli* and *Staphylococcus aureus*. New findings about the activities of these plant extracts could



**Figure 1:** Chromatogram profile of *S. arvensis* L. ethyl acetate fractions by thin layer chromatography (TLC). Mobile phase *n*-hexane:ethyl acetate (4:1). Spot staining: *p*-anisaldehyde sulfuric acid. 1-16: fractions number, I-V: fraction groups number. 17. ethanol extract, 18. ethyl acetate extract, 19. *n*-hexane extract.

**Table 1:** The yield of *S. arvensis* L. ethyl acetate fractions.

Fractions	Weight (g)	Group Fractions	Final weight (g)	Yield (%)
1	0.08			
2	0.19		0.53	10.66%
3	0.26			
4	0.27			
5	0.29	I	0.56	11.14%
6	0.25			
7	0.38	II	0.64	12.88 %
8	0.14			
9	0.36			
10	0.29	III	1.01	20.18%
11	0.36			
12	0.39			
13	0.46	IV	1.21	24.28%
14	0.37			
15	0.33	V	0.82	16.44%
16	0.50			

Note: Five gram of ethyl acetate extract was fractionated to be 16 fractions (Latin number) and then grouped into five fractions (Rome number).

**Table 2:** The  $IC_{50}$  of antioxidant activities of *S. arvensis* L. ethyl acetate fractions.

Sample	$IC_{50}$ ( $\mu$ g/mL)	Activity
I	145.07	moderate
II	110.52	moderate
III	46.62	active
IV	22.56	active
V	22.82	active

**Table 3:** Diameter of imbibition zone of *S. arvensis* L. ethyl acetate fractions against *Escherichia coli* and *Staphylococcus aureus* (mm).

Fractions	<i>Escherichia coli</i> (mm)				<i>Staphylococcus aureus</i> (mm)			
	Diameter of inhibition zone (mm)		Percentage of inhibition (%)		Diameter of inhibition zone (mm)		Percentage of inhibition (%)	
	25%	50%	25%	50%	25%	50%	25%	50%
I	0	0	0	0	0	0	0	0
II	3.17±3.01	4.50±4.50	29.15±11.88	41.56±19.71	0	0	0	0
III	10.33±2.17	12.10±2.33	61.81±22.61	72.19±25.11	3.28±3.40	6.00±1.00	22.16±16.07	29.86±16.50
IV	16.78±0.70	19.22±0.07	102.05±15.34	117.38±23.11	10.37±0.55	15.17±1.04	49.35±13.30	83.77±17.32
V	9.53±5.24	13.01±5.22	45.95±24.17	63.17±23.16	5.09±5.31	10.08±3.02	31.87±30.56	60.93±19.25
Chloramphenicol (1000 mg/L)	20.43±1.11				20.67±0.31			

**Table 4: Growth and inhibition percentage of *S. arvensis* L. ethyl acetate fractions against *Plasmodium falciparum* strain 3D7.**

Concentration (µg/mL)	Fractions	% Parasitemia		Growth percentage	Inhibition percentage
		0h	48h		
control (-)	I	1.02	6.37	5.35	-
	II	1.02	6.48	5.46	-
	III	1.02		5.68	-
	IV	1.02		5.58	-
	V	0.62		3.10	-
100	I	1.02	5.01	3.99	25.42
	II	1.02	3.66	2.64	51.65
	III	1.02		3.06	46.13
	IV	1.02		0.62	88.89
	V	0.62		0.01	99.68
10	I	1.02	5.85	4.83	09.72
	II	1.02	4.29	3.27	40.11
	III	1.02		5.07	10.74
	IV	1.02		3.64	34.77
	V	0.62		1.99	35.81
1	I	1.02	6.59	5.57	00.00
	II	1.02	5.96	4.49	09.52
	III	1.02		5.21	08.27
	IV	1.02		4.54	18.64
	V	0.62		3.15	00.00
0.1	I	1.02	6.82	5.80	00.00
	II	1.02	6.41	5.41	00.92
	III	1.02		5.43	04.40
	IV	1.02		5.95	00.00
	V	0.62		3.22	00.00
0.01	I	1.02	8.12	6.10	00.00
	II	1.02	8.08	6.00	00.00
	III	1.02		6.72	00.00
	IV	1.02		7.16	00.00
	V	0.62		3.52	00.00

**Table 5: The IC<sub>50</sub> value of antiplasmodial activities of *S. arvensis* L. ethyl acetate fractions.**

Fractions	IC <sub>50</sub> (µg/mL)	Activities
I	494.947	Not Active
II	51.323	Not Active
III	153.664	Not Active
IV	12.068	Active
V	13.343	Active

lead to the isolation and identification of active compounds for further pharmaceutical applications.

## ACKNOWLEDGMENTS

The authors would like to thank The Ministry of Education of the Republic of Indonesia that was funded this research (grant no.714/UN3.14/PT/2020).

## REFERENCES

1. WHO. World Health Organization. Publishes List of Bacteria for Which New Antibiotics are Urgently Needed. 2017.
2. Kemenkes RI. Info DATIN Pusat Data dan Informasi Kesehatan Kementerian Kesehatan RI : Malaria. Kementerian Kesehatan Republik Indonesia, Jakarta. 2016.
3. Muharani, Fitrya, Farida S. Antibacterial Activity Assay of Ethanolic Extract of Medicinal Plant Musi Tripe in District of Musi Banyuasin, South Sumatera. *Ind Pharm J.* 2017;7(2):127-35.
4. WHO. World Health Organization: World Malaria Report 2021. WHO Press. Geneva. 2021.
5. Ganapathy S, Karpagam S. *In Vitro* Evaluation of Antibacterial Potential of *Andrographis paniculata* Against Resistant Bacterial Pathogens Methicillin-Resistant *Staphylococcus aureus* (MRSA) and Multiple Drug Resistant *Escherichia coli* (MDR *E. coli*). *Int J Bioassays.* 2016;5(3):4879-81.
6. Wahyuni DK, Rahayu S, Purnama PR, Saputro TB, Suharyanto, Wijayanti N, *et al.* Morpho-anatomical structure and DNA barcode of *Sonchus arvensis* L. *Biodiversitas.* 2019;20(1):2417-26.
7. Wahyuni DK, Lestari S, Kuncoro EP, Purnobasuki H. Callus induction and its metabolite profiles of *Sonchus arvensis* L. under temperature treatment. *Ann Biol.* 2020;36(2):299-303.
8. Wahyuni DK, Purnobasuki H, Kuncoro EP, Ekasari W. Callus Induction of *Sonchus arvensis* L. and Its Antiplasmodial Activity. *African J Inf Dis.* 2020;14(1):1-7.
9. Wahyuni DK, Rahayu S, Zaidan AH, Ekasari W, Prasongsuk S, Purnobasuki H. Growth, Secondary Metabolite Production, and *In Vitro* Antiplasmodial Activity of *Sonchus arvensis* L. Callus Under Dolomite [CaMg(CO<sub>3</sub>)<sub>2</sub>] Treatment. *PLoS ONE.*

- 2021;16(8):e0254804.
10. Rachmawati F, Nuria MC, Sumantri. Antibacterial Activity Assay of Chloroform Fraction of Ethanolic Extract of pegagan (*Centella asiatica* (L.) Urb.) and Its Bioactive Compound Identification. Faculty of Pharmacy, University of Wahid Hasyim, Semarang. 2011.
  11. Tapan S. Quantitative HPLC Analysis of Phenolic Acids, Flavonoids, and Ascorbic Acid in Four Different Solvent Extracts of Two Wild Edible Leaves, *Sonchus arvensis* L. and *Oenanthe linearis* of North-Eastern Region in India. *J Appl Pharma Sci.* 2016;6(2):157-66.
  12. Xia Z, Qu W, Lu H, Fu J, Ren Y, Liang J. Sesquiterpene Lactones from *Sonchus arvensis* L. and Their Antibacterial Activity Against *Streptococcus mutans* ATCC 25175. *Fitoterapia.* 2009;81(5):424-8.
  13. Kanaani S, Sani AM. Chemical Composition of Essential Oils and *In Vitro* Antibacterial Activity of Methanolic Extract of *Sonchus arvensis* L. and *Eremurus spectabilis* Against Food-Borne Pathogenic Bacteria. *J Essent Oil-Bear Plants.* 2015;18(5):1093-9.
  14. Seal T. Quantitative HPLC Analysis of Phenolic Acids, Flavonoids, and Ascorbic Acid in Four Different Solvent Extracts of Two Wild Edible Leaves, *Sonchus arvensis* and *Oenanthe linearis* of North-Eastern Region in India. *J Appl Pharma Sci.* 2016;6(1):157-66.
  15. Marston A, Hostettmann A. Bioassay Methods in Natural Product Research and Drug Development: Biological and Chemical Evaluation of Plant Extracts and Subsequent Isolation Strategy. Proceedings of the Phytochemical Society of Europe. Springer Science Business Media Dordrecht. 1999;43:67-80.
  16. Fu R, Zhang Y, Guo Y, Liu F, Chen F. Determination of Phenolic Contents and Antioxidant Activities of Extracts of *Jatropha curcas* L. Seed Shell, A By-Product, A New Source of Natural Antioxidant. *Industrial Crops Prod.* 2014;58(1):265-70.
  17. Ekasari W, Pratiwi DW, Amanda Z, Suciati WA, Arwati H. Various Parts of *Helianthus annuus* Plants as New Sources of Antimalarial Drugs. *Evid Based Complementary Alternat Med.* 2019;2019:7390385.
  18. Nasr A, Zhou X, Huang S, Wang Y, Li X, Guo-Ping Zhu G. Comparative Effects of Some Extraction Solvents on The Antimicrobial Activity of *Eucalyptus camaldulensis* Leaf, Bud, Capsule, and Seed Crude Extracts. *Nat Prod Res.* 2019;33(17):2560-5.
  19. Dhanani T, Shah S, Gajbhiye NA, Kumar S. Effect of Extraction Methods on Yield, Phytochemical Constituents, and Antioxidant Activity of *Withania somnifera*. *Arabian J Chem.* 2017;10(1):S1193-9.
  20. Adam OAO, Abadi RSM, Ayoub SMH. The Effect of Extraction Method and Solvents on Yield and Antioxidant Activity of Certain Sudanese Medicinal Plant Extracts. *J Phytopharmacol.* 2019;8(5):248-52.
  21. D'Auria M, Mecca M, Bruno MR, Todaro L. Extraction Methods and Their Influence on Yield When Extracting Thermo-Vacuum Modified Chestnut Wood. *Forests.* 2021;12:73.
  22. Ali KS, Mohammed ASA, Munayem RT. Phytochemical Screening and Thin Layer Chromatography of *Acacia etbaica* ssp. *Uncinata* leaves. *World J Pharma Res.* 2017;6(12):1278-83.
  23. Ahamed T, Rahman SKM, Shohae AM. Thin Layer Chromatographic Profiling and Phytochemical Screening of Six Medicinal Plants in Bangladesh. *IJB.* 2017;11(1):131-40.
  24. Prieto JM. Procedure: Preparation of DPPH Radical, and Antioxidant Scavenging Assay. Prieto's DPPH Microplate Protocol. <https://www.researchgate.net/file.PostFileloader.html?id=503cd1c9e39d5ead11000043&assetKey=AS%3A271744332435456%401441800305338>, March, 30th, 17.30 Western Indonesian Time, 2021. 2012.
  25. Esmaeili AK, Taha RM, Mohajer S, Banisalam B. Antioxidant Activity and Total Phenolic and Flavonoid Content of Various Solvent Extracts from *In Vivo* and *In Vitro* grown *Trifolium pratense* L. (Red Clover). *BioMed Res Int.* 2015;2015:643285.
  26. Olennikov DN, Zilfikarov IN, Toropova AA, Ibragimov TA. Chemical Composition of *Callisia fragrans* Wood Juice and Its Antioxidative Activity (*In Vitro*). *Chem Plant Raw Mat J.* 2008;4(1):95-100.
  27. Yahya MA, Nurrosyidah IH. Antioxidant activity Ethanol Extract of gotu kola (*Centella asiatica* L.) with DPPH Method (2,2-diphenyl-1-picrylhydrazil). *J Halal Prod Res.* 2020;3:106-12.
  28. Serri A, Mahboubi A, Moghimi H. Investigating The Antimicrobial Efficacy of Liposomal Vancomycin in Gram-Positive and Gram-Negative Bacteria: A Preliminary Mechanistic Study. *Iranian Pharma Sci.* 2018;2018:203946010.
  29. Gonelimali FD, Lin J, Miao W, Xuan J, Charles F, Chen M, et al. Antimicrobial Properties and Mechanism of Action of Some Plant Extracts Against Food Pathogens and Spoilage Microorganisms. *Front Microbiol.* 2018;9:1639.
  30. Owusu E, Ahorlu MM, Afutu E, Akumwena A, Asare GA. Antimicrobial Activity of Selected Medicinal Plants from a Sub-Saharan African Country Against Bacterial Pathogens from Post-Operative Wound Infections. *Med Sci.* 2021;9(2):23.
  31. Khameneh B, Iranshahy M, Sohelli V, Bazzaz BSF. Review on Plant Antimicrobials: A Mechanistic Viewpoint. *Antimicrob Resist Infect Control.* 2019;8:118.
  32. Lima RB, Rocha e Silva LF, Melo MR, Costa JS, Picanço NS, Lima ES. *In vitro* and *In Vivo* Antimalarial Activity of Plants from The Brazilian Amazon. *Malar J.* 2015;14:508.
  33. Tajuddeen N, van Heerden N. Antiplasmodial Natural Products: An update. *Malar J.* 2019;18(1):404.
  34. Widyawaruyanti A, Harwiningtias N, Tumewu L, Hafid AF, Soetjpto. Effect of formulated *Artocarpus champeden* Extract on Parasite Growth and Immune Response of *Plasmodium berghei*-infected Ice. *Evid Based Complementary Alternat Med.* 2020;2020:4678634.
  35. Bordignon A, Ciekiewicz E, Campos PE, Frederich M. Study of The Antiplasmodial Potential of *Vernonia fimbriifera* Less. (Asteraceae) Leaves, an Endemic Plant from Reunion Island. *Planta Medica.* 2016;81(S01):S1-S381.

**Cite this article:** Wahyuni DK, Nariswari A, Supriyanto A, Purnobasuki H, Punnapayak H, Bankeeree W, et al. Antioxidant, Antimicrobial, and Antiplasmodial Activities of *Sonchus arvensis* L. Leaf Ethyl Acetate Fractions. *Pharmacogn J.* 2022;14(6)Suppl: 993-998.

# Antioxidant, Antimicrobial, and Antiplasmodial Activities of *Sonchus arvensis* L. Leaf Ethyl Acetate Fractions

## ORIGINALITY REPORT

12%

SIMILARITY INDEX

9%

INTERNET SOURCES

8%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	2%
2	<a href="http://www.science.gov">www.science.gov</a> Internet Source	2%
3	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	1%
4	Sameh Sobhy El - Hadad, Natali Aleksandrovna Tikhomirova, Mahmoud Abd El - Aziz. "Biological activities of dihydroquercetin and its effect on the oxidative stability of butter oil", <i>Journal of Food Processing and Preservation</i> , 2020 Publication	1%
5	Kabouche, A.. "Antioxidant abietane diterpenoids from <i>Salvia barrelieri</i> ", <i>Food Chemistry</i> , 2007 Publication	<1%
6	<a href="http://www.pyiephyomaung.yolasite.com">www.pyiephyomaung.yolasite.com</a> Internet Source	<1%



7	<a href="http://cer.ihtm.bg.ac.rs">cer.ihtm.bg.ac.rs</a> Internet Source	<1 %
8	<a href="http://pesquisa.bvsalud.org">pesquisa.bvsalud.org</a> Internet Source	<1 %
9	<a href="http://link.springer.com">link.springer.com</a> Internet Source	<1 %
10	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	<1 %
11	Mahfuz Elmastaş, Ibrahim Dermirtas, Omer Isildak, Hassan Y. Aboul - Enein. " Antioxidant Activity of S - Carvone Isolated from Spearmint ( ) ", Journal of Liquid Chromatography & Related Technologies, 2007 Publication	<1 %
12	<a href="http://www.jmedsoc.org">www.jmedsoc.org</a> Internet Source	<1 %
13	M. Gajendiran, S. Divakar, N. Raaman, S. Balasubramanian. "In Vitro Drug Release Behavior, Mechanism and Antimicrobial Activity of Rifampicin Loaded Low Molecular Weight PLGA-PEG-PLGA Triblock Copolymeric Nanospheres", Current Drug Delivery, 2013 Publication	<1 %
14	<a href="http://id.123dok.com">id.123dok.com</a> Internet Source	<1 %

- |    |   |      |
|----|---|------|
| 15 | <a href="https://repository.icr.ac.uk">repository.icr.ac.uk</a><br>Internet Source  | <1 % |
| 16 | <a href="http://www.jove.com">www.jove.com</a><br>Internet Source   | <1 % |
| 17 | Caroline Issler Rodrigues, Diego Mota da Costa, Acidália Carine Vieira Santos, Maria José Moreira Batatinha et al. "Assessment of in vitro anthelmintic activity and bio-guided chemical analysis of BRS Boyrá pineapple leaf extracts", <i>Veterinary Parasitology</i> , 2020<br>Publication | <1 % |
| 18 | Roberto Lanna Filho, Reginaldo da Silva Romeiro, Eduardo Alves. "Bacterial spot and early blight biocontrol by epiphytic bacteria in tomato plants", <i>Pesquisa Agropecuária Brasileira</i> , 2010<br>Publication  | <1 % |
| 19 | Sagorika Panda, Jyoti Ranjan Rout, Pallabi Pati, Manoranjan Ranjit, Santi Lata Sahoo. "Antimalarial activity of <i>Artemisia nilagirica</i> against <i>Plasmodium falciparum</i> ", <i>Journal of Parasitic Diseases</i> , 2017<br>Publication  | <1 % |
| 20 | Van Tang Nguyen, Michael C. Bowyer, Quan Van Vuong, Ian A. Van Altena, Christopher J. Scarlett. "Phytochemicals and antioxidant capacity of Xao tam phan ( <i>Paramignya</i>  | <1 % |

trimer) root as affected by various solvents and extraction methods", Industrial Crops and Products, 2015

Publication

21

[ejournal.umm.ac.id](http://ejournal.umm.ac.id)

Internet Source

<1 %

22

[scholar.sun.ac.za](http://scholar.sun.ac.za)

Internet Source

<1 %

23

[www.biorxiv.org](http://www.biorxiv.org)

Internet Source

<1 %

24

B J Thomas, R T Evans, D A Hawkins, D Taylor-Robinson. "Sensitivity of detecting Chlamydia trachomatis elementary bodies in smears by use of a fluorescein labelled monoclonal antibody: comparison with conventional chlamydial isolation.", Journal of Clinical Pathology, 1984

Publication

<1 %

25

Gülsen Tel-Çayan, Mehmet Öztürk, Mehmet E. Duru, Mujeeb Ur Rehman, Achyut Adhikari, Aziz Türkoğlu, M. Iqbal Choudhary.

"Phytochemical investigation, antioxidant and anticholinesterase activities of Ganoderma adspersum", Industrial Crops and Products, 2015

Publication

<1 %

- 26 Uttpal Anand, Samapika Nandy, Avinash Mundhra, Neela Das, Devendra Kumar Pandey, Abhijit Dey. "A review on antimicrobial botanicals, phytochemicals and natural resistance modifying agents from Apocynaceae family: Possible therapeutic approaches against multidrug resistance in pathogenic microorganisms", Drug Resistance Updates, 2020  
Publication <1 %
- 
- 27 Zahra Hasanpour, Peyman Salehi, Atusa Aliahmadi, Mahdieh Hoseinpour, Hossein Behboodi, Dan Staerk, Morteza Bararjanian. "Semi-synthesis of Novel Thebaine Derivatives and their Anti-bacterial and Wound Healing Properties", Research Square Platform LLC, 2023  
Publication <1 %
- 
- 28 [orbi.uliege.be](https://orbi.uliege.be)  
Internet Source <1 %
- 
- 29 [rjptonline.org](https://rjptonline.org)  
Internet Source <1 %
- 
- 30 [worldwidescience.org](https://worldwidescience.org)  
Internet Source <1 %
- 
- 31 Kyriacos C Nicolaou, Stephan Rigol. "A brief history of antibiotics and select advances in

their synthesis", The Journal of Antibiotics,  
2017

Publication

---

32

S.F. Mbatchi, B. Mbatchi, J.T. Banzouzi, T.  
Bansimba, G.F. Nsonde Ntandou, J.-M.  
Ouamba, A. Berry, F. Benoit-Vical. "In vitro  
antiplasmodial activity of 18 plants used in  
Congo Brazzaville traditional medicine",  
Journal of Ethnopharmacology, 2006

<1 %

Publication

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Antioxidant, Antimicrobial, and Antiplasmodial Activities of *Sonchus arvensis* L. Leaf Ethyl Acetate Fractions

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---