

p-ISSN: 0854-4263

e-ISSN: 2477-4685

INDONESIAN JOURNAL OF
**Clinical Pathology and
Medical Laboratory**

Majalah Patologi Klinik Indonesia dan Laboratorium Medik

Diterbitkan oleh Perhimpunan Dokter Spesialis Patologi Klinik Indonesia

Published by Indonesian Association of Clinical Pathologists

Accredited No : 66b/DIKTI/KEP/2011, 9 September 2011



INDONESIAN JOURNAL OF Clinical Pathology and Medical Laboratory

Majalah Patologi Klinik Indonesia dan Laboratorium Medik

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [ANNOUNCEMENTS](#) [AU](#)

[Home](#) > [About the Journal](#) > [Editorial Team](#)

Editorial Team

Editor-in-chief

[Puspa Wardhani](#), Editor-in-chief Indonesian Journal of Clinical Pathology and Medical Laboratory, Indonesia, Indonesia

Editorial Boards

[Ida Parwati](#), Department of Clinical Pathology, Faculty of Medicine, Padjajaran University, Bandung, Indonesia, Indonesia

[Anak Agung Wiradewi Lestari](#), Department of Clinical Pathology, Faculty of Medicine, Udayana University/Sanglah Hospital, Bali, Indonesia, Indonesia

[Yulia Nadar Indrasari](#), Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia

[Nuri Diah Indrasari](#), Department of Clinical Pathology, Faculty of Medicine, Indonesia University/Dr. Cipto Mangunkusumo, Jakarta, Indonesia, Indonesia

[Efrida Efrida](#), Department of Clinical Pathology, Faculty of Medicine, Andalas Univesity/M.Djamil Hospital, Padang, Indonesia, Indonesia

[Puspa Wardhani](#), Editor-in-chief Indonesian Journal of Clinical Pathology and Medical Laboratory, Indonesia, Indonesia

[Nyoman Suci Widiastuti](#), Department of Clinical Pathology, Faculty of Medicine, Diponegoro University/Dr. Kariadi, Semarang, Indonesia, Indonesia

[Uleng Bahrun](#), Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Dr. Wahidin Sudirohusodo Hospital, Makassar, Indonesia, Indonesia

[Budi Mulyono](#), Department of Clinical Pathology, Faculty of Medicine, Gajah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia

Editorial Assistant

[Dian Wahyu Utami](#), Indonesian Journal of Clinical Pathology and Medical Laboratory, Indonesia, Indonesia

Peer-Reviewers

[Tony Badrick](#), Royal College of Pathologists of Australia Quality Assurance Programs, Australia

[Che Maraina Bt Che Hussin](#), Universiti Sains Malaysia, Malaysia

[Zilfalil Bin Alwi](#), Universiti Sains Malaysia, Malaysia

[Teguh Triyono](#), Department of Clinical Pathology, Faculty of Medicine, Gadjah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia

[Wendy Erber](#), The University of Western Australia, Australia

[Hans Vrielink](#), Sanquin Blood Supply, Plesmanlaan 125, 1066 CX Amsterdam, the Netherlands, Netherlands

[Ni Kadek Mulyantari](#), Department of Clinical Pathology, Faculty of Medicine, Udayana University/Sanglah General Hospital Bali, Indonesia, Indonesia

[Adi Koesoema Aman](#), Department of Clinical Pathology, Faculty of Medicine, Sumatera Utara University/Adam Malik

Hospital, Medan, Indonesia, Indonesia
Basti Andriyoko, Department of Clinical Pathology, Faculty of Medicine, Padjajaran University/Dr. Hasan Sadikin Hospital, Bandung, Indonesia, Indonesia
Paulus Budiono Notopuro, Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia
Tonang Dwi Ardyanto, Department of Clinical Pathology, Faculty of Medicine, Sebelas Maret University/Dr. Moewardi Hospital, Surakarta, Indonesia, Indonesia
Agustin Iskandar, Faculty of Medicine, Brawijaya University, Indonesia
Tenri Esa, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Dr. Wahidin Sudirohusodo, Makassar, Indonesia, Indonesia
Rachmawati Adiputri Muhiddin, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Dr. Wahidin Sudirohusodo, Makassar, Indonesia, Indonesia
Liong Boy Kurniawan, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Wahidin Sudirohusodo Hospital, Makassar, Indonesia, Indonesia
Andaru Dahesih Dewi, Department of Clinical Pathology, Faculty of Medicine, Gajah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia
Umi Solekhah Intansari, Department of Clinical Pathology, Faculty of Medicine, Gajah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia
Banundari Rachmawati, Department of Clinical Pathology, Faculty of Medicine, Diponegoro University/Dr. Kariadi Hospital, Semarang, Indonesia, Indonesia
Agnes Rengga Indrati, Department of Clinical Pathology, Faculty of Medicine, Padjajaran University/Hasan Sadikin Hospital, Bandung, Indonesia, Indonesia
Rikarni Rikarni, Department of Clinical Pathology, Faculty of Medicine, Andalas University/M.Djamil Hospital, Padang, Indonesia, Indonesia
Hani Susianti, Department of Clinical Pathology, Faculty of Medicine, Brawijaya University/Saiful Anwar Hospital, Malang, Indonesia, Indonesia
Adhi Kristianto Sugianli, Department of Clinical Pathology, Faculty of Medicine, Padjajaran University/Dr. Hasan Sadikin Hospital, Bandung, Indonesia, Indonesia
Yetti Hernaningsih, Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia
Yuyun Widaningsih, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Wahidin Sudirohusodo Hospital, Makassar, Indonesia, Indonesia
Osman Sianipar, Department of Clinical Pathology, Faculty of Medicine, Gajah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia
Rismawati Yaswir, Department of Clinical Pathology, Faculty of Medicine, Andalas University/M. Djamil Hospital, Padang, Indonesia, Indonesia
Maimun Zulhaidah Arthamin, Department of Clinical Pathology, Faculty of Medicine, Brawijaya University/Saiful Anwar Hospital, Malang, Indonesia, Indonesia
Aryati Aryati, Department of Clinical Pathology, Faculty of Medicine, Airlangga University/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia
Rahajuningsih Dharma, Department of Clinical Pathology, Faculty of Medicine, Indonesia University/Ciptomangunkusumo Hospital, Jakarta, Indonesia
Mansyur Arif, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Wahidin Sudirohusodo Hospital, Makassar, Indonesia
July Kumalawati, Department of Clinical Pathology, Faculty of Medicine, Indonesia University/Ciptomangunkusumo Hospital, Jakarta, Indonesia, Indonesia
Purwanto AP, Department of Clinical Pathology, Faculty of Medicine, Diponegoro University/Dr. Kariadi Hospital, Semarang, Indonesia, Indonesia
Jusak Nugraha, Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia
Edi Widjayanto, Department of Clinical Pathology, Faculty of Medicine, Brawijaya University/Saiful Anwar Hospital, Malang, Indonesia, Indonesia
Ninik Sukartini, Department of Clinical Pathology, Faculty of Medicine, Indonesia University/Dr. Ciptomangunkusumo Hospital, Jakarta, Indonesia, Indonesia
Kusworini Handono, Department of Clinical Pathology, Faculty of Medicine, Brawijaya University/Dr. Saiful Anwar Hospital, Malang, Indonesia, Indonesia
Ranti Permatasari, Department of Clinical Pathology, Faculty of Medicine, Sumatera Utara University/Adam Malik Hospital, Medan, Indonesia, Indonesia



INDONESIAN JOURNAL OF Clinical Pathology and Medical Laboratory

Majalah Patologi Klinik Indonesia dan Laboratorium Medik

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [ANNOUNCEMENTS](#) [AU](#)

[Home](#) > [Archives](#) > **Vol 26, No 2 (2020)**

Vol 26, No 2 (2020)

Table of Contents

Articles

Comparison of Hepcidin Levels in Children with and without Soil-Transmitted Helminths Infection	PDF <i>Pg. 129-133</i>
<i>Dewi Saputri, Yunilda Andriyani, Almaycano Ginting</i>	
Human Sperm Cells After Purification Using SCLB Can Be Stored at 4o, -20o, or -80oC Before Small RNA Isolation	PDF <i>Pg. 134-137</i>
<i>Berliana Hamidah, Ashon Sa'adi, Rina Yudiwati</i>	
Diagnostic Value of Encode TB IgG and IgM Rapid Test to Support Pulmonary Tuberculosis Diagnosis	PDF <i>Pg. 138-141</i>
<i>Notrisia Rachmayanti, Aryati Aryati, Tutik Kusmiati</i>	
Diagnostic Test of PIVKA-II as A Tumor Marker for Hepatocellular Carcinoma	PDF <i>Pg. 142-145</i>
<i>Dwi Priyadi Djatmiko, I Putu Adi Santosa, Elvin Richela Lawanto, Bogi Pratomo, Hani Susianti</i>	
Correlation of Serum Interleukin-6, TNF-α, Procalcitonin and Leukocyte Count in Patients with Suspected Sepsis	PDF <i>Pg. 146-150</i>
<i>Erfina Lim, Jusak Nugraha</i>	

<p>Evaluation of the Progressivity Parameters of Chronic Kidney Disease after Branched-Chain Amino Acid Supplementation in Children</p> <p><i>Esthy Poespitaningtyas, Roedi Irawan, Ninik Asmaningsih Soemyarso, Jusak Nugraha</i></p>	<p>PDF</p> <p>Pg. 151-157</p>
<p>Analysis of Liver Fibrosis Degree with APRI Score and FIB-4 Index on Patients with Non-Alcoholic Fatty Liver Disease</p> <p><i>Gillian Elvira Seipalla, Nurahmi Nurahmi, Ibrahim Abd Samad</i></p>	<p>PDF</p> <p>Pg. 158-161</p>
<p>Comparison of the Profile and TSH Levels from Several Types of Blood Collection Tubes</p> <p><i>Gunawan Eka Putra, Ninik Sukartini, Suzanna Immanuel, Fify Henrika, Nuri Dyah Indrasari</i></p>	<p>PDF</p> <p>Pg. 162-167</p>
<p>Correlation between Plasma Osteopontin and Alkaline Phosphatase in Type 2 Diabetes Mellitus Patients</p> <p><i>Josua TH Sinambela, M.I Diah Pramudianti, Dian Ariningrum</i></p>	<p>PDF</p> <p>Pg. 168-174</p>
<p>Infection of Cytomegalovirus in Cholestasis Infant with Biliary Atresia</p> <p><i>Lasmauli Situmorang, Bagus Setyoboedi, Sjamsul Arief, Gondo Mastutik</i></p>	<p>PDF</p> <p>Pg. 175-181</p>
<p>Seropositivity of Anti-Rubella Antibodies as A Marker for Rubella Infection in Infants at High Risk of Congenital Deafness</p> <p><i>Nyilo Purnami, Risa Etika, Martono Martono, Puspa Wardhani</i></p>	<p>PDF</p> <p>Pg. 182-186</p>
<p>Soluble Suppression of Tumorigenicity-2 Levels As Prognostic Marker in Non-ST-segment Elevation Myocardial Infarction</p> <p><i>Sherly Purnamawaty, Tenri Esa, Ibrahim Abd Samad</i></p>	<p>PDF</p> <p>Pg. 187-192</p>
<p>Description of Fecal Culture Results in Diarrhea Patients Due To Antibiotic Use</p> <p><i>Suci Tresna, I.G.A.A Putri Sri Rejeki, Puspa Wardhani</i></p>	<p>PDF</p> <p>Pg. 193-197</p>
<p>Analysis of D-dimer Levels in Deep Vein Thrombosis Patients</p> <p><i>Anton Triyadi, Rachmawati A. Muhiddin, Agus Alim Abdullah</i></p>	<p>PDF</p> <p>Pg. 198-202</p>



<p>The Relation of 25-Hydroxyvitamin D Level with Metabolic Syndrome in Type 2 Diabetes Mellitus Patients</p> <p><i>M.I. Diah Pramudianti, Dian Ariningrum, Medityas Winda Krissinta</i></p>	<p>PDF</p> <p>Pg. 203-209</p>
<p>Diagnostic Value of Plasmotec Malaria-3 Antigen Detection on Gold Standard Microscopy</p> <p><i>Trieva Verawaty Butarbutar, Puspa Wardhani, Aryati Aryati</i></p>	<p>PDF</p> <p>Pg. 210-216</p>
<p>Differences of Bone Marrow Features and BCR-ABL Variants in Chronic Granulocytic Leukemia Post Tyrosine Kinase Inhibitor Therapy</p> <p><i>Wivina Riza Devi, M Darwin Prenggono, Purwanto AP, Imam B</i></p>	<p>PDF</p> <p>Pg. 217-222</p>
<p>Profile of Rapid Molecular Test of Tuberculosis Using Xpert MTB/RIF</p> <p><i>Faigah Aprilia Sy Faraid, Irda Handayani, Tenri Esa</i></p>	<p>PDF</p> <p>Pg. 223-228</p>
<p>Analysis of Hematologic Parameters and Serum Bilirubin Levels in Complicated and Uncomplicated Acute Appendicitis Patients</p> <p><i>Rini Rahman, Ani Kartini, Yuyun Widaningsih, Agus Alim Abdullah</i></p>	<p>PDF</p> <p>Pg. 229-234</p>
<hr/>	
<h2>Literature Review</h2>	
<p>Mean Platelet Volume as A Marker of Thrombosis Event in Antiphospholipid Syndrome Patients</p> <p><i>Michael Dwinata, Jonathan H. Haposan, Inolyn Pandjaitan</i></p>	<p>PDF</p> <p>Pg. 235-240</p>
<hr/>	
<h2>Case Report</h2>	
<p>Gestational Trophoblastic Neoplasia with Hyperthyroidism</p> <p><i>Devi Rahmadhona, Betty Agustina Tambunan</i></p>	<p>PDF</p> <p>Pg. 241-248</p>
<p>A 24-Year-Old Male with Gigantism, Growth Hormone Deficiency, Suspected Clivus Chordoma, Primary Hypothyroidism, Hypogonadism and Pancytopenia</p> <p><i>W.A. Arsana, M.I. Diah Pramudianti</i></p>	<p>PDF</p> <p>Pg. 249-256</p>

Cover	PDF
	<i>Pg.</i>

Contents	PDF
	<i>Pg.</i>

Back Matter

Guidelines for Author and Subscribes Form	PDF
	<i>Pg.</i>

Contact Us:

Laboratorium Patologi Klinik RSUD Dr. Soetomo

Jl. Mayjend. Prof. Dr. Moestopo 6-8 Surabaya.

Telp/fax (031-5042113, 085733220600

E-mail: majalah.ijcp@yahoo.com

Website: <https://indonesianjournalofclinicalpathology.org>

Indonesian Journal of Clinic
licensed under a [Creative Commons](#)

ACCREDITED No. 36a/E/KPT/2016, 23 Mei 2016

OAI Address: <https://indonesianjournalofclinicalpathology.org/>

Find us on:



Infection of Cytomegalovirus in Cholestasis Infant with Biliary Atresia

Lasmauli Situmorang,¹ Bagus Setyoboedi,¹ Gondo Mastutik,² Sjamsul Arief¹

¹ Division of Hepatology, Department of Child Health, Faculty of Medicine, Airlangga University/Dr. Soetomo Hospital, Surabaya, Indonesia.
E-mail: baguzze@gmail.com

² Department of Anatomic Pathology, Faculty of Medicine, Airlangga University/Dr. Soetomo Hospital, Surabaya, Indonesia

ABSTRACT

Biliary Atresia (BA) is extrahepatic cholestasis that results in death within the first two years if the diagnosis and intervention are delayed. The etiology and pathogenesis of BA are still undetermined. Viral infections, including Cytomegalovirus (CMV), are presumed to be one of the causes. Cytomegalovirus infection is more common in intrahepatic than extrahepatic cholestasis such as BA. There are limited data about Cytomegalovirus infection in cholestatic infants with BA. This study compared the incidence of CMV infection in cholestatic infants with biliary atresia and non-biliary atresia. A cross-sectional study was performed in December 2017 - August 2018 in cholestatic infants aged 1-6 months. Liver biopsy, histopathological examination followed by PCR CMV examination were performed on cholestatic infants. The results of the PCR examination were compared between BA and non-BA infants. Statistical analysis of Chi-Square, t-test independent and Mann-Whitney U resulting in $p < 0.05$ were stated as significant. Thirty-seven children were obtained during the study period, consisting of sixteen children with BA and twenty-one children with non-BA. Biliary atresia was predominantly found in female than male children, despite no differences were found between the groups ($p = 0.163$). There were differences in body weight ($p = 0.002$) age ($p = 0.009$), birth weight ($p = 0.02$) and gestational age ($p = 0.03$) between children with BA and non-BA. There was no significant difference in the incidence of CMV infection in cholestatic infants with BA and non-BA ($p = 0.338$). Cytomegalovirus infection in cholestatic infants with BA was less than non-BA cholestatic infants.

Keywords: Biliary atresia, cholestasis, cytomegalovirus, polymerase chain reaction

INTRODUCTION

Biliary Atresia (BA) remains a great challenge for clinicians because it has poor clinical outcomes if not early diagnosed and intervened.^{1,2} Biliary atresia is a type of extrahepatic cholestasis which is frequently found in infants. However; the etiology and pathogenesis of BA are still undetermined. Particular viruses have been suggested to play a role in pathogenesis BA, including group C Rotavirus, Reovirus and Cytomegalovirus (CMV).³

Cytomegalovirus infection is often found in intrahepatic cholestasis; however, currently, there are several studies which showed CMV infection in extrahepatic cholestasis, including BA.^{4,5} Cholestasis is classified as intrahepatic and extrahepatic cholestasis, and there are several methods for diagnosing CMV infection, including Polymerase Chain Reaction (PCR). There is no single examination that is 100% accurate in diagnosing CMV infection. Polymerase Chain Reaction examination can use blood, urine and tissue specimens, which are ideally carried out at the age of three weeks after birth.⁶ The PCR examination cannot be performed at the age of

three weeks because most cases of CMV infection are asymptomatic and high in cost; however, PCR can detect viral DNA with low amounts of sample and time-efficient.⁷

The gold standard of diagnosis of BA is intraoperative cholangiography; however, liver biopsy and following histopathological examination have quite high sensitivity of approximately 96.9%.⁸ Several studies showed positive CMV results in PCR examination of liver biopsies of patients with AB.^{9,10} However, there is no data on the incidence of CMV infection in cholestatic infants with BA in Dr. Soetomo Hospital, Surabaya. Therefore, a preliminary study is needed to determine the incidence of CMV infection in cholestatic patients with BA and without BA by using PCR in liver tissue in Dr. Soetomo Hospital, Surabaya.

METHODS

A cross-sectional study was performed from December 2017 to August 2018. Cholestasis infants aged 1-6 months old who were treated at Hepatology division were included. Cholestasis infants associated with severe infections (sepsis) or

severe multi-organ abnormalities, a history of ganciclovir treatment, and immunodeficient patients were excluded. Each subject underwent a laboratory test (complete blood count, total bilirubin, direct bilirubin, AST, ALT, albumin) and liver tissue biopsy conducted by the Pediatric Hepatology consultant. Biopsy of liver tissue was stored in a tube then sent to the Pathology Department of Dr. Soetomo Hospital Surabaya and Tropical Disease Airlangga University. The study protocol was approved by the Ethical Commission of Health Research of Dr. Soetomo Hospital with number No.729/Panke.KKE/XII/2017.

Polymerase chain reaction examination was carried out by extracting DNA using QIAampDNA Mini Kit (Qiagen) from a liver biopsy and based on manual according to the kit. Beta-globin genes were identified using PC03 + and PC04 + primers with the ability to produce 110 bp products with certain sequences (Table 1). Cytomegalovirus was identified by nested PCR with primer MIE4 and MIE5 for first-cycle which produced 435bp; while IE1 and IE2 products were used for the second cycle which produced 161bp.

Polymerization chain reaction mixtures required for β globin were mastermix (Promega): 10 μ L/reaction, FWD primer (PC03 +) 10pmol: 1 μ L/reaction, primary REV (PC04 +) 10pmol: 1 μ L/reaction, ddH₂O (water): 5 μ L/reaction, DNA template: 3 μ L/reaction, with PCR conditions as follows: initial denaturation at 94°C for 5 minutes, denaturation at 94°C for 30 sec, annealing at 55°C for 30 seconds, elongation at 72°C for 45 seconds, final elongation at 72°C for 7 minutes. All of this process was performed 40 cycles.

Four microliters were taken from first-cycle products for second-cycle PCR. Polymerase chain reaction mixture in second-cycle was the same as first-cycle, the differences were only in the product used. The PCR product was visualized by electrophoresis in 2% agarose gel, stained with ethidium bromide, and viewed under ultraviolet light.

Data were collected and presented as a written

explanation, tabulation, and diagrams. Descriptive analysis was used to calculate the number of BA and non-BA cases, the number of CMV infection in BA and non-BA cholestatic infants, and compare the number of CMV infection between BA and BA patients with the Chi-Square test.

RESULTS AND DISCUSSION

There were 37 cholestasis infants involved in this study, dominated by 21 male infants an average age of 2.9 (SD 1.28) months and an average body weight of 4632 (SD 1070) gram. Most patients live outside of Surabaya. Table 2 showed the basic characteristics of pediatric patients with cholestasis. It can be seen that the number of cholestatic infants with BA was smaller (43.2%) compared to without BA (56.8%).

Table 2. Baseline characteristic of cholestasis infant

Characteristic	
Age, mean (\pm SD)	2.9 (\pm 1.28)
Age, n (%)	
1 month	4 (10.8)
2 month	14 (37.9)
3 month	7 (18.9)
4 month	6 (16.2)
5 month	6 (16.2)
6 month	0 (0)
Birth weight, mean (\pm SD)	4632,4 (\pm 1070,06)
Gender (%)	
Girl	16 (43.2)
Boy	21(56.8)
Gestationalage, n (%)	
Aterm	25 (67.62)
Preterm	12 (32.4)
Type of cholestasis, n (%)	
Biliary atresia	16 (43.2)
Non-biliary atresia	21 (56.8)
Residence, n (%)	
Surabaya	11 (29.7)
Outside of Surabaya	26 (70.3)

Table 1. The primer used with the sequence and its product

Primer	Sequence	Product
MIE4	5'-CCA AGC GGC CTC TGA TAA CCA AGC C-3'	435bp
MIE5	5'-CAG CAC CAT CCT CCT CTT CCT CTG G-3'	435bp
IE1	5'- CCA CCC GTG GTG CCA GCT CC-3'	161bp
IE2	5'-CCC GCT CCT CCT GAG GAC CC-3'	161bp
PC03+	5'-CCT CTG ACA CAA CTG TGT TCA CTA GC-3'	110bp
PC04+	5'-TCA CCA CCA ACT TCA TCC ACG TTC ACC-3'	110bp

This study aimed to compare the incidence of CMV infection in BA and non-BA cholestatic infants. In addition to clinical manifestation, histopathological examination of liver tissue biopsy was used to distinguish between BA and non-BA cholestatic infants.¹¹ Biliary atresia is typically characterized by biliary duct proliferation, bile plugs, and portal tract edema/fibrosis in biopsy liver. A study showed that liver biopsy had a sensitivity, specificity, and accuracy of 88.2%.¹² Similar to study by Lee and Looi, biliary duct proliferation in BA showed 95% sensitivity and 88% specificity, while bile plugs showed 68% sensitivity and 86 % specificity.¹³ Russo *et al.* found significant differences between BA and non-BA, indicated by more severe biliary duct proliferation, bile plugs in the ductus and canaliculi and portal fibrosis in BA cases.¹⁴

In this study, BA cases were predominantly found in female infants compared to male infants. Contrastingly, non-BA cases in male infants were

higher compared to female infants. There were significant differences in body weight, age, birth weight and gestational age (Table 3).

The study by Bellomo-Brandao *et al.* found that from 165 infants, intrahepatic cholestasis was found in 62.64% male infants, while extrahepatic cholestasis was found in 55.25% female infants with p-value = 0.026.¹⁵ This finding was similar to this study; despite no significant differences were found, this study showed that BA or extrahepatic cholestasis was commonly found in female infants.

This study showed significant differences in birth weight between BA and non-BA cholestatic infants. The birth weight of BA infants was greater than non-BA infants, indicated by birth weight > 2500 grams was more commonly found in BA infants. This finding was similar to previous research suggesting that higher body weight and greater length at birth were found in children with extrahepatic cholestasis.¹⁵ A study by Fischler *et al.*

Table 3. Characteristics of BA and non-BA infants

Characteristic	Biliary Atresia (BA) (n=16)	Non-Biliary Atresia (Non-BA) (n=21)	P
Age, n (%)			
Boy	7 (33.3)	14(66.7)	0
Girl	9 (56.3)	7 (33.7)	0.163*
Birth weight, mean (± SD)	5218.7 (±926.08)	4185(±966.07)	
Age (month), mean (± SD)	3.5 (±1.15)	2.43 (±1.21)	0.002**
Age (month), n (%)			
One	0 (0)	4 (100.0)	0.009***
Two	4 (28.6)	10 (71.4)	
Three	4 (57.1)	3 (42.9)	
Four	4 (66.7)	2 (33.3)	
Five	4 (66.7)	2 (33.3)	0.111*
Gestational age, n (%)			
Aterm	15 (60.0)	10(40.0)	
Preterm	1 (8.3)	11 (91.7)	0.003*
Birth weight, Mean (± SD)	2953.1 (±295.22)	2542.4 (±684.16)	0.020**
Birth weight, n (%)			
< 2500	1 (9.1)	10 (90.9)	
>2500	15 (57.7)	11 (42.3)	0.006*
IgM CMV n (%)			
Positive	9 (69.2)	4 (30.8)	0
Negative	7 (29.2)	17 (70.8)	.019*
IgG CMV n (%)			
Positive	15 (45.5)	18 (54.5)	0.435*
Negative	1 (25.0)	3 (75)	

SD= Standard Deviation *Chi-Square, ** independent t-test, *** Mann-Whitney U

found that preterm birth was superior in children with BA 3/30 (10%) compared to those without BA 5/55 (9%). However, this difference was not significant. Also, this study found that preterm birth was more frequently found in non-BA infants.¹⁶

The older age was found in BA infants compared to non-BA infants when patients were admitted to the hospital. This was different from the previous study which found that older infants were found in cholestatic patients without BA, supported by other studies which found that there were no significant differences between both groups.^{15,16}

There were nostatistically significant differences in the onset of jaundice between cholestatic infants with BA and without BA, the onset of the yellow appearance of BA infants was longer than non-BA infants (Table 4).This was a contrast to previous

studies which found the remarkably quick onset of jaundice in patients with BA.¹⁷ The difference of the onset of jaundice in these two groups could have been different based on parents' perspective and knowledge. Parents/families sometimes do not know that the children have pathologic icteric and patients with BA still show good nutritional status at the onset of the disease.

On laboratory examination, significant differences in leukocytes count were found between the two groups (Table 4); whereas there were no significant differences in laboratory results, such as complete blood count (hemoglobin, platelets), liver function (ALT/AST), albumin, direct bilirubin, and total bilirubin levels. Higher leukocyte count was found in patients with BA and in accordance with these findings, Wibowo reported comparable

Table 4. Clinical manifestation of BA and non-BA infants

Clinical manifestation	Mean (±SD)		p
	Biliary atresia	Non-biliary atresia	
Onset of jaundice	3.4 (±0.44)	3.7 (±0.39)	0.259**
Direct bilirubin	31.4(±26.62)	25.1(±29.29)	0.646*
Total bilirubin	9.1 (±4.59)	9.6 (±4.86)	0.968**
Hemoglobin	8.4 (±6.2)	8.5 (±6.11)	0.963**
Leukocyte	10.5 (±2.07)	11.2 (±2.88)	0.034**
Platelet	14640 (±5844)	10469 (3682)	0.304*
AST	355.583(±145.052)	371.444 (156.949)	0.63**
ALT	246.2 (±95.39)	215.8 (±161.66)	0.101**
Albumin	213.9 (±133.73)	164.9(±114.27)	0.063*




Figure 1. Histopathological features of liver tissue biopsy with a 400x magnification of a microscope. **Figure A** shows a picture of the portal track of intrahepatic cholestasis as indicated by the presence of giant cell hepatitis (red arrow) and no bile duct proliferation. **Figure B** shows a picture of BA (extrahepatic cholestasis) as biliary ducts proliferation that contains a bile plug (blue arrow) in the portal tract.




Figure 2. (A) Electrophoresis of β globin PCR gene using PCO3 and PCO4 primers (in 8 samples) which produced 110bp products followed by (B) electrophoresis PCR results using primers MIE4 and MIE5 which produced 435 bp

Table 5. Comparison of CMV infection in cholestatic infants with BA and without BA based on the results of PCR in liver tissue

Liver Biopsy PCR CMV, n (%)	Biliary Atresia	Non-Biliary Atresia	p
Positive	9 (37.5)	15(62.5)	0.338*
Negative	7 (53.8)	6 (46.2)	

results, although the increase in leukocytes in BA remained unexplained.¹⁸

The diagnosis of BA was based on clinical manifestations (yellowing of the eyes and whole body, a cholic stool) and anatomical pathology examination (histopathological features such as bile plug, ductular proliferation, and portal edema with and/or fibrosis of liver biopsy tissue (Figure 1).

Biopsy samples were taken and extracted from liver tissue, then PCR was carried out with β globin using PCO3 primers to determine the quality of the samples. If a positive result (yielding a product of 100-200 bp) is obtained, the examination must be continued by PCR examination using primers MIE4 and MIE5. A positive result is reported if the product produces 400-500bp. In this study, positive results were obtained for all β globin effects; therefore, PCR was performed (Figures 2).

The detection of CMV by PCR showed positive results in 24 infants and negative results in 13 patients. The incidence of CMV infection in cholestatic infants with BA and without BA was 56.2% and 71.4%, respectively. The polymerase chain reaction is a diagnostic instrument that has high sensitivity and ability to detect the presence of CMV, despite low specificity and low CMV infection.¹⁹

Cytomegalovirus infection is initially more common in intrahepatic cholestasis (without BA);

however, several studies have shown that CMV infection can be found in extrahepatic cholestasis (BA). The study found that viruses including CMV can be a trigger leading to dysregulation of immune mechanisms with genetic influences and eventually cause BA.²⁰ Cytomegalovirus infection has the ability to replicate both in hepatocytes and cholangiocytes. This virus can directly induce damage to the liver and biliary duct system and induce damage to the immune system in infected cells, leading to the formation of inclusion of bodies in hepatocyte and vascular cells of epithelial cells, especially along with biliary duct epithelial cells.²¹

There were no significant differences between BA and non-BA cholestatic infants based on the PCR of CMV in liver tissue (Table 5). This study showed that positive PCR results of CMV were only found in 9 (37.5%) BA patients. The study about PCR CMV was begun by conducting several studies on animals, and subsequently was carried out on humans.²²⁻²⁵ In this study, there were no significant differences in the number of CMV infection in infants with BA and without BA. This was because BA could be caused by other viruses such as Rotavirus, Reovirus, Epstein-Barr Virus (3.5%) and Adenovirus (5.8%).^{26,27,9} Presumed role of Rotavirus and Reovirus in BA have also been studied for a long time.

Fjaer *et al.* found 4 cases of CMV infection from a

total of 9 cholestatic patients. However, positive PCR of CMV from liver tissue was only found in 2 patients from 4 cases of CMV infection. Cytomegalovirus infection in the other 2 patients was caused by Epstein-Barr Virus.²⁸ The presence of the Human Herpes virus 6 in liver tissue was also demonstrated by Domiati *et al.* in their study about Human Herpes six virus in BA patients. Cytomegalovirus infection; however, was not found in the study subjects or controls.²⁹

CONCLUSION AND SUGGESTION

Cytomegalovirus infection is found in intrahepatic and extrahepatic cholestatic infants like BA. In this study, the lower incidence of CMV infection in cholestatic infants with BA was found compared to non-BA. However, there was no significant difference in the incidence of CMV infection in cholestatic infants with BA or without BA. Future research with longer research time and PCR was needed to determine the causal virus of BA.

REFERENCES

- Santos JL, Carvalho E, Bezerra JA. Advances in biliary atresia: From patient care to research. *Braz. J. Med. Biol. Res.* 2010; 43(6): 522-527.
- Bassett MD, Murray KF. Biliary atresia: Recent progress. *J. Clin. Gastroenterol.* 2008; 42(6): 720-729.
- Moreira RK, Cabral R, Cowles RA, Lobritto SJ. Biliary atresia: A multi disciplinary approach to diagnosis and management. *Arch. Pathol. Lab. Med.* 2012; 136(7): 746-760.
- Sira MM, Salem TAH, Sira AM. Biliary atresia: A challenging diagnosis. *Global Journal of Gastroenterology.* 2013; 1: 24-35.
- Oliveira NL, Kanawaty FR, Costa SC, Hessel G. Infection by Cytomegalovirus in patients with neonatal cholestasis. *Arq. Gastroenterol.* 2002; 39(2): 132-136.
- Buonsenso D, Serranti D, Gargiullo L, Ceccarelli M, Ranno O, Valentini P. Congenital Cytomegalovirus infection: Current strategies and future perspectives. *Eur Rev Med Pharmacol Sci.* 2012; 16(17): 919-935.
- Caliendo AM, Schuurman R, Yen-Lieberman B, Spector SA, Andersen J, *et al.* Comparison of quantitative and qualitative PCR assays for Cytomegalovirus DNA in plasma. *J Clin Microbiol.* 2001; 39(4): 1334-1338.
- Mack CL, RJ Sokol. Unraveling the pathogenesis and etiology of biliary atresia. *Pediatr Res.* 2005; 57: 87-94.
- XuY, YuJ, ZhangR, Yin Y, Ye J, *et al.* The perinatal infection of Cytomegalovirus is an important etiology for biliary atresia in China. *Clin. Pediatr (Phila).* 2012; 51(2): 109-113.
- Rauschenfels S, Krassmann M, Al-Masri AN, Verhagen W, Leonhardt J, *et al.* Incidence of hepatotropic viruses in biliary atresia. *Eur.J. Pediatr.* 2009; 168(4): 469-476.
- Rashed YK, Saber MA, Tawfik M, Mourad WS. Histopathological features and accuracy for diagnosing biliary atresia by prelaparotomy liver biopsy in Egypt. *Egyptian. Pediatric Association Gazette.* 2013; 61: 42-45.
- Rastogi AN, Krishnani K, Yachha V, Khanna U, Poddar, *et al.* Histopathological features and accuracy for diagnosing biliary atresia by prelaparotomy liver biopsy in developing countries. *J. Gastroenterol Hepatol.* 2009; 24(1): 97-102.
- Lee WS, Looi LM. Usefulness of a scoring system in the interpretation of histology in neonatal cholestasis. *World J. Gastroenterol.* 2009; 15(42): 5326-5333.
- Russo P, Magee JC, Boitnott J, Bove KE, Raghunathan T, *et al.* Design and validation of the biliary atresia research consortium histologic assessment system for cholestasis in infancy. *Clinical Gastroenterology and Hepatology: American Gastroenterological Association.* 2011; 9(4): 357-362.
- Bellomo-Brandao MA, Arnaut LT, Tommaso AM, Hessel G. Differential diagnosis of neonatal cholestasis: Clinical and laboratory parameters. *J. Pediatr.* 2010; 86(1): 40-44.
- Fischler B, Woxenius S, Nemeth A, Papadogiannakis N. Immunoglobulin deposits in liver tissue from infants with biliary atresia and the correlation to cytomegalovirus infection. *J. Pediatr. Surg.* 2005; 40(3): 541-546.
- Bazlul Karim AS, Kamal M. Cholestasis jaundice during infancy: experience at a tertiary-care center in Bangladesh. *Indian. J. Gastroenterol.* 2005; 24(2): 52-54.
- Wibowo S, Santoso NB. Karakteristik klinik dan laboratorik kolestasis intrahepatal dan ekstrahepatal di bangsal perawatan anak RSUD Dr. Saiful Anwar Malang. *M. Med. Indones.* 2012; 46(2): 108-114.
- Jahan M. Laboratory diagnosis of CMV infection: A review. *Bangladesh J. Med. Microbiol.* 2010; 4(2): 39-44.
- Mack CL. The pathogenesis of biliary atresia: Evidence for a virus-induced autoimmune disease. *Semin. Liver Dis.* 2007; 27(3): 233-242.
- Lazim HH, Kadhim HS, Arif H, Al Khafaji KR. The association between biliary atresia and Cytomegalovirus hepatitis. *J.Nepal Paediatr. Soc.* 2015; 35(3): 269-274.
- Wang W, Zheng S, Shong Z, Zhao R. Developmental of a guinea pig model of perinatal Cytomegalovirus induced hepatobiliary injury. *Fetal. Pediatr. Pathol.* 2011; 30(5): 301-311.
- Wang W, Donnelly B, Bondoc A. The rhesus rotavirus gene encoding VP4 is a major determinant in the pathogenesis of biliary atresia in newborn mice. *J. Virol.* 2011; 85(17): 9069-9077.
- Soomro GB, Abbas Z, Hassan M, Luck N, Memon Y, Khan AW. Is there any association of extrahepatic biliary atresia with cytomegalovirus or other

- infections?. J. Pak. Med. Assoc, 2011; 61(3): 281-283.
25. Yaghobi R, Didari M, Gramizadeh B. Study of viral infections in infants with biliary atresia. Indian J. Pediatr, 2011; 78(4): 478-481.
 26. Von Sochaczewski CO, Pintelon I, Brouns I, Dreier A, Klemann C, *et al.* Rotavirus particles in the extrahepatic bile duct in experimental biliary atresia. J. Pediatr. Surg, 2014; 49(4): 520-524.
 27. Tyler KL, RJ Sokol, SM Oberhaus. Detection of reovirus RNA in hepatobiliary tissues from patients with extrahepatic biliary atresia and choledochal cysts. Hepatology, 1998; 27(6): 1475-1482.
 28. Fjaer RB, Bruu AL, Nordbo. Extrahepatic bile duct atresia and viral involvement. Pediatr. Transplant, 2005; 9(1): 68-73.
 29. Domiati-Saad R, Dawson DB, Margraf LR. Cytomegalovirus and human herpesvirus 6, but not human papillomavirus, are present in neonatal giant cell hepatitis and extrahepatic biliary atresia. Pediatr. Dev. Pathol, 2000; 3(4): 367-373.



INDONESIAN JOURNAL OF CLINICAL PATHOLOGY AND MEDICAL LABORATORY (IJCPML)

📍 [PERHIMPUNAN DOKTER SPESIALIS PATOLOGI KLINIK INDONESIA](#)

🌟 P-ISSN : 08544263 <> E-ISSN : 24774685



0

Impact Factor



601

Google Citations



Sinta 2

Current Accreditation

[🔍 Google Scholar](#) [📖 Garuda](#) [🌐 Website](#) [🔗 Editor URL](#)

History Accreditation

2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026

Garuda [Google Scholar](#)

[Immunogenicity Assessment on Clinical Trials of SARS-CoV-2 Vaccines](#)

Indonesian Association of Clinical Pathologist and Medical laboratory

[INDONESIAN JOURNAL OF CLINICAL PATHOLOGY AND MEDICAL LABORATORY Vol](#)

[28, No 2 \(2022\) 202-208](#)

📅 2022

📄 DOI: [10.24293/ijcpml.v28i2.1975](#)

🏆 [Accred : Sinta 2](#)

[Concordance Test of Various Erythrocyte Indices for Screening of Beta Thalassemia Carrier](#)

Indonesian Association of Clinical Pathologist and Medical laboratory

[INDONESIAN JOURNAL OF CLINICAL PATHOLOGY AND MEDICAL LABORATORY Vol](#)

[28, No 2 \(2022\) 137-142](#)

📅 2022

📄 DOI: [10.24293/ijcpml.v28i2.1842](#)

🏆 [Accred : Sinta 2](#)

[Gene Expression of SOX2, OCT4, and Nanog by Small Molecule Compound VC6TFZ on Peripheral Blood Mononuclear Cell](#)

Indonesian Association of Clinical Pathologist and Medical laboratory

[INDONESIAN JOURNAL OF CLINICAL PATHOLOGY AND MEDICAL LABORATORY Vol](#)

[28, No 2 \(2022\) 115-120](#)

📅 2022

📄 DOI: [10.24293/ijcpml.v28i2.1759](#)

🏆 [Accred : Sinta 2](#)

[Correlation between Inflammatory Markers of Platelet Index and Vitamin D with Body Mass Index](#)

Indonesian Association of Clinical Pathologist and Medical laboratory

[INDONESIAN JOURNAL OF CLINICAL PATHOLOGY AND MEDICAL LABORATORY Vol](#)

[28, No 2 \(2022\) 161-164](#)

📅 2022

📄 DOI: [10.24293/ijcpml.v28i2.1816](#)

🏆 [Accred : Sinta 2](#)



**KOMITE ETIK PENELITIAN KESEHATAN
RSUD Dr. SOETOMO SURABAYA**

**KETERANGAN KELAIKAN ETIK
(" ETHICAL CLEARANCE ")**

729 / Panke.KKE / XII / 2017

KOMITE ETIK RSUD Dr. SOETOMO SURABAYA TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA PENELITIAN DENGAN JUDUL :

" Kesesuaian Pemeriksaan Serologi CMV dan PCR CMV Jaringan Hati pada Kolestasis Bayi "

PENELITI UTAMA : Dr. Bagus Setyoedi, dr., Sp.A (K)

**PENELITI LAIN : 1. Sjamsul Arief, dr., Sp.A (K), MARS
2. Dwiyanti Puspitasari, dr., DTM&H, MCTM (TP), Sp.A (K)
3. Dr. Hari Basuki N, dr., M.Kes
4. Dr. Gondo Mastutik, drh., M.Kes
5. Reny Widayanti, dr**

UNIT / LEMBAGA / TEMPAT PENELITIAN : RSUD Dr. Soetomo Surabaya

DINYATAKAN LAIK ETIK

**Berlaku dari : 19 DEC 2017 s.d 19 DEC 2018
SURABAYA, 19 DEC 2017**

KETUA

(Dr. Elizeus Hanindito, dr., Sp.An, KIC,KAP)
NIP. 19511007 197903 1 002

