

Effect of Corrots (Daucus carota L.) on Gastric Histopathology of Piroxicam-Induced Mice as a Peptic Ulcer Prevention

by Jennifer Susanto

Submission date: 04-Oct-2022 05:06PM (UTC+0800)

Submission ID: 1916295297

File name: ology_of_Piroxicam-Induced_Mice_as_a_Peptic_Ulcer_Prevention.pdf (379.4K)

Word count: 2434

Character count: 13457

Effect of Carrots (*Daucus carota* L.) on Gastric Histopathology of Piroxicam-Induced Mice as a Peptic Ulcer Prevention

Jennifer Susanto¹, Danti Nur Indiatuti², Gondo Mastutik³

¹Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

²Department of Anatomy, Histology, and Pharmacology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

³Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

ABSTRACT

Introduction: Non-steroidal anti-inflammatory drugs (NSAIDs) consumption contributes significantly to gastrointestinal morbidity, including peptic ulcers. NSAIDs cause gastric mucosal damage by inhibiting PGE2 and PGI2 secretion, increasing HCl secretion, and also causing local irritation and neutrophil adhesion. Flavonoids and chlorogenic acid are gastroprotective agents found in carrots (*Daucus carota* L.). Therefore, this study aimed to validate the effect of carrots on gastric histopathology of piroxicam-induced mice.

Methods: This was an experimental study using a post-test-only control group design. 35 mice were divided into 5 groups and were allowed to adapt for 1 week. The negative control group received 0.25 mL aqua dest, while the positive control group received 0.104 mg famotidine. Group A, B, and C received carrot extract with each dosage was 200 mg/kgBW, 300 mg/kgBW, and 400 mg/kgBW. The mice were induced 2 hours later with 0.052 mg piroxicam. The treatments lasted 14 days. Each stomach was taken and examined on the 15th day. Afterward, histopathological preparations were made.

Results: The results of the histopathological signs of inflammation using the Kruskal-Wallis method showed $p = 0.000$, indicating that there was an effect of the carrot extract in preventing inflammation on the gastric histopathology ($p < 0.05$).

Conclusion: In conclusion, this study confirmed that carrot extract is more effective in avoiding peptic ulcers due to NSAID consumption.

* Correspondence: danti-n-i@fk.unair.ac.id

JUXTA: Jurnal Ilmiah Mahasiswa Kedokteran Universitas Airlangga

p-ISSN: 1907-3623; e-ISSN: 2684-9453

DOI: 10.20473/juxta.V13i12022.88-92

Open access under Creative Commons Attribution-ShareAlike 4.0 International License (CC-BY-SA)

ARTICLE INFO

Article history:

Received 2 June 2022

Received in revised form
28 June 2022

Accepted 27 July 2022

Available online 10 August 2022

Keywords:

Carrot,
Famotidine,
Health risk,
Peptic ulcer,
Piroxicam.



Introduction

Non-steroidal anti-inflammatory drugs (NSAID) have common side effects that damage the digestive tract mucosa through inhibition of cyclooxygenase-1 (COX-1) enzyme which can reduce the secretion of gastric mucosal cytoprotective materials, especially PGE2 and PGI2, through local irritation due to direct contact with gastric mucosa,¹ through inhibition of cyclooxygenase-2 (COX-2) enzyme which can increase neutrophil adhesion,² through reactive oxygen species (ROS) production,^{3,4} and also through its prominent acidic nature.⁴ Based on 2013 Basic Health Research in Indonesia, there were 24,496 NSAIDs stored by 20,516 households purchased without a prescription. One of them is piroxicam which has side effects of peptic ulcers and gastrointestinal bleeding up to 9.5 times higher than others.^{5,6}

Peptic ulcers are ulceration of the gastric or duodenum mucosa.⁷ The best therapy is proton pump inhibitor (PPI). However, PPI is an acid-labile drug and only works after being absorbed in the intestine,⁵ thus the H2 receptor antagonist (H2RA) was the drug used in this study because it directly inhibits the H2 receptor. Famotidine, for example, was effective in preventing peptic ulcers in indomethacin-induced rats by reduction of gastric degenerative changes, gastric acidity, and ROS production.⁸ However, PPI and H2RA have side effects such as diarrhea and headache.⁵ Therefore, carrots (*Daucus carota L.*) which contain anti-inflammatory substances, known as phenolic substances, can be used as an alternative to prevent peptic ulcers.⁸ The most dominant phenolic compound in carrots is chlorogenic acid which can reduce the release of inflammatory mediators, especially TNF- α .^{9,10} Carrots also contain flavonoids that inhibit acid secretion, reduce pepsin level and activity, and also increase gastric mucous and bicarbonate secretion. Flavonoids also have antioxidative, anti-inflammatory, and cytoprotective effect.^{11,12} Therefore, this study aimed to prove the effect of carrot extract as a peptic ulcer prevention on piroxicam-induced mice using histopathological signs of inflammation as the measurement.

Methods

This study used an experimental method with a post-test-only control group design that randomized mice into groups consisting of 2 control groups and 3 experimental groups.

Drugs

The dosage of the piroxicam used in this study was 20 mg \times 0.0026 = 0.052 mg/treatment/mice (2 tablets of piroxicam were crushed). 25.48 mg of piroxicam were taken and dissolved in 122.5 ml aqua dest. The dosage of the famotidine used in this study was 40 mg \times 0.026 = 0.052 mg/treatment/mice (1 tablet of famotidine was crushed). 10.192 mg of famotidine were taken and dissolved in 24.5 ml aqua dest. 0.0026 was the constant surface-area ratio from man to mouse.¹³

Carrot Extract

Fresh carrots were pureed with methanol using a blender. Then the juice was put in a place to be macerated for 3 days. Afterward, the juice was filtered to separate the pulp and the filtrate. The filtrate was evaporated using a rotary evaporator to evaporate the methanol. Carrot extract was dissolved in aqua dest.

Animals

The sample population of this study was male mice (*Mus musculus*) with an average weight of 25 grams, aged 2-3 months, and healthy. Samples that died during the trial period were excluded. The animals were provided by Department of Anatomy, Histology, and Pharmacology Faculty of Medicine Universitas Airlangga Surabaya. The animals were housed in standard cages and supplemented with food and water ad libitum. After 7 days of adaptation, the mice were randomly divided into 5 groups and were treated as follows: (negative control) aqua dest 0.25 ml; (positive control) famotidine 0.104 mg; (A) 200 mg/kg BW carrot extract; (B) 300 mg/kg BW carrot extract; and (C) 400 mg/kg BW carrot extract. 2 hours later, all mice received 0.052 mg piroxicam. All treatments were administered orally. The treatments were given for 14 days. On the 15th day, the mice were examined and the stomachs were removed. All procedures were approved by the ethics committee of Faculty of Medicine Universitas Airlangga Surabaya

Histology Preparation

The stomach was opened along the greater curvature. Then, the part that had rugae (corpus to antrum) was given 10% formalin solution, dehydrated through increasing the grades of alcohol, printed into a paraffin block, and stored in the refrigerator. Paraffin blocks were cut to a thickness of 5 μ m. The pieces floated in warm water with a temperature of 60°C for 24 hours. The preparation was then removed and stained with hematoxylin-eosin (HE) solution.

Determination of Histopathological Signs of Inflammation

Each preparation was viewed in 3 fields per specimen (400x).

Score 0: normal gastric mucosa

Score 1: signs of inflammation of the gastric mucosa, such as hyperemia and edema, accompanied by neutrophils on the basement membrane, but the epithelial surface was still intact and there were no lymphocytes or plasma cells in the lamina propria

Score 2: erosion or bleeding on the mucosal surface

Score 3: erosions that penetrate the mucosa or transmural bleeding

Score 4: necrotic tissue in the basal lamina formed by granulation tissue¹⁴

Statistical Analysis

For histopathology the inflammatory sign, this study used the Kruskal-Wallis method because the data measurement scale was an ordinal. The results were displayed in tables.

Results

Table 1. Mean value and the Kruskal-Wallis test result of histopathological score

Group	Histopathological Score	Asymp. sig. (p)
Negative Control	1.95 ± 0.26	0.000
Positive Control	1.67 ± 0.29	
A	0.84 ± 0.07	
B	0.50 ± 0.14	
C	0.39 ± 0.10	

Table 1 shows that the severity of the inflammation decreased as the dosage increasing. There was an effect of treatment given on the histopathological score of inflammation signs. Similarly, this study used post hoc Mann-Whitney test to know which groups were significantly different. The results showed that group A and group B were not significantly different, the same with group B and group C.

Below are some pictures from the histopathological score of inflammation signs measurement:




Figure 1. Negative Control (Score 3)




Figure 2. Positive Control (Score 2)




Figure 3. Group A (Score 1)




Figure 4. Group B (Score 1)




Figure 5. Group C (Score 0)

Discussion

On the histopathological signs of inflammation, all doses of carrot extract were more effective than famotidine. However, the difference in histopathological scores between group A and B and group B and C was less significant because of the close difference in dosage. The dosage used in the previous study were 150 mg/kg BW and 300 mg/kg BW, the results showed that ulcer index and

curative ratio percentage were significantly different.¹⁵ Therefore, carrot extract is more effective at preventing peptic ulcers than famotidine. This is caused by the antioxidant and gastroprotection compounds found in carrots such as flavonoids and chlorogenic acid.^{8,16}

Flavonoids can increase gastric prostaglandins, reduce H⁺ secretion, reduce free radicals, and regulate both pro and anti-inflammatory cytokines.^{12,16,17} One of the flavonoids in carrots is flavonols. There are several substances that are included in the flavonol class, one of which is quercetin. Quercetin can inhibit gastric acid secretion and lipid peroxidation. It also can increase mucous bicarbonate secretion.¹⁸

On the other hand, chlorogenic acid can also reduce gastric tissue lesions through several ways. First, it can decrease neutrophil migration and infiltration to the lesion area. Neutrophils are the first leukocytes to enter the area of inflammation and secrete chemicals to destroy the damaging agent. Thus, it can damage the host tissue.^{10,19} Second, it can decrease local secretion of pro-inflammatory mediators by modulating the key transcription factors.^{10,20} Lastly, it can decrease free radicals production. It can restore enzyme function and glutathione levels that can be reduced during inflammation.¹⁰

In comparison to carrot extract, famotidine is the most potent H₂RA. In previous studies, famotidine could reduce the side effect of NSAID consumption.^{5,8} However, the results showed that it was not as effective as carrot extract due to its only mechanism of action that reduce aggressive factors but does not increase defensive factors.^{8,21} If the use of NSAIDs should be continued, then PPI is a better choice than H₂RA.⁸

Conclusion

Carrot extract at a dose of 200 mg/kg, 300 mg/kg, and 400 mg/kg of body weight can prevent inflammation better than famotidine 0.104 mg for the histopathological signs of inflammation. In conclusion, carrot extract is more effective in preventing peptic ulcers due to NSAIDs consumption than famotidine.

Acknowledgments

The authors are extremely thankful to all staff in Department of Anatomy, Histology, and Pharmacology and Department of Pathology Anatomy Faculty of Medicine Universitas Airlangga for the help in this study.

Conflict of Interest

The authors declared there is no conflict of interest.

References

- Grosser T, Smyth E, FitzGerald GA. Anti-inflammatory, Antipyretic, and Analgesic Agents; Pharmacotherapy of Gout. In: Brunton LL, Chabner BA, Knollmann BC (eds) *Goodman & Gilman's: The Pharmacological Basis of Therapeutics, 12e*. New York, NY: McGraw-Hill Education, <http://accessmedicine.mhmedical.com/content.aspx?aid=1127868620> (2015).
- Habeeb A, Tiwari SK, Bardia A, et al. Peptic Ulcer Disease: Descriptive Epidemiology, Risk Factors, Management and Prevention. *SMGroup*, <http://www.smgebooks.com/peptic-ulcer-disease/chapters/PUD-16-06.pdf> (2016).
- Firdaus SB, Ghosh D, Chattopadhyay A, et al. Protective Effect of Antioxidant Rich Aqueous Curry Leaf (*Murraya koenigii*) Extract Against Gastro-Toxic Effects of Piroxicam in Male Wistar Rats. *Toxicol Reports* 2014; 1: 987–1003. [PubMed] [CrossRef]
- Nabil M, El Raey MA, Abdo W, et al. Gastro-Protective Effects of Albizia anthelmintica Leaf Extract on Indomethacin-Induced Gastric Ulcer in Wistar Rats: In Silico and In Vivo Studies. *Antioxidants* 2021; 10: 176. [PubMed] [CrossRef]
- Soleha M, Isnawati A, Fitri N, et al. Profil Penggunaan Obat Antiinflamasi Nonstereoid di Indonesia. *J Kefarmasian Indones* 2018; 109–117. [CrossRef]
- Katzung BG, Masters SB, Trevor AJ. *Farmakologi Dasar & Klinik*. 12th ed. Jakarta: EGC, 2012.
- Chan F, Lau J. *Peptic Ulcer Disease*. 10th ed. Philadelphia: Elsevier Saunders, 2016.
- Leja M, Kamińska I, Kramer M, et al. The Content of Phenolic Compounds and Radical Scavenging Activity Varies with Carrot Origin and Root Color. *Plant Foods Hum Nutr* 2013; 68: 163–170. [PubMed] [CrossRef]
- Arscott SA, Tanumihardjo SA. Carrots of Many Colors Provide Basic Nutrition and Bioavailable Phytochemicals Acting as a Functional Food. *Compr Rev Food Sci Food Saf* 2010; 9: 223–239. [CrossRef]
- López-Gámez G, Elez-Martínez P, Martín-Belloso O, et al. Applying Pulsed Electric Fields to Whole Carrots Enhances the Bioaccessibility of Carotenoid and Phenolic Compounds in Derived Products. *Foods* 2021; 10: 1321. [PubMed] [CrossRef]
- Shimoyama AT, Santin JR, Machado ID, et al. Antiulcerogenic Activity of Chlorogenic Acid in Different Models of Gastric Ulcer. *Naunyn Schmiedebergs Arch Pharmacol* 2013; 386: 5–14. [PubMed] [CrossRef]
- Zhang W, Lian Y, Li Q, et al. Preventative and Therapeutic Potential of Flavonoids in Peptic Ulcers. *Molecules* 2020; 25: 4626. [PubMed] [CrossRef]
- Paget G, Barnes J. *Evaluation of Drug Activities: Pharmacometrics*. London: Academic Press, 1964.
- Kumar V, Abbas A, Aster J. *Robbins Basic Pathology*. Philadelphia: Elsevier Saunders, 2013.
- Wehbe K, Mroueh M, Daher CF. The Potential Role of Daucus carota Aqueous and Methanolic Extracts on Inflammation and Gastric Ulcers in Rats. *J Complement Integr Med*; 6. Epub ahead of print 16 January 2009. [CrossRef]
- da Silva Dias JC. Nutritional and Health Benefits of Carrots and Their Seed Extracts. *Food Nutr Sci* 2014; 05: 2147–2156. [CrossRef]
- Serafim C, Araruna ME, Júnior EA, et al. A Review of the Role of Flavonoids in Peptic Ulcer (2010–2020). *Molecules* 2020; 25: 5431. [PubMed] [CrossRef]
- Anand David A, Arulmoli R, Parasuraman S. Overviews of Biological Importance of Quercetin: A



- Bioactive Flavonoid. *Pharmacogn Rev* 2016; 10: 84. [PubMed]
19. Zatorski H, Salaga M, Zielińska M, *et al.* Experimental Colitis in Mice is Attenuated by Topical Administration of Chlorogenic Acid. *Naunyn Schmiedebergs Arch Pharmacol* 2015; 388: 643–651. [PubMed] [CrossRef]
20. Liang N, Kitts D. Role of Chlorogenic Acids in Controlling Oxidative and Inflammatory Stress Conditions. *Nutrients* 2015; 8: 16. [PubMed] [CrossRef]
21. Tjokropawiro A. *Buku ajar ilmu penyakit dalam. Ed.2: Fakultas Kedokteran Universitas Airlangga Rumah Sakit Pendidikan Dr. Soetomo Surabaya.* Surabaya: Airlangga University Press, 2015.

Effect of Corrots (Daucus carota L.) on Gastric Histopathology of Piroxicam-Induced Mice as a Peptic Ulcer Prevention

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

11%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | phcogj.com
Internet Source | 1% |
| 2 | Iqbal Syahputra, Eko Purnanto, Ade Utia Detty, Indra Kumala. "Hubungan Diabetes Melitus Tipe II Dengan Kejadian Benigna Prostat Hiperplasia Di Rumah Sakit Pertamina Bintang Amin Bandar Lampung", MAHESA : Malahayati Health Student Journal, 2022
Publication | 1% |
| 3 | H Anthony, F.C. Nwinyi, A Mohammed. "Effects of Aqueous-Methanol Bulb Extract of Allium sativum on Gastric Ulcer and Gastrointestinal Motility", Journal of Veterinary and Biomedical Sciences, 2022
Publication | 1% |
| 4 | S Ilyas, M Tanjung, I Nurwahyuni, F Y Pasaribu. " Effect of Mahkota Dewa Ethanolic Extracts (Scheff.) to Hepar Histology of Female Rat () In Preeclampsia ", IOP | 1% |

Conference Series: Earth and Environmental Science, 2019

Publication

5	jih.ejournal.unri.ac.id Internet Source	1 %
6	medrech.com Internet Source	1 %
7	rua.ua.es Internet Source	1 %
8	garuda.kemdikbud.go.id Internet Source	1 %
9	medcraveonline.com Internet Source	1 %
10	www.jphres.org Internet Source	1 %
11	www.ncbi.nlm.nih.gov Internet Source	1 %
12	Satish Bhoge, Prabal P. Singh, Deepak K. Das, Yakub Ali, Abhijeet Dhulap. "Design, synthesis, anti-inflammatory evaluation and in silico molecular docking of novel furan-based derivatives with promising TNF- α inhibitory activity", Research Square Platform LLC, 2022 Publication	1 %
13	journal2.unusa.ac.id Internet Source	1 %

14	www.jidmr.com Internet Source	1 %
15	f1000research.com Internet Source	1 %
16	www.researchgate.net Internet Source	1 %
17	André T. Shimoyama, José Roberto Santin, Isabel D. Machado, Ana Mara de Oliveira e Silva et al. "Antiulcerogenic activity of chlorogenic acid in different models of gastric ulcer", <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2012 Publication	<1 %
18	Mohamed A. Lebda, Rabab E. Mostafa, Nabil M. Taha, Eman M. Abd El-Maksoud et al. "Commiphora myrrh Supplementation Protects and Cures Ethanol-Induced Oxidative Alterations of Gastric Ulceration in Rats", <i>Antioxidants</i> , 2021 Publication	<1 %
19	coek.info Internet Source	<1 %
20	jurnal.unpad.ac.id Internet Source	<1 %
21	www.europeanreview.org Internet Source	<1 %

22 www.quantobasta.com <1 %
Internet Source

23 www.veterinaryworld.org <1 %
Internet Source

24 es.scribd.com <1 %
Internet Source

25 riset.unisma.ac.id <1 %
Internet Source

26 "OS02: Tuberculosis 1 : Oral Sessions", <1 %
Respirology, 2013.
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

C23 Effect of Corrots (Daucus carota L.) on Gastric Histopathology of Piroxicam-Induced Mice as a Peptic Ulcer Prevention

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
