

ISSN (printed) : 2088-4559

ISSN (online) : 2477-0256

Pharmaci ana

Accredited by Ministry of Research, Technology and Higher Education, Republic of Indonesia (Decree No. 32a/E/KPT/2017)

Jurnal Kefarmasian

Vol 12. No 2. July 2022



UNIVERSITAS AHMAD DAHLAN
IN COLLABORATION WITH
IKATAN APOTEKER INDONESIA (IAI)



Home > About the Journal > **Editorial Team**

Editorial Team

Editor in Chief

Prof.Dr. Nurkhasanah Mahfudh, M.Si., Apt, Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

Editorial Board

apt., Phebe Hendra, M.Si., Ph.D, Sanata Dharma University, Indonesia

Prof. Dr. apt. Dyah Aryani Perwitasari, M.Si., Ph.D., Fakultas Farmasi, Universitas Ahmad Dahlan, Yogyakarta, Indonesia

Dr. Kenny Chitcholtan, Christchurch School of Medicine and Health Sciences, New Zealand

Prof. Dr. Irwandi Jaswir, International Islamic University Malaysia (IIUM), Malaysia

Dr. drh Sapto Yuliani, MP, Fakultas Farmasi, Universitas Ahmad Dahlan, Yogyakarta, Indonesia

Sukanya Dej-adisai Ph.D, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat-Yai, Songkhla 90110 Thailand, Thailand

Dr. Fezah Binti Othman, Universiti Putra Malaysia, Malaysia

Prof. Dr. apt. Zullies Ikawati, Fakultas Farmasi, Universitas Gadjah Mada, Yogyakarta, Indonesia

Prof. Dr. apt. Abdul Rohman, M.Si., Fakultas Farmasi, Universitas Gadjah Mada Yogyakarta, Indonesia

Prof. Dr. apt. Akhmad Kharis Nugroho, M.Si., Fakultas Farmasi, Universitas Gadjah Mada, Indonesia

Dadan Hermawan, M.Si., Ph.D, Fakultas MIPA, Universitas Jenderal Soedirman, Indonesia

Dr. Moch. Saiful Bachri, Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

Prof. Dr. Achmad Mursyidi, Universitas Ahmad Dahlan, Indonesia

Dr Isnaeni Yudi Haryanto, Fakultas Farmasi, Universitas Airlangga, Indonesia

Dr. apt. Keri Lestari Dandan, M.Si., Fakultas Farmasi, Universitas Padjadjaran, Indonesia

Dr. apt. Laela Hayu Nurani, M.Si., Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

apt., Anita Sukmawati, Ph.D, Fakultas Farmasi, Universitas Muhammadiyah Surakarta, Indonesia

Dr. apt. Tedjo Yuwono, Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

Dr. apt. Nining Sugihartini, M.Si, Fakultas Farmasi, Universitas Ahmad Dahlan, Yogyakarta, Indonesia

Dr. apt. Lannie Hadisoewignyo, M.Si, Fakultas Farmasi, Universitas Katolik Widya Mandala Surabaya, Indonesia

lin Narwanti, MSc., Fakultas Farmasi Universitas Ahmad Dahlan, Indonesia

apt. Nina Salamah, M.Sc, Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

apt Faridah Baroroh, MSc, Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

apt. Susan Fitria Candradewi, M.Sc, Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

apt. Citra Ariani Edityaningrum, M.Si, Fakultas Farmasi, Universitas Ahmad Dahlan

Copy Editor

Aning Setiya Rini, S.T, Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia

Pharmaciana

ISSN Print: 2088-4559 | ISSN Online: 2477-0256

Website: <http://journal.uad.ac.id/index.php/PHARMACIANA>

Office: Faculty of Pharmacy, Universitas Ahmad Dahlan

Jl. Prof. Dr. Soepomo, S.H., Janturan, Warungboto, Umbulharjo, Yogyakarta, Indonesia

Kode pos 55164

Email: pharmaciana@pharm.uad.ac.id



Home > Archives > **Vol 12, No 2 (2022)**

Vol 12, No 2 (2022)

Pharmaciana

Table of Contents

Pandan Wangi rice as reference food and the use of 25 g portion in glycemic index test	PDF
<i>Didah Nur Faridah, Puspo Edi Giriwono, Ratna Sari Listyaningrum, Masayu Nur Ulfa</i>	147-155
In vitro antidiabetic activity of Peperomia pellucida extract and fraction by alpha-amylase inhibition pathway	PDF
<i>Sholihatil Hidayati, Shinta Shinta Mayasari, Lindawati Setyaningrum, Ayu Dwi Wardani, Qurrotul Aini</i>	156-163
The effectiveness of avocado (Persea americana M.) leaves infusion as urinary stones dissolving in rats	PDF
<i>Ni Made Dwi Sandhiutami, Rika Sari Dewi, Alya Fariha</i>	164-172
Incision wound healing test ethanolic extract gel from Ekor Naga (Rhaphidophora pinnata (L.f) Schott) leaves in white male rats	PDF
<i>Fathnur Sanik, Havizur Rahman, Ave Olivia Rahman, Agung Giri Samudra, Cassa de Floris, Muhaimin Muhaimin</i>	173-180
Burn wound healing activity of ethanol extract gel of Green Algae (Ulva lactuca L) in mice	PDF
<i>Wahyu Widyaningsih, Supto Yuliani, Vivi Sofia, Reka Rukmiati, Khozanatul Ulwy</i>	181-187
Free fatty acids regulating action of Capparis decidua fruit on dyslipidemia in rats	PDF
<i>Avijit Saha, Sangeeta Ghosh, Alok K. Hazra, Sandip Ghosh, Tapas Kumar Sur</i>	188-195
Correlation of antibiotics prescribing appropriateness based on the pharmaceutical care network europe (PCNE) method and clinical response in Community-Acquired Pneumonia	PDF
<i>Cut Fatia Ulfa, Woro Supadmi, Dyah Aryani Perwitasari, Endang Yuniarti</i>	196-207
Sleep quality and duration during pandemic uninvolved to impaired fasting glucose and hyperuricemia among health care practitioners	PDF
<i>Phebe Hendra, Fenty Fenty, Christianus Heru Setiawan, Leonardo Susanto Utomo, Gregorius Bhaskara Wikanendra, Putu Dyana Christasani, Dita Maria Virginia</i>	208-217
The comparison between combination of candesartan-amlodipine and candesartan-furosemide on blood pressure in hypertensive patients with chronic kidney disease	PDF
<i>Fitri Rachmaini, Dian Ayu Juwita, Almahdy A, Rahmad Abdillah, Yolanda Mayestika Wati</i>	218-226
The association between knowledge and cardiovascular risk in high school students in Bali Indonesia	PDF
<i>Preiffer Agus Prasajo, Yohana Helenora Desy Lopez, Christianus Heru Setiawan</i>	227-234

Causality assessment, severity and preventability of adverse drug reactions due to first-line antitubercular agents	PDF
<i>Gunjan Upadhyay, Apexa B Shukla, Darshan J Dave</i>	235-244
Effect of consumption of vegetables and fruit on nitric oxide levels and T-glutathione levels in type 2 diabetes mellitus patients	PDF
<i>Prita Anggraini Kartika Sari, Endang Darmawan, Akrom Akrom</i>	245-252
<hr/>	
The antioxidant activity of several antidiabetic herbal products	PDF
<i>Ichwan Ridwan Rais, Andhika Septiawan, Meta Ayuni, Dhega Agung Wichaksono, Nanik Sulistyani</i>	253-262
Standardization of <i>Cassia spectabilis</i> DC leaves and antimalarial activities of ethyl acetate extract	PDF
<i>Wiwied Ekasari, Heny Arwati, Nindya Tresiana Putri, Dewi Hariyani, Rosalia Friska Ananda, Eko Suhartono</i>	263-274

Pharmaciana

ISSN Print: 2088-4559 | ISSN Online: 2477-0256

Website: <http://journal.uad.ac.id/index.php/PHARMACIANA>

Office: Faculty of Pharmacy, Universitas Ahmad Dahlan

Jl. Prof. Dr. Soepomo, S.H., Janturan, Warungboto, Umbulharjo, Yogyakarta, Indonesia

Kode pos 55164

Email: pharmaciana@pharm.uad.ac.id

Standardization of *Cassia spectabilis* DC leaves and antimalarial activities of ethyl acetate extract

Wiwied Ekasari^{1*}, Heny Arwati², Nindya Tresiana Putri¹, Dewi Hariyani¹,
Rosalia Friska Ananda¹, Eko Suhartono³

¹Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga,
Jl. Mulyorejo, Surabaya, East Java, Indonesia

²Department of Parasitology, Faculty of Medicine, Universitas Airlangga,
Jl. Mayjen Prof. Dr. Moestopo No.47, Surabaya, East Java, Indonesia

³Department of Biochemistry and Molecular Biology, Faculty of Medicine, Universitas Lambung Mangkurat,
Jl. Veteran Sungai Bilu No.128, Banjarmasin, South Kalimantan, Indonesia

Submitted: 17-03-2022

Reviewed: 04-04-2022

Accepted: 15-06-2022

ABSTRACT

One of the Indonesian medicinal plants, *Cassia spectabilis* is traditionally used to treat several diseases, including malaria. The quality of raw materials influences the quality of drug derived from a plant. Standardizing the raw materials followed by the antimalarial activity test are needed to assure the quality of product prepared from plants. The aim of this study was to standardize the quality of *C. spectabilis* leaves and to evaluate the antimalarial activities of ethyl acetate extract. The fresh material of *C. spectabilis* leaves was observed in its specific and non-specific parameters. *In vitro* test was done by using *Plasmodium falciparum* 3D7. *In vivo* test was done using a 4-day suppressive test method against mice infected with *P. berghei* for four consecutive days. Heme detoxification inhibitory activity test was carried out using the modified Basilico method. The leaves of *C. spectabilis* meet the quality requirement for raw materials of traditional medicine. The ethyl acetate extract showed *in vitro* antiplasmodial activity against *P. falciparum* 3D7 and *in vivo* antimalarial activity against *P. berghei* infection with an IC₅₀ value of 27.28 µg/mL and ED₅₀ value of 1.74 mg/kg, respectively. The extract also showed heme detoxification inhibitory activity with an IC₅₀ value of 0.33±0.01 mg/mL. The leaves of *C. spectabilis* meet the quality requirement, and the ethyl acetate extract from standardized *C. spectabilis* leaves possessed a potential antimalarial activity that deserves to be further developed.

Keywords: malaria, *Cassia spectabilis*, heme detoxification, *Plasmodium berghei*, *Plasmodium falciparum*, standardization

*Corresponding author:

Wiwied Ekasari

Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga

Jl. Mulyorejo, Surabaya, East Java, Indonesia

Email: wiwied-e@ff.unair.ac.id



INTRODUCTION

Malaria is an infectious disease caused by parasite from the genus of *Plasmodium*, and transmitted by the bite of female *Anopheles* mosquito. Malaria remains the leading cause of death worldwide, most commonly in Africa and some Asian countries. Meanwhile, malaria in developed country caused by the imported cases from endemic areas (Talapko et al., 2019). The reduction in mortality and morbidity by at least 40% in 2020, as the important goals of the Global Technical Strategy for malaria is difficult to achieve (World Health Organization, 2018). Since 2010, a significant reduction in the burden of malaria has occurred, however, a slowdown and an increase in the number of cases between 2015 and 2017 have been observed. Reducing the number of cases in countries with the highest-burden is the most critical step in the global eradication of malaria (Talapko et al., 2019).

Drug resistance has been reported as severe global problem, even to all antimalarial drugs, however, artemisinin-based combination therapy (ACT) remains an effective therapy (Alonso & Noor, 2017). New drugs are needed to overcome the problem, where a number of new antimalarial discoveries and developments are ongoing, and some of them come from medicinal plants.

The species of *Cassia* (Caesalpiniaceae) is a well-known medicinal plant found in India and other tropical countries. One of the known species for the treatment of malaria is *Cassia siamea* which traditionally is consumed by drinking the decoction of leaves or flowers. The decoction of the flowers is also used as a body bath to treat malaria and liver disorders (Kamagaté et al., 2014). Several scientific studies have also proven the antimalarial activity of *C. siamea* along with its active compounds, either *in vitro* or *in vivo* (Ekasari et al., 2009). Apart from that, several other species of *Cassia* with clear antimalarial properties functioned throughout Africa, such as *C. occidentalis*, *C. africana*, *C. floribunda*, and *C. hirsuta*. Some of them have been shown to have *in vivo* antimalarial activity (Grace et al., 2012).

Previously, an *in vitro* antimalarial activity of methanolic extracts of leaves from *C. spectabilis* has been reported to show the highest inhibition against *P. falciparum* with an IC₅₀ value of 2.66 µg/mL (Ekasari et al., 2018). The ethanol extract of the leaves also showed antimalarial activity, both in the *in vitro* and *in vivo* test, with an IC₅₀ value of 12.52 µg/mL, and an ED₅₀ value of 131.5 mg/kg, respectively (Ekasari et al., 2018). The results proved the *C. spectabilis* leaves extract can be further potentially developed into antimalarial drug.

Medicinal plants play an important role in promoting health. They are widely spread all over the world but mostly grow in tropical countries. Recently, 25% of modern medicines, directly or indirectly were derived from plants (Patwekar et al., 2015). The quality of herbal drugs is highly affected by the quality of the raw materials. Further, the various factors such as cultivation, harvesting, and production influence the quality of raw materials. Quality assurance has to meet the specified standard. Raw materials of poor quality can affect the pharmacological activity of the product, therefore standardization of raw materials is the main key in obtaining a qualified drug (Yadav & Prajapati, 2011), because standardization is an essential step in obtaining a safe, effective, and qualified pharmaceutical product (Purwantiningsih et al., 2011). The determination of the specific and non-specific standard parameters of *C. spectabilis* leaves are reported herein. The improvement of the results will make these Indonesian plants as the material for traditional medicine, especially for antimalarial medicament.

Further investigation showed active antimalarial alkaloid compounds that reveals a structural pattern identical to (-)-7-hydroxycassine (Ekasari et al., 2021). The first antimalarial drug which contains alkaloid is quinine isolated from the *Cinchona* bark and is still used to treat malaria that is resistant to several drugs. Alkaloids of *C. spectabilis* leaves present in higher amounts in ethanol, methanol, and ethyl acetate extracts (Veerachari & Bopaiah, 2011). Antimalarial activity tests of methanol and ethanol extract of *C. spectabilis* leaves have been done before. Therefore, this study used ethyl acetate leaf extract of *C. spectabilis* to determine its antimalarial activity.

Assessment of antimalarial activity can be done *in vitro* to *P. falciparum* and *in vivo* to rodent plasmodia (Fidock et al., 2004). *In vitro* test includes IC₅₀ determination against *P. falciparum* type, both drug-resistant and drug-sensitive strains. *P. falciparum* culture can be used to study how parasites enter erythrocytes, screen for new drugs, isolate and characterize strains and clones, and identify immunogenic

antigens and parasite genomes (Kaira et al., 2006). As for the advantages of the *in vitro* method, there are precise and efficient; fast; a large number of compounds can be evaluated at the same time; synergism or antagonism with drug combinations can be studied; and a better assessment of the intrinsic activity of a drug.

In vivo test includes a four-day primary test for suppression of parasitaemia or inhibition of parasite growth in mice. This is the most widely used preliminary test, in which the efficacy of a compound is assessed by comparing blood parasitaemia and survival time of mice in treated and untreated mice (Kaira et al., 2006).

In addition to testing for *in vitro* and *in vivo* antimalarial activities, the identification of compounds that inhibit β -hematin formation is an approach for detecting antimalarial drugs (Mosaddegh et al., 2018). Malaria parasites digest hemoglobin in vacuoles into amino acids and heme, where the free heme formed can be toxic to the parasites themselves. To protect the body from the poisonous heme, *Plasmodium* is known to have several detoxification mechanisms such as the formation of hemozoin, heme-binding protein, and degradation of free heme by H_2O_2 (Slater & Cerami, 1992). Some drugs show an antimalarial effect through the inhibition of hemozoin formation, such as quinoline and xanthenes and their derivatives, which are the most important mechanism for detoxification (Fong & Wright, 2013). For *in vitro* antimalarial assessment, a synthetic polymer that is identical to hemozoin, namely β -hematin, is used.

Based on the description above, the aim of this study is to observe the antimalarial activity of ethyl acetate extract of *C. spectabilis* leaves, both *in vitro* against *P. falciparum* and *in vivo* against *P. berghei* infected mice, as well as its heme detoxification inhibitory activity.

MATERIALS AND METHODS

Plant Materials

Fresh *C. spectabilis* leaves were obtained and determined at Purwodadi Botanical Garden – Indonesian Institute of Sciences (LIPI), Pasuruan, Indonesia (B-160/IPH.06/KS.02/III/2019). The dried leaves were produced using a standard guideline *How to make simplicia* (Ministry of Health Republic of Indonesia, 1985) and were ground into powder using a pollinating machine.

Standardization of *C. spectabilis* Leaves

Standardization of *C. spectabilis* leaves were carried out based on standard guideline *Parameter Standar Umum Ekstrak* (Ministry of Health Republic of Indonesia, 2000) by determining the specific and non-specific parameters. The specific parameters including identification, macroscopic, microscopic, organoleptic, water- and ethanol-soluble extract content, as well as chemical content such as essential oils and total flavonoids were determined. Essential oil content was determined using xylol distillation method, while total flavonoid content was determined using aluminum chloride spectrophotometric method.

The non-specific parameters including loss on drying, total water content, total ash, total acid insoluble ash, as well as residue of pesticides, heavy metals, and microbial contamination were determined. Determination of total water content was carried out using toluene distillation method, while total ash and total acid insoluble ash were determined using thermogravimetric analysis method.

Preparation of ethyl acetate crude extracts

C. spectabilis leaves were extracted at room temperature in the research room of the Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia. The extract was made by macerating 500 g of dry leaf powder in 2500 mL of ethyl acetate for 3x24 h. The extract was then evaporated on a rotary evaporator.

Experimental animals

Adult BALB/c male mice (20–30 g) were obtained from the Faculty of Veterinary Medicine, Universitas Airlangga. Animals were acclimatized to the laboratory environment (room temperature

24±1°C, humidity 55±5%, 12 h light/dark cycle) for two weeks before the test and provided free access to food and water sources. The research protocol was approved by the Animal Use and Care Committee of Faculty of Veterinary Medicine, Universitas Airlangga (2.KE.181.10.2018).

In Vitro antimalarial activity

The stock solution of extract was prepared in dimethyl sulfoxide (DMSO) and diluted with complete RPMI 1640 culture media until the final concentrations of the solutions were ranging from 0.001 to 100 µg/mL. The test was carried out in duplicate. The plates were incubated in a candle jar for 48 h under CO₂ conditions at 37°C. The contents of the well were harvested by making a thin smear of blood on clean microscopic slides after the incubation process. The blood smear was then fixed using methanol and stained for 10 min in 10% Giemsa solution (pH 7.3). Each IC₅₀ values was calculated using probit analysis based on the percentage of *P. falciparum* growth inhibition.

In Vivo antimalarial activity

In vivo testing of ethyl acetate leaf extract of *C. spectabilis* was carried out on rodent malaria parasites (*Plasmodium berghei*) using a 4-day suppressive test method after the inoculation of 2x10⁷ parasites/mice intraperitoneally (Fidock et al., 2004). A group of eight mice (BALB/c, male, eight weeks old) was used for each dose. Five other mice were kept as donors who were inoculated but untreated. The negative and positive control groups each received a suspension of 0.5% sodium carboxymethylcellulose (Na CMC) solution and dihydroartemisinin + piperazine (20.8 + 166.4 mg/kg/day, p.o.) (standard drug). Four doses of ethyl acetate leaf extract of *C. spectabilis* tested 1, 10, 100, and 200 mg/kg/day. Each mouse was given a 0.5% Na CMC solution, plant extracts, and standard drugs orally. The first administration was started after three hours of parasite inoculation (D₀) and then continued at the same time for three additional days (D₁–D₃). The percentage of parasitaemia was determined on D₄ with a thin blood smear using the formula described by Fidock et al. (2004). ED₅₀ was rated as a dose causing 50% inhibition of parasite growth compared to growth in negative controls.

Heme detoxification inhibitory activity

Heme detoxification inhibitory activity was carried out on ethyl acetate extract of *C. spectabilis* leaves based on the modified Basilico method (Basilico et al., 1998). The test materials were plant extracts and positive control (chloroquine diphosphate) with various levels, which were 4; 2; 1; 0.5; 0.25; and 0.1 mg/mL. Each level was replicated three times. The dissolution process of the test material used 10% DMSO and was carried out by dissolving the test material in 100 µL of DMSO and added with distilled water to a concentration of 10% DMSO. Each microtube was added as much as 50 µL of the test material, 100 µL of 1 mM hematin solution in 0.2 M NaOH, and 50 µL of glacial acetic acid and incubated for 24 h at 37°C. Then the solution was centrifuged at 8000 rpm for 10 min, the supernatant was removed, and the precipitate was washed with 200 µL of DMSO. Washing was performed three times. At the final stage of washing, the precipitate was dissolved in 200 µL of 0.1 M NaOH as much as 100 µL of the solution was distributed into 96-well microplates to measure the absorbance using ELISA reader at a wavelength of 405 nm. Inhibition percentage was calculated, and IC₅₀ (the content of the test compound which was able to inhibit the formation of β-hematin by up to 50%) was calculated using probit analysis.

Statistical analysis

The data were expressed as mean ± standard error of the mean (S.E.M). Statistical significance of means was determined by one-way analysis of variance (ANOVA), followed by Dunnett's (posthoc test) to compare the measured parameters (parasitemia and level of β-hematin) with negative controls. The analysis was performed with a 95% confidence interval, and *p*-values less than 0.05 was considered to be statistically significant.

RESULT AND DISCUSSION

Determining specific standard parameters of *C. spectabilis* includes macroscopic and microscopic observation, organoleptic test, water-soluble essence, ethanol-soluble essence, volatile oil, as well as total flavonoid level. Macroscopic observation results are shown in [Figure 1](#) and [Table 1](#).



Figure 1. *Cassia spectabilis* leaves and flowers

Table 1. Morphological observation of *Cassia spectabilis* leaves

Characteristic	Result
Leaf	
Shape	Oval-oblong
Leaf tip	Blunt with a shallow notch at the tip
Leaf base	Blunt or quite rounded
Surface	Upper surface: glabrous, quite glossy Lower surface: Pubescent
Edge	Even
Leaf vein	Pinnate
Size	
Length	3 – 7.5 cm
Width	1 – 2.5 cm
Color	Green to brownish-green
Stipule	Long

Microscopic observations, including fragments of mesophyll, crystal fibers, mesophyll with calcium oxalate crystal, and trichome, are shown in [Figure 2](#).




Figure 2. Fragments of (A) mesophyll; (B) crystal fibers (prism shape, blue arrow); (C) mesophyll with calcium oxalate crystal (blue arrow); and (D) trichome

Other results of the specific and non-specific parameters were summarized in [Table 2](#). For determination of non-specific parameters, such as loss on drying, water level, total ash, and acid-insoluble ash of dried leaves powder of *C. spectabilis* it meets the quality required to be used as the raw material of traditional medicine.

In the determination of microbes, the dried leaves powder of *C. spectabilis* already met the requirements regulated by the [National Agency of Drug and Food Control \(2014\)](#), as well as the determination of residue of pesticides and heavy metals [Table 2](#). Pesticide is a chemical substance still used in agriculture, whereas it can cause damage to health because of its carcinogenic, mutagenic, and teratogenic effects ([El-Nahhal & Radwan, 2013](#)). Heavy metal is a chemical element with high molecular weight. It is in solid form at room temperature. Heavy metal is essential for living creatures in small amounts, but in large amounts, it causes a dangerous effect on the human.

In vitro antimalarial test results from ethyl acetate leaf extract of *C. spectabilis* are shown in [Table 3](#), where the IC_{50} value obtained was 27.28 $\mu\text{g/mL}$.

Table 2. Summary of results of determination of specific and non-specific parameters of dried leaves powder of *C. spectabilis*

Parameter	Result	Limit allowed for dried herbal raw materials
Specific parameters:		
Organoleptic		
Color	Slightly brownish	
Taste	Fresh	
Smell	Weak	
Water-soluble essence	3.72 ± 0.11 % w/w	-
Ethanol soluble essence	1.61 ± 0.09 % w/w	-
Volatile oil	0.17 % v/w	-
Total flavonoid content	2.22 ± 0.20 % w/w	-
Non-specific parameters:		
Loss on drying	7.81 ± 0.11 % w/w	-
Water level	7.54 ± 0.38 % w/w	-
Total ash	8.56 ± 0.09 % w/w	-
Acid-insoluble ash	2.26 ± 1.14 % w/w	-
Microbial contamination:		
Mold	60 colonies/g	≤ 104 colonies/g
Yeast	< 10 colonies/g	≤ 104 colonies/g
APM <i>Coliform</i>	7.4 colonies/g	≤ 106 colonies/g
<i>Escherichia coli</i>	Negative	Negative/g
<i>Salmonella sp.</i>	Negative	Negative/g
<i>Staphylococcus aureus</i>	Negative	Negative/g
<i>Pseudomonas aeruginosa</i>	Negative	Negative/g
Pesticide residues:		
Organochlorine	Negative	
Carbamate	Negative	
Heavy metal residues		
Lead (Pb)	0.250 mg/kg	≤ 10 mg/kg
Cadmium (Cd)	0.199 mg/kg	≤ 0.3 mg/kg
Zinc (Zn)	4.679 mg/kg	-
Copper (Cu)	0.346 mg/kg	-

Note: Data are expressed as mean ± S.E.M. ($n = 3$)


Based on the classification according to Gessler et al. (1994), the antimalarial activity of extracts with an IC_{50} value of $<10 \mu\text{g/mL}$ is considered very good; $10\text{--}50 \mu\text{g/mL}$ is moderate, and $>50 \mu\text{g/mL}$ is low. This suggests that *in vitro* antimalarial activity from ethyl acetate leaf extract of *C. spectabilis* was moderate because the IC_{50} value was in the range of 10 to $50 \mu\text{g/mL}$.

In vivo antimalarial activity test results from ethyl acetate leaf extract of *C. spectabilis* against *P. berghei* infected mice can be seen in Figure 3 and Table 4. Figure 3 shows that each mouse has a different parasitic growth profile. This difference is due to variations in the immune system of each mouse.

Table 3. The average inhibition percentage of ethyl acetate extract of *C. spectabilis* leaves against *P. falciparum* 3D7 strain *in vitro*

Sample	Concentration (µg/mL)	Parasitemia (%)	Inhibition (%)	IC ₅₀ (µg/mL)
EACS	100	0.36 ± 0.03	100.00 ± 0.00	27.28
	10	2.54 ± 0.05	41.18 ± 6.00	
	1	2.69 ± 0.07	35.66 ± 6.88	
	0.1	3.20 ± 0.11	16.91 ± 8.95	
	0.01	4.44 ± 0.30	0.00 ± 0.00	
Negative control	-	3.66 ± 0.13	-	-

Note: Data are expressed as mean ± S.E.M. ($n = 2$). No significantly different in the percentage of parasitemia and inhibition compared to the negative control ($p > 0.05$). EACS = ethyl acetate extract of *C. spectabilis* leaves. DMSO was used for solubilizing the extracts and as a negative control

**Figure 3.** The effect of ethyl acetate extract of *C. spectabilis* leaves on parasitemia percentage of *P. berghei* infected mice on 4-day suppression test; data are mean ± S.E.M. ($n = 8$); Na CMC = sodium carboxymethylcellulose, DHP = dihydroartemisinin + piperazine, EACS = ethyl acetate extract of *C. spectabilis* leaves; numbers refer to doses in mg/kg/day**Table 4.** Percentage of parasitemia result treated *in vivo* of negative control, positive control, and ethyl acetate extract of *C. spectabilis* leaves

Treatment	Dose (mg/kg/day, p.o.)	Parasitemia (%)		Suppression (%)	ED ₅₀ (mg/kg)
		Day 0	Day 4		
EACS	1	1.46 ± 0.11	8.24 ± 0.20*	46.23	1.74
	10	1.32 ± 0.14	6.01 ± 0.21*	62.81	
	100	1.52 ± 0.11	4.09 ± 0.19*	79.62	
	200	1.22 ± 0.10	3.20 ± 0.17*	84.30	
DHP	20.8 + 166.4	1.23 ± 0.22	0.00 ± 0.00*	100.00	ND
Negative control	-	1.63 ± 0.07	14.24 ± 0.51	-	-

Note: Data are expressed as mean ± S.E.M. ($n = 8$). *Significantly different compared to negative control ($p < 0.05$). EACS = ethyl acetate extract of *C. spectabilis* leaves; DHP = dihydroartemisinin + piperazine. Na CMC were used as a negative control. ND = Not determined

In vivo antimalarial activity test of ethyl acetate leaf extract of *C. spectabilis* against *P. berghei*-infected mice resulted in the inhibition of parasite growth was a dose-dependent manner, with the greatest inhibition of 84.30% at a dose of 200 mg/kg. *In vivo* antimalarial activity is calculated from the ED₅₀ value, which was based on the probit analysis that revealed the dose which inhibits the growth of parasites by 50% of the total population. The analysis resulted in an ED₅₀ value of 1.74 mg/kg, which revealed the ethyl acetate leaf extract of *C. spectabilis* to be a very active antimalarial (Muñoz et al., 2000). This value was lower than that of 90% ethanol extract of *C. spectabilis* leaves, which was 131.5 mg/kg (Ekasari et al., 2018). On the other hand, ethanol extract, chloroform extract, and aqueous leaf extract of *C. siamea* have an ED₅₀ of 34.70 mg/kg, 19.59 mg/kg, and 83.77 mg/kg, respectively (Ekasari et al., 2009). Ethyl acetate extract from other plant species, such as *Garcinia husor* (Clusiaceae) bark, showed antimalarial activity against *P. berghei*-infected mice with an ED₅₀ value of 22.30 mg/kg (Kainama et al., 2019), which was much higher than that of ethyl acetate leaf extract of *C. spectabilis* in this current study.

The heme detoxification inhibitory activity test of ethyl acetate leaf extract of *C. spectabilis* using Basilico method and read by ELISA reader at a wavelength of 405 nm resulted in heme α -chlorohemin reacted with acetate in a 24 h incubation formed β -hematin. The use of DMSO solution was aimed to remove the residual hematin that was still mixed with β -hematin crystals that were insoluble in DMSO washing solution. The application of NaOH to the formed-hematin precipitate would convert it to alkaline hematin, which could be measured by ELISA reader. The heme detoxification inhibitory activity was expressed in IC₅₀, which was the level of the test compound capable of inhibiting β -hematin formation by up to 50%. IC₅₀ calculation results using probit analysis can be seen in Table 5.

Table 5. IC₅₀ values of chloroquine diphosphate and ethyl acetate extract of *C. spectabilis* leaves

Sample	Concentration (mg/mL)	Level of β -hematin (mM)	Inhibition (%)	IC ₅₀ (mg/mL)
EACS	4.00	15.71 \pm 4.01*	88.92 \pm 1.78	0.33 \pm 0.01
	2.00	29.51 \pm 2.92*	78.54 \pm 0.36	
	1.00	45.26 \pm 5.60*	67.25 \pm 1.10	
	0.50	61.06 \pm 7.04*	55.75 \pm 0.62	
	0.25	74.34 \pm 7.71*	46.01 \pm 0.67	
	0.10	95.93 \pm 7.87*	30.02 \pm 1.70	
	CQ	4.00	32.19 \pm 9.51*	
2.00		43.98 \pm 8.62*	68.64 \pm 2.86	
1.00		54.89 \pm 8.55*	60.54 \pm 1.90	
0.50		69.17 \pm 7.77*	49.85 \pm 0.47	
0.25		88.02 \pm 7.03*	35.76 \pm 1.65	
0.10		102.62 \pm 5.85*	24.78 \pm 3.35	
Negative control		-	137.72 \pm 14.48	-

Note: Data are expressed as mean \pm S.E.M. ($n = 3$). *Significantly different compared to negative control ($p < 0.05$). EACS = ethyl acetate extract of *C. spectabilis* leaves; CQ = chloroquine diphosphate. DMSO was used as a negative control

The mechanism of action of antimalarial compounds in inhibiting heme detoxification consists of two mechanisms, the first interaction occurs between terpenoid compounds, phenols, and sterols with the heme electronic system, and the second mechanism is the interaction between the test compound (extract) which has a hydroxyl group that can bind heme iron ions (Turalely et al., 2011). Based on these facts, the mechanism of action of ethyl acetate leaf extract of *C. spectabilis* inhibiting heme detoxification probably by interacting with the alkaloid hydroxyl group in the extracted compound, which is bound to the heme iron ion.

According to Frölich et al. (2005), the compounds that have a formation resistance in inhibiting β -hematin formation greater than 60% was a good potential as inhibitors of β -hematin formation, while smaller than 40% was as weak inhibitors in inhibitory effect of ethyl acetate leaf extract of *C. spectabilis* in β -hematin formation was a good potential inhibitor, where was greater than 60% at the concentrations of 4; 2; and 1 mg/mL. In the result of inhibition percentage in β -hematin extract and positive control formation, it was shown that higher levels in the test material would increase the percentage inhibition in β -hematin formation.

The IC₅₀ value of heme detoxification inhibitory activity from ethyl acetate leaf extract of *C. spectabilis* was 0.33 mg/mL. It is lower than the IC₅₀ value of positive control with a value of 0.56 mg/mL, indicating that ethyl acetate leaf extract of *C. spectabilis* was an active extract in inhibiting heme detoxification (Baelmans et al., 2000). Based on this result, heme detoxification inhibitory activity could be a mechanism of action of ethyl acetate leaf extract of *C. spectabilis* as an antiplasmodial.

CONCLUSION

Overall, this study was concluded that the dried leaves powder of *C. spectabilis* meets the quality requirements and recommended to be used as the raw material of traditional medicine. The results of antiplasmodial activity showed that the ethyl acetate leaf extract of *C. spectabilis* has good activity as an antiplasmodial, either *in vitro* or *in vivo*, and showed heme detoxification inhibitory activity. Further research is needed on the compounds that play a role in antimalarial activity and their mechanism of action so that they can be used as an alternative as anti-malarial drugs or as a combination drug with other antimalarials.

ACKNOWLEDGEMENT

We would like to thank The Ministry of Research, Technology, and Higher Education of the Republic of Indonesia for its financial support.

REFERENCES

- Alonso, P., & Noor, A. M. (2017). The global fight against malaria is at crossroads. *The Lancet*, 390(10112), 2532–2534. [https://doi.org/10.1016/S0140-6736\(17\)33080-5](https://doi.org/10.1016/S0140-6736(17)33080-5)
- Baelmans, R., Deharo, E., Muoz, V., Sauvain, M., & Ginsburg, H. (2000). Experimental conditions for testing the inhibitory activity of chloroquine on the formation of β -hematin. *Experimental Parasitology*, 96(4), 243–248. <https://doi.org/10.1006/expr.2000.4558>
- Basilico, N., Pagani, E., Monti, D., Olliaro, P., & Taramelli, D. (1998). A microtitre-based method for measuring the haem polymerization inhibitory activity (HPIA) of antimalarial drugs. *Journal of Antimicrobial Chemotherapy*, 42(1), 55–60. <https://doi.org/10.1093/jac/42.1.55>
- Ekasari, W., Basuki, D. R., Arwati, H., & Wahyuni, T. S. (2021). Antiplasmodial activity of Ethanolic extract of *Cassia spectabilis* DC leaf and its inhibition effect in heme detoxification. *BMC Complementary Medicine and Therapies*, 21(1), 1–12. <https://doi.org/10.1186/s12906-021-03239-9>
- Ekasari, W., Wahyuni, T. S., Arwati, H., & Putri, N. T. (2018). Determination of effective dose of antimalarial from *Cassia spectabilis* leaf ethanol extract in *Plasmodium berghei*-infected mice. *African Journal of Infectious Diseases*, 12(Special Issue 1), 110–115. <https://doi.org/10.2101/Ajid.12v1S.16>
- Ekasari, W., Widyawaruyanti, A., Zaini, N. C., Syafruddin, D., Honda, T., & Morita, H. (2009). Antimalarial activity of cassiarin A from the leaves of *Cassia siamea*. *Heterocycles*, 78(7), 1831–1836. <https://doi.org/10.3987/COM-09-11680>
- El-Nahhal, Y., & Radwan, A. A. (2013). Human health risks: Impact of pesticide application. *Journal of Environment and Earth Science*, 3(7), 199–210.
- Fidock, D. A., Rosenthal, P. J., Croft, S. L., Brun, R., & Nwaka, S. (2004). Antimalarial drug discovery: efficacy models for compound screening. *Nature Reviews. Drug Discovery*, 3(6), 509–520.

- <https://doi.org/10.1038/nrd1416>
- Fong, K. Y., & Wright, D. W. (2013). Hemozoin and antimalarial drug discovery. *Future Medicinal Chemistry*, 5(12), 1437–1450. <https://doi.org/10.4155/fmc.13.113>
- Frölich, S., Schubert, C., Bienzle, U., & Jenett-Siems, K. (2005). In vitro antiplasmodial activity of prenylated chalcone derivatives of hops (*Humulus lupulus*) and their interaction with haemin. *Journal of Antimicrobial Chemotherapy*, 55(6), 883–887. <https://doi.org/10.1093/jac/dki099>
- Gessler, M. C., Nkunya, M. H. H., Mwasumbi, L. B., Heinrich, M., & Tanner, M. (1994). Screening Tanzanian medicinal plants for antimalarial activity. *Acta Tropica*, 56(1), 65–77. [https://doi.org/10.1016/0001-706x\(94\)90041-8](https://doi.org/10.1016/0001-706x(94)90041-8)
- Grace, M. H., Lategan, C., Grazioplene, R., Smith, P. J., Raskin, I., & Lila, M. A. (2012). Antiplasmodial activity of the ethnobotanical plant *Cassia fistula*. *Natural Product Communications*, 7(10), 1263–1266. <https://doi.org/10.1177/1934578x1200701002>
- Kainama, H., Fatmawati, S., Santoso, M., Kakisina, P., & Ersam, T. (2019). In vitro and in vivo antiplasmodial of stem bark extract of *Garcinia husor*. *HAYATI Journal of Biosciences*, 26(2), 81–89. <https://doi.org/10.4308/hjb.26.2.81>
- Kaira, B. S., Chawla, S., Gupta, P., & Valecha, N. (2006). Screening of antimalarial drugs: An overview. *Indian Journal of Pharmacology*, 38(1), 5–12. <https://doi.org/10.4103/0253-7613.19846>
- Kamagaté, M., Koffi, C., Kouamé, N. M., Akoubet, A., Yao, N. A. R., & Die-Kakou, H. M. (2014). Ethnobotany, phytochemistry, pharmacology and toxicology profiles of *Cassia siamea* Lam. *The Journal of Phytopharmacology*, 3(1), 57–76.
- Ministry of Health Republic of Indonesia. (1985). *How to make simplicia*. Ministry of Health Republic of Indonesia.
- Ministry of Health Republic of Indonesia. (2000). *General standard parameters of medicinal plant extract*. Ministry of Health Republic of Indonesia.
- Mosaddegh, M., Irani, M., & Esmaeili, S. (2018). Inhibition test of heme detoxification (ITHD) as an approach for detecting antimalarial agents in medicinal plants. *Research Journal of Pharmacognosy*, 5(1), 5–11.
- Muñoz, V., Sauvain, M., Bourdy, G., Callapa, J., Bergeron, S., Rojas, I., Bravo, J. A., Balderrama, L., Ortiz, B., Gimenez, A., & Deharo, E. (2000). A search for natural bioactive compounds in Bolivia through a multidisciplinary approach. Part I. Evaluation of the antimalarial activity of plants used by the Chacobo Indians. *Journal of Ethnopharmacology*, 69(2), 127–137. [https://doi.org/10.1016/S0378-8741\(99\)00148-8](https://doi.org/10.1016/S0378-8741(99)00148-8)
- National Agency of Drug and Food Control. (2014). Regulation of head of national agency of drug and food control republic Indonesia number 12 about quality requirements of traditional medicine. National Agency of Drug and Food Control.
- Patwekar, S. L., Suryawanshi, A. B., Gaikwad, M. S., Pedewad, S. R., & Potulwar, A. P. (2015). Standardization of herbal drugs: An overview. *The Pharma Innovation Journal*, 4(9), 100–104.
- Purwantiningsih, Purwantini, I., & Santoso, D. (2011). Identification of standard parameters of Kepel leaves [*Stelechocarpus Burahol* (BL.) Hook. F. & TH.] and the extract as raw material for anti-hyperuricemic medicaments. *Asian Journal of Pharmaceutical and Clinical Research*, 4(Suppl. 1), 149–153.
- Slater, A. F. G., & Cerami, A. (1992). Inhibition by chloroquine of a novel haem polymerase enzyme activity in malaria trophozoites. *Letters to Nature*, 355(6356), 167–169. <https://doi.org/10.1038/355167a0>
- Talapko, J., Škrlec, I., Alebić, T., Jukić, M., & Včev, A. (2019). Malaria: The past and the present. *Microorganisms*, 7(6), 179. <https://doi.org/10.3390/microorganisms7060179>
- Turalely, R., Susidarti, R. A., & Wijayanti, M. A. (2011). In vivo antiplasmodial of the most active fraction and its compound of Kapur leaves (*Harmsioplanax aculeatus* Harms) extract against *Plasmodium berghei*. *Tropical Medicine Journal*, 1(2), 131–140. <https://doi.org/10.22146/tmj.4575>
- Veerachari, U., & Bopaiah, A. K. (2011). Preliminary phyto-chemical evaluation of the leaf extract of

- five Cassia species. *Journal of Chemical and Pharmaceutical Research*, 3(5), 574–583.
- World Health Organization. (2018). *World malaria report 2018*. WHO Press.
- Yadav, P., & Prajapati, P. K. (2011). Quality control parameters for medicinal plants, an overview. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 1(5), 12–16.



**KOMISI ETIK PENELITIAN
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA
*Animal Care and Use Committee (ACUC)***

**KETERANGAN KELAIKAN ETIK
“ ETHICAL CLEARANCE ”**

No : 2.KE.181.10.2018

**KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE)
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA,
TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG
DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :**

PENELITIAN BERJUDUL : Efikasi Ekstrak Etanol Daun *Cassia spectabilis* Sebagai Obat Antimalaria dari Bahan Alami

PENELITI UTAMA : Wiwied Ekasari

UNIT/LEMBAGA/TEMPAT PENELITIAN : Fakultas Farmasi Universitas Airlangga

DINYATAKAN : LAIK ETIK

Surabaya, 30 Oktober 2018

Ketua,

Dr. Nusdianto Triakoso, M.P.,Drh.
NIP. 196805051997021001

Mengetahui,
Dekan FKH-Unair,

Prof. Dr. Rudi Sianto, M.Kes.,Drh.
NIP. 195601051986011001





KEMENTERIAN PENDIDIKAN, KEBUDAYAAN,
RISET, DAN TEKNOLOGI
DIREKTORAT JENDERAL PENDIDIKAN TINGGI, RISET,
DAN TEKNOLOGI

Jalan Jenderal Sudirman, Senayan, Jakarta 10270
Telepon (021) 57946104, Pusat Panggilan ULT Dikti 126
Laman www.dikti.kemdikbud.go.id

SALINAN

KEPUTUSAN DIREKTUR JENDERAL PENDIDIKAN TINGGI,
RISET, DAN TEKNOLOGI
KEMENTERIAN PENDIDIKAN, KEBUDAYAAN, RISET, DAN TEKNOLOGI
REPUBLIK INDONESIA

NOMOR 105/E/KPT/2022
TENTANG
PERINGKAT AKREDITASI JURNAL ILMIAH PERIODE I TAHUN 2022

DIREKTUR JENDERAL PENDIDIKAN TINGGI, RISET, DAN TEKNOLOGI,

- Menimbang : a. bahwa untuk melaksanakan ketentuan Pasal 6 ayat (5) Peraturan Menteri Riset, Teknologi dan Pendidikan Tinggi Nomor 9 Tahun 2018 tentang Akreditasi Jurnal Ilmiah dan berdasarkan Berita Acara Penetapan Hasil Akreditasi Jurnal Periode 1 Tahun 2022 pada tanggal 28 Maret 2022, perlu menetapkan peringkat akreditasi jurnal ilmiah periode I tahun 2022;
- b. bahwa berdasarkan pertimbangan sebagaimana dimaksud dalam huruf a, perlu menetapkan Keputusan Direktur Jenderal Pendidikan Tinggi, Riset, dan Teknologi tentang Peringkat Akreditasi Jurnal Ilmiah Periode I Tahun 2022;

- Mengingat : 1. Undang-Undang Nomor 12 Tahun 2012 tentang Pendidikan Tinggi (Lembaran Negara Republik Indonesia Tahun 2012 Nomor 158, Tambahan Lembaran Negara Republik Indonesia Nomor 5336);
2. Peraturan Pemerintah Nomor 4 Tahun 2014 tentang Penyelenggaraan Pendidikan Tinggi dan Pengelolaan Perguruan Tinggi (Lembaran Negara Republik Indonesia Tahun 2014 Nomor 16, Tambahan Lembaran Negara Republik Indonesia Nomor 5500);

3. Peraturan Presiden Nomor 62 Tahun 2021 tentang Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi (Lembaran Negara Republik Indonesia Tahun 2021 Nomor 156);
4. Peraturan Menteri Riset, Teknologi, dan Pendidikan Tinggi Nomor 9 Tahun 2018 tentang Akreditasi Jurnal Ilmiah (Berita Negara Republik Indonesia Tahun 2018 Nomor 428);
5. Peraturan Menteri Pendidikan, Kebudayaan, Riset, dan Teknologi Nomor 28 Tahun 2021 tentang Organisasi dan Tata Kerja Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi (Berita Negara Republik Indonesia Tahun 2021 Nomor 963);
6. Keputusan Direktur Jenderal Pendidikan Tinggi, Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi Nomor 105/E/KPT/2021 tentang Asesor Akreditasi Jurnal Ilmiah Nasional;
7. Keputusan Direktur Jenderal Pendidikan Tinggi, Riset, dan Teknologi Kementerian Pendidikan Tinggi, Riset, dan Teknologi Nomor 134/E/KPT/2021 tentang Pedoman Akreditasi Jurnal Ilmiah;

MEMUTUSKAN:

Menetapkan : KEPUTUSAN DIREKTUR JENDERAL PENDIDIKAN TINGGI, RISET, DAN TEKNOLOGI TENTANG PERINGKAT AKREDITASI JURNAL ILMIAH PERIODE I TAHUN 2022.

KESATU : Menetapkan peringkat akreditasi jurnal ilmiah periode I tahun 2022 sebagaimana tercantum dalam Lampiran yang merupakan bagian yang tidak terpisahkan dari Keputusan Direktur Jenderal ini.

KEDUA : Peringkat akreditasi Jurnal Ilmiah sebagaimana dimaksud dalam Diktum KESATU berlaku selama 5 (lima) tahun mulai Volume, Nomor dan Tahun Terbitan sampai Volume, Nomor dan Tahun terbitan sesuai Lampiran Keputusan Direktur Jenderal ini.

- KETIGA : Jurnal Ilmiah sebagaimana dimaksud dalam Diktum KESATU dapat mengajukan kembali kenaikan peringkat akreditasi setelah menerbitkan paling sedikit 4 (empat) nomor penerbitan.
- KEEMPAT : Jurnal ilmiah yang telah memiliki peringkat akreditasi sebagaimana dimaksud dalam Diktum KESATU wajib:
- mencantumkan masa berlaku akreditasi, dengan menuliskan tanggal penetapan dan tanggal akhir masa berlaku akreditasi; dan
 - menampilkan sertifikat akreditasi.
- KELIMA : Apabila setelah ditetapkannya Keputusan Direktur Jenderal ini ditemukan ketidaksesuaian antara jurnal ilmiah sebagaimana Diktum KESATU dengan Pedoman Akreditasi Jurnal Ilmiah, peringkat akreditasi jurnal ilmiah dapat diturunkan atau dicabut.
- KEENAM : Keputusan Direktur Jenderal ini mulai berlaku pada tanggal ditetapkan.

Ditetapkan di Jakarta
pada tanggal 7 April 2022

Plt. DIREKTUR JENDERAL PENDIDIKAN
TINGGI, RISET, DAN TEKNOLOGI,

TTD.

NIZAM
NIP 196107061987101001

Salinan sesuai dengan aslinya

Plt. Sekretaris Direktorat Jenderal Pendidikan Tinggi, Riset, dan Teknologi
Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi,

Tjitjik Srije Tjahjandarie
NIP 196502061988102001



57	Lentera Hukum	26213710	Universitas Jember	Reakreditasi Tetap di Peringkat 2 mulai Volume 8 Nomor 3 Tahun 2021 sampai Volume 13 Nomor 2 Tahun 2026
58	LIMNOTEK - Perairan Darat Tropis di Indonesia	25498029	Pusat Penelitian Limnologi dan Masyarakat Limnologi Indonesia	Reakreditasi Tetap di Peringkat 2 mulai Volume 27 Nomor 2 Tahun 2020 sampai Volume 32 Nomor 1 Tahun 2025
59	Majalah Obstetri & Ginekologi	25981013	Departemen Obstetri dan Ginekologi, Fakultas Kedokteran Universitas Airlangga	Reakreditasi Tetap di Peringkat 2 mulai Volume 29 Nomor 2 Tahun 2021 sampai Volume 34 Nomor 1 Tahun 2026
60	Marine Research in Indonesia	24432008	Pusat Penelitian Oseanografi - Lembaga Ilmu Pengetahuan Indonesia	Reakreditasi Tetap di Peringkat 2 mulai Volume 45 Nomor 2 Tahun 2020 sampai Volume 50 Nomor 1 Tahun 2025
61	Mimbar Ilmu	26859033	Universitas Pendidikan Ganesha	Reakreditasi Naik Peringkat dari Peringkat 3 ke Peringkat 2 mulai Volume 26 Nomor 3 Tahun 2021 sampai Volume 31 Nomor 2 Tahun 2026
62	Mosharafa: Jurnal Pendidikan Matematika	25278827	Program Studi Pendidikan Matematika, Institut Pendidikan Indonesia	Reakreditasi Naik Peringkat dari Peringkat 3 ke Peringkat 2 mulai Volume 10 Nomor 2 Tahun 2021 sampai Volume 15 Nomor 1 Tahun 2026
63	Nadwa : Jurnal Pendidikan Islam	25028057	Fakultas Ilmu Tarbiyah dan Keguruan UIN Walisongo Semarang	Reakreditasi Tetap di Peringkat 2 mulai Volume 15 Nomor 1 Tahun 2021 sampai Volume 19 Nomor 2 Tahun 2025
64	Pharmaciana: Jurnal Kefarmasian	24770256	Fakultas Farmasi, Universitas Ahmad Dahlan	Reakreditasi Tetap di Peringkat 2 mulai Volume 11 Nomor 3 Tahun 2021 sampai Volume 16 Nomor 2 Tahun 2026