

# Effect of low oxygen tension on transcriptional factor OCT4 and SOX2 expression in New Zealand rabbit bone marrow-derived mesenchymal stem cells

*by* Erma Erma Safitri

---

**Submission date:** 08-Aug-2022 05:17PM (UTC+0800)

**Submission ID:** 1880215203

**File name:** Vet\_World\_Erma\_First\_Corresponding.pdf (868.01K)

**Word count:** 5321

**Character count:** 28481

## Effect of low oxygen tension on transcriptional factor OCT4 and SOX2 expression in New Zealand rabbit bone marrow-derived mesenchymal stem cells

Erma Safitri<sup>1,2</sup>

<sup>19</sup> 1. Department of Veterinary Reproduction, Faculty <sup>45</sup> Veterinary Medicine, Universitas Airlangga, Surabaya 60115, Indonesia; 2. Stem Cells Research Division, Institute Tropical Disease, Universitas Airlangga, Surabaya 60115, Indonesia.  
**Corresponding author:** Erma Safitri, e-mail: erma-s@fkh.unair.ac.id

**Received:** 22-06-2020, **Accepted:** 14-10-2020, **Published online:** 18-11-2020

<sup>9</sup> **doi:** www.doi.org/10.14202/vetworld.2020.2469-2476 **How to cite this article:** Safitri E (2020) Effect of low oxygen tension on transcriptional factor OCT4 and SOX2 expression in New Zealand rabbit bone marrow-derived mesenchymal stem cells, *Veterinary World*, 13(11): 2469-2476.

### Abstract

**Background and Aim:** Octamer-binding transcription factor 4 (OCT4) and sex-determining region Y-box 2 (SOX2) are transcription factors whose functions are essential to maintain the pluripotency of embryonic stem cells. The purpose of this study was to derive stem cells for *in vitro* culture and to maintain their viability and pluripotency, with the goal to <sup>42</sup>tain a cell line for transplantation in patients with degenerative diseases or injuries. This research focused on examining the effect of low oxygen tension on the ability of bone marrow-derived mesenchymal stem cells (BM-MSCs) to express OCT4 and SOX2 *in vitro*.

**Materials and Methods:** BM-MSCs were obtained from femurs of 2000 to 3000 g New Zealand male rabbits. BM-MSCs were divided into three groups to test different culture conditions: A control group under hyperoxia condition (21% O<sub>2</sub>) and two treatment groups with low oxygen tension (1% and 3% O<sub>2</sub>). We characterized the BM-MSCs using flow cytometric measurement of cluster differentiation 44 (CD44) and cluster differentiation 90 (CD90) expression. The expression of OCT4 and SOX2 was measured by immunofluorescence staining after 48 h of incubation in chambers with normal or low oxygen tension with controlled internal atmosphere consisting of 95% N<sub>2</sub>, 5% CO<sub>2</sub>, and 1% O<sub>2</sub> (T1) and 3% O<sub>2</sub> (T2). We considered OCT4 and SOX2 as two markers of pluripotency induction. All immunofluorescence data were subjected to a *post hoc* normality Tukey's honestly significant difference test; all differences with p<5% were considered significant.

**Results:** BM-MSCs were positive for CD44 and CD90 expression after isolation. Oxygen tension culture conditions of 1% and 3% O<sub>2</sub> led to OCT4 and SOX2 expression on culture days 2 and 4 (p<0.05), respectively, as compared to the hyperoxia condition (21% O<sub>2</sub>).

**Conclusion:** Based on the OCT4 and SOX2 immunofluorescence data, we conclude that the stem cells were pluripotent at low O<sub>2</sub> tension (at 1% O<sub>2</sub> on day 2 and at 3% O<sub>2</sub> on day 4), whereas under 21% O<sub>2</sub> the OCT4 and SOX2 were <sup>43</sup>not expressed.

**Keywords:** bone marrow, cluster differentiation 44, cluster differentiation 90, culture *in vitro*, low O<sub>2</sub> tension, mesenchymal stem cells, octamer-binding transcription factor 4, pluripotency, sex-determining region Y-box 2.

### Introduction

Pluripotency is the ability of stem cells to differentiate into all cell types <sup>27</sup> constitute the adult body. Multipotent of rabbit's bone marrow-derived mesenchymal stem cells (BM-MSCs) can be induced to become pluripotent *in vitro* when cultured under low O<sub>2</sub> tension conditions [1]. Low O<sub>2</sub> tension is required to induce a favorable microenvironment to maintain MSCs *in vitro*, mimicking their BM niche [2], to ensure their viability for transplantation [3]. The low O<sub>2</sub> tension in the BM helps to maintain stem cells adjusted to the body's normal physiological condition [4]. This main <sup>47</sup>ance would likely occur if the stem cells are in the G0 phase and not in a cycling state

(G1/S/G2/M) [5]. However, MSCs non-proliferative and undifferentiated *in vivo* if there is no injury [6].

MSCs require physiological O<sub>2</sub> tension levels to be 1% in the BM [7], 10-15% in the adipose tissue, and 2-9% in almost all other tissues [8]. Therefore, to obtain pluripotent characteristics *in vitro*, a low O<sub>2</sub> tension has to be guaranteed during cell culture procedures to achieve their suitable physiological microenvironment. However, the optimal concentration and duration of low O<sub>2</sub> tension to induce or maintain pluripotency have not yet been defined. Low O<sub>2</sub> tension has been used in *in vitro* microenvironments to support several stem cell types: 0-5% O<sub>2</sub> for hematopoietic stem cells [5]; 2% for adipose stem cells [9], 1-5% for neural stem cells (NSCs) [10], 3% during 7 days for human cord blood cells [11], and 2% for MSCs [12]. Recent studies have focused on searching for the *in vitro* factor combination necessary to control the proliferation of stem cells that would enable them to remain viable and undifferentiated [13] and to prevent apoptosis, senescence, or gene mutations.

<sup>1</sup> Copyright: Safitri, E. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.



In light of the potential therapeutic benefits that can derive from stem cell transplantation in patients with a degenerative disease or injury, understanding how to keep cell viability and pluripotency characteristics *in vitro* are of primordial importance [14]. Common *in vitro* culture conditions rely on hyperoxia (high O<sub>2</sub> levels), which represents non-physiological environment for stem cells that are used to hypoxia (low O<sub>2</sub> levels). Moreover, the pluripotency of stem cells *in vitro* is also affected by the cultivation time under low O<sub>2</sub> condition [12, 15]. For example, after 48 h of low O<sub>2</sub> tension, the role of the transcription factor hypoxia-inducible factor (HIF)-1 $\alpha$  is replaced by HIF-2 $\alpha$ , which has different target genes [15]. The expression of HIF target genes *in vitro* was suggested to influence the expression of the pluripotency genes [16] octamer-binding transcription factor 4 (OCT4), sex-determining region Y-box 2 (SOX2), NANOG [17], reduced expression (REX-1) [18], Kruppel-like factor 4, and c-MYC [19].

The exact procedure to induce pluripotency in MSCs using low O<sub>2</sub> tension *in vitro* before stem cell transplantation still remains unexplored. Therefore, this study aimed to determine the O<sub>2</sub> concentration (comparing 1%, 3%, and 21%) and the culture time (1, 2, 4, and 8 days) required to achieve pluripotency of MSCs *in vitro*.

## Materials and Methods

### Ethical approval

All experimental protocols were reviewed and approved by the Animal Care and Use Committee, the Institutional Animal Ethics Committee (Veterinary Ethical Clearance), Faculty of Veterinary Medicine, Universitas Airlangga.

### Study period and location

All experiments were conducted at Stem Cells Research Laboratory, Institute Tropical Disease, Universitas Airlangga, Surabaya 60115, Indonesia from January to April 2020.

### Animal source of MSCs

Seven-month-old New Zealand male rabbits (1800 g) were housed in polycarbonate cages with 12-h light-dark cycle at a constant temperature of 21°C. Rabbits were provided from Laboratory animal trial, Institute Tropical Disease, Universitas Airlangga, Surabaya 60115. Adult rabbits were provided a food pellet diet and fresh hay, vegetables, and water *ad libitum*. For the isolation of BM-MSCs, we used one femur from one male rabbit. The sampling technique of the BM-MSCs from the rabbit's femur was based on Lemeshow's method [20]. Animals were anesthetized before MSCs isolation.

### Isolation and culture of BM-MSCs

The femur from a male rabbit ( $\pm$ 3000 g) was excised and its BM cells were slowly flushed out from the bone cavity. The BM aspirate was transferred to a 15 mL Heparin tube (Thermo Fisher®,

USA) previously filled with 3 mL  $\alpha$ -minimum essential medium (Sigma-Aldrich®, USA) supplemented with 10% inactivated fetal bovine serum (FBS), 2 mL L-glutamine (Sigma-Aldrich®, USA), and 100 U/mL penicillin and streptomycin (Thermo Fisher, USA). The final concentration was 10 mL. The aspirate was then transferred to a 15 mL sterile tube and diluted with 1 $\times$  sterile phosphate-buffered saline (PBS) (Sigma-Aldrich, USA) to a total volume of 8 mL maintaining a neutral pH. The tube was then rinsed twice with 5 mL PBS and then the aspirate was transferred to Ficoll solution (Sigma-Aldrich®, USA) in a separated 15 mL tube. The aspirate was then coated with Ficoll before being centrifuged (Sorvall™ MX Mic 30 Ultracentrifuge, Thermo Fisher, USA) at 1600 rpm for 15 min at room temperature. After centrifugation, the collection was done from the "buffy coat" location on the surface of the Ficoll-PBS solution using a sterile Pasteur pipette and placed in a 15 mL tube (Thermo Fisher®, USA). Each sample was diluted with 1 $\times$  PBS to a total volume of 15 mL and the tube was turned 3-5 times to achieve an even mix. Next, the sample was centrifuged at 1600 rpm for 10 min, the supernatant was discarded and cells were suspended in 8 mL of cell culture medium before heating. After seeding between 5 cm<sup>2</sup> and 8 cm<sup>2</sup> (appropriate 200 million) of cells on the plate, cells were incubated at 37°C and 5% CO<sub>2</sub> and allowed to attach. Approximately 24 h later, the media and cells that were not attached were discarded and the dish was washed twice with 5 mL pre-heated 1 $\times$  PBS before 10 mL of fresh cell culture medium was added to the dish for further incubation. The cells were incubated at 37°C and 5% CO<sub>2</sub> and observed daily using an inverted microscope. Culture media were changed every 3 days. This process was continued until the cells reached a confluency of 80-90%. Forty-eight hours before cell collection, cells were balanced in the incubator at 37°C in a humidified tissue culture incubator containing 5% CO<sub>2</sub> and 95% air. Non-adherent cells were removed after 24 h and the presence of MSCs was confirmed under an inverted microscope [20].

### Identification of surface markers of BM-MSCs by flow cytometry

BM-MSCs were harvested by centrifugation at the end of the cell culture period and a single-cell suspension was prepared using trypsinization before flow cytometry analysis. Approximately 2 $\times$ 10<sup>5</sup> cells were washed twice with PBS. Cells were then incubated in test tubes or microtiter plates with unlabeled or fluorescently conjugated antibodies and analyzed by flow cytometry (Becton Dickinson FACSVerser, San Diego, USA). The antibodies for surface markers were a PerCP-Cy5.5 anti-human cross anti-rabbit CD45 (Cat # 45-0459-42, Bioscience, Thermo Fisher, USA) and an anti-cluster differentiation 90 (CD90) FITC anti-rabbit CD90 (Bs-0778R, Biossusa, San Diego, USA).



### Low O<sub>2</sub> tension in MSCs culture

At the third cell culture passage, BM-MSCs were collected and placed on culture dishes with diameter of 5 cm at a density of  $2 \times 10^7$  cells/cm<sup>2</sup>. Iscove's Modified Dulbecco's medium was supplemented with 15% FBS (BM-MSCs stimulator supplement), and antibiotics (100 IU penicillin and 100 µg/mL streptomycin) and was used to culture cells at a temperature of 37°C and with a mixture of 5% CO<sub>2</sub> and 95% air [21]. Low O<sub>2</sub> tensions (1 and 3%) [22,23] were achieved by integrating the culture flask into a specialized incubator for low O<sub>2</sub> tension (Modular Incubator Chamber).

The BM-MSCs at passage three post-isolation were divided into three treatment groups [22,23], each with ten replicates: Control group with BM-MSCs under hyperoxic conditions (O<sub>2</sub>=21%); treatment Group 1 with BM-MSCs under 1% O<sub>2</sub> tension; and treatment Group 2 with BM-MSCs under 3% O<sub>2</sub> tension. All groups were cultured for 1, 2, 4, or 8 days.

### Analysis of pluripotency induction


The analysis of the pluripotency induction was based on the identification of OCT4 and SOX2 proteins by immunofluorescence staining [24,25]. For this, MSCs were harvested, collected in a 15 mL tube, and fixed with methanol. After 15 min, rabbit FITC-conjugated anti-OCT4/POU5F1 polyclonal antibody Cat # 45-0459-42, Bioscience, Thermo Fisher, USA) and rabbit FITC-conjugated anti-SOX2 polyclonal antibody (Cat. No bs-0523R-FITC; Biossusa) were added. The samples were washed with PBS, placed on glass slide and then incubated in incubator at 37°C for 1 h so that it adheres and finally analyzed under a fluorescence microscope [21,25].

### Statistical analysis

The immunofluorescence data were analyzed using the IBM SPSS Statistics 20.0 (IBM Corp., NY, USA) and subjected to a *post hoc* normality Tukey's honestly significant difference test; all differences with  $p < 5\%$  were considered significant.

### Results

BM-MSCs are multipotent stem cells and thus do not express pluripotent cell-specific transcription factors such as OCT4 and SOX2. We tested whether low O<sub>2</sub> tension could induce rabbit BM-MSCs to become pluripotent *in vitro*. After being cultured under hypoxic (1% or 3% O<sub>2</sub>) or normoxic (21% O<sub>2</sub>) conditions, we characterized the OCT4 and SOX2 expression in BM-MSCs by immunofluorescence. We observed that OCT4 and SOX2 positive cells appeared after 2 and 4 days under 1% and 3% O<sub>2</sub> tension conditions (Figures-1-4), respectively, whereas under normal O<sub>2</sub> tension OCT4 and SOX2 expression was not identified ( $p < 0.05$ ) (Figures-5 and 6). Of note, the fluorescence intensity on day 4 at 3% O<sub>2</sub> tension was lower than on day 2 at 1% O<sub>2</sub> tension ( $p < 0.05$ ) (Tables-1 and 2 or Figures-7 and 8 for complete immunofluorescence quantification). This suggests that the



**Figure-1:** The positive expression on day 2 of the octamer-binding transcription factor 4 (OCT4) is 1% O<sub>2</sub> in hypoxic precondition treatment group during which cultivation time is 1, 2, 4, and 8 days using fluorescence microscope; (A) without a filter and microscope (400×); (B) containing a green filter, OCT4 positively shows green fluorescence (red arrow).

hypoxic conditioning led to the expression of transcription factors potentially enabling the BM-MSCs to become pluripotent. Furthermore, this observation indicates that the cultivation time under a specific O<sub>2</sub> tension also plays an important role in the timing of OCT4 and SOX2 expression after isolation.

Overall, this result shows that the treatment with low O<sub>2</sub> tension leads to the expression of the pluripotency transcription factors OCT4 and SOX2 in otherwise multipotent BM-MSCs, which may indicate that some cells were induced to become pluripotent.

### Discussion

Our analysis of pluripotency based on OCT4 and SOX2 immunofluorescence. The study found that human ESC cultured for 48 h under low O<sub>2</sub> conditions expressed HIF2-α, which was shown to be an upstream regulator of OCT4 and therefore essential for the maintenance of pluripotency. Interestingly, other transcription factors, such as SOX2 and NANOG, are also regulated by HIF2-α [15]. Furthermore, OCT4, SOX2, and NANOG maintain stem cell characteristics and suppress the expression of differentiation genes.


The relationship between hypoxia, HIF2-α, and pluripotency genes was previously highlighted by a




**Table-1:** Score of OCT4 in 1, 2, 4, and 8 days in some treatments.

Treatments	Average OCT4 expression score±SD in 1 day	Average OCT4 expression score±SD in 2 days	Average OCT4 expression score±SD in 4 days	Average OCT4 expression score±SD in 8 days
Conventional/ Hyperoxia O <sub>2</sub> 21% (C)	0.13 <sup>a</sup> ±0.50	0.14 <sup>a</sup> ±0.35	0.15 <sup>a</sup> ±0.60	0.15 <sup>a</sup> ±0.50
Low O <sub>2</sub> tension O <sub>2</sub> 1% (T1)	0.40 <sup>a</sup> ±0.35	1.75 <sup>b</sup> ±0.55	3.75 <sup>c</sup> ±0.45	3.70 <sup>c</sup> ±0.40
Low O <sub>2</sub> tension O <sub>2</sub> 3% (T2)	0.25 <sup>a</sup> ±0.50	0.50 <sup>a</sup> ±0.35	1.50 <sup>b</sup> ±0.30	1.45 <sup>b</sup> ±0.50

<sup>a,b,c,d</sup>Different superscripts in the same column or row were significantly different (p<0.005). OCT4=Octamer-binding transcription factor 4, SD=Standard deviation



**Figure-2:** The positive expression on day 2 of sex-determining region Y-box 2 (SOX2) is 1% O<sub>2</sub> in hypoxic precondition treatment group during which cultivation time is 1, 2, 4, and 8 days using the fluorescence microscope; (A) without filter and microscope (400×); (B) with a green filter, SOX2 positively shows green fluorescence (red arrow).



**Figure-3:** Positive expression on day 4 of the octamer-binding transcription factor 4 (OCT4) is 3% O<sub>2</sub> in hypoxic precondition treatment group during which cultivation time is 1, 2, 4, and 8 days using the fluorescence microscope; (A) without filter and microscope (400×); (B) with a green filter, OCT4 positively shows green fluorescence (red arrow).

study showing that the reprogramming efficiency during iPS cell generation from mouse embryonic fibroblasts was higher under hypoxia than under normoxia [26,27]. Furthermore, a study describing NSCs transplantation proposed that preconditioning under a hypoxic environment is necessary during cells *in vitro* maintenance [28]. This is in line with the high reliance of NSCs on low O<sub>2</sub> during early development [29]. The therapeutic efficacy of stem cell transplantation requires adequate knowledge of the mechanisms that affect stem cells during their *in vitro* maintenance, such as migration, proliferation, commitment, and dependence on oxygen levels in the appropriate culturing conditions.


Hypoxic culture conditions were previously linked to the intracellular stabilization pathway of HIF

transcription factors. Under low oxygen tension, the  $\alpha$  subunit of the HIF (HIF- $\alpha$ ) becomes stable and dominant due to the absence of Proline hydroxylase activity [30], which under normoxia acts as a HIF inhibitor through oxygen-dependent hydroxylation of a conserved asparagine in the COOH-terminal transactivation domain (CAD) of HIF- $\alpha$ . When oxygen is not available, the asparagine residue is not hydroxylated and transcriptional co-activators such as p300/CBP-CREB (cAMP-response element-binding protein) can interact with CAD, triggering the complete activity of the cell's response to hypoxia through the utilization of HIF1- $\alpha$ . After HIF1- $\alpha$  is stabilized, it translocates from the cytoplasm to the nucleus and undergoes a dimerization with HIF1- $\beta$  [31], also called ARNT. The dimerization between HIF1- $\alpha$  and HIF1- $\beta$  forms the HIF complex and requires the bHLH domain

**Table-2:** Score of SOX2 in 1, 2, 4, and 8 days in some treatments.

Treatments	Average SOX2 expression score±SD in 1 day	Average SOX2 expression score±SD in 2 days	Average SOX2 expression score±SD in 4 days	Average SOX2 expression score±SD in 8 days
Conventional/Hyperoxia O <sub>2</sub> 21% (C)	0.12 <sup>a</sup> ±0.40	0.14 <sup>a</sup> ±0.50	0.15 <sup>a</sup> ±0.40	0.15 <sup>a</sup> ±0.30
Low O <sub>2</sub> tension O <sub>2</sub> 1% (T1)	0.30 <sup>a</sup> ±0.40	1.95 <sup>b</sup> ±0.45	2.75 <sup>c</sup> ±0.35	2.95 <sup>c</sup> ±0.50
Low O <sub>2</sub> tension O <sub>2</sub> 3% (T2)	0.20 <sup>a</sup> ±0.50	0.40 <sup>a</sup> ±0.40	1.25 <sup>b</sup> ±0.40	1.40 <sup>b</sup> ±0.55


<sup>a,b,c,d</sup>Different superscripts in the same column or row were significantly different ( $p < 0.005$ ). SOX2=Sex-determining region Y-box 2, SD=Standard deviation



**Figure-4:** Positive expression on day 4<sup>th</sup> of the sex-determining region Y-box 2 (SOX2) is 3% O<sub>2</sub> in hypoxic precondition treatment group during which cultivation time is 1, 2, 4, and 8 days using fluorescence microscope; (A) without filter and microscope (400×); (B) with a green filter, SOX2 positively shows green fluorescence (red arrow).

and parts of the periodic acid-Schiff (PAS) domain. The bHLH domain was described to be also important for the binding of the HIF complex to DNA [32] specific DNA sequences, which are referred to as hypoxia response elements (HREs; 5'-TACGTG-12 activating the expression of target genes. Of note, the N-terminal transactivating domain confers target gene specificity to HIF [33].

Conversely, under normox<sup>51</sup> conditions, cells trigger the degradation pathway of HIF-1 $\alpha$  through the oxygen-sensitive hydroxylation of the HIF1- $\alpha$  subunit [34]. HIF1- $\alpha$  contains many proline (Pro) amino acid residues, namely, in the oxygen-dependent degradation domains, which are hydroxylated by prolyl hydroxylases (PHDs). PHDs catalyze this reaction




**Figure-5:** The negative expression of octamer-binding transcription factor 4 (OCT4) in 21% O<sub>2</sub> normoxia during which cultivation time is 1, 2, 4, and 8 days using fluorescence microscope; (A) without filter and microscope (400×); (B) with green filter, OCT4 negatively does not show green fluorescence.

through the addition of oxygen in the Pro-402 and Pro-564 residues turning them into 4-hydroxyproline. Oxygen and 2-oxoglutarate are required for the activity of three PHD isoforms (PHD1, PHD2, and PHD3), which have an equally high dependence on oxygen [34]. Following hydroxylation, HIF-1 $\alpha$  is destabilized, ubiquitinated, and then degraded by the proteasome [35]. According to Vriend and Reitter [36], the von-Hippel-Lindau E3 ubiquitin ligase, tumor suppression, plays an important role in the regulation of the ubiquitination process of HIF-1 $\alpha$  by binding to its hydroxylation subunit. Moreover, this study showed that HIF-1 $\alpha$  degradation culminates in the absence of OCT4 and SOX2 expression in culture of BM-MSCs.

Our study infers the pluripotency activity in BM-MSCs after low O<sub>2</sub> treatment, which is one of the main research focuses on the adult stem cells field. The balance between differentiation, apoptosis, and






**Figure-6:** Negative expression of sex-determining region Y-box 2 (SOX2) in 21% O<sub>2</sub> normoxia during which cultivation time is 1, 2, 4, and 8 days using fluorescence microscope; (A) without filter and microscope (400×); (B) with green filter, SOX24 negatively does not show green fluorescence.

self-renewal, characteristic of stem cells, needs to be achieved through the regulation by the microenvironmental niche where the stem cells are located. According to Mohyeldin *et al.* [37], in feeder-free *in vitro* culture conditions, the fate of the stem cells is affected by growth factors, interleukins, and serum composition, but is also influenced by the oxygen levels used for cell culture.


Pluripotency, the ability of a cell to differentiate into any cell belonging to the three embryonic layers (ectoderm, mesoderm, and endoderm), is generally an exclusive characteristic of ESCs. Compared to more committed progenitor cells, stem cells, in particular, ESCs possess a higher differentiation potential [13]. ESCs are derived from the inner cell mass of the blastocyst and are easy to differentiate *in vitro* into various types of cells, including nerve, blood, cardiac, and immune cells. In this study, we show that MSCs, adult stem cells that are generally defined as multipotent, can express OCT4 and SOX2 genes after being cultured under low O<sub>2</sub> tension. In the future, we expect to confirm that MSCs cultured under these *in vitro* conditions have a pluripotent potential and are also fully capable of differentiating into different cell types. This hypothesis is based on other studies, which show high efficiency of cell reprogramming to iPS cell state under low O<sub>2</sub> tension conditions [26].

### Conclusion

Based on immunofluorescence measurements, we suggest that some BM-MSCs have pluripotency



**Figure-7:** Score of octamer-binding transcription factor 4 in 1, 2, 4, and 8 days in some treatments.



**Figure-8:** Score of sex-determining region Y-box 2 in 1, 2, 4, and 8 days in some treatments.

characteristics since they express OCT4 and SOX2 at low O<sub>2</sub> tension (after 2 days under 1% O<sub>2</sub> and after 4 days under 3% O<sub>2</sub> tension), whereas at 21% O<sub>2</sub> tension OCT4 and SOX2 were not expressed. Thus, we conclude that *in vitro* low oxygen tension conditions increase OCT4 and SOX2 expression ( $p < 0.05$ ) as compared to conventional or hyperoxic conditions.

### Author's Contributions

ES: Conceptualization, methodology, data analysis, research and ethical clearance preparation of the research equipment, observation of Immunofluorescence method (OCT4 and SOX2), draft for manuscript preparation. Preparation of formal experimental and corresponding author. She has read and approved the final manuscript.

### Acknowledgments

The study was supported by funding from the Directorate General of Higher Education, Indonesia (DIKTI) 2020, The National Education Ministry, Republic of Indonesia, based on Decree number: 27/E1/KPT/2020 and agreement Contract number 4/AMD/E1/KP.PTNBH/2020 and 765/UN3.14/PT2020.

### Competing Interests

The author declares that she has no competing interests.

### Publisher's Note

Veterinary World remains neutral with regard to jurisdictional claims in published institutional affiliation.

## References

1. Safitri, E., Prasetyo, R.H., Hariadi, M. and Srianto, P. (2018) Hypoxic preconditioning for inducing pluripotency of rabbit's bone marrow-derived mesenchymal stem cells. *Biochem. Cell. Arch.*, 18(2): 1509-1517.
2. Lambertini, E., Penolazzi, L., Angelozzi, M., Bergamin, L.S., Manferdini, C., Viecelli, D. S.F., Paolella, F., Lisignoli, G. and Piva, R. (2018) Hypoxia preconditioning of Human MSCs: a Direct Evidence of HIF-1 $\alpha$  and Collagen Type XV Correlation. *Cell Phys. Biochem.*, 51:2237-2249.
3. Safitri, E. and Hariadi, E. (2019) Comparison of biotechnological culture of hypoxia-conditioned rat mesenchymal stem cells with conventional *in vitro* culture of normoxia-conditioned rat mesenchymal stem cells for testicular failure therapy with low libido in rats. *Vet. World*, 12(6): 916-924.
4. Safitri, E., Srianto, P., Widiyatno, T.V., Sandhika, W. and Prasetyo, R.H. (2018) Viability of rabbit adipocyte stem cells cultured under different oxygen concentrations *in vitro*. *Philipp. J. Vet. Med.*, 55(1): 1-8.
5. Daignan-Fornier, B. and Sagot, I. (2011) Proliferation/quiescence: When to start? Where to stop? What to stock? *Cell Div.*, 6(10): 1-5.
6. Rumman, M., Dhawan, J. and Kassem, M. (2015) Concise review: Quiescence in adult stem cells: Biological significance and relevance to tissue regeneration. *Stem Cells*, 33(10): 2903-2912.
7. Kim, D.S., Ko, Y.J., Lee, M.W., Park, H.J., Kim, Y.J., Sung, K.W., Koo, H.K. and Yoo, K.H. (2016) Effect of low oxygen tension on the biological characteristics of human bone marrow mesenchymal stem cells. *Cell Stress Chaperones*, 21(6): 1089-1099.
8. Gruber, M., Mathew, L.K., Runge, A.C., Garcia, J.A. and Simon, M.C. (2010) EPAS1 is required for spermatogenesis in the postnatal mouse testis. *Biol. Reprod.*, 82(6): 1227-1236.
9. Shi, S., Xie, J., Zhong, J., Lin, S., Zhang, T., Sun, K., Fu, N., Shao, X. and Lin, Y. (2016) Effects of low oxygen tension on gene profile of soluble growth factors in co-cultured adipose-derived stromal cells and chondrocytes. *Cell Prolif.*, 49(3): 341-351.
10. Mannello, F., Medda, V. and Tonti, G.A. (2011) Hypoxia and neural stem cells: From invertebrates to brain cancer stem cells. *Int. J. Dev. Biol.*, 55(6): 569-581.
11. Ivanovic, Z., Hermite, F., de la Grange, P.B., Dazey, B., Belloc, F., Lacombe, F., Vezon, G. and Praloran, V. (2004) Simultaneous maintenance of human cord blood SCID-repopulating cells and expansion of committed progenitors at low O<sub>2</sub> concentration (3%). *Stem Cells*, 22(5): 716-724.
12. Fan, L., Liu, R., Li, L., Shi, Z., Dang, X. and Wang, K. (2015) Low oxygen tension enhances osteogenic potential of bone marrow-derived mesenchymal stem cells with osteonecrosis-related functional impairment. *Stem Cell Int.*, 2015: 1-8.
13. Halim, D.H., Murty, H., Sandra, F., Boediono, A., Djuwantono, T. and Setiawan, B. (2010) Stem Cell Dasar Teori Dan Aplikasi Klinis. 1<sup>st</sup> ed. Erlangga, Indonesia. p154-196.
14. Masarani, M., Wazait, H. and Dinneen, M. (2006) Mumps orchitis. *J. R. Soc. Med.*, 99(11): 573-575.
15. Forristal, C.E., Wright, K.L., Hanley, N.A., Oreffo, O. and Houghton, F.D. (2010) Hypoxia inducible factors regulate pluripotency and proliferation in human embryonic stem cells cultured at reduced oxygen tensions. *Reproduction*, 139(1): 85-97.
16. Youssef, A. and Han, V.K.M. (2016) Low oxygen tension modulates the insulin-like growth factor-1 or -2 signaling via both insulin-like growth factor-1 receptor and insulin receptor to maintain stem cell identity in placental mesenchymal stem cells. *Endocrinology*, 157(3): 1163-1174.
17. Seymour, T., Twigger, A.J. and Kakulas, F. (2015) Pluripotency genes and their functions in the normal and aberrant breast and brain. *Int. J. Mol. Sci.*, 16(11): 27288-27301.
18. Festuccia, N., Osomo, R., Wilson, V. and Chambers, I. (2013) The role of pluripotency gene regulatory network components in mediating transitions between pluripotent cell states. *Curr. Opin. Genet. Dev.*, 23(5): 504-511.
19. Neganova, I., Shmeleva, E., Munkley, J., Chichagova, C., Anyfantis, G., Anderson, R., Passos, J., Elliott, D.J., Armstrong, L. and Lako, M. (2016) JNK/SAPK signaling is essential for efficient reprogramming of human fibroblasts to induced pluripotent stem cells. *Stem Cells*, 34(5): 1198-1212.
20. Hussain, S., Tamizhselvi, R., George, L. and Manickam, V. (2016) Assessment of the role of noni (*Morinda citrifolia*) juice for inducing osteoblast differentiation in isolated rat bone marrow derived mesenchymal stem cells. *Int. J. Stem Cells*, 9(2): 221-229.
21. Eleotério, R.B., Sepulveda, R.V., Reis, E.C.C., Valente, F.L. and Borges, A.P.B. (2016) Isolation, expansion and differentiation of mesenchymal stromal cells from rabbits' bone marrow. *Pesq. Vet. Bras.*, 36(5): 423-430.
22. Safitri, E., Utama, S., Widiyatno, T.V., Sandhika, W. and Prasetyo, R.H. (2016) Auto-regeneration of mice testicle seminiferous tubules due to malnutrition based on stem cells mobilization using honey. *Asian Pac. J. Reprod.*, 5(1): 30-34.
23. Prasetyo, R.H. and Safitri, E. (2016) Effects of honey to mobilize endogenous stem cells in efforts intestinal and ovarian tissue regeneration in rats with protein energy malnutrition. *Asian Pac. J. Reprod.*, 5(3): 198-203.
24. Safitri E., Utama, S., Bumi, C., Aulani'am, A. and Rantam, F.A. (2014) Hypoxic preconditioning for the regeneration of spermatogenesis process. *Adv. Nat. App. Sci.*, 8(8): 42-46.
25. De Miguel, M.P., Alcaina, Y., de la Maza, D.S. and Lopez-Iglesias, P. (2015) Cell metabolism under microenvironmental low oxygen tension levels in stemness, proliferation and pluripotency. *Curr. Mol. Med.*, 15(4): 343.
26. Rizzino, A. (2015) Concise review: The Sox2-Oct4 connection: Critical players in a much larger interdependent network integrated at multiple levels. *Stem Cells*, 31(6): 1033-1039.
27. Yoshida, Y., Takahashi, K., Okita, K., Ichisaka, T. and Yamanaka, S. (2009) Hypoxia enhances the generation of induced pluripotent stem cells. *Cell Stem Cell*, 5(3): 237-241.
28. Moreno, M., Fernandez, V., Monllau, J.M., Borrell, V., Lerin, C. and de la Iglesia, N. (2015) Transcriptional profiling of hypoxic neural stem cells identifies calcineurin-NEATc4 signaling as a major regulator of neural stem cell biology. *Stem Cell Reports*, 5(2): 157-165.
29. Wu, J., Sun, Z., Sun, S., Wu, J., Weisel, R.D. and Keating, A. (2008) Intravenously administered bone marrow cells migrate to damaged brain tissue and improve neural function in ischemic rats. *Cell Transplant.*, 16(10): 993-1005.
30. Ivan, M., Kond, K., Yang, H., Kim, W., Valindo, J., Ohh, M., Saic, A., Asara, J.M., Lane, W.S. and Kaelin, W.G. Jr. (2001) HIF1 $\alpha$  targeted for VHL-mediated destruction by proline hydroxylation: Implications for O<sub>2</sub> sensing. *Science*, 292(5516): 464-468.
31. Ku, Q. and Costa, M. (2006) Hypoxia-inducible factor-1 (HIF-1). *Mol. Pharmacol.*, 70(5): 1469-1480.
32. Zagorska, A. and Dulak, J. (2004) HIF-1: The knowns and unknowns of hypoxia sensing. *Acta Biochim. Pol.*, 51(3): 563-585.
33. Tuckerman, J.R., Zhao, Y., Hewitson, K.S., Tian, Y.M., Pugh, C.W., Ratcliffe, P.J. and Mole, D.R. (2004) Determination and comparison of specific activity of the HIF-prolyl hydroxylases. *FEBS Lett.*, 576(1-2): 145-150.
34. Ciria, M., Garcia, N.A., Ontoria-Oviedo, I.,



- Gonzalez-King, H., Carrero, R., De La Pompa, J.L., Montero, J.A. and Sepulveda, P. (2017) Mesenchymal stem cell migration and proliferation are mediated by hypoxia-inducible factor-1 $\alpha$  upstream of notch and SUMO pathways. *Stem Cells Dev.*, 26(13): 973-985.
35. Fandrey, J., Gorr, T.A. and Gassmann, M. (2006) Regulating cellular oxygen sensing by hydroxylation. *Cardiovasc. Res.*, 71(4): 642-651.
36. Vriend, J. and Reitter, R.J. (2016) Melatonin and the von hippel-Lindau/HIF-1 oxygen sensing mechanism: A review. *Biochim. Biophys. Acta*, 1865(2): 176-183.
37. Mohyeldin, A., Muvdi, T.G. and Hiojosa, A.Q. (2010) Oxygen in stem cell biology: A critical component of the stem cell niche. *Cell Stem Cell*, 7(7): 150-161.

\*\*\*\*\*

# Effect of low oxygen tension on transcriptional factor OCT4 and SOX2 expression in New Zealand rabbit bone marrow-derived mesenchymal stem cells

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://nutritionandmetabolism.biomedcentral.com">nutritionandmetabolism.biomedcentral.com</a> Internet Source	2%
2	Soyolmaa Gurdorj, Nyamdolgor Uranbileg, Mungun-Ochir B, Baararjargal P et al. "Study results of diagnosis of Maedi-visna disease in sheep", Mongolian Journal of Agricultural Sciences, 2020 Publication	1%
3	<a href="http://www.ijstemcell.com">www.ijstemcell.com</a> Internet Source	1%
4	<a href="http://oak.kribb.re.kr">oak.kribb.re.kr</a> Internet Source	1%
5	<a href="http://bmchealthservres.biomedcentral.com">bmchealthservres.biomedcentral.com</a> Internet Source	1%
6	<a href="http://downloads.hindawi.com">downloads.hindawi.com</a> Internet Source	1%
7	<a href="http://www.zora.uzh.ch">www.zora.uzh.ch</a> Internet Source	1%



8	<a href="http://vir.sgmjournals.org">vir.sgmjournals.org</a> Internet Source	<1 %
9	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	<1 %
10	M. Ivan Ariful Fathoni, Gunardi, Fajar Adi-Kusumo, Susanna Hilda Hutajulu. "Survival analysis of breast cancer patients in Yogyakarta", <i>Journal of Physics: Conference Series</i> , 2021 Publication	<1 %
11	<a href="http://microbenotes.com">microbenotes.com</a> Internet Source	<1 %
12	<a href="http://thesesups.ups-tlse.fr">thesesups.ups-tlse.fr</a> Internet Source	<1 %
13	Yongjin Hu. "Characterization of fermented black soybean natto inoculated with <i>Bacillus natto</i> during fermentation", <i>Journal of the Science of Food and Agriculture</i> , 2010 Publication	<1 %
14	<a href="http://imrpress.com">imrpress.com</a> Internet Source	<1 %
15	<a href="http://www.researchsquare.com">www.researchsquare.com</a> Internet Source	<1 %
16	<a href="http://www.scribd.com">www.scribd.com</a> Internet Source	<1 %

17

Alessandro Prigione. "The Senescence-Related Mitochondrial/Oxidative Stress Pathway is Repressed in Human Induced Pluripotent Stem Cells", *Stem Cells*, 2010

Publication

---

&lt;1 %

18

[bmcmusculoskeletdisord.biomedcentral.com](http://bmcmusculoskeletdisord.biomedcentral.com)

Internet Source

---

&lt;1 %

19

Lailatul Muniroh, Triska Susila Nindya, Santi Martini, Rondius Solfaine. "Effect of Plectranthus amboinicus Extract on BUN and Creatinine Levels and Cellular Response Proinflammatory Factors TNF- $\alpha$  and IL-1 $\beta$  on Gout Arthritis Patients", *Makara Journal of Health Research*, 2014

Publication

---

&lt;1 %

20

bai, hua, wei zhang, Xujun Qin, Tao Zhang, Hao Wu, Jangzheng Liu, and Chun X Hai. "Hydrogen peroxide modulates the proliferation/quiescence switch in the liver during embryonic development and post-hepatectomy regeneration", *Antioxidants & Redox Signaling*, 2015.

Publication

---

&lt;1 %

21

"Therapeutic Neovascularization—Quo Vadis?", Springer Science and Business Media LLC, 2007

Publication

---

&lt;1 %



22	<a href="http://d-nb.info">d-nb.info</a> Internet Source	<1 %
23	<a href="http://pdffox.com">pdffox.com</a> Internet Source	<1 %
24	<a href="http://www.microbiologyresearch.org">www.microbiologyresearch.org</a> Internet Source	<1 %
25	Furness, S.G.B.. "The dioxin (aryl hydrocarbon) receptor as a model for adaptive responses of bHLH/PAS transcription factors", FEBS Letters, 20070731 Publication	<1 %
26	Huiyuan Sun, Shukuan Ling, Dingsheng Zhao, Jianwei Li et al. "Ginsenoside Re Treatment Attenuates Myocardial Hypoxia/Reoxygenation Injury by Inhibiting HIF-1 $\alpha$ Ubiquitination", Frontiers in Pharmacology, 2020 Publication	<1 %
27	<a href="http://ecronicon.com">ecronicon.com</a> Internet Source	<1 %
28	<a href="http://www.lib.kobe-u.ac.jp">www.lib.kobe-u.ac.jp</a> Internet Source	<1 %
29	"Targeting Proteins for Destruction by the Ubiquitin System: Implications for Human Pathobiology", Annual Review of Pharmacology and Toxicology, 02/2009	<1 %

30

Elisabetta Lambertini, Letizia Penolazzi, Giulia Pelliello, Caterina Pipino et al. "Pro-Osteogenic Properties of Violina pumpkin (*Cucurbita moschata*) Leaf Extracts: Data from In Vitro Human Primary Cell Cultures", *Nutrients*, 2021

Publication

---

<1 %

31

Purwani, Ni Nyoman, Handoko Darmokoesoemo, and Ni Nyoman Tri Puspaningsih. "Hydrolysis of Corncob Xylan using  $\beta$ -xylosidase GbtXyl43B from *Geobacillus Thermoleovorans* IT-08 Containing Carbohydrate Binding Module (CBM)", *Procedia Chemistry*, 2016.

Publication

---

<1 %

32

Rosa Carbó, Marta Ginovart, Akatibu Carta, Xavier Portell, Luis J. del Valle. "Effect of aerobic and microaerophilic culture in the growth dynamics of *Saccharomyces cerevisiae* and in training of quiescent and non-quiescent subpopulations", *Archives of Microbiology*, 2015

Publication

---

<1 %

33

Vito Antonio Baldassarro, Alessandra Marchesini, Luciana Giardino, Laura Calzà. "Differential effects of glucose deprivation on the survival of fetal versus adult neural stem

<1 %



# cells - derived oligodendrocyte precursor cells", Glia, 2019

Publication

---

34	<a href="http://acikerisim.mu.edu.tr">acikerisim.mu.edu.tr</a> Internet Source	<1 %
35	<a href="http://ajnd.us">ajnd.us</a> Internet Source	<1 %
36	<a href="http://cob.silverchair-cdn.com">cob.silverchair-cdn.com</a> Internet Source	<1 %
37	<a href="http://dukespace.lib.duke.edu">dukespace.lib.duke.edu</a> Internet Source	<1 %
38	<a href="http://findresearcher.sdu.dk:8443">findresearcher.sdu.dk:8443</a> Internet Source	<1 %
39	<a href="http://in.umh-csic.es">in.umh-csic.es</a> Internet Source	<1 %
40	<a href="http://jcs.biologists.org">jcs.biologists.org</a> Internet Source	<1 %
41	<a href="http://liu.se">liu.se</a> Internet Source	<1 %
42	<a href="http://rep.bioscientifica.com">rep.bioscientifica.com</a> Internet Source	<1 %
43	<a href="http://www.deepdyve.com">www.deepdyve.com</a> Internet Source	<1 %
44	<a href="http://www.diss.fu-berlin.de">www.diss.fu-berlin.de</a> Internet Source	<1 %

---

---

45 "Treated Plasmodium berghei infected pregnant mice by Andrographis paniculata tablet (AS201-01) decreasing the TLR-4 expression and apoptosis index of placental tissue", Journal of Applied Pharmaceutical Science, 2018

Publication

<1 %

---

46 Johanna Myllyharju. "Prolyl 4-hydroxylases, key enzymes in the synthesis of collagens and regulation of the response to hypoxia, and their roles as treatment targets", Annals of Medicine, 2009

Publication

<1 %

---

47 Keiyo Takubo. "Chapter 18 The Hypoxia Regulatory System in Hematopoietic Stem Cells", IntechOpen, 2012

Publication

<1 %

---

48 Sophie V eriter, Najima Aouassar, Pierre-Yves Adnet, Marie-Sophie Paridaens et al. "The impact of hyperglycemia and the presence of encapsulated islets on oxygenation within a bioartificial pancreas in the presence of mesenchymal stem cells in a diabetic Wistar rat model", Biomaterials, 2011

Publication

<1 %

---

49 Suherni Susilowati, Imam Mustofa, Wurlina Wurlina, Indah Norma Triana, Suzanita Utama, Budi Utomo. "A 150 kDa Protein

<1 %



Derived from Bull Seminal Plasma Extended the Survival Time of Kacang Goat Sperm Stored at 5°C", Veterinary Medicine International, 2021

Publication

---

50

Szymon Grochans, Jan Korbecki, Donata Simińska, Wojciech Żwieręto et al. "CCL18 Expression Is Higher in a Glioblastoma Multiforme Tumor than in the Peritumoral Area and Causes the Migration of Tumor Cells Sensitized by Hypoxia", International Journal of Molecular Sciences, 2022

Publication

---

<1 %

51

K. Lundgren, C. Holm, G. Landberg. "Common Molecular Mechanisms of Mammary Gland Development and Breast Cancer", Cellular and Molecular Life Sciences, 2007

Publication

---

<1 %

52

Oihana Iriondo, Desirea Mecnas, Yilin Li, Christopher R. Chin et al. "Hypoxic memory of tumor intrinsic type I interferon suppression promotes breast cancer metastasis", Cold Spring Harbor Laboratory, 2022

Publication

---

<1 %

53

Takashi Semba, Rachel Sammons, Xiaoping Wang, Xuemei Xie, Kevin N. Dalby, Naoto T. Ueno. "JNK Signaling in Stem Cell Self-Renewal

<1 %

# and Differentiation", International Journal of Molecular Sciences, 2020

Publication

---

---

Exclude quotes      Off

Exclude matches      Off

Exclude bibliography      On