

Veterinary World

Editor-in-Chief

Anjum V. Sherasiya - Ex-Veterinary Officer, Department of Animal Husbandry, Gujarat State, India.

Editorial board

- Shambhunath Choudhary - Veterinary Pathologist II, Pathology | Charles River Laboratories, 640 North Elizabeth Street, Spencerville, OH 45887, USA.
- Suresh H. Basagoudanavar - FMD Vaccine Research Laboratory, Indian Veterinary Research Institute, Bangalore- 560024, Karnataka, India.
- R. G. Jani - Ex-Coordinator of Wildlife Health, Western Region Centre, Indo-US Project, Department of Veterinary Medicine, Veterinary College, Anand Agricultural University, Anand - 388001, Gujarat, India.
- G. N. Gongal - WHO South -East Asia Regional Office, New Delhi -110002, India.
- Md. Tanvir Rahman Department of Microbiology and Hygiene, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.
- Deepmala Agarwal - Cancer Prevention Laboratory, Pennington Biomedical Research Center, Baton Rouge, LA, USA.
- Fouad Kasim Mohammad - Department of Pharmacology & Toxicology, Vice President for Administrative & Financial Affairs, University of Mosul, P.O. Box 11136, Mosul, Iraq.
- Nicole Borel - Department of Pathology, Vetsuisse Faculty, University of Zurich, CH-8057 Zurich, Switzerland.
- B. A. Lubisi - Virology, MED Programme, ARC - Onderstepoort Veterinary Institute, No. 100 Old Soutpan Road, Onderstepoort, Tshwane, 0110, South Africa.
- Kumar Venkitanarayanan - Graduate Programs Chair, Honors and Pre-Vet Programs Advisor, Department of Animal Science, University of Connecticut, Storrs, CT 06269, USA.
- Kemin Xu - Department of Veterinary Medicine, University of Maryland, College Park College Park, MD, 20742, USA.
- Vassilis Papatsiros - Faculty of Veterinary Medicine, Department of Medicine (Porcine Medicine), University of Thessaly, Thessaly, Greece.
- K. P. Singh - School of Medicine and Dentistry, University of Rochester, Department of Environmental Medicine, Room: 4-6820, 601 Elmwood Avenue, Box-EHSC, Rochester, New York-14620, USA.
- Ashok K. Chockalingam - Division of Applied Regulatory Science, U.S. Food and Drug Administration, 10903, New Hampshire Avenue, Silver Spring, Maryland 20993, USA.
- Ashutosh Wadhwa - Center for Global Safe Water, Sanitation and Hygiene at Emory University, Hubert Department of Global Health, Rollins School of Public Health, Emory University, 1518 Clifton Rd. NE, Atlanta, GA 30322, USA.
- Luiz Otavio Pereira Carvalho - Laboratory of Immunomodulation and Protozoology, Oswaldo Cruz Institute, Ministry of health (Brazil), Pavilhao "108" - Sala: 09, Av. Brasil, 4365 - Manguinhos, Rio de Janeiro - RJ, CEP: 21040-360, Brazil.

- Mallikarjun Bidarimath - Cornell Stem Cell Program, Department of Biomedical Sciences, T2-012 Veterinary Research Tower, Cornell University, College of Veterinary Medicine, Ithaca, NY 14853-6401, USA.
- Chyer Kim - Virginia State University, Petersburg, VA, USA.
- Ionel D. Bondoc - Department of Public Health, Faculty of Veterinary Medicine Iasi, University of Agricultural Sciences and Veterinary Medicine Iasi, Romania.
- Filippo Giarratana - Department of Veterinary Medicine, University of Messina, Polo Universitario dell'Annunziata, 98168 Messina, Italy.
- Abdelaziz ED-DRA - Department of Biology, Faculty of Science, Moulay Ismail University, BP. 11201 Zitoune, Meknes, Morocco.
- Eduardo Jorge Boeri - Institute of Zoonosis Luis Pasteur, Buenos Aires, Argentina.
- Liliana Aguilar-Marcelino - CENID-PARASITOLOGIA VETERINARIA, Instituto Nacional de Investigaciones Forestales Agrícolas y Pecuarias: Jiutepec, Morelos, Mexico.
- Guilherme Dias de Melo - Institut Pasteur, Paris, Ile-de-France, France.
- Anut Chantiratikul - Department of Agricultural Technology, Faculty of Technology, Mahasarakham University, Muang, Mahasarakham Province 44150 Thailand.
- Panagiotis E Simitzis - Department of Animal Breeding and Husbandry, Faculty of Animal Science and Aquaculture, Agricultural University of Athens, 75 Iera Odos, 11855, Athens, Greece.
- Bartosz Kieronczyk - Poznan University of Life Sciences, Poznan, Greater Poland, Poland.
- Mario Manuel Dinis Ginja - Department of Veterinary Sciences, Center for Research and Agro-Environmental and Biological Technologies, University of Tras-os-Montes and Alto Douro, Portugal
- Nuh Kilic - Department of Surgery, Faculty of Veterinary Medicine, Adnan Menderes University, Turkey.
- Hanna Markiewicz - Milk Examination Laboratory, Kazimierz Wielki University in Bydgoszcz, Poland.
- Kai Huang - University of Texas Medical Branch at Galveston, Galveston, TX, USA.
- N. De Briyne - Federation of Veterinarians of Europe, Brussels, Belgium.
- Hasan Meydan - Akdeniz University, Faculty of Agriculture, Antalya, Turkey.
- Suleyman Cilek - Kirikkale Universitesi, Kirikkale, kirikkale, Turkey.
- Rodrigo Alberto Jerez Ebensperger - University of Zaragoza, Spain.
- Joao Simoes - Universidade de Tras-os-Montes e Alto Douro, Vila Real, Portugal.
- Alberto Elmi - University Of Bologna, Ozzano dell'Emilia, Bologna, Italy.
- Parag Nigam - Department of Wildlife Health Management, Wildlife Institute of India, Dehradun, India.
- Ali Aygun - Selcuk Universitesi, Konya, Turkey.

Volume - 14

No.11

November-2021

ISSN: 0972-8988

EISSN: 2231-0916



Scopus: Citescore-2.6, SJR-0.550, SNIP-1.387

Veterinary World is indexed in

Academic Journals Database, AGORA, AGRICOLA, AGRIS, CABI, CAS, DOAJ, EBSCO, ESCI - Thomson Reuters, Gale, Google Scholar, HINARI, Index Scholar, Indian Animal Science Abstracts, Indian Science Abstracts, JournalSeek, Open J-gate, ProQuest, PubMed, PubMed Central, SCOPUS, TEEAL

Veterinary World

Open access and peer reviewed journal



Editorial office

Veterinary World,
Star, Gulshan Park,
NH-8A, Chandrapur Road,
Wankaner - 363621,
Dist. Morbi, Gujarat, India
Website: www.veterinaryworld.org
E-mail: editorveterinaryworld@gmail.com

Printed and Published by Dr. Anjum V. Sherasiya on behalf of Veterinary World. Printed and Published at Star, Gulshan Park, N.H. 8A, Chandrapur Road, Wankaner-363621, Dist. Morbi, Gujarat, India.
Editor-in-Chief: Dr. Anjum V. Sherasiya

Veterinary World

ISSN: 0972-8988, EISSN: 2231-0916, www.veterinaryworld.org

Volume-14

No.11

November-2021

The articles in Veterinary World are open access articles licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Review (Published online: 03-11-2021)

1. Advances in nanotechnology and the benefits of using cellulose nanofibers in animal nutrition
Geovane Rosa de Oliveira, Carla de Andrade, Cristina Santos Sotomaior and Leandro Batista Costa
Veterinary World, 14(11): 2843-2850

Research (Published online: 06-11-2021)

2. Emergence of *Dirofilaria repens* (Spirurida: Onchocercidae) in dogs in Eastern Thailand
Wanarit Jitsamai, Patchana Kamkong, Sariya Asawakarn and Piyanan Taweethavonsawat
Veterinary World, 14(11): 2851-2854

Research (Published online: 09-11-2021)

3. Seroprevalence and risk factors for brucellosis in small ruminant flocks in Karnataka in the Southern Province of India
Krithiga Natesan, Triveni Kalleshamurthy, Mangadevi Nookala, Chaitra Yadav, Nagalingam Mohandoss, Somy Skariah, Swati Sahay, Bibek Ranjan Shome, Obli Rajendran Vinodh Kumar, Habibur Rahman and Rajeswari Shome
Veterinary World, 14(11): 2855-2862

Research (Published online: 10-11-2021)

4. Comparative study utilizing different post-breeding treatment regimens in cyclic Arabian mares
Khalid Mohammed Karam, Ahmed Saed Alebady, Haitham O. Alhilfi and Dhia Hussain Al-Delemi
Veterinary World, 14(11): 2863-2868

Research (Published online: 10-11-2021)

5. Multidrug resistance, biofilm formation, and virulence genes of *Escherichia coli* from backyard poultry farms
Theeb Al-Marri, Abdulla Al-Marri, Reham Al-Zanbaqi, Ahmad Al Ajmi and Mahmoud Fayez
Veterinary World, 14(11): 2869-2877

Research (Published online: 10-11-2021)

6. Comparison of polymerase chain reaction and microscopy for the detection of *Fasciola* spp. in the fecal matter of domestic bovines in Kalasin Province, Thailand
Sirikanda Thanasuwan and Anupong Tankrathok
Veterinary World, 14(11): 2878-2882

Research (Published online: 11-11-2021)

7. Prebiotic and synbiotic effect on rumen papilla length development and rumen pH in 12-week-old calves
A. Arne and A. Ilgaza
Veterinary World, 14(11): 2883-2888

Review (Published online: 11-11-2021)

8. Review of herbal medicine works in the avian species
Tyagita Hartady, Mas Rizky A. A. Syamsunarno, Bambang Pontjo Priosoeryanto, S. Jasni and Roostita L. Balia
Veterinary World, 14(11): 2889-2906

Research (Published online: 12-11-2021)

9. Phenytoin: A promising non-antibiotic drug for the topical treatment of digital dermatitis in dairy cows
El-Sayed El-Shafaey, Mohamed A. Hamed, Eman Abo Elfadl, Naglaa A. Gomaa and Mohamed Abdo Rizk
Veterinary World, 14(11): 2907-2912

Research (Published online: 12-11-2021)

10. Expression of the recombinant C-terminal of the S1 domain and N-terminal of the S2 domain of the spike protein of porcine epidemic diarrhea virus

Jiraporn Sritun, Natnaree Inthong, Siriluk Jala, Sakuna Phatthanakunanan, Khomson Satchasataporn, Kaitkanoke Sirinarumitr, Preeda Lertwatcharasarakul and Theerapol Sirinarumitr
Veterinary World, 14(11): 2913-2918

Research (Published online: 16-11-2021)

11. Efficacy of *Dipterocarpus alatus* oil combination with *Rhinacanthus nasutus* leaf and *Garcinia mangostana* pericarps against canine demodicosis

Atchara Artchayasawat, Parichart Boueroy, Thidarut Boonmars, Benjamabhorn Pumhirunroj, Pranee Sriraj, Ratchadawan Aukkanimart, Sirintip Boonjaraspinyo, Opal Pitaksakulrat, Panaratana Ratanasuwan, Apiporn Suwannatrai, Chatanun Eamudomkarn, Porntip Laummaunwai and Wu Zhiliang
Veterinary World, 14(11): 2919-2928

Research (Published online: 21-11-2021)

12. Isolation and antibiotic sensitivity of *Campylobacter* species from fecal samples of broiler chickens in North West Province, South Africa

Kealeboga Mileng, Tsepo A. Ramatla, Rendani V. Ndou, Oriel M. M. Thekiso and Michelo Syakalima
Veterinary World, 14(11): 2929-2935

Research (Published online: 21-11-2021)

13. In vitro cytotoxicity and virucidal efficacy of potassium hydrogen peroxymonosulfate compared to quaternary ammonium compound under various concentrations, exposure times and temperatures against African swine fever virus

Watcharee Sovijit, Machimaporn Taesuji, Khate Rattanamas, Darsaniya Punyadarsaniya, Thanongsak Mamom, Hoa Thi Nguyen and Sakchai Ruenphet
Veterinary World, 14(11): 2936-2940

Research (Published online: 24-11-2021)

14. A survey of the usage (frequency and pattern) of antibiotics at the University of Maiduguri Veterinary Teaching Hospital, Maiduguri, Borno State, Nigeria

Nubwa Daniel, Kefas David Malgwi, Bukar Umaru, Isaac John Omeh and Ladi Sanya
Veterinary World, 14(11): 2941-2946

Research (Published online: 24-11-2021)

15. Seasonal content of heavy metals in the "soil-feed-milk-manure" system in horse husbandry in Kazakhstan

Nazym Kozhanova, Nurzhan Sarsembayeva, Bozena Lozowicka and Zhassulan Kozhanov
Veterinary World, 14(11): 2947-2956

Research (Published online: 24-11-2021)

16. Inactivated vaccine against Aujeszky's disease

Zhanat B. Kondibaeva, Bolat A. Yespembetov, Khairulla B. Abeuov, Assiya K. Mussayeva, Sarsenbek T. Siyabekov, Saltanat T. Nussupova, Elmira K. Akmatova, Yerlan K. Pazylov, Kydyrbay T. Maikhin and Nazym S. Syrym

Veterinary World, 14(11): 2957-2963

Research (Published online: 25-11-2021)

17. The ability of probiotics to ameliorate blood and gonad damage caused by copper toxicity in Nile tilapia (*Oreochromis niloticus*)

Alfiah Hayati, Manikya Pramudya and Hari Soepriandono
Veterinary World, 14(11): 2964-2970

Research (Published online: 25-11-2021)

18. Evaluation of secondary metabolites of herbal plant extracts as an antiviral effect on infectious bursal disease virus isolates in embryonated chicken eggs

Rawaa Saladdin Jumaa, Dhuha Ismael Abdulmajeed and Abdulkarim Jafar Karim
Veterinary World, 14(11): 2971-2978

Research (Published online: 26-11-2021)

19. Preliminary study of coronavirus disease 2019 on pets in pandemic in Denpasar, Bali, Indonesia

Hamong Suharsono, Ali Ghufon Mukti, Ketut Suryana, I. Wayan Masa Tenaya, Dilasdita Kartika Pradana, Guy Daly and Mochamad Panji Pujasakti
Veterinary World, 14(11): 2979-2983

Review (Published online: 26-11-2021)

20. Nociceptive pain and anxiety in equines: Physiological and behavioral alterations

I. Hernández-Avalos, D. Mota-Rojas, J. E. Mendoza-Flores, A. Casas-Alvarado, K. Flores-Padilla, A. E. Miranda-Cortes, F. Torres-Bernal, J. Gómez-Prado and P. Mora-Medina

Veterinary World, 14(11): 2984-2995

Research (Published online: 27-11-2021)

21. Cross-sectional seasonal prevalence and relative risk of ectoparasitic infestations of rodents in North Sinai, Egypt

Doaa S. Farid, Nahla H. Sallam, Ahmed M. Salah Eldein and Essam S. Soliman

Veterinary World, 14(11): 2996-3006

Research (Published online: 27-11-2021)

22. Effect of novel natural feed additive containing Averrhoa bilimbi L. fruit filtrate, wheat bran, and Saccharomyces cerevisiae on growth performance and meat characteristics of broilers

Sugiharto Sugiharto, Anugrah R. Pratama, Turrini Yudiarti and Tugay Ayasan

Veterinary World, 14(11): 3007-3014

Research (Published online: 27-11-2021)

23. Prevalence of the MDR1 gene mutation in herding dog breeds and Thai Ridgebacks in Thailand

Chommanad Lerdkrai and Nuch Phungphosop

Veterinary World, 14(11): 3015-3020

Research (Published online: 28-11-2021)

24. Occurrence of Marek's disease in vaccinated Algerian broiler breeder flocks: A histopathological survey

Abdelaziz Lounas, Mohamed Besbaci, Madjid Akkou and Oumennoune Tali

Veterinary World, 14(11): 3021-3027

Research (Published online: 28-11-2021)

25. Increasing the efficiency of the herd reproduction system by introducing innovative technologies into dairy farming in Northern Kazakhstan

Vitaly Anatolevich Raketsky, Askar Myrzakhmetovich Nametov, Vasily Arkadyevich Sozinov and Abdrakhman Abdybekuly Baisakalov

Veterinary World, 14(11): 3028-3037

Research (Published online: 29-11-2021)

26. Microbiological, sensory, and physicochemical quality of curd cheeses originating from direct sales

Ewa Januś, Piotr Sablik, Małgorzata Jakubowska and Klaudia Wróbel

Veterinary World, 14(11): 3038-3047

Research (Published online: 29-11-2021)

27. Evaluation of the phytochemical content, antimicrobial and antioxidant activity of Cocos nucifera liquid smoke, Garcinia mangostana pericarp, Syzygium aromaticum leaf, and Phyllanthus niruri L. extracts

Tiurma Pasaribu, Arnold P. Sinurat, Elizabeth Wina and Triwardhani Cahyaningsih

Veterinary World, 14(11): 3048-3055

Research (Published online: 30-11-2021)

28. Effectiveness of mesenchymal stem cells cultured under hypoxia to increase the fertility rate in rats (*Rattus norvegicus*)

Erma Safitri and Hery Purnobasuki

Veterinary World, 14(11): 3056-3064

Research (Published online: 30-11-2021)

29. In vitro experiments of *Pediculus humanus capitis* (Phthiraptera: Pediculidae) resistance to permethrin and 6-paradol in East Jakarta: Detoxification enzyme activity and electron microscopic changes in lice

Rizal Subahar, Lisawati Susanto, Rachmanin Aidilla, Annisa Putri Aulia, Yulhasri Yulhasri, Rawina Winita, Nadar S. Lubis and Ika Puspa Sari

Veterinary World, 14(11): 3065-3075

Effectiveness of mesenchymal stem cells cultured under hypoxia to increase the fertility rate in rats (*Rattus norvegicus*)

Erma Safitri¹ and Hery Purnobasuki²

1. Department of Veterinary Science, Division of Veterinary Reproduction, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya 60115, Indonesia; 2. Department of Biology, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60115, Indonesia.

Corresponding author: Erma Safitri, e-mail: erma-s@fkh.unair.ac.id

Co-author: HP: hery-p@fst.unair.ac.id

Received: 22-06-2021, **Accepted:** 20-10-2021, **Published online:** 30-11-2021

doi: www.doi.org/10.14202/vetworld.2021.3056-3064 **How to cite this article:** Safitri E, Purnobasuki H (2021) Effectiveness of mesenchymal stem cells cultured under hypoxia to increase the fertility rate in rats (*Rattus norvegicus*), *Veterinary World*, 14(11): 3056-3064.

Abstract

Background and Aim: Mesenchymal stem cells (MSCs) transplanted into the testes of rats with testicular failure can help rescue fertility. However, the low viability of transplanted MSCs limits the success of this treatment. This study aimed to determine the effectiveness of MSCs cultured under hypoxia to increase the fertility rate in rats (*Rattus norvegicus*).

Materials and Methods: Bone marrow-derived MSCs (200 million cells/rat) were transplanted into male rat models with induced infertility (10 rats/treatment group) after 4 days of culture in 21% O₂ (normoxia) and 1% O₂ (hypoxia). Ten fertile and 10 untreated infertile rats served as controls. In the infertile male rats that had been fasted from food for 5 days, the fasting condition induced malnutrition and then resulted in testicular failure.

Results: The results indicated that the MSCs cultured under hypoxic conditions were more effective than those cultured in normoxic conditions as a treatment for testicular failure in infertile male rats based on the increased number of cells expressing p63 as a quiescent cell marker and ETV5 as a transcription factor expressed in Sertoli and germ cells. Furthermore, the structure of the seminiferous tubules, which contain spermatogonia, primary and secondary spermatocytes, and spermatid, Sertoli, and Leydig cells, was improved in infertile male rats treated with the MSCs cultured under hypoxic conditions.

Conclusion: The testicular transplantation of MSCs cultured under hypoxic conditions was an effective treatment for testicular failure in rats.

Keywords: fertility, hypoxia culture, malnutrition, stem cell, testicular failure.

Introduction

Mesenchymal stem cell (MSC) transplantation using rabbit [1] and rat [2] bone marrow and later rat [3] and rabbit [4] adipose tissue was shown to be effective in rebuilding the tissues supporting the endogenous stem cells, allowing them to multiply and mature into sperm cells. The viability of the transplanted MSCs from the bone marrow [2,5], adipose tissue [3], or umbilical cord blood [6] is low; hence, this treatment has limited efficacy. The reduced viability of MSCs is thought to be caused by a normoxia culture with a high oxygen concentration (O₂>20%). Cell senescence [7], cell apoptosis [8], and gene mutation [9] can all be caused by normoxia culture. As a result, the low viability of MSCs restricts the success of cell transplant therapy. It is hypothesized that the efficiency of MSC transplantation is influenced by apoptosis [10-15]. To achieve therapeutic impact, substantial doses of MSCs are necessary. Several

researchers are attempting to acquire adequate dosage without the use of boosters, hence reducing the influence on rising expenses. Due to these issues, the effectiveness of the treatment remains unclear. Thus, alternative investigations are still required to learn more about the efficacy of more relevant treatments.

Other studies have indicated the critical significance of stem cell cultivation under hypoxic conditions as follows: To keep transplanted MSCs alive and adaptable, they were grown in a hypoxic environment (1-3% O₂) [2,4]. This condition was induced by cells in the quiescent state [4,16,17], allowing the cells to live longer [18,19]. Hypoxia-inducible factor 2 (HIF2), a critical regulator of progenitor stem cell function, may influence the expression of p63 as a definite marker in quiescent cells. Quiescent MSCs are self-renewing stem cells that remain in gap 0 and do not cycle (i.e., gap 1/synthesis/gap 2/mitosis) [20] or in undifferentiated states [21]. However, there is still a high potential for cell renewal [22]. At the indifferent stem stage, self-renewal is a symptom of the biological process and defense mechanism [23]. The homing signal based on the vascular endothelial growth factor (VEGF) expression following transplantation is critical for culture hypoxia-conditioned rat MSCs (1% O₂ concentration) [2]. VEGF is a component of the stem cell extracellular matrix that helps

Copyright: Safitri and Purnobasuki. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

maintain a favorable milieu for stem cells to survive after transplantation.

Scientific evidence on the effectiveness of MSCs cultured under hypoxia for testicular failure is still lacking. Therefore, we determined whether HIF2 (with HIF2 alpha [HIF2 α] monoclonal antibody [ep190b] as a marker) regulated the transplantation of MSCs in the form of quiescent MSCs (with p63/TP73L monoclonal antibody as a marker) derived from rat bone marrow and whether HIF2 played a role crucial for spermatogonial stem cells (SSCs). Infertile males with testis failure can be treated with transplantation of MSCs cultured under hypoxic conditions. The findings of this study are relevant in the area of male reproductive health.

This study aimed to determine the effectiveness of MSCs cultured under hypoxia to increase the fertility rate in rats (*Rattus norvegicus*).

Materials and Methods

Ethical approval

The study was approved by Animal Care and Use Committee (No: 239-KE; *Komisi Etik Penelitian*) of the Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia.

Study period and location

The study was conducted from March 2018 to 2020 at Department of Veterinary Science, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia.

Stem cell isolation

Stem cells were harvested from the bone marrow through aspiration of the femur, tibia, and ulna [24] of 3-month-old rats (*Rattus norvegicus*) [25]. The aspirate was placed in heparinized tubes (Z181099, Sigma-Aldrich®, Burlington, Massachusetts, USA) and stored at 4°C to be transported to the laboratory [26].

Stem cell culture

The aspirate from the rat bone marrow was transferred into 15 mL sterile tubes (SIAL0790-500EA, Sigma centrifuge tubes, Sigma-Aldrich®), rinsed twice with 5 mL sterile phosphate-buffered saline (PBS), (MFCD00131855, Sigma-Aldrich®), and filled up to a total volume of 10 mL. The diluted sample was added with the same volume of Ficoll (Biowest, Nuaille, France) in a separate 15 mL tube. Centrifugation was performed for 15 min at room temperature (37°C) at 1600 rpm. After centrifugation, the cells were collected from the Ficoll-PBS interface using a sterile Pasteur pipette (Corning™ C7095BNMR, Thermo Fisher Scientific, Waltham, MA, USA) and transferred into a 15 mL tube. The cells were resuspended in PBS up to a total volume of 15 mL. The tube was gently inverted and shaken (CLS6791 Sigma, Corning LSE Benchtop Shaking Incubator with Platform, Sigma-Aldrich®) 5 times to homogenize the suspension.

The suspension was centrifuged again for 10 min. The supernatant and floating cells were discarded, and the cell pellet was resuspended in 6 mL of

alpha-modified essential medium (α -MEM) (M0894; Sigma-Aldrich). Mononucleated cells were placed on a plate in 10 cm² with approximately 2 \times 10⁷ cells and incubated at 37°C in a humidified atmosphere with 5% CO₂ (BioSpherix, Florida, USA) for 24 h to let the cells adhere (sticking in a plate). After 24 h, the media and non-adherent cells were discarded. The adherent cells were rinsed twice with 5 mL PBS, and 10 mL fresh α -MEM medium was then added into the dish, which was returned into the incubator. The culture was observed daily under an inverted microscope. The medium was changed every 4 days, preceded by a rinse with 10 mL PBS, after which 10 mL fresh α -MEM medium was added. The culture was continued until approximately 75-80% confluence was achieved. After confluence, the cells were passaged into several other dishes to cultivate subcultures [26]. Passaging was performed three times, and then, the cells were assigned to two conditions: Hypoxic precondition treatments of 1% in a hypoxia chamber (BioSpherix, Florida, USA) inside a 5% CO₂ incubator and another treatment using 21% O₂ concentration (normoxia) over 4 days. The MSCs were observed under a microscope before being transferred into the testes.

Infertility rat model

This study used male rats with testicular failure that had been fasted from food for 5 days but provided with drinking water *ad libitum* [2,27]. The fasting condition for 5 days induced malnutrition and then resulted in testicular failure. The malnutrition condition caused the adrenal cortex to function suboptimally in producing dehydroepiandrosterone (DHEA). Low levels of DHEA in the blood can cause fatigue and decrease sperm concentration. DHEA is the most potent precursor of steroid hormones, such as testosterone, which is produced by the renal adrenal cortex [28] and Leydig cells of the testis [29]. Low testosterone production can lead to decrease spermatogenesis and, thus, testicular failure. The animal models used in this study were healthy 8-10-week-old Wistar strain male rats (*R. norvegicus*) with a body weight of 250-300 g. The rats were placed in individual plastic cages in the experimental animal laboratory at the Faculty of Veterinary Medicine, Universitas Airlangga.

MSC transplantation methods

The MSCs were transplanted into male rats with testicular failure, which were then compared with the negative and positive control rats. The T1 group consisted of 10 infertile male rats that were transplanted with stem cells cultured under normoxia (21% O₂ concentration) for 4 days with a dose of 200 million cells/rat [1]. The T2 group consisted of 10 infertile male rats that were transplanted with stem cells cultured under hypoxia (1% O₂ concentration) for 4 days with a dose of 200 million cells/rat [1]. The positive (fertile) control group was composed of 10 normal male rats (fertile) injected with 0.1 mL PBS. The negative

(infertile) control group consisted of 10 infertile male rats injected with 0.1 mL PBS.

The testes of the male rats were surgically excised after 54 days (one cycle of spermatogenesis) [30] to collect testicular tissue. The improvement in the testicular tissue was observed by histopathological preparations using hematoxylin and eosin (H&E) stain (B8438, Sigma-Aldrich®). Immunohistochemical (IHC) observation was performed to determine the expression of p63 (with p63/TP73L monoclonal antibody) as a marker of quiescent cells, HIF2 α (with HIF2 α monoclonal antibody [ep190b]) as a marker that is crucial for endogenous cells (SSCs, Sertoli cells, etc.), and ETV5 (with ETV5 monoclonal antibody) as a marker for the transcriptional factor to improve testicular failure and infertility. *In vitro* fertilization between the ovum and sperm cells was performed to observe the fertility of the male rats.

Histopathological assessment

Histopathological examination of the testicular tissues for the presence of Sertoli cells, Leydig cells, spermatogonia, spermatocytes, and primary and secondary and spermatid cells was performed on the testicular tissues that were fixed with 10% formalin. The testes were then dehydrated through a series of increasing alcohol concentrations, cleared with xylol, and embedded in paraffin. Thin sections mounted on slides were processed for H&E staining [31].

Histopathological examination was performed using a light microscope with a magnification of 200 \times . Five fields of view were assessed for each slide. Observations and identification of the spermatogonia and Sertoli and Leydig cells and regeneration of seminiferous tubules were based on the existing histological description [27].

IHC observation

IHC observation was performed to determine the expressions of p63, HIF2 α , and ETV5. The samples were prepared for histopathological examination of the testicular tissue. After deparaffinization of the preparation (paraffin block) with xylene 3 times each for 3 min, rehydration of the preparation with 100%, 95%, and 70% ethanol each for 2, 2, and 1 min and finally with water for 1 min was performed. Subsequently, the preparation was soaked in peroxidase blocking solution at 37°C for 10 min and then incubated in pre-diluted blocking serum at 25°C for 10 min.

In the next procedure, the IHC technique used monoclonal antibodies to determine the expressions of p63, HIF2 α , and ETV5. The preparation was soaked in p63 (MAB1916, Monoclonal Anti-p63/TP73L antibody, R&D Systems Inc., Minneapolis, USA), HIF2 α , (MA1-16519, HIF-2 α Monoclonal Antibody (ep190b), Thermo Fisher Scientific), and ETV5 (MBS831454, Mouse anti-Human ETV5 Monoclonal Antibody, MyBioSource, San Diego, USA) monoclonal antibodies for 10 min and then washed with PBS for 5 min.

Subsequently, the preparation was incubated with a secondary antibody (conjugated to horseradish peroxidase) at 25°C for 10 min, washed with PBS for 5 min, and then incubated again with peroxidase at 25°C for 10 min. Next, the preparation was washed with PBS for 5 min and then incubated with chromogen diaminobenzidine at 25°C for 10 min.

Furthermore, the preparation was incubated with H&E for 3 min and washed with water. Finally, the preparation was cleaned, dropped with mounting media, and closed with a coverslip. Then, the expressions of p63, HIF2 α , and ETV5 (brown color) were observed on the cells using a light microscope with 400 \times . Five fields of view (one tubule/field view) were assessed for each slide through the scoring system. The following IHC scoring system [32] was used: IHC score = A \times B, where A denotes the wide percentage of expressions and B is the intensity of the chromogen color (Table-1).

Medium preparation, sperm and oocyte collection, and *in vitro* fertilization

The media, M16 (MR-016 EMD Millipore, Sigma-Aldrich Inc., Darmstadt, Germany) and PBS (10010031 PBS, pH 7.4; Thermo Fisher Scientific), were manufactured in accordance with the established procedure for producing these two media. Before use for *in vitro* fertilization, a droplet medium was prepared in a Petri dish (Sterilin™ 100 mm Petri Dishes, Thermo Fisher Scientific) with a volume of 50 μ L as a washing medium and 25 μ L as an *in vitro* culture medium. The droplet medium was then incubated for 3 h in a 5% CO₂ incubator at 37°C before being used for *in vitro* fertilization [33].

Sperm cells were collected from male rats after being sacrificed by dislocation of the fourth cervical spine and then disinfected with 70% alcohol. A Y-shaped incision was made in the abdomen, the stomach contents were removed, and the left testicle was pulled out. The fats were separated, and then part of the cauda epididymis, the mature sperm storage, was taken. The obtained cauda epididymis was washed with PBS twice, cut into small pieces to free

Table-1: Semi-quantitative IHC scale taking into account both percentage of positive cells (A) and intensity of reaction color (B) with the final score representing product of the two variables (A \times B).

A	B
0 patients no cells with positive reaction	0 patients no color reaction
1 patient 10% cells with positive reaction	1 patient low intensity of color reaction
2 patients 11-50% cells with positive reaction	2 patients moderate intensity of color reaction
3 patients 51-80% cells with positive reaction	3 patients intense color reaction
4 patients >80% cells with positive reaction	

IHC=Immunohistochemical

the spermatozoa, placed on M16, and then incubated in a 5% CO₂ incubator at 37°C [34].

Before the oocytes were collected, PMSG and hCG hormones were injected subcutaneously to stimulate superovulation: On the 0th h on the 1st day, female rats were injected with 5 IU of 0.1 mL PMSG to stimulate the process of folliculogenesis and left to stand for 48 h. The female rats were injected with 5 IU of 0.1 mL hCG 48 h after the PMSG was injected and mated directly with a single vasectomized male to stimulate ovulation. After 17 h, a vaginal plug was used to confirm that the female had mated. Then, the oocytes were flushed. The female rats were killed by dislocation of the fourth cervical spine. A Y-shaped incision was then made on the abdomen, and the uterus was removed and separated from the fallopian tube section and then rinsed with M16. The oocytes from the ampulla of the fallopian tube were flushed using an inverted microscope [35].

Fertility rate observations

The fertility of spermatozoa was determined by examining *in vitro* fertilization. Semen analysis was conducted according to the guidelines of the World Health Organization. Semen was processed over a two-layer discontinuous density gradient, formed by a top layer of 40% (v/v) PureSperm (Nidacon Lab AB, Gothenburg, Sweden) and a lower layer of 80% (v/v) PureSperm, by centrifugation at 1500 g for 15 min at 37°C. The pellet was resuspended in 3 µL SAGE fertilization medium with 5% HSA and spun down at 200 g for 10 min at 37°C [36]. The oocyte and sperm were both placed in the Petri dishes containing M16 drops under a mineral oil overlay and incubated in 5% CO₂ incubators at 37°C for 5 h for *in vitro* fertilization [37]. Fertilization was confirmed by the presence of the second polar body of oocyte [36] through an inverted microscope (Nikon Eclipse TE 2000S; Nikon, Tokyo, Japan) at 400× with Hoffman modulation optics. In rats, the first polar body is known to degenerate.

The embryo quality was evaluated on days 2-3. On days 2-3, embryo development was assessed, including blastomere number, size, regularity, presence, and percentage of fragmentation. All embryos were graded on a scale of 1-5, with 1 being the best. Grade 1 embryos had symmetrical blastomeres of equal size with no cytoplasmic fragmentation. Grade 2 embryos had blastomeres of equal size and minor cytoplasmic fragmentation covering <10% of the embryo surface. Grade 3 embryos had even or uneven blastomeres and minor cytoplasmic fragmentation covering 10-25% of the embryo surface. Grade 4 embryos had blastomeres of equal or unequal size and moderate to high cytoplasmic fragmentation covering 25-50% of the embryo surface. Grade 5 embryos contained few blastomeres of any size and severe fragmentation covering >50% of the volume of the embryo. Embryos with good quality were defined as those with six to

eight equal-sized cells and <10% fragmentation on day 3 [36]. The fertility rate of sperm formula in rats was calculated as follows: Number of good-quality embryos/the number of mature oocytes × 100% [35].

Statistical analysis

The expressions of p63, HIF2α, and ETV5 and the fertility rate of the sperm cells were statistically analyzed using the SPSS software (v. 17 for Windows XP; SPSS Inc., Chicago, IL, USA) with a 99% confidence level ($\alpha=0.01$) and 0.05 significant difference ($p<0.05$). The comparative steps for hypothesis testing were as follows: Normality data test, Kolmogorov-Smirnov test, homogeneity of variance test, analyses of variance, and Tukey's HSD *post hoc* test with 5% least significant difference.

Results

Data were collected from 40 male rats, which were divided into four treatment groups: Normal males in the positive (fertile) control, infertile males treated with PBS in the negative (infertile) control, infertile males transplanted with stem cells cultured under normoxia (21% O₂ concentration) for 4 days in the first treatment (T1) group, and infertile males transplanted with stem cells cultured under hypoxia (1% O₂ concentration) for 4 days in the second treatment (T2) group. The results indicated that transplantation with MSCs from hypoxic precondition culture improves testicular function by decreasing the extent of damage and increasing fertility. The expression levels of p63, HIF2α, and ETV5 increased, as were the regeneration of the testicular tissue, as described in more detail in the following, such as intact tubular seminiferous tissue; formation of Sertoli cells, Leydig cells, spermatogonia, spermatocytes, and primary-secondary and spermatid cells; and improvement of the fertility rate of sperm *in vitro*.

Expression of p63

The average score of the p63 expression in the T2 group was 6.0^b±0.50, although the score was lower than that in the positive (fertile) control group, in which p63 was expressed (9.60^a±0.44), but the score was still much higher than that in the T1 (2.5^c±0.25) and negative (infertile) control groups, in which p63 was not expressed (0.41^d±0.22) (Figure-1 and Table-2).

Expression of HIF2α

The average score of the HIF2α expression in the T2 group was 7.0^a±0.75. This was the highest among those in the other groups, T1 (3.9^b±0.44), positive (fertile) control (0.6^c±0.34), and negative (infertile) control (0.3^c±0.33) (Figure-2 and Table-2). An increase in the score of the HIF2α expression could occur if they were given a low O₂ concentration.

Expression of ETV5

The average score of the ETV5 expression in the T2 group was 7.2^b±0.34, although the score was lower in the positive (fertile) control group (10.5^a±0.25), but the score was still higher than that in the T1 (2.2^c±0.15)

Table-2: The average score of p63, HIF2 α , and ETV5 expression in some treatments as shown by the IHC method in testicular tissue of rat.

Treatments	Sample (n)	Score \pm SD		
		Average score p63 expression	Average score HIF2 α expression	Average score ETV5 expression
Positive (fertile) control group	10	9.6 ^a \pm 0.44	0.6 ^c \pm 0.34	10.5 ^a \pm 0.25
Negative (infertile) control group	10	0.41 ^d \pm 0.22	0.3 ^c \pm 0.33	0.3 ^d \pm 0.23
Infertile male transplanted with stem cells from normoxia culture (21% O ₂ concentration) (T1 group)	10	2.5 ^c \pm 0.25	3.9 ^b \pm 0.44	2.2 ^c \pm 0.15
Infertile male transplanted with stem cells from hypoxia culture (1% O ₂ concentration) (T2 group)	10	6.0 ^b \pm 0.50	7.0 ^a \pm 0.75	7.2 ^b \pm 0.34

^{a-d}Different superscripts in the same column were significantly different ($p < 0.005$). IHC=Immunohistochemical, HIF=Hypoxia-inducible factor




Figure-1: Average score of p63 expression (brown): (a) Positive (fertile) control group=9.6^a \pm 0.44; (b) T2 group=6.0^b \pm 0.50; (c) T1 group=2.5^c \pm 0.25; (d) negative (infertile) control group=0.41^d \pm 0.22. (a-d) 200 \times with the IHC method. IHC=Immunohistochemical.




Figure-3: Average score of ETV5 expression (brown): (a) Positive (fertile) control group=10.5^a \pm 0.25; (b) T2 group=7.2^b \pm 0.34; (c) T1 group=2.2^c \pm 0.15; (d) negative (infertile) control group=0.3^d \pm 0.23. (a-d) 400 \times with the IHC method. IHC=Immunohistochemical.




Figure-2: Average score of HIF2 α expression (brown): (a) Positive (fertile) control group=0.6^c \pm 0.34; (b) T2 group=7.0^a \pm 0.75; (c) T1 group=3.9^b \pm 0.44; (d) negative (infertile) control group=0.3^c \pm 0.33. (a-d) 400 \times with the IHC method. IHC=Immunohistochemical, HIF=Hypoxia-inducible factor.

and negative (infertile) control groups, in which ETV5 was not expressed (0.3^d \pm 0.23) (Figure-3 and Table-2).

Regeneration of testicular tissue

Microscopic examinations of five different fields of view revealed that the T2 group experienced repaired testicular tissue. The improvements were identified based on the regeneration of Sertoli cells, Leydig cells, spermatogonia, spermatocytes, primary-secondary and spermatid cells, and seminiferous tubules. An overview of these improvements could be compared with the positive (fertile) control, which did not experience testicular degeneration. The T2 group remained under the normal condition (Figures-4a and d, Table-3). The T1 group did not exhibit testicular tissue repair, indicating that intact seminiferous tubules were not observed, and spermatogonia, Sertoli cells, and Leydig cells were degenerated. This amount of damage was comparable to that of the negative (infertile) control (Figures-4b and c, Table-3).

The number of different types of cells was counted based on the characteristics of each cell, as in the following:

Spermatogonium: It has a round shape and is located near the basement membrane, and the nucleus has an oval shape with fine chromatin and a thin nuclear membrane.

Table-3: The average number of Sertoli cells, Leydig cells, spermatogonium, spermatocyte primary-secondary, and spermatid cells in some treatments.

Treatments	Sample (n)	Average number of Sertoli cells \pm SD	Average number of Leydig cells \pm SD	Average number of spermatogonium cells \pm SD	Average number of spermatocyte primary-secondary cells \pm SD	Average number of spermatid cells \pm SD
Positive (fertile) control group	10	11.5 ^d \pm 0.45	29.56 ^d \pm 0.85	30.65 ^d \pm 0.70	74.25 ^d \pm 1.65	85.55 ^d \pm 1.50
Negative (infertile) control group	10	2.1 ^a \pm 0.47	5.15 ^a \pm 0.75	5.35 ^a \pm 0.75	30.15 ^a \pm 1.70	12.25 ^a \pm 1.75
Infertile male transplanted with stem cells from normoxia culture (21% O ₂ concentration) (T1 group)	10	4.2 ^b \pm 0.30	10.35 ^b \pm 0.65	12.95 ^b \pm 0.60	43.45 ^b \pm 1.45	30.50 ^b \pm 1.30
Infertile male transplanted with stem cells from hypoxia culture (1% O ₂ concentration) (T2 group)	10	7.8 ^c \pm 0.53	17.85 ^c \pm 0.75	20.75 ^c \pm 0.80	69.85 ^{cd} \pm 1.50	55.25 ^c \pm 1.25

^{a-d}Different superscripts in the same column were significantly different ($p < 0.005$)




Figure-4: Rat testicular tissue with hematoxylin and eosin (H&E) staining: (A) Positive (fertile) control group: (a) spermatogonia cells, (b) Sertoli cells, (c) Leydig cells, and d. intact tubules seminiferous; (B) T2 group: (a) Spermatogonia cells, (b) Sertoli cells, (c) Leydig cells, and (d) intact tubules seminiferous; (C) T1 group: (a) Spermatogonia cells, (b) Sertoli cells, (c) Leydig cells, and (d) damaged tubules seminiferous; (D) negative (infertile) control group: Loss of (a) spermatogonia cells, (b) Sertoli cells, (c) Leydig cells, and (d) damaged tubules seminiferous. (A-D) 200 \times . H&E: Hematoxylin and eosin.

Primary spermatocyte: It has the largest size among gamete cells, with heterochromatin in the nucleus, and is located between the basal membrane and the tubular lumen.

Secondary spermatocyte: It is rarely observed in the seminiferous tubules because it quickly divides into spermatids.

Spermatid: It has a round shape and is smaller than spermatocytes, and the nucleus is round, pale, and bright.

Sertoli cell: It has a slim, irregular shape, and the base attaches to the basement membrane of the seminiferous tubules, having one nucleus located at the center.

Leydig/interstitial cell: It is located in the loose connective tissue between the tubules; this is a large

cell, polygonal in shape, and the nucleus is clearly visible and also polygonal in shape.

Improvement of the fertility rate of the sperm

The T2 group exhibited a significant improvement in the fertility rates compared with the T1 group, although the rate did not reach that of the positive (fertile) control group (Table-4).

Discussion

In this study, we determined whether MSCs exposed to hypoxic conditions could repair testicular function more effectively than MSC cultured under normoxic conditions. Hypoxia, in this study, was adjusted to the normal and physiological conditions in which the stem cells were *in situ* (*in vivo*). Therefore, an *ex situ* (*in vitro*) study was conducted by inducing hypoxic conditions so that the conditions of the stem cells cultured are the same as the *in situ* physiological conditions (*in vivo*). In *in vivo* conditions, the MSCs are in the form of quiescent.

The induction of quiescent MSCs through p63 quiescent expression increased the life expectancy of stem cells by maintaining their viability and the adaptive condition for stem cell transplantation. The role of hypoxia in maintaining quiescence stem cells begins with the induction of HIF2 α . Subsequently, HIF2 α activated the gene pluripotency after it had been preceded by an initial adaptation time by HIF1 α . The pluripotency of these stem cells may prolong the lifetime of quiescent cells so that the function of stem cells (stemness) is maintained. Furthermore, after transplantation, it mobilized the endogenous stem cells toward the defect area (testicular tissue). The process of mobilization can occur through several ways: (a) Induction of proteolysis (protein degradation) from the microenvironment of bone marrow, such as induction of pharmacological agents (granulocyte-colony stimulating factor or cyclophosphamide) or induction of quiescent stem cells that were transplanted (p63 marker); (b) blockade of CXCR4 or VLA-4 by specific blocking molecules, such as

Table-4: Fertility rate of sperm of male rat.

S. No.	Treatment	Number of mature oocyte	Good quality embryos	Fertility rate (%)
1.	Positive (fertile) control group	50	43	86
2.	Negative (infertile) control group	50	2	4
3.	Infertile male transplanted with stem cells from normoxia culture (21% O ₂ concentration) (T1 group)	50	9	18
4.	Infertile male transplanted with stem cells from hypoxia culture (1% O ₂ concentration) (T2 group)	50	31	62

AMD3100 or BIO4860; (c) effect from neural mediators, such as dopamine and beta-2 adrenergic receptors; (d) element modulation from the coagulation of cascade apoptosis; (e) inflammatory reaction causing injury signals or immune response induction, such as NF- κ B, cytokines, or β -catenin, through Wnt from the tissue damage; and (f) homing signals, such as SDF1, CXCL12, VEGF, HGF, PDGF, and integrin, or transcriptional factors, such as HIF2 α and ETV5, which appeared and recruited endogenous stem cells [27]. The mobilization of endogenous stem cells will further move the stem cells to the testicular tissue, resulting in spermatogenesis processes and rescue of testicular failure and infertility repair.

This study demonstrated that the MSCs from the hypoxic precondition culture were effective for therapy in male rats with testicular failure and infertility based on the increased of p63 expression as a quiescent cell marker as a crucial for progenitor of stem cell function and ETV5 expression as a transcriptional factor for regeneration of testicular tissue and improvement of the fertility of sperm.

The regenerative efforts of stem cells were differentiated by the decreased expression of p63 compared with that in the positive (fertile) control group. This study demonstrated that the score of p63 in the T2 group was lower than that in the positive (fertile) control group but was still better than those in the T1 and negative (infertile) control groups, which is slightly expressed (Figure-1 and Table-1).

The p63 gene can maintain the viability of stem cells and regenerate stem cells from various tissue cells, which are known as ringmasters. In the previous study [4], the absence of p63 was showed a decreased of proliferation ability of cells, indicating that p63 is a key function in increasing the division of stem cells because the p63 gene directly promotes and control the stem cell environment and maintain undifferentiation.

Hypoxic precondition led to HIF1 α release; therefore, so that not bound by the van Hippel-Lindau factor as a factor inhibited for HIF1 α action furthermore[38]. Furthermore, HIF1 α would be bound to HIF1 β so that complex HIF1 was formed [9]. The HIF1 α +HIF1 β bond occurred in specific DNA sequences known as the hypoxia response element (HRE) 5'-TACGC-3'. Complex HIF1 α and HIF1 β bonds on HRE occurred at the start of exposure to hypoxia [39], thus causing cell cycle arrest and gene expression [40]. This inhibited the p21 expression,

resulting in cell cycle inactivation and resistance to senescence and exhausting cells [7]. This is thought to slow down the proliferation of cultured stem cells; thus, quiescent cells can still be maintained [41].

Long-term maintenance of quiescent cells was also thought to be influenced by cultivation time-dependent hypoxic preconditions. After 48 h under low O₂ tension, the role of HIF1 α would be replaced by that of HIF2 α with different target genes [39]. The target genes in *in vitro* culture were expected to induce the expression of pluripotency genes [42], such as OCT4, SOX2, NANOG [9,43], and REX-1 [23]. A hypoxic precondition is an effort to change the ability of multipotent stem cells to become pluripotent.

The mean identification of the HIF2 α score in the T2 group was the highest among those of the other groups. *In vitro* cultures, low oxygen tension (hypoxia), and cultivation time-dependent administration of oxygen induce expressions of pluripotency genes [37], such as OCT4, SOX2 [1,44], REX-1 [44], and NANOG [45]. Pluripotency genes are activated by HIF2 α [46] after preceding initial adaptation time by HIF1 α [9]. The pluripotency of these stem cells can retain quiescent cells; therefore, the function of stem cells is maintained. The quiescent cells with p63 as a marker, which is regulated by HIF2 α from hypoxic precondition culture with 1% O₂ concentration for 4 days, are crucial for conducive niche *in vivo*, so that after transplantation, the stem cells could be transdifferentiated through spermatogenesis in the seminiferous tubules of the testis.

Furthermore, stem cells from the hypoxic precondition culture were found to be effective based on ETV5 formation in the testicular tissue, with an average score of 2.95^b±0.50. The ETV5 as a marker of SSC function that can increase and improve the testicular environment and support endogenous stem cells, so that stem cells can be mobilized to the testicular tissue that has failed, resulting in improvement and rescue for fertility.

In this study, IHC methods were employed to identify ETV5. The score of the ETV5 expression in the T2 group was approximately 3. Although this score was below that of the positive (fertile) control group, it was still well above those of the T1 and negative (infertile) control groups, which was slightly expressed (Figure-3 and Table-1). Previous research demonstrated that bone marrow-derived MSCs are adult stem cells that quickly grow and differentiate

into cells that are needed in response to the presence of defects [40].

Regeneration of the testicular tissue was identified as an intact seminiferous tubule tissue; formation of Sertoli cells, Leydig cells, spermatogonia, spermatocytes, and primary and secondary and spermatid cells. The viability of stem cells that differentiate into cells is necessary. In infertile conditions, the degenerative testicular tissue can be regenerated if the stem cells are viable. If they are not viable, then the testicular tissue will remain degenerated.

The survival of stem cells in the animal model of degenerative tissue, such as testicular failure, is beyond the scope of the therapeutic effect of MSC treatment. In addition, poor survival following cell transplantation is a crucial factor [5]. This study demonstrated that stem cells from the hypoxic precondition culture survived based on the effectiveness of therapy in male rats with testicular failure and infertility through the regeneration of their testes, which can be observed using IHC methods, H&E staining, and a light microscope [31]. Testicular tissue repair was confirmed by the regeneration of the seminiferous tubules, based on observing of seminiferous tubules become intact and compact again [27].

In this study, light microscopy examination revealed that testicular tissue repair occurred in the T2 group. The overview of the testicular tissue repair can be compared with that in the positive (fertile) control group, which did not experience testicular degeneration and remained in the normal condition (Figure-4 and Table-2), whereas the T1 group did not exhibit improvement in their testicular tissue. The tissue damage was comparable with that in the negative (infertile) control group with testicular degeneration (Figure-4 and Table-2).

Conclusion

Transplantation of MSCs cultured under hypoxic conditions is an effective treatment for testicular failure in a rat infertility model.

Authors' Contributions

ES: Supervised the whole study. ES and HP: Designed the study, analyzed the data, wrote and edited the article. Both authors read and approved the final manuscript.

Acknowledgments

The authors greatly acknowledge The Indonesian Directorate General Higher Education (DIKTI) for funding this study (Grant number : 4/AMD/E1/KP.PTNBH/2020). The authors also acknowledge chairman and staff of LIPJHKI, who assisted in the grammar check and Prof. Dr. R. Heru Prasetyo, dr., MS., SpParK, for his assistance in the study.

Competing Interests

The authors declare that they have no competing interests.

Publisher's Note

Veterinary World remains neutral with regard to jurisdictional claims in published institutional affiliation.

References

- Safitri, E., Utama, S., Bumi, C. and Mulyani, S.W.M. (2014) Hypoxic preconditioning for viable and self-renewing mesenchymal stem cells (MSCs) as the regeneration of spermatogenesis process. *Adv. Nat. Appl. Sci.*, 8(8): 42-46.
- Safitri, E. and Hariadi, M. (2019) Comparison of biotechnological culture of hypoxia-conditioned rat mesenchymal stem cells with conventional *in vitro* culture of normoxia-conditioned rat mesenchymal stem cells for testicular failure therapy with low libido in rats. *Vet. World*, 12(6): 916-924.
- Cakici, C., Buyrukcu, B. and Duruksu, G. (2013) Recovery of fertility in azoospermia rats after injection of adipose-tissue-derived mesenchymal stem cells: Tha sperm generation. *Bio. Med. Res. Inter.*, 10: 1-18.
- Safitri, E., Srianto, P., Widiyatno, T.V., Sandhika, W. and Prasetyo, R.H. (2018) Viability of rabbit adipocyte stem cells cultured under different oxygen concentrations *in vitro*. *Philippine J. Vet. Med.*, 55(SI): 1-8.
- Li, L., Chen, X., Wang, W.E. and Zeng, C. (2016) Review article: How to improve the survival of transplanted mesenchymal stem cell in ischemic heart? *Stem. Cells Inter.*, 2016(SI): 1-14.
- Scaradavou, A., Smith, K.M., Hawke, R., Schaible, A., Abboud, M., Keman, N.A., Young, J.W. and Barker, J.N. (2015) Cord blood units with low CD34+ cell viability have a low probability of engraftment after double unit transplantation. *Biol. Blood Marrow Transplant.*, 16(4): 500-508.
- Tsai, C.C., Chen, Y.J., Yew, T.L., Chan, L.L., Wang J.Y., Chiu, C.H. and Hung, S.C. (2011) Hypoxia inhibits senescence and maintains mesenchymal stem cell properties through down-regulation of E2A-p21 by HIF-TWIST. *Blood J. Hematol.*, 117(2): 459-469.
- Mulyani, S.W.M., Setiawati, E.M., Safitri, E. and Astuti, E.R. (2014) The role of heat shock protein 27(HSP 27) as inhibitor apoptosis in hypoxic conditions of bone marrow stem cell culture. *Dent. J. Maj. Ked. Gigi.*, 47(1): 41-44.
- Szablowska-Gadomska, I., Zayat, V. and Buzanska, L. (2017) Influence of low oxygen tension on expression of pluripotency genes in stem cells. *Acta Neuroiol. Exp.*, 71(1): 86-93.
- Cai, B., Li, X., Wang, L.Y., Liu, Y., Yang, F., Chen, H., Yin, K., Tan, X., Tan, X., Zhu, J., Pan, Z., Wang, B. and Lu, Y. (2013) Apoptosis of bone marrow mesenchymal stem cells caused by homocysteine via activating JNK signal. *PLoS One.*, 8(5): 1-9.
- Chen, T.L., Zhu, G.L., Wang, J.A., Wang, Y., He, X.L. and Jiang, J. (2014) Apoptosis of bone marrow mesenchymal stem cells caused by hypoxia/reoxygenation via multiple pathways. *Int. J. Clin Exp. Med.*, 7(12): 4686-4697.
- Teixeira, F.G., Panchalingam, K.M., Anjo, S.I., Manadas, B., Pereira, R., Sousa, N., Salgado, A.J. and Behie, L.A. (2015) Do hypoxia/normoxia culturing conditions change the neuroregulatory profile of Wharton Jelly mesenchymal stem cell secretome? *Stem Cell Res. Ther.*, 6(1): 133-147.
- Qin, H.H., Filippi, C., Sun, S., Lehec, S., Dhawan, A. and Hughes, R.D. (2015) Hypoxic preconditioning potentiates the trophic effects of mesenchymal stem cells on co-cultured human primary hepatocytes. *Stem Cell Res. Ther.*, 2015(1): 1-12.
- Zhou, H., Li, D., Shi, C. and Xin, C. (2015) Effects of exendin-4 on bone marrow mesenchymal stem cell proliferation, migration and apoptosis *in vitro*. *Sci. Rep.*, 5: 1-12.
- Galleu, A., Riffo-Vasquez, Y., Trento, C., Lomas, C., Dolcetti, L., Cheung, T.S., Bono, N.V., Barbiene, R.,

- Halai, K., Ward, S., Weng, L., Chikraverty, R., Lombardy, B., Watt, F.M., Orchard, K., Marks, D.J., Apperly, J., Bornhouser, M., Welezak, H., Bennet, C. and Dazzi, F. (2017) Apoptosis in mesenchymal stromal cells induces *in vivo* recipient-mediated immunomodulation. *Sci. Transl. Med.*, 9(416): 7828.
16. Suda, T., Takubo, K. and Semenza, G.L. (2011) Cell stem cell review. Metabolic regulation of hematopoietic stem cells in hypoxic niche. *Cell Stem Cell.*, 9(4): 296-310.
17. Nakamura-Ishizu, A., Takizawa, H. and Suda, T. (2014) Reviews: The analysis, roles and regulation of quiescence in hematopoietic stem cells. *Development*, 141(24): 4656-4666.
18. Takubo, K. (2011) The hypoxia regulatory system in hematopoietic stem cells. *Adv. Hemato. Stem Cell Res.*, 20(12): 133-146.
19. Yue, F., Bi1, P., Wang, C., Shan, T., Nie, W., Ratliff, T. and Gavin, T.P. (2017) Pten is necessary for the quiescence and maintenance of adult muscle stem cells. *Nat Comm.*, 10: 1-13.
20. Cheung, T.H. and Rando, T.A. (2013) Molecular regulation of stem cell quiescence. *Nat. Rev. Mol. Cell Biol.*, 14(6): 1-26.
21. Rumman, M., Dhawan, J. and Kassem, M. (2015) Concise review: Quiescence in adult stem cells: Biological significance and relevance to tissue regeneration. *Stem. Cells*, 33(10): 2903-2912.
22. Daignan-Fornier, B. and Sagot, I. (2011) Proliferation/quiescence: When to start? Where to stop? What to stock? *Cell Div.*, 6(20): 1-5.
23. Kolf, C.M., Cho, E. and Tuan, R.S. (2007) Review mesenchymal stromal. Biology of adult mesenchymal stem cells: Regulation of niche, self-renewal, and differentiation. *Arthritis Res. Ther.*, 9(1): 204.
24. Kang, X.Q., Zang, W.J., Song, T.S., Xu, X.L., Yu, X.J., Li, D.L., Meng, K.W., Wu, S.L. and Zhao, Z.Y. (2005) Rat bone marrow mesenchymal stem cells differentiate into hepatocytes *in vitro*. *World J. Gastroenterol.*, 11(22): 3479-3484.
25. Gala, K., Burdzińska, A., Idziak, M., Makula, J. and Paczek, L. (2011) Characterization of bone-marrow-derived rat mesenchymal stem cells depending on donor age. *Cell Biol. Int.*, 35(10): 1055-1062.
26. Eleotério, R.B., Sepúlveda, R.V., Reis, E.C.C., Falente, F.L. and Borges, A.P.B. (2016) Isolation, expansion and differentiation of mesenchymal stromal cells from rabbits' bone marrow. *Pesq. Vet. Bras.*, 36(5): 423-430.
27. Safitri, E., Utama, S., Widiyatno, T.V., Sandhika, W. and Prasetyo, R.H. (2016) Auto-regeneration of mice testicle seminiferous tubules due to malnutrition based on stem cells mobilization using honey. *Asian Pac. J. Reprod.*, 5(1): 31-35.
28. Hackbert, L. and Heiman, J.R. (2002) Acute dehydroepiandrosterone (DHEA) effect on sexual arousal in postmenopausal women. *J. Women Health Gend. Based Med.*, 11(2): 155-162.
29. Hafez, E.S.E. and Hafez, B. (2010) Reproduction in Farm Animals. 7th ed. Wiley-Blackwell, Philadelphia, PA, USA. p157-198.
30. França, L.R., Ogawa, T., Avarbock, M.R., Brinstel, R.L. and Russell, L.D. (1998) Germ cell genotype controls cell cycle during spermatogenesis in the rat. *Biol. Reprod.*, 59(1): 1371-1377.
31. Razi, M., Najafi, G., Feyzi, S., Karimi, A., Shahmohammadloo, S. and Nejati, V. (2012) Histological and histochemical effects of glyphosate on testicular tissue and function. *Iran. J. Reprod. Med.*, 10(3): 181-192.
32. Nowak, M., Madej, J. and Dziegiel, P. (2007) Intensity of COX2 expression in cells of soft tissue fibrosarcomas in dogs as related to grade of tumour malignancy. *Bull Vet. Inst. Pulawy.*, 51(2): 275-279.
33. Widjiati, W., Rachmawati, A. and Sosiawati, S.M. (2012) Identification of epidermal growth factor (EGF) protein in 46 kDa from oocyte maturation *in vitro* of cow with. *J. Vet. Med.*, 6(1): 32-35.
34. Nabavi, N., Todehdehghan, F. and Shiravi, A. (2013) Effect of caffeine on motility and vitality of sperm and *in vitro* fertilization of outbreed mouse in T6 and M16 media. *Iran J. Reprod. Med.*, 11(9): 741-746.
35. Widjiati, W., Pusporini, S.E. and Arifin, M.Z. (2012) Comparative of fertility rate and barriers to development of mice embryos that cultured between M16 medium and human tubal fluid. *J Vet Med.*, 13(3): 227-234.
36. Jin, H., Shu, Y. and Dai, S. (2014) The value of second polar body detection 4 hours after insemination and early rescue ICSI in preventing complete fertilisation failure in patients with borderline semen. *Reprod. Fertil. Dev.*, 26(2): 346-350.
37. Youssef, A. and Han, V.K.M. (2016) Low oxygen tension modulates the insulin-like growth factor-1 or -2 signaling via both insulin-like growth factor-1 receptor and insulin receptor to maintain stem cell identity in placental mesenchymal stem cells. *Endocrinology*, 157(3): 1163-1174.
38. Takubo, K. and Suda, T. (2012) Roles of the hypoxia response system in hematopoietic and leukemic stem cells. *Int. J. Hematol.*, 95(47): 483.
39. Forristal, C.E., Wright, K.L., Hanley, N.A., Oreffo, R.O.C. and Houghton, F.D. (2010) Hypoxia inducible factor regulate pluripotency and proliferation in human embryonic stem cells cultured at reduced oxygen tensions. *Reproduction*, 139(1): 85-97.
40. Li, L. and Bhatia, R. (2011) Stem cell quiescence. *Clin. Cancer Res.*, 7(15): 57-69.
41. Arai, F. and Suda, T. (2008) Quiescent Stem Cells in the Niche. The Stem Cell Research Community, StemBook, London. p382.
42. Grskovic, M. and Santos, M.R. (2008) The Pluripotent Transcriptome. The Stem Cell Research Community, StemBook, London. p24.
43. Yamanaka, S. (2007) Strategies and new developments in the generation of patient-specific pluripotent stem cells. *Stem Cell Rev.*, 1(1): 39-49.
44. Lavagnoli, T., Gupta, P., Hormanseder, E., Botenbal, H.M., Dharmalenghan, G., Carroll, T., Gurdon, J.B., Fisher, A.G. and Markensclanger, M. (2014) Initiation and maintenance of pluripotency gene expression in the absence of cohesion. *Genes Dev.*, 29(1): 23-38.
45. Festuccia, N., Osorno, R., Wilson, V. and Chambers, I. (2013) The role of pluripotency gene regulatory network components in mediating transitions between pluripotent cell states. *Curr. Opin. Genet. Dev.*, 23(5): 504-511.
46. Dengler, V.L., Galbraitmh, M. and Espinosa, J.M. (2014) Transcriptional regulation by hypoxia inducible factors. *Crit. Rev. Biochem. Mol. Biol.*, 49(1): 1-15.
