

Treatment with Mammalian Ste-20-like Kinase 1,2 (MST1,2) Inhibitor XMU-MP-1 Improves Glucose Tolerance in Streptozotocin-Induced Diabetes Mice

by Zakiyatul Faizah

Submission date: 21-Nov-2022 05:30PM (UTC+0800)

Submission ID: 1960195496

File name: es_Glucose_Tolerance_in_Streptozotocin-Induced_Diabetes_Mice.pdf (2.53M)

Word count: 7763

Character count: 51082

Treatment with Mammalian Ste-20-like Kinase 1/2 (MST1/2) Inhibitor XMU-MP-1 Improves Glucose Tolerance in Streptozotocin-Induced Diabetes Mice

Zakiyatul Faizah ^{1,†}, Bella Amanda ^{1,†}, Faisal Yusuf Ashari ^{1,†}, Efta Triastuti ^{2,3}, Rebecca Oxtoby ², Anny Setijo Rahaju ⁴, M. Aminudin Aziz ¹, Maria Inge Lusida ^{5,6} and Delvac Oceandy ^{2,*}

¹ Department of Biomedical Sciences, Faculty of Medicine, Universitas Airlangga, Surabaya 60132, Indonesia; zakiyatul-f@fk.unair.ac.id (Z.F.); bella.amanda@fk.unair.ac.id (B.A.); faisalyusufashari@gmail.com (F.Y.A.); azizandro@gmail.com (M.A.A.)

² Division of Cardiovascular Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester M13 9PT, UK; efta.triastuti@postgrad.manchester.ac.uk (E.T.); b.oxtoby@hotmail.com (R.O.)

³ Department of Pharmacy, Faculty of Medicine, Universitas Brawijaya, Malang 65145, Indonesia

⁴ Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya 60132, Indonesia; anny_sr@fk.unair.ac.id

⁵ Institute of Tropical Disease, Universitas Airlangga, Surabaya 60286, Indonesia; ingelusida@itd.unair.ac.id

⁶ Department of Microbiology, Faculty of Medicine, Universitas Airlangga, Surabaya 60132, Indonesia

* Correspondence: delvac.oceandy@manchester.ac.uk; Tel.: +44-161-275-1772

† These authors contributed equally to this work.

Academic Editor: Luigi Milella

Received: 28 August 2020; Accepted: 22 September 2020; Published: 24 September 2020



Abstract: Diabetes mellitus (DM) is one of the major causes of death in the world. There are two types of DM—type 1 DM and type 2 DM. Type 1 DM can only be treated by insulin injection whereas type 2 DM is commonly treated using anti-hyperglycemic agents. Despite its effectiveness in controlling blood glucose level, this therapeutic approach is not able to reduce the decline in the number of functional pancreatic β cells. MST1 is a strong pro-apoptotic kinase that is expressed in pancreatic β cells. It induces β cell death and impairs insulin secretion. Recently, a potent and specific inhibitor for MST1, called XMU-MP-1, was identified and characterized. We hypothesized that treatment with XMU-MP-1 would produce beneficial effects by improving the survival and function of the pancreatic β cells. We used INS-1 cells and STZ-induced diabetic mice as in vitro and in vivo models to test the effect of XMU-MP-1 treatment. We found that XMU-MP-1 inhibited MST1/2 activity in INS-1 cells. Moreover, treatment with XMU-MP-1 produced a beneficial effect in improving glucose tolerance in the STZ-induced diabetic mouse model. Histological analysis indicated that XMU-MP-1 increased the number of pancreatic β cells and enhanced Langerhans islet area in the severe diabetic mice. Overall, this study showed that MST1 could become a promising therapeutic target for diabetes mellitus.

Keywords: diabetes mellitus; MST1/2; Hippo pathway; XMU-MP-1

1. Introduction

Diabetes mellitus (DM) is one of the major causes of death in the world [1]. A total of 463 million (9.3%) people worldwide were estimated to suffer from diabetes in 2019. This number is estimated to increase to 578 million (10.2%) in 2030 and 700 million (10.9%) in 2045 [2]. DM might induce multi-organ complications, such as retinopathy, neuropathy, cerebrovascular diseases, nephropathy, and cardiovascular diseases [3]. At present, the most commonly used treatment for DM is anti-hyperglycemic agents, that can reduce the detrimental effects of high blood glucose level [4]. However, this therapeutic strategy cannot

56
stop the decline in the number of functional pancreatic β cells, which are important in the production and secretion of insulin. Discovery of new therapeutic strategies that can prevent the damage of pancreatic β cells due to DM will be useful in tackling this disease in the future.

There are two types of DM—type 1 DM (T1DM) and type 2 DM (T2DM). T1DM is caused by pancreatic β cell destruction mainly due to the autoimmune process. Its onset often occurs in children and young adults. Since the cause of T1DM is pancreatic β cell damage, the main treatment strategy for this condition is injection with insulin [5]. On the other hand, T2DM is caused by insulin resistance in the peripheral target organs as well as defective insulin secretion [6]. T2DM is normally associated with metabolic syndrome (obesity, hypertension, dyslipidemia). These conditions can change the molecular properties of the insulin receptor in the target tissues, such as adipose and skeletal muscle tissues. In contrast to T1DM, the main therapeutic strategy for T2DM is using anti-hyperglycemic agents that work by inducing insulin sensitivity (glitazones), reducing glucose production in liver (metformin) or enhancing insulin release by pancreatic β cells (sulfonylureas) [7].

Pancreatic β cell damage due to apoptosis plays an important role in both types of DM. In T1DM, β cell destruction occurs following an autoimmune reaction, which triggers activation of inflammatory cells and secretion of cytokines [8]. This will eventually activate the apoptotic pathways in the pancreatic β cells [9]. In T2DM, metabolic stress might lead to the activation of proinflammatory cytokines such as tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6) [10]. These cytokines might induce insulin resistance in the peripheral target organs, as well as contribute to the apoptotic cell death of pancreatic β cells [10].

59
One of the major signaling pathways that is heavily involved in the regulation of cell apoptosis, survival, and regeneration is the Hippo pathway [11]. The core components of the Hippo pathway in mammals include kinases—mammalian sterile 20-like kinase 1/2 (MST1/2) and large tumor suppressor 1/2 (LATS1/2), adaptor molecules (salvador (SAV1) and MOB kinase activator 1 (MOB1)), the transcriptional co-activators Yes-associated protein (YAP), and WW domain containing transcription regulator 1 (WWTR1/TAZ) [8,11]. Activation of the Hippo pathway by phosphorylation of the kinases results in YAP inactivation and hence reduces the expression of its target genes, which include anti-apoptotic and pro-survival genes. On the other hand, inhibition of the Hippo pathway result in YAP activation and eventually in induction of target genes expression [8,11].

Beside regulating the cell function via the canonical pathway, several members of the Hippo pathway might regulate apoptosis through alternative pathways. For example, MST1 was shown to be a powerful pro-apoptotic molecule through the modulation of B-cell lymphoma-extra-large (Bcl-xL) phosphorylation [12], regulation of the Forkhead box O1-NOXA (FoxO1-NOXA) pathway [13], and phosphorylation of histone H2AX [14]. MST1 is expressed in pancreatic β cells, making this kinase important in the pathophysiology of DM [15]. MST1 induces β cell death and impairs insulin secretion by initiating proteasomal degradation of pancreatic and duodenal homeobox 1 (PDX1), and β cell transcription factor [16]. In keeping with this finding, genetic inhibition of MST1 protected against β cell apoptosis and improved the diabetic phenotype in streptozotocin (STZ)-induced diabetes model in mice [17]. This finding prompted a notion that MST1 could be a novel therapeutic target for DM. Furthermore, a study using a cancer drug, neratinib, which was found to inhibit MST1 activity, indicated that pharmacological inhibition of MST1 was protective against the development of STZ-induced diabetes mellitus [17].

Recently, a novel molecule XMU-MP-1 was identified and characterized as a potent and specific inhibitor for MST1/2 [18]. It was demonstrated that treatment with XMU MP-1 in mice enhanced the proliferation of hepatocytes and small intestinal epithelial cells [18]. XMU MP-1 could also improve heart function and remodeling in a model of chronic cardiac pressure overload in mice, by decreasing apoptosis, fibrosis, and pathological hypertrophy [19]. Given the crucial role of MST1 in the pancreatic β cells and in the pathogenesis of DM, we hypothesized that treatment with XMU-MP-1 would produce beneficial effects by improving the survival and function of the pancreatic β cells. In this study, we tested this idea by performing in vitro and in vivo studies using the insulinoma cell line

(INS-1) and an STZ-induced diabetes model in mice. We used the rat insulinoma cell line INS-1 832/13 in this study. We refer to this cell line as INS-1 cells, thereafter.

2. Results

2.1. The Effects of XMU-MP-1 Treatment on INS-1 Cells

To examine whether treatment with XMU-MP-1 (Figure 1A) modulates the Hippo signaling pathway in pancreatic β cells, we treated INS-1 cells with increasing doses of XMU-MP-1 (1–5 μ M). As XMU-MP-1 targets the MST1/2 activity, we first looked at the phosphorylation of LATS1 and MOB1, which are known as downstream substrates of MST1/2. Western blot analysis showed reduction in LATS1 and MOB1 phosphorylation, as shown in Figure 1B. Further downstream, we explored if inhibition of MST1/2 in INS-1 cells would activate YAP, a major target effector of the Hippo Pathway. Using an established adenoviral-driven luciferase reporter system [19], we detected a significant and dose-dependent increase in YAP activity, following XMU-MP-1 treatment (Figure 1C). Together, these findings suggest that XMU-MP-1 treatment could inhibit MST1/2 activity in INS-1 cells, which subsequently enhanced the YAP activity.




Figure 1. Effects of XMU-MP-1 treatment on INS-1 cells. (A) Molecular structure of XMU-MP-1. (B) XMU-MP-1 treatment reduced the phosphorylation of MOB1 and LATS1 in INS-1 cells. (C) Using an adenoviral-based YAP-luciferase reporter system, we detected a significant increase of YAP activity in INS-1 cells treated with XMU-MP-1, at doses of 3 μ M and 5 μ M for 24 h ($n = 6$ in each group). (D) MTT assay indicated that there was a reduction in cell survival following treatment with 3 μ M and 5 μ M of XMU-MP-1 for 24 h ($n = 10$ in each group). ** $p < 0.01$, *** $p < 0.001$.

In order to determine if the XMU-MP-1 treatment produced toxic effects in INS-1 cells, we performed an MTT assay to assess the level of INS-1 cell viability, following treatment with XMU-MP-1. Data shown in 1D indicated that at a lower dose (1 μ M) of XMU-MP-1 did not induce significant cell death, however, cell viability was decreased after treatment with higher doses of XMU-MP-1 (3–5 μ M). It is important to note that despite statistically significant reduction in INS-1 cell

viability, there were still more than ~70% INS-1 cells that survived at 24 h, after treatment with 5 μ M XMU-MP-1 (Figure 1C).

2.2. XMU-MP-1 Treatment Reduces STZ-Mediated Cell Death of INS-1 Cells

To investigate the effects of XMU-MP-1 treatment on STZ-induced toxicity in INS-1 cells, we assessed cell viability using an MTT assay. First, we performed experiments to determine the toxic dose of STZ in INS-1 cells. As expected, we found a dose dependent reduction in INS-1 cell viability following STZ treatment (1–2 mM for 24 h), with a statistical significance only observed, following treatment with 2 mM STZ (Figure 2A). We next analyzed the effects of XMU-MP-1 treatment, at doses of 1–3 μ M, in addition to STZ administration. We observed more cell survival following treatment with 1 μ M XMU-MP-1 on INS-1 treated with 1 mM STZ (Figure 2B) and in cells treated with 2 mM STZ (Figure 2C), although the differences did not reach statistical significance.




Figure 2. Effects of XMU-MP-1 treatment on INS-1 cell survival following streptozotocin treatment. (A) Treatment with STZ at 1–2 mM for 24 h resulted in a reduction in cell survival ($n = 22–25$ in each group, $*** p < 0.001$). (B) There was slight but non-significant increase in cell survival following XMU-MP-1 treatment in INS-1 cells, upon addition of 1 mM STZ ($n = 18–32$ in each group) or (C) 2 mM STZ ($n = 11–21$ in each group).

2.3. Effects of XMU-MP-1 Treatment in STZ-Induced Diabetic Mice

Next, we conducted *in vivo* experiments using the STZ-induced diabetic mouse model, to examine the effects of XMU-MP-1 treatment. Figure 3A describes the design of the *in vivo* experiments. Wild-type Balb/c mice were injected with STZ at a dose of 50 mg/kg body weight (BW) per day, for 5 consecutive days. We analyzed the random blood glucose level before the STZ induction (day 0), and at day 7, 14, 21, and 28, following the first STZ injection. We treated diabetic mice with either XMU-MP-1 at 1 mg/kg BW/day or an equivalent volume of DMSO, for consecutive 21 days, starting from day 15 after the first STZ injection. At the end of the experiments (day 35), we measured the fasting blood glucose and performed glucose tolerance test (GTT).

We found that the blood glucose level was significantly elevated at two weeks, following the first STZ injection (Figure 3B). We classified these diabetic mice into two groups—mild diabetes group (random blood glucose 200–300 mg/dL) and severe diabetes group (random blood glucose >300 mg/dL). All diabetic mice in each group were randomly assigned to the experimental group (received XMU-MP-1 treatment) or the control diabetic group (injected with DMSO).

Random blood glucose was assessed every 7 days after the first STZ injection. We did not observe any difference in random blood glucose levels between the XMU-treated mice and the DMSO-treated mice, at any time points before and after the XMU/DMSO treatment (Figure 3C). Similarly, when we evaluated random blood glucose levels in the mild and severe diabetes groups, we did not find any difference between the XMU-treated and the DMSO-treated mice (Figure 3D,E).

Body weight was also monitored during the course of the experiments. Similar to the random blood glucose data, we did not observe any difference in body weight between the XMU-treated diabetic mice, DMSO-treated diabetic mice, and the normal controls (Figure 3F–H).




Figure 3. The effect of XMU-MP-1 treatment in STZ-induced diabetes model in mice. (A) Schematic diagram explaining the design of the experiments. (B) Analysis of random blood glucose (RBG) at day 14 after the first STZ injection (before XMU-MP-1 treatment); control, $n = 20$; STZ, $n = 40$. **** $p < 0.0001$. (C) RBG levels in all mice during the course of experiment (control, $n = 10$; STZ + DMSO, $n = 20$; STZ + XMU-MP-1, $n = 20$). (D) RBG levels in mice with mild diabetes (RBG 200–300 mg/dL at day 14) (control, $n = 10$; STZ + DMSO, $n = 10$; STZ + XMU-MP-1, $n = 10$). (E) RBG levels in mice with severe diabetes (RBG > 300 mg/dL at day 14) (control, $n = 10$; STZ + DMSO, $n = 10$; STZ + XMU-MP-1, $n = 10$). Body weight in (F) all mice, (G) mild diabetes mice, and (H) severe diabetes mice, during the course of the experiments.

At the end of the experiments (day 35), we analyzed the fasting blood glucose and performed a glucose tolerance test. We found that the fasting blood glucose level was decreased in the XMU-treated group, compared to the DMSO-treated mice (Figure 4A). Furthermore, the glucose tolerance test (GTT) revealed that XMU-treated diabetic mice displayed a better response to bolus glucose injection (1g/kg body weight) compared to DMSO-treated diabetic mice (Figure 4B). In XMU-treated mice, the blood glucose started to decrease at 15 min after peritoneal glucose injection, whereas in the DMSO-treated diabetic group, the blood glucose stabilized only after 120 min after glucose injection.

These data suggest that XMU-MP-1 might improve the response against glucose load in STZ-induced diabetic mice.




Figure 4. Analysis of blood glucose phenotype at the end of the experiments (day 35). **(A)** Analysis of fasting blood glucose (FBG) in all diabetic mice showed that mice treated with XMU-MP-1 exhibited lower FBG level (STZ + DMSO, $n = 10$; STZ + XMU-MP-1, $n = 10$). $* p < 0.05$. **(B)** Intraperitoneal glucose tolerance test (GTT) was performed by injecting mice with 1 g/kg BW glucose, followed by analysis of blood glucose at several time points as indicated in the graph. In all diabetic mice combined, we found that XMU-MP-1 treatment improved the blood glucose levels. (STZ + DMSO, $n = 11$; STZ + XMU, $n = 11$; $* p < 0.05$, $*** p < 0.001$). **(C)** Results of GTT in mild diabetes mice (RBG = 200–300 mg/dL at the start of treatment) (STZ + DMSO, $n = 5$; STZ + XMU, $n = 5$). **(D)** Results of GTT analysis in severe diabetes mice (RBG > 300 mg/dL at the start of treatment) (STZ + DMSO, $n = 6$; STZ + XMU, $n = 6$). $* p < 0.05$, $** p < 0.01$, $*** p < 0.001$.

Next, we assessed the GTT results separately in the mild and severe diabetes groups. Evaluation of GTT results in these groups showed that the beneficial effects of XMU-MP-1 were observed only in the severe diabetic group (Figure 4C,D).

2.4. Protective Effect of XMU-MP-1 Treatment in Pancreatic Tissues

To study the effects of XMU-MP-1 treatment on the histological structure of the pancreas, we examined hematoxylin-and-eosin-stained pancreatic tissue sections (Figure 5A). We focused our analysis on the structure of Langerhans islets by calculating the average cell number within one islet, and the average surface area of the Langerhans islets in the whole pancreatic sections. In the control mice (non-diabetic/no STZ), we found a trend of increased cell number and Langerhans islet area following XMU-MP-1 treatment, although the difference did not reach statistical significance (Figure 5B,C). We then analyzed the effects of XMU-MP-1 treatment in all diabetic mice (combined group

of mild and severe diabetes). We did not find any significant effects of the XMU-MP-1 treatment in terms of the Langerhans islet surface area and the average cell number within the islet (Figure 5B,C).




Figure 5. Histological analysis of pancreatic tissue sections. (A) Representative images of hematoxylin-eosin-stained pancreatic tissue sections showing Langerhans islets in all groups of experimental animals (scale bar = 100 μm). (B) The average number of cells in the Langerhans islets was examined in all groups. No significant difference in cell number was observed. (C) Analysis of the Langerhans islets surface area in all groups of mice. No significant difference in cell number was observed. (Control (non-diabetic), $n = 6$; control + XMU, $n = 6$; diabetic, $n = 19$; diabetic + XMU, $n = 19$).

However, further analysis in each group of diabetic mice showed beneficial effects of XMU-MP-1 treatment, when given to the severe diabetic mice. Whilst there was no significant difference regarding the cell number and Langerhans islets surface area in the mild diabetic group (Figure 6A,B), we found a significantly enhanced cell number and islets area following the XMU-MP-1 treatment in the severe diabetic group (Figure 6C,D). These findings were in line with the GTT analysis and supported the idea that XMU-MP-1 was effective to improve the phenotype in the severe diabetic group but not in the mild diabetic mice.




Figure 6. Analysis of Langerhans islets average cell number and surface area in specific diabetes groups. (A) Average cell number in the Langerhans islets and (B) the Langerhans islets surface area in mild diabetic mice. No significant difference was observed in this group of diabetic mice between the DMSO and XMU-MP-1 treatment (mild diabetes + DMSO, $n = 10$; mild diabetes + XMU MP-1, $n = 11$). (C) However, in the severe diabetic group there was a trend of a higher cell number and (D) a significant increase in the Langerhans islets surface area in the XMU-MP-1-treated mice, compared to the DMSO-treated diabetic mice (severe diabetes + DMSO, $n = 9$, severe diabetes + XMU MP-1, $n = 8$); * $p < 0.05$.

3. Discussion

The main finding of this study was that treatment with the MST1/2 inhibitor XMU-MP-1 produced a beneficial effect in improving glucose tolerance in the STZ-induced diabetic mouse model. XMU-MP-1 treatment increased the number of pancreatic β cells and enhanced the Langerhans islet area in severe diabetic mice.

XMU-MP-1 is a potent and specific inhibitor of MST1 and MST2, which are key components of the Hippo pathway [11,20]. XMU-MP-1 is able to efficiently inhibit MST1/2 activities in various cell types including hepatocytes, the macrophage cell line, osteosarcoma cell line [18], rat primary cardiomyocytes [19], microglial cells [20], and breast cancer cell lines [21].

As for other kinase inhibitors, it is important to consider the selectivity of XMU-MP-1. A previous study reported XMU-MP-1 selectivity by performing a KINOMEscan analysis against a panel of 468 kinases [18]. The authors reported only very few kinases that displayed a high affinity to XMU-MP-1. It was reported that at the concentration of 1 μ M, XMU-MP-1 inhibited two kinases to the same extent as the inhibition of MST1/2. These were aurora kinase A (AURKA) and phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit gamma (PIK3CG). AURKA is involved in regulating β cell proliferation [22], whereas PIK3CG was shown to positively modulate insulin secretion [23]. Based on these data, the expected phenotypes of AURKA and PIK3CG inhibition would be a reduction in pancreatic β cell function, which is contrary to our findings. Thus, we believe that the phenotypes of the diabetic mice following XMU-MP-1 in our model were more likely due to inhibition of MST1/2 kinases.

Here, we found that XMU-MP-1 was also effective in inhibiting MST1/2 and hence induced YAP activity in the pancreatic beta cell line INS-1. This finding is important because previous reports showed

that genetic ablation of MST1 resulted in the protection against the development of STZ-induced diabetes mellitus in mice [16] via improvement of pancreatic β cell survival, and thus it is essential for studying the effects of XMU-MP-1 treatment in pancreatic β cells, as well as in the STZ-induced diabetes model.

Our *in vitro* data using the INS-1 cell line suggested that XMU-MP-1 treatment did not produce a significant effect in enhancing cell survival, following STZ treatment. However, we observed a slight trend of increasing cell survival against STZ-induced toxicity when XMU-MP-1 was added at a lower dose (1 μ M). The absence of the protective effects of XMU-MP-1 in the *in vitro* model might be due to a possible toxic effect of XMU-MP-1 to INS-1 cells. When added at higher doses (3–5 μ M), XMU-MP-1 caused a slight but statistically significant reduction of INS-1 cell survival. These effects might be cell-specific as toxicity was not reported when XMU-MP-1 was used in different cell types [18,19]. Although XMU-MP-1 is considered to be one of the most specific inhibitors of MST1/2, at higher doses, it also inhibits several other kinases such as AURKA, ABL1, DCAMKL1, and MAP3K2/3 [18]. Thus, the toxic effects of XMU-MP-1 on INS-1 cells might be due to an alteration of other signaling pathways and not solely due to the inhibition of the Hippo pathway.

In this study, we tested the effects of XMU-MP-1 treatment on an STZ-induced diabetes model in mice. STZ is widely used to induce experimental β -pancreatic cell damage, both *in vitro* and *in vivo*. STZ is a naturally occurring nitrosourea that is capable of inducing irreversible cell damage, through the production of free radicals and induction of DNA damage [24]. Its effects are specific to pancreatic β -cells, since it enters the cell via the glucose transporter 2 (GLUT2), which is abundantly expressed in the β - cells [25]. Since it targets the pancreatic β cells, the STZ-induced diabetes is often considered to be the most suitable model for T1DM. Although there were several limitations, such as possible toxic effects to other organs/cell types, STZ is still the most widely used model for T1DM [26].

In this study, we chose this model, because a previous report demonstrated that MST1 was involved in regulating pancreatic β cell apoptosis [16]. Active MST1 promotes pancreatic β cell apoptosis, a molecular process that resembles the progression of type-1 diabetes mellitus. Thus, inhibition of MST1 using a genetic or pharmacological approach should protect against the development of type 1 DM [16]. This was the main reason for the use of the STZ-induced diabetes model in this study. However, we believe that it is also important to study the effects of MST1 inhibition in the other type of DM, i.e., the type 2 DM. Several rodent models of T2DM are established including diet-induced T2DM (in which the animals were subjected to a high-fat and high-calorie diet for a period of time) and the genetically modified mouse models of T2DM [27].

A study using another inhibitor of MST1, Neratinib—which also showed MST1 inhibition to be protective—was also performed using STZ-induced diabetes mellitus [17]. Our data were consistent with these previous studies. We observed a significant improvement of glucose tolerance test (GTT) and fasting blood glucose in the XMU-MP-1-treated mice at the end of the experiments. However, analysis of random blood glucose (RBG) at various time-points did not show any changes between the XMU-MP-1 treated mice, compared to those treated with DMSO. The difference might be caused by the timing of the RBG test, which were carried out at random time-points. The GTT is considered to be a more sensitive assay to assess the physiological function of β cells in Langerhans islets, to respond to an acute increase of the blood glucose level. Analysis of the fasting blood glucose level supports the GTT data that XMU-MP-1 might produce beneficial effects in the STZ diabetes model. Interestingly, our data indicated that the positive effect of XMU-MP-1 treatment was observed in the severe diabetes group of mice, suggesting that the protective and possibly regenerative effects of XMU-MP-1 was more prominent if there was a significant damage of the pancreatic cells. Taken together, our data might suggest an improvement of pancreatic β cell function after XMU-MP-1 treatment, although it was only limited to the severe diabetic mice, when they were challenged to a high glucose load during GTT.

Our finding, at least in part, was in line with previous studies that used gene deletion and MST1 inhibitor neratinib to inhibit MST1 function [16,17]. However, the beneficial effects of XMU-MP-1 in preventing the extent of STZ-induced diabetes did not seem to be as strong as those of neratinib or

in the genetic knockout models. This was probably due to some degree of toxicity of XMU-MP-1 in pancreatic β cells.

So, how does XMU-MP-1 improve pancreatic β cell function in the severe STZ-induced diabetic mice? Histological analysis revealed an increased number of cells within the Langerhans islets that eventually expanded the Langerhans islet area in mice treated with XMU-MP-1. There are several possibilities to explain this phenotype. XMU-MP-1 induces cell proliferation; it reduces cell death by inhibition of apoptosis; or a combination of both. In other cell types, e.g., hepatocytes and intestinal cells, it is understood that both genetic ablation of MST1/2 and pharmacological inhibition using XMU-MP-1, resulted in increased cell proliferation [18,28,29]. In neonatal cardiomyocytes, XMU-MP-1 also induced cell proliferation, although this effect was not observed in adult cardiomyocytes, which are known to be terminally differentiated and mitotically inactive cells [19]. In contrast, tissue-specific knock out of MST1 and MST2 in pancreatic tissue resulted in the opposite phenotype, i.e., reduction of pancreatic size, which was associated with morphological defects [30,31], suggesting a critical role of the Hippo/YAP pathway in the pancreas during embryonic development. These findings showed that the role of the Hippo pathway might be organ- and context-specific and the effects of the MST1/2 pharmacological inhibition in the pancreas might not simply be due to induction of cell proliferation. Indeed treatment with MST1 inhibitor, neratinib, resulted in β cell mass recovery and attenuation of cell apoptosis, but it did not affect cell proliferation [17]. In contrast, following STZ stimulation, MST1 KO mice showed both a reduction of apoptosis and an increase in cell proliferation that eventually enhanced the β cell mass [16]. Thus, it is possible that XMU-MP-1, besides promoting β cell survival (as indicated in our in vitro model), also induced proliferation of the pancreatic β cells in vivo.

One of the most important factors in drug development is to understand the safety, toxicity, and possible side effects. It is known that inhibition of the core components of the Hippo pathway, which results in induction of the YAP activity, is associated with increased cell proliferation and survival. Previous observations using MST1/2 inhibitor (XMU-MP-1) and YAP/TAZ activator (TT-10) suggested increases in hepatocytes, cardiomyocytes, and small intestinal epithelial cell proliferation [18,32]. These reports might suggest possible carcinogenic effects of these compounds, when used for a longer term. However, our own observation on using XMU-MP-1 to treat cardiac hypertrophy in mice showed that there were no adverse effects in the heart, kidney, or liver [19]. Histological analysis as well as evaluation of serum markers of liver and kidney function indicated no significant damage in these major organs, following XMU-MP-1 treatment [19].

Overall, our study adds to the growing body of evidence that targeting pancreatic β cell survival and regeneration could become a new strategy for the treatment of diabetes mellitus. There are several molecular targets and compounds that were studied in this field [17,33,34]. Our data support the idea that the key components of the Hippo signaling pathway, in this case MST1/2, could become a target for diabetes therapy. Further studies are required to define the exact mechanism(s) as well as evaluate the toxicity and the off-target effects of the inhibitor. Equally important, studies using different models of diabetes, e.g., the high-fat-diet-induced diabetes model, also need to be conducted to understand if MST1/2 inhibition is also effective in controlling this type of pathological condition.

4. Materials and Methods

4.1. INS-1 Cell Culture

The INS-1 cells were obtained from Sigma and are widely used as a model to study pancreatic beta-cell functions in vitro [35]. INS-1 cells were cultured and maintained in a T-175 culture flask with a growth medium containing 500 mL of RPMI 1640 with no L-Glutamine (Gibco #11534446), 10% fetal bovine serum (Gibco #11550356), 100 U/mL of penicillin/streptomycin (Gibco #11528876), 50 μ M of 2-mercaptoethanol (Gibco #11528926), 1 mM of sodium pyruvate (Gibco #12539059), 2 mM of L-Glutamine (Gibco #15410314), and 10 mM of HEPES (Gibco #12509079). The medium was changed

every two days or when the cells reached 80–85% confluency. After that, the INS-1 cells were passaged to several T-175 flasks and maintained in the 5% CO₂ incubator at 37 °C. For experimental purposes, we seeded the INS-1 cells on 24-well plates (10⁵ cells per well for YAP luciferase and MTT assays) and 6-well plates (5 × 10⁵ cells per well for protein extraction).

4.2. Yes-Associated Protein (YAP) Luciferase Assay

The measurement of the transcriptional activity of YAP was conducted by utilizing the GAL4-TEAD-UAS luciferase reporter system, as described previously [19]. The INS-1 cells were seeded at a density of 10⁵ cells/well on 24-well plates and were kept in the 5% CO₂ incubator at 37 °C for 24 h. On the following day, the cells were transfected with both adenoviruses expressing GAL4-TEAD and UAS-luciferase. The treatments (DMSO as control, 1 μM, 3 μM, and 5 μM of XMU-MP-1) were administered concomitantly with the adenoviruses. After 24 h of incubation, the cells were washed with PBS (Phosphate Buffer Saline) and lysed with 1× cell culture lysis buffer (Promega #E1531), at room temperature for 20 min. A total 20 μL of the cell lysate was mixed with 50 μL of luciferase substrate (Promega #E151A) in a luminometer tube. The luciferase signal was detected using the LB9507 luminometer.

4.3. Western Blot

Protein expression levels were examined by Western blot. RIPA buffer was used to extract total protein from the INS-1 cells. Cells were seeded on 6-well plates, at a density of 5 × 10⁵ cells/well, then treated with DMSO, 1 μM, 3 μM, or 5 μM of XMU-MP-1 for 24 h. The protein concentration was quantified using the Pierce BCA assay kit system (Thermo Scientific #23225). A total of 20 μg of protein samples with 6× Laemmli buffer were loaded into SDS-polyacrylamide separating gel (8% for LATS1 detection and 12% for MOB1 detection). An electrophoresis system was used to separate the proteins based on the molecular size. After transferring the proteins onto nitrocellulose membrane and blocking the membrane with 5% skim milk, the membrane was incubated with primary antibodies—anti-phospho-MOB1 (Cell Signaling Technology, #8699S), anti-total-MOB1 (Cell Signaling Technology, #13730S), anti-phospho-LATS1 (Cell Signaling Technology, #9157S), and anti-total-LATS1 (Protein Tech #17049-1-AP), overnight at 4 °C. After washing in TBST buffer, the membrane was incubated with the secondary antibody (anti-rabbit IgG, HRP-linked antibody from Cell Signaling Technology #7074S), for an hour, at room temperature. The ECL Western blotting detection reagent (Amersham Biosciences #RPN2232) was used to visualize the protein bands in a ChemiDoc XRS+ Imaging system. The band intensities were quantified using the Image Lab software (3.0, Bio-Rad Laboratories, Inc., Hercules, CA, USA).

4.4. INS-1 Cell Survival Experiments

The INS-1 cells were treated with 1–2 mM of STZ (Sigma Aldrich #18883-66-4) for 24 h, to induce cell damage. DMSO-treated cells were used as a negative control. In the treatment groups, XMU-MP-1 (1–3 μM) was added concomitantly with STZ. Cells were then incubated at 37 °C for 24 h. MTT solution (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, Sigma) at a concentration of 0.5 mg/mL was used to detect the normal cell metabolites, by producing purple crystals at the bottom of the plates. The cells were incubated with the MTT solution for two hours at 37 °C. When the crystals appeared, they were dissolved using a solubilization solution containing isopropanol and 0.1 M HCl (100 μL/well). The absorbance of the dissolved crystals in the solubilization solution was measured using a Multiscan Ascent microplate reader (Thermo LabSystems Inc., Philadelphia, PA, USA), at 500 nm wavelength.

4.5. In Vivo Experiments

Animal experiments were performed in compliance with the principles for the ethical treatment of animals in research, as described in the Government Regulation of Republic of Indonesia No. 95/2012 and in accordance with the guidelines for the care and use of animals in biomedical

research (National Guidelines on Health Research Ethics 2011, Ministry of Health, Indonesia). The experimental procedures were approved by the Research Ethics Committee, Faculty of Medicine, Universitas Airlangga.

We used wild type male Balb/c mice, which were obtained from UD Wistar, Yogyakarta, a certified laboratory animal breeder supplier. We used 12-weeks-old mice with an average body weight (BW) of 24.22 ± 1.53 g, at the start of the experiments. Mice were housed in the experimental animal facility within the Institute of Tropical Disease (Universitas Airlangga) in standard housing conditions for laboratory animals. Mice were maintained on a 12-h light/dark cycle in a controlled temperature of 19–22 °C and humidity of 40–65% for 1 week, before the experiments were started. Mice were fed with standard chow diet BR501.

To generate a model of diabetes mellitus, mice were injected with STZ at a dose of 50 mg/kg BW/day, for 5 consecutive days. We analyzed random blood glucose (RBG) at basal levels before STZ injection (day 0), and at day 7, 14, 21, and 28, following the first STZ injection. The criterion used for diabetes was RBG level more than 200 mg/dL, at day 14 after the first STZ injection. At day 14, the mice were divided into 5 groups: (i) control group; (ii) diabetic mice with RBG less than 300 mg/dL and treated with XMU-MP-1; (iii) diabetic mice with RBG less than 300 mg/dL and treated with vehicle (DMSO); (iv) diabetic mice with RBG more than 300 mg/dL and treated with XMU-MP-1; and (v) diabetic mice with RBG more than 300 mg/dL and treated with vehicle (DMSO). XMU-MP-1 at a dose of 1 mg/kg BW/day or equal volume of vehicle (DMSO) were administered intraperitoneally, starting at day 15 after the first STZ injection, for a total of 21 days. At the end of the experiments (day 35), mice were sacrificed following intraperitoneal glucose tolerance test (GTT) and fasting blood glucose measurement.

4.6. Random Blood Glucose Measurement

Random blood glucose measurement was performed on blood samples obtained from the mouse tail vein. Glucose level was measured using a glucometer kit (One Touch Select; Lifescan; Malvern, PA, USA), following the protocol recommended by the manufacturer.

4.7. Intraperitoneal Glucose Tolerance Test

For the intraperitoneal GTT, mice were fasted for 12 h overnight. The following morning, mice were injected intraperitoneally with 40% glucose solution (Otsuka, Otsu, Japan), at a dose of 1 g glucose/kg body weight. Blood glucose levels were measured using a glucometer test, before glucose injection, and at 15, 30, 60, 90, and 120 min after glucose injection.

4.8. Histological Analysis of Pancreatic Tissues

Pancreas tissues collected at the end of the experiment were fixed using 4% normal buffered formalin for 24 h. Tissues were processed overnight using a Krime automated tissue processor and were then embedded in paraffin wax. The histological sections were prepared at 5 µm thickness, using a rotary microtome (Leica 2125, Chicago, IL, USA). Analysis of the Langerhans islets surface area and cell number were conducted using an Olympus BX-41 microscope with 400× magnification. Images were analyzed using the ImageJ software (v1.52, NIH, Bethesda, MD, USA).

4.9. Data Analysis

Data are presented as mean \pm standard error of the mean (SEM). Statistical analysis was performed using the GraphPad Prism software (v8.4.3, GraphPad Software, San Diego, CA, USA). One-way analysis of variance (ANOVA) was used to compare significance among groups. If ANOVA produced a significant value of F ($p < 0.05$) and there was no significant variance in homogeneity then the post-hoc multi comparison analysis (Tukey's) were conducted. For comparison between two groups, Student's t -test was used. Multiple t -test analysis was used to compare IPGTT data at each specific time-point; p value < 0.05 was considered to be significant.

Author Contributions: Z.F. designed and performed experiments, analyzed data, and wrote manuscript; B.A., performed experiments, analyzed data; F.Y.A., performed experiments, analyzed data; E.T., performed experiments; R.O., performed experiments; A.S.R. validation and analysis of histological data; M.A.A., designed and performed experiments, analyzed data, and supervised the project; M.I.L., project supervision and funding management; D.O., conceived the idea, designed experiments, analyzed data, wrote, and edited manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by a Mandate Research Grant from Universitas Airlangga, grant number 1408/UN3/2019.

Acknowledgments: We gratefully acknowledge the embryology laboratory assistants in the Faculty of Veterinary Medicine, Universitas Airlangga, and the NPMRD laboratory staff of the Institute of Tropical Disease, Universitas Airlangga, for the technical support during this study. We thank Dr. Nicholas Stafford (University of Manchester) for proofreading this article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. World Health Organization. *Global Health Estimates 2016: Disease burden by Cause, Age, Sex, by Country and by Region, 2000–2016*; World Health Organization: Geneva, Switzerland, 2018.
2. Saeedi, P.; Petersohn, I.; Salpea, P.; Malanda, B.; Karuranga, S.; Unwin, N.; Colagiuri, S.; Guariguata, L.; Motala, A.A.; Ogurtsova, K.; et al. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9(th) edition. *Diabetes Res Clin. Pract.* **2019**, *157*, 107843. [[CrossRef](#)] [[PubMed](#)]
3. Orasanu, G.; Plutzky, J. The pathologic continuum of diabetic vascular disease. *J. Am. Coll. Cardiol.* **2009**, *53*, S35–S42. [[CrossRef](#)] [[PubMed](#)]
4. Matthaedi, S.; Stumvoll, M.; Kellerer, M.; Haring, H.U. Pathophysiology and pharmacological treatment of insulin resistance. *Endocr. Rev.* **2000**, *21*, 585–618. [[CrossRef](#)] [[PubMed](#)]
5. Ilonen, J.; Lempainen, J.; Veijola, R. The heterogeneous pathogenesis of type 1 diabetes mellitus. *Nat. Rev. Endocrinol.* **2019**, *15*, 635–650. [[CrossRef](#)] [[PubMed](#)]
6. Skyler, J.S.; Bakris, G.L.; Bonifacio, E.; Darsow, T.; Eckel, R.H.; Groop, L.; Groop, P.H.; Handelsman, Y.; Insel, R.A.; Mathieu, C.; et al. Differentiation of Diabetes by Pathophysiology, Natural History, and Prognosis. *Diabetes* **2017**, *66*, 241–255. [[CrossRef](#)] [[PubMed](#)]
7. Chaudhury, A.; Duvoor, C.; Reddy Dendi, V.S.; Kraleti, S.; Chada, A.; Ravilla, R.; Marco, A.; Shekhawat, N.S.; Montales, M.T.; Kuriakose, K.; et al. Clinical Review of Antidiabetic Drugs: Implications for Type 2 Diabetes Mellitus Management. *Front. Endocrinol.* **2017**, *8*, 6. [[CrossRef](#)] [[PubMed](#)]
8. Lu, J.; Liu, J.; Li, L.; Lan, Y.; Liang, Y. Cytokines in type 1 diabetes: Mechanisms of action and immunotherapeutic targets. *Clin. Transl. Immunol.* **2020**, *9*, e1122. [[CrossRef](#)]
9. Kim, K.A.; Lee, M.S. Recent progress in research on beta-cell apoptosis by cytokines. *Front. Biosci.* **2009**, *14*, 657–664. [[CrossRef](#)]
10. Akash, M.S.; Rehman, K.; Chen, S. Role of inflammatory mechanisms in pathogenesis of type 2 diabetes mellitus. *J. Cell. Biochem.* **2013**, *114*, 525–531. [[CrossRef](#)]
11. Misra, J.R.; Irvine, K.D. The Hippo Signaling Network and Its Biological Functions. *Annu. Rev. Genet.* **2018**, *52*, 65–87. [[CrossRef](#)]
12. Del Re, D.P.; Matsuda, T.; Zhai, P.; Maejima, Y.; Jain, M.R.; Liu, T.; Li, H.; Hsu, C.P.; Sadoshima, J. Mst1 promotes cardiac myocyte apoptosis through phosphorylation and inhibition of Bcl-xL. *Mol. Cell* **2014**, *54*, 639–650. [[CrossRef](#)]
13. Valis, K.; Prochazka, L.; Boura, E.; Chladova, J.; Obsil, T.; Rohlena, J.; Truksa, J.; Dong, L.F.; Ralph, S.J.; Neuzil, J. Hippo/Mst1 stimulates transcription of the proapoptotic mediator NOXA in a FoxO1-dependent manner. *Cancer Res.* **2011**, *71*, 946–954. [[CrossRef](#)]
14. Wen, W.; Zhu, F.; Zhang, J.; Keum, Y.S.; Zykova, T.; Yao, K.; Peng, C.; Zheng, D.; Cho, Y.Y.; Ma, W.; et al. MST1 promotes apoptosis through phosphorylation of histone H2AX. *J. Biol. Chem.* **2010**, *285*, 39108–39116. [[CrossRef](#)] [[PubMed](#)]
15. Ardestani, A.; Maedler, K. MST1: A promising therapeutic target to restore functional beta cell mass in diabetes. *Diabetologia* **2016**, *59*, 1843–1849. [[CrossRef](#)] [[PubMed](#)]

16. Ardestani, A.; Paroni, F.; Azizi, Z.; Kaur, S.; Khobragade, V.; Yuan, T.; Frogne, T.; Tao, W.; Oberholzer, J.; Pattou, F.; et al. MST1 is a key regulator of beta cell apoptosis and dysfunction in diabetes. *Nat. Med.* **2014**, *20*, 385–397. [[CrossRef](#)]
17. Ardestani, A.; Li, S.; Annamalai, K.; Lupse, B.; Geravandi, S.; Dobrowolski, A.; Yu, S.; Zhu, S.; Baguley, T.D.; Surakattula, M.; et al. Neratinib protects pancreatic beta cells in diabetes. *Nat. Commun.* **2019**, *10*, 1–17. [[CrossRef](#)] [[PubMed](#)]
18. Fan, F.; He, Z.; Kong, L.L.; Chen, Q.; Yuan, Q.; Zhang, S.; Ye, J.; Liu, H.; Sun, X.; Geng, J.; et al. Pharmacological targeting of kinases MST1 and MST2 augments tissue repair and regeneration. *Sci. Transl. Med.* **2016**, *8*, 352ra108. [[CrossRef](#)]
19. Triastuti, E.; Nugroho, A.B.; Zi, M.; Prehar, S.; Kohar, Y.S.; Bui, T.A.; Stafford, N.; Cartwright, E.J.; Abraham, S.; Oceandy, D. Pharmacological inhibition of Hippo pathway, with the novel kinase inhibitor XMU-MP-1, protects the heart against adverse effects during pressure overload. *Br. J. Pharmacol.* **2019**, *176*, 3956–3971. [[CrossRef](#)]
20. Qing, J.; Liu, X.; Wu, Q.; Zhou, M.; Zhang, Y.; Mazhar, M.; Huang, X.; Wang, L.; He, F. Hippo/YAP Pathway Plays a Critical Role in Effect of GDNF Against Abeta-Induced Inflammation in Microglial Cells. *DNA Cell Biol.* **2020**, *39*, 1064–1071. [[CrossRef](#)]
21. Sun, H.L.; Men, J.R.; Liu, H.Y.; Liu, M.Y.; Zhang, H.S. FOXM1 facilitates breast cancer cell stemness and migration in Yap1-dependent manner. *Arch. Biochem. Biophys.* **2020**, *685*, 108349. [[CrossRef](#)]
22. Hobson, A.; Draney, C.; Stratford, A.; Becker, T.C.; Lu, D.; Arlotto, M.; Tessem, J.S. Aurora Kinase A is critical for the Nkx6.1 mediated beta-cell proliferation pathway. *Islets* **2015**, *7*, e1027854. [[CrossRef](#)] [[PubMed](#)]
23. Li, L.X.; MacDonald, P.E.; Ahn, D.S.; Oudit, G.Y.; Backx, P.H.; Brubaker, P.L. Role of phosphatidylinositol 3-kinasegamma in the beta-cell: Interactions with glucagon-like peptide-1. *Endocrinology* **2006**, *147*, 3318–3325. [[CrossRef](#)] [[PubMed](#)]
24. Szkudelski, T. The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiol. Res.* **2001**, *50*, 537–546. [[PubMed](#)]
25. Schnedl, W.J.; Ferber, S.; Johnson, J.H.; Newgard, C.B. STZ transport and cytotoxicity. Specific enhancement in GLUT2-expressing cells. *Diabetes* **1994**, *43*, 1326–1333. [[CrossRef](#)] [[PubMed](#)]
26. Gvazava, I.G.; Rogovaya, O.S.; Borisov, M.A.; Vorotelyak, E.A.; Vasiliev, A.V. Pathogenesis of Type 1 Diabetes Mellitus and Rodent Experimental Models. *Acta Nat.* **2018**, *10*, 24–33. [[CrossRef](#)]
27. Fang, J.Y.; Lin, C.H.; Huang, T.H.; Chuang, S.Y. In Vivo Rodent Models of Type 2 Diabetes and Their Usefulness for Evaluating Flavonoid Bioactivity. *Nutrients* **2019**, *11*, 530. [[CrossRef](#)]
28. Zhou, D.; Conrad, C.; Xia, F.; Park, J.S.; Payer, B.; Yin, Y.; Lauwers, G.Y.; Thasler, W.; Lee, J.T.; Avruch, J.; et al. Mst1 and Mst2 maintain hepatocyte quiescence and suppress hepatocellular carcinoma development through inactivation of the Yap1 oncogene. *Cancer Cell* **2009**, *16*, 425–438. [[CrossRef](#)]
29. Zhou, D.; Zhang, Y.; Wu, H.; Barry, E.; Yin, Y.; Lawrence, E.; Dawson, D.; Willis, J.E.; Markowitz, S.D.; Camargo, F.D.; et al. Mst1 and Mst2 protein kinases restrain intestinal stem cell proliferation and colonic tumorigenesis by inhibition of Yes-associated protein (Yap) overabundance. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, E1312–E1320. [[CrossRef](#)]
30. Gao, T.; Zhou, D.; Yang, C.; Singh, T.; Penzo-Mendez, A.; Maddipati, R.; Tzatsos, A.; Bardeesy, N.; Avruch, J.; Stanger, B.Z. Hippo signaling regulates differentiation and maintenance in the exocrine pancreas. *Gastroenterology* **2013**, *144*, 1543–1553.e1. [[CrossRef](#)]
31. George, N.M.; Day, C.E.; Boerner, B.P.; Johnson, R.L.; Sarvetnick, N.E. Hippo signaling regulates pancreas development through inactivation of Yap. *Mol. Cell. Biol.* **2012**, *32*, 5116–5128. [[CrossRef](#)]
32. Hara, H.; Takeda, N.; Kondo, M.; Kubota, M.; Saito, T.; Maruyama, J.; Fujiwara, T.; Maemura, S.; Ito, M.; Naito, A.T.; et al. Discovery of a Small Molecule to Increase Cardiomyocytes and Protect the Heart After Ischemic Injury. *JACC Basic Transl. Sci.* **2018**, *3*, 639–653. [[CrossRef](#)] [[PubMed](#)]
33. Lee, D.; Kim, Y.M.; Jung, K.; Chin, Y.W.; Kang, K.S. Alpha-Mangostin Improves Insulin Secretion and Protects INS-1 Cells from Streptozotocin-Induced Damage. *Int. J. Mol. Sci.* **2018**, *19*, 1484. [[CrossRef](#)] [[PubMed](#)]

34. Zheng, S.; Zhao, M.; Ren, Y.; Wu, Y.; Yang, J. Sesamin suppresses STZ induced INS-1 cell apoptosis through inhibition of NF-kappaB activation and regulation of Bcl-2 family protein expression. *Eur. J. Pharmacol.* **2015**, *750*, 52–58. [[CrossRef](#)] [[PubMed](#)]
35. Hohmeier, H.E.; Mulder, H.; Chen, G.; Henkel-Rieger, R.; Prentki, M.; Newgard, C.B. Isolation of INS-1-derived cell lines with robust ATP-sensitive K⁺ channel-dependent and -independent glucose-stimulated insulin secretion. *Diabetes* **2000**, *49*, 424–430. [[CrossRef](#)] [[PubMed](#)]

Sample Availability: Samples of the compounds are not available from the authors.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Treatment with Mammalian Ste-20-like Kinase 1,2 (MST1,2) Inhibitor XMU-MP-1 Improves Glucose Tolerance in Streptozotocin-Induced Diabetes Mice

ORIGINALITY REPORT

17%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|---|-----|
| 1 | nutritionandmetabolism.biomedcentral.com Internet Source | <1% |
| 2 | uknowledge.uky.edu Internet Source | <1% |
| 3 | www.atsjournals.org Internet Source | <1% |
| 4 | bnrc.springeropen.com Internet Source | <1% |
| 5 | coek.info Internet Source | <1% |
| 6 | www.frontiersin.org Internet Source | <1% |
| 7 | www.omicsdi.org Internet Source | <1% |
| 8 | Chien-Hsun Huang, Fu-Ting Wang, Wen-Hsiung Chan. "Role of caspase-3-cleaved/activated PAK2 in brusatol-triggered | <1% |

apoptosis of human lung cancer A549 cells",
Toxicology Research, 2022

Publication

9

Kyung Kim, . "Radix clematidis extract protects against cytokine- and streptozotocin-induced \hat{I}^2 -cell damage by suppressing the NF- \hat{I}° B pathway", International Journal of Molecular Medicine, 1998.

Publication

<1 %

10

Xiaoling Lu, Huiqian Yu, Jiaoyao Ma, Kunkun Wang, Luo Guo, Yanping Zhang, Boan Li, Zehang Zhao, Huawei Li, Shan Sun. "Loss of Mst1/2 activity promotes non-mitotic hair cell generation in the neonatal organ of Corti", npj Regenerative Medicine, 2022

Publication

<1 %

11

dspace.knust.edu.gh

Internet Source

<1 %

12

thno.org

Internet Source

<1 %

13

www.jbc.org

Internet Source

<1 %

14

Abigail Robertson, Tamer M.A. Mohamed, Zeinab El Maadawi, Nicholas Stafford et al. "Genetic ablation of the mammalian sterile-20 like kinase 1 (Mst1) improves cell reprogramming efficiency and increases

<1 %

induced pluripotent stem cell proliferation and survival", Stem Cell Research, 2017

Publication

15

Cristiane A Dalla-Torre, Maisa Yoshimoto, Chung-Hae Lee, Anthony M Joshua et al. "Effects of THBS3, SPARC and SPP1 expression on biological behavior and survival in patients with osteosarcoma", BMC Cancer, 2006

Publication

<1 %

16

Konstantin N. Belosludtsev, Natalia V. Belosludtseva, Mikhail V. Dubinin. "Diabetes Mellitus, Mitochondrial Dysfunction and Ca²⁺-Dependent Permeability Transition Pore", International Journal of Molecular Sciences, 2020

Publication

<1 %

17

Lidan Sun, Chuandong Wang, Yuxuan Dai, Yingying Chu, Jing Han, Jie Zhou, Xingguang Cai, Wenlong Huang, Hai Qian. "Coumaglutide, a novel long-acting GLP-1 analog, inhibits β -cell apoptosis in vitro and invokes sustained glycemic control in vivo", European Journal of Pharmacology, 2015

Publication

<1 %

18

Xu Li, Shu Zhuo, Ting Zhuang, Yong Suk Cho et al. "YAP inhibits ER α and ER⁺ breast cancer

<1 %

growth by disrupting a TEAD-ER α signaling axis", Nature Communications, 2022

Publication

19 catalog.lib.kyushu-u.ac.jp <1 %
Internet Source

20 core.ac.uk <1 %
Internet Source

21 docksci.com <1 %
Internet Source

22 publications.aston.ac.uk <1 %
Internet Source

23 "Poster Sessions", Journal of Diabetes, 04/2011 <1 %
Publication

24 "The Diabetes Textbook", Springer Science and Business Media LLC, 2019 <1 %
Publication

25 Shu Zhuo, Xu Li, Yong Suk Cho, Yuchen Liu, Yingzi Yang, Jian Zhu, Jin Jiang. "YAP antagonizes TEAD-mediated AR signaling and prostate cancer growth", Cold Spring Harbor Laboratory, 2022 <1 %
Publication

26 downloads.hindawi.com <1 %
Internet Source

27 opac.ll.chiba-u.jp

Internet Source

<1 %

28

www.biorxiv.org

Internet Source

<1 %

29

Deepmala Singh, Ramprasad Regar, Pushpanjali Soppina, Virupakshi Soppina, Sriram Kanvah. "Imaging of lipid droplets using coumarin fluorophores in live cells and *C. elegans*", *Journal of Photochemistry and Photobiology B: Biology*, 2022

Publication

<1 %

30

Tercel, M, R N Marnane, M A Tatnell, R J Stevenson, A Halim, G-L Lu, C Duchesnes, M Truong, W A Denny, W R Wilson, and K G Mountjoy. "An indoline-derived compound that markedly reduces mouse body weight", *International Journal of Obesity*, 2012.

Publication

<1 %

31

Tomohiro Imamura, Yuki T. Yanagihara, Yasumasa Ohyagi, Norimichi Nakamura et al. "Insulin deficiency promotes formation of toxic amyloid- β 42 conformer co-aggregating with hyper-phosphorylated tau oligomer in an Alzheimer's disease model", *Neurobiology of Disease*, 2020

Publication

<1 %

32

actanaturae.ru

Internet Source

<1 %

33

elifesciences.org

Internet Source

<1 %

34

pubs.rsc.org

Internet Source

<1 %

35

scholar.ufs.ac.za:8080

Internet Source

<1 %

36

services.rmh.med.sa

Internet Source

<1 %

37

Hyunjung Park, Hyerin Park, Jiyeon Baek, Hyuk Moon, Simon Weonsang Ro. "Target Therapy for Hepatocellular Carcinoma: Beyond Receptor Tyrosine Kinase Inhibitors and Immune Checkpoint Inhibitors", *Biology*, 2022

Publication

<1 %

38

Vasileios Angelis, Stephen R. D. Johnston, Amin Ardestani, Kathrin Maedler. "Case Report: Neratinib Therapy Improves Glycemic Control in a Patient With Type 2 Diabetes and Breast Cancer", *Frontiers in Endocrinology*, 2022

Publication

<1 %

39

Z Salah. "WW domain interactions regulate the Hippo tumor suppressor pathway", *Cell Death and Disease*, 06/2011

Publication

<1 %

40 Floriane Gibault, Manon Sturbaut, Fabrice Baily, Patricia Melnyk, Philippe Cotelle. "Targeting Transcriptional Enhanced Associate Domains (TEADs)", Journal of Medicinal Chemistry, 2017
Publication <1 %

41 Lirong Ma, Jianning Li, Hui Song, Yan Li, Yue Wang, Yansong Luan, Yuhan Li, Hui Qi, Yi Yang, Lei Cai. "Lipotoxicity induced pancreatic beta cell damage is associated with GPR119/MST1/FoxO1 Pathway", Archives of Medical Science, 2021
Publication <1 %

42 Sri Umijati, Sri Kardjati, Ismudijanto, Sunarjo. "Empowering Mothers through Mentoring on 6-60 Months Children's Nutrition Care: An Effort to Prevent Child Malnutrition", Electronic Journal of General Medicine, 2021
Publication <1 %

43 Yoon Keun Cho, Yeonho Son, Abhirup Saha, Doeun Kim et al. "STK3/STK4 signalling in adipocytes regulates mitophagy and energy expenditure", Nature Metabolism, 2021
Publication <1 %

44 eurjther.com
Internet Source <1 %

45 hull-repository.worktribe.com
Internet Source

<1 %

46

journals.plos.org

Internet Source

<1 %

47

www.thno.org

Internet Source

<1 %

48

Kim, H.S.. "IFN- γ sensitizes MIN6N8 insulinoma cells to TNF- α -induced apoptosis by inhibiting NF- κ B-mediated XIAP upregulation", *Biochemical and Biophysical Research Communications*, 20051028

Publication

<1 %

49

Timea Frosch, Elisabeth Wyrwich, Di Yan, Juergen Popp, Torsten Frosch. "Fiber-Array-Based Raman Hyperspectral Imaging for Simultaneous, Chemically-Selective Monitoring of Particle Size and Shape of Active Ingredients in Analgesic Tablets", *Molecules*, 2019

Publication

<1 %

50

static-site-aging-prod2.impactaging.com

Internet Source

<1 %

51

www.science.gov

Internet Source

<1 %

52

Amin Ardestani, Blaz Lupse, Kathrin Maedler. "Hippo Signaling: Key Emerging Pathway in

<1 %

Cellular and Whole-Body Metabolism", Trends in Endocrinology & Metabolism, 2018

Publication

53

C. Uluoglu, D. B. Durakoglugil, C. Karasu, G. Ozbey, A. Gunes, H. Zengil. "The Effect of Experimental Diabetes on the Twenty - Four - Hour Pattern of the Vasodilator Responses to Acetylcholine and Isoprenaline in the Rat Aorta", Chronobiology International, 2009

Publication

<1 %

54

Chanyue Wang, Ye Wang, Hao Yang, Zirun Tian, Manli Zhu, Xiaoting Sha, Ju Ran, Linlin Li. "Uygur type 2 diabetes patient fecal microbiota transplantation disrupts blood glucose and bile acid levels by changing the ability of the intestinal flora to metabolize bile acids in C57BL/6 mice", BMC Endocrine Disorders, 2022

Publication

<1 %

55

Dahae Lee, Da Lee, Sungyoul Choi, Jin Lee, Dae Jang, Ki Kang. "Identification and Isolation of Active Compounds from Astragalus membranaceus that Improve Insulin Secretion by Regulating Pancreatic β -Cell Metabolism", Biomolecules, 2019

Publication

<1 %

56

Deng, Yaoting, Yurika Matsui, Wenfei Pan, Qiu Li, and Zhi-Chun Lai. "Yap1 plays a protective role in suppressing free fatty acid-induced apoptosis and promoting beta-cell survival", *Protein & Cell*, 2016.

Publication

<1 %

57

Huayu Shang, Trisha A. VanDusseldorp, Ranggui Ma, Yan Zhao, Jason Cholewa, Nelo Eidy Zanchi, Zhi Xia. "Role of MST1 in the regulation of autophagy and mitophagy: implications for aging-related diseases", *Journal of Physiology and Biochemistry*, 2022

Publication

<1 %

58

Jason D. Ray, Kyle B. Kener, Benjamin F. Bitner, Brent J. Wright et al. "Nkx6.1-mediated insulin secretion and β -cell proliferation is dependent on upregulation of c-Fos", *FEBS Letters*, 2016

Publication

<1 %

59

Jing Wang, Linwu Zhuang, Yan Ding, Zhenzhong Wang, Wei Xiao, Jingbo Zhu. "A RNA-seq approach for exploring the protective effect of Ginkgolide B on Glutamate-induced astrocytes injury", *Journal of Ethnopharmacology*, 2021

Publication

<1 %

60

Koji Takayama, Yohei Kawakami, Makoto Kobayashi, Nick Greco et al. "Local intra-

<1 %

articular injection of rapamycin delays articular cartilage degeneration in a murine model of osteoarthritis", Arthritis Research & Therapy, 2014

Publication

61

Minyang Fu, Yuan Hu, Tianxia Lan, Kun-Liang Guan, Ting Luo, Min Luo. "The Hippo signalling pathway and its implications in human health and diseases", Signal Transduction and Targeted Therapy, 2022

Publication

62

Shibu Marthandam Asokan, Ting Wang, Wei-Ting Su, Wan-Teng Lin. "Antidiabetic Effects of a Short Peptide of Potato Protein Hydrolysate in STZ-Induced Diabetic Mice", Nutrients, 2019

Publication

63

Tai L. Guo, Dori R. Germolec, Jian Feng Zheng, Linda Kooistra et al. "Genistein Protects Female Nonobese Diabetic Mice from Developing Type 1 Diabetes When Fed a Soy- and Alfalfa-free Diet", Toxicologic Pathology, 2014

Publication

64

Weibin Shi, Jing Li, Kelly Bao, Mei-Hua Chen, Zhenqi Liu. "Ldlr-Deficient Mice with an Atherosclerosis-Resistant Background Develop Severe Hyperglycemia and Type 2

<1 %

<1 %

<1 %

<1 %

Diabetes on a Western-Type Diet", Biomedicines, 2022

Publication

65

Xiaolei Cao, Chenliang Wang, Jiyang Liu, Bin Zhao. "Regulation and functions of the Hippo pathway in stemness and differentiation", Acta Biochimica et Biophysica Sinica, 2020

Publication

<1 %

66

dspace.cuni.cz

Internet Source

<1 %

67

repositori.irta.cat

Internet Source

<1 %

68

repository.ajou.ac.kr

Internet Source

<1 %

69

research.vu.nl

Internet Source

<1 %

70

thieme-connect.de

Internet Source

<1 %

71

www.spandidos-publications.com

Internet Source

<1 %

72

www.thieme-connect.com

Internet Source

<1 %

73

"Probiotic Research in Therapeutics", Springer Science and Business Media LLC, 2022

Publication

<1 %

74

Amelia U. Schirmer, Lucy M. Driver, Megan T. Zhao, Carrow I. Wells et al. "Non-canonical role of Hippo tumor suppressor serine/threonine kinase 3 STK3 in prostate cancer", *Molecular Therapy*, 2022

Publication

<1 %

75

Amin Ardestani, Matthew S. Tremblay, Weijun Shen, Kathrin Maedler. "Neratinib is an MST1 inhibitor and restores pancreatic β -cells in diabetes", *Cell Death Discovery*, 2019

Publication

<1 %

76

H. J. Welters. "Conditional expression of hepatocyte nuclear factor-1, the maturity-onset diabetes of the young-5 gene product, influences the viability and functional competence of pancreatic β -cells", *Journal of Endocrinology*, 07/01/2006

Publication

<1 %

77

Lusida, Maria Inge. "Preface", *Procedia Chemistry*, 2016.

Publication

<1 %

78

Mingming Liu, Meng Yan, Jinlong He, Huizhen Lv et al. "Macrophage MST1/2 Disruption Impairs Post-Infarction Cardiac Repair via LTB₄", *Circulation Research*, 2021

Publication

<1 %

79

Máire E. Doyle, Josephine M. Egan.
"Mechanisms of action of glucagon-like
peptide 1 in the pancreas", *Pharmacology &
Therapeutics*, 2007

Publication

<1 %

80

Nur Raihan Abdullah, Mohd Hamzah Mohd
Nasir, Nur Hafizah Azizan, Wan Abd Al Qadr
Imad Wan-Mohtar, Faez Sharif. "Bioreactor-
grown exo- and endo- β -glucan from
Malaysian *Ganoderma lucidum*: An in vitro
and in vivo study for potential antidiabetic
treatment", *Frontiers in Bioengineering and
Biotechnology*, 2022

Publication

<1 %

81

Shoshy A. Brinch, Enya Amundsen-Isaksen,
Sandra Espada, Clara Hammarström et al.
"The Tankyrase Inhibitor OM-153
Demonstrates Antitumor Efficacy and a
Therapeutic Window in Mouse Models",
Cancer Research Communications, 2022

Publication

<1 %

82

Toshikatsu Matsui, Tadahiro Shinozawa.
"Spontaneous recovery from sunitinib-
induced disruption of sarcomere in human
iPSC-cardiomyocytes and possible
involvement of the Hippo pathway", *BMC
Pharmacology and Toxicology*, 2021

Publication

<1 %

83 Zhang, Yaqin, Hongliang Mei, Wei Shan, Li Shi, Xiaoi Chang, Yunxia Zhu, Fang Chen, and Xiao Han. "Lentinan protects pancreatic β cells from STZ-induced damage", Journal of Cellular and Molecular Medicine, 2016.
Publication

84 assets.researchsquare.com
Internet Source

85 bsdwebstorage.blob.core.windows.net
Internet Source

86 dev.biologists.org
Internet Source

87 elib.suub.uni-bremen.de
Internet Source

88 iovs.arvojournals.org
Internet Source

89 jnanobiotechnology.biomedcentral.com
Internet Source

90 pancreapedia.org
Internet Source

91 preview-cellandbioscience.biomedcentral.com
Internet Source

92 spandidos-publications.com
Internet Source

| | | |
|-----|--|------|
| 93 | staff.najah.edu Internet Source | <1 % |
| 94 | www.hindawi.com Internet Source | <1 % |
| 95 | www.phcog.com Internet Source | <1 % |
| 96 | www.preprints.org Internet Source | <1 % |
| 97 | www.tandfonline.com Internet Source | <1 % |
| 98 | "Posters", <i>Diabetic Medicine</i> , 12/2006 Publication | <1 % |
| 99 | Anjana Sharma, Veera Ganesh Yerra, Ashutosh Kumar. "Emerging role of Hippo signalling in pancreatic biology: YAP re-expression and plausible link to islet cell apoptosis and replication", <i>Biochimie</i> , 2017 Publication | <1 % |
| 100 | Ardiansah Bayu Nugroho, Nicholas Stafford, Min Zi, Sukhpal Prehar et al. "MicroRNA-411 Expression Improves Cardiac Phenotype Following Myocardial Infarction in Mice", <i>JACC: Basic to Translational Science</i> , 2022 Publication | <1 % |
| 101 | Burkart, Volker. "Role of Poly(ADP-Ribose) Polymerase in the Pathogenesis of Pancreatic | <1 % |

Islet Cell Death and Type 1 Diabetes",
Handbooks in Pharmacology and Toxicology,
2000.

Publication

102 The Hippo Signaling Pathway and Cancer,
2013. <1 %

Publication

103 Yan Xia, Ting Chang, Yingmei Wang, Yixiong
Liu, Wenhui Li, Ming Li, Heng-Yu Fan. "YAP
Promotes Ovarian Cancer Cell Tumorigenesis
and Is Indicative of a Poor Prognosis for
Ovarian Cancer Patients", PLoS ONE, 2014 <1 %

Publication

104 "The Hippo Pathway", Springer Science and
Business Media LLC, 2019 <1 %

Publication

105 Hikari Hirakida, Shinsuke Nakamura, Satoshi
Inagaki, Shohei Tsuji, Masahiro Hayashi,
Masamitsu Shimazawa, Hideaki Hara. "Anti-
diabetic effects of astaxanthin-rich extract
derived from Paracoccus carotinifaciens on
pancreatic β cells", Journal of Functional
Foods, 2022 <1 %

Publication

106 Mohd Iqbal Yattoo, Archana Saxena,
Arumugam Gopalakris, Mahmoud Alagawany,
Kuldeep Dhama. "Promising Antidiabetic
Drugs, Medicinal Plants and Herbs: An <1 %

Update", International Journal of
Pharmacology, 2017

Publication

107 Pasquale Sileo, Clémence Simonin, Patricia Melnyk, Marie-Christine Chartier-Harlin, Philippe Cotelle. "Crosstalk between the Hippo Pathway and the Wnt Pathway in Huntington's Disease and Other Neurodegenerative Disorders", Cells, 2022

Publication

108 Qian Wu, Xuemeng Miao, Jingjing Zhang, Ludan Xiang et al. "Astrocytic YAP protects the optic nerve and retina in an experimental autoimmune encephalomyelitis model through TGF- β signaling", Theranostics, 2021

Publication

109 Rachel M McQuade, Vanesa Stojanovska, Rhian Stavely, Cara Timpani et al. "Oxaliplatin-induced enteric neuronal loss and intestinal dysfunction is prevented by co-treatment with BGP-15", British Journal of Pharmacology, 2018

Publication

110 Yoojin Seo, So-Yeon Park, Hyung-Sik Kim, Jeong-Seok Nam. "The Hippo-YAP Signaling as Guardian in the Pool of Intestinal Stem Cells", Biomedicines, 2020

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Treatment with Mammalian Ste-20-like Kinase 1,2 (MST1,2) Inhibitor XMU-MP-1 Improves Glucose Tolerance in Streptozotocin-Induced Diabetes Mice

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15
