



**KOMISI ETIK PENELITIAN
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA
*Animal Care and Use Committee (ACUC)***

**KETERANGAN KELAIKAN ETIK
“ ETHICAL CLEARANCE ”**

No : 239-KE

**KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE)
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA,
TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG
DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :**

PENELITIAN BERJUDUL : Quiscent Bone Marrow Mesenchymal Stem Cells
Melalui Hypoxia Precondition Sebagai Upaya
Mempertahankan Viabilitas Kultur Stem Cells

PENELITI UTAMA : Erma Safitri

**UNIT/LEMBAGA/TEMPAT
PENELITIAN** : Program Studi Ilmu Kedokteran
Fakultas Kedokteran Universitas Airlangga

DINYATAKAN : LAIK ETIK

Surabaya, 14 Januari 2013

Mengetahui,
Dekan FKH-Unair

Prof. Romziah Sidik, Ph.D., drh.
NIP. 195312161978062001

Ketua,

Dr. E. Bimo Aksono, M.Kes., Drh.
NIP. 196609201992031003

C.33
C

The PHILIPPINE JOURNAL OF

Veterinary Medicine

Volume 55

Special Issue

December 2018

**Published by the College of Veterinary Medicine
University of the Philippines Los Baños**

ISSN 0031-7705



The 2nd Veterinary Medicine
International Conference
Surabaya, Indonesia
4-5 July 2018

The Philippine Journal of Veterinary Medicine

Volume 55

Special Issue

December 2018

The Philippine Journal of Veterinary Medicine is a peer-reviewed international journal of basic and applied research in veterinary medicine and science. It is published semi-annually, for the periods January-June and July-December each year, by the College of Veterinary Medicine, University of the Philippines Los Baños. All articles are subjected to double-blind review.

Authors of articles appearing in the journal are solely responsible for opinions expressed therein.

All rights reserved. No article of the journal may be reproduced in any form and by any means without a written permission from the publisher or the Editor-in-Chief.

EDITORIAL BOARD:

Jezie A. Acorda, DVM, MAgr, PhD
Editor-in-Chief

Dennis V. Umali, DVM, PhD
Associate Editor

Joseph F. dela Cruz, DVM, MS, PhD
Remil L. Galay, DVM, PhD
Technical Editors

Jesalyn L. Constante, DVM, MS
Business Manager

SUPPORT STAFF:

Ms. Jocelyn E. Arcinas
Mr. Fernando P. Micoso

The annual subscription price is US\$100.00 (net) for foreign subscribers (inclusive of mailing cost) and Philippine PhP1,500.00 plus mailing cost for local subscribers. Prices for current single issue and back issues are available on request. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by air delivery to foreign subscribers.

All communications should be addressed to:

The Editor-in-Chief

Philippine Journal of Veterinary Medicine

College of Veterinary Medicine

University of the Philippines Los Baños

Laguna, Philippines 4031

Telefax Nos. +63-49-536-2727, +63-49-536-2730

Email: pjvm1964@gmail.com, vetmed_uplb@yahoo.com.ph

This journal is Abstracted/Indexed by: SCOPUS, Biological Abstracts, Focus on: Veterinary Science & Medicine, Zoological Records, CAB Abstracts, Index Veterinarius, Veterinary Bulletin, Parasitology Database, Helminthological Abstracts, Protozoological Abstracts, Review of Medical and Veterinary Entomology, EBSCO, ASEAN Citation Index, Prescopus Russia, *i-journals* (www.ijournals.my), *i-focus* (www.ifocus.my), *i-future* (www.ifocus.my), Philippine E-Journals (<https://ejournals.ph>) and UPLB Journals Online (<http://journals.uplb.edu.ph/index.php/PJVM>).

© 2018 College of Veterinary Medicine, University of the Philippines Los Baños

The Philippine Journal of Veterinary Medicine

Volume 55

Special Issue

December 2018

CONTENTS

Original Articles

Medicine

- Viability of Rabbit Adipocyte Stem Cells Cultured Under Different Oxygen Concentrations *In Vitro*. 1
E Safitri, P Srianto, TV Widiyatno, W Sandhika and RH Prasetyo

Microbiology

- Antigenic Site of Glycoprotein Encoding Gene in Rabies Virus Isolate from Indonesia. 9
J Rahmahani, S Suwarno and FA Rantam
- Characterization of Newcastle Disease Virus Lentogenic Strain Infected Native Chickens from Surabaya, Indonesia. 17
FA Rantam, R Ernawati, AP Rahardjo, IL Rahmawati, D Kartika, NS Widjaja and J Rahmahani

Nutrition

- Effect of Concentrate to Forage Ratio on Milk Urea Nitrogen, Milk Production and Reproductive Performance of Dairy Cows. 25
S Utama, S Mulyati, W Wurlina and I Mustofa

Pathology

- Toxicity, Stability and Renal Histopathology of Alkaloid of Jarong (*Achyranthes aspera* Linn.) (Caryophyllales: Amaranthaceae) Leaf on Mice. 35
DK Meles, W Wurlina, I Mustofa, S Zakaria, A Basori, M Hariadi, E Safitri, DKSC Putri and N Suwasanti
- Histochemical Expression of Transforming Growth Factor Beta and Tumor Necrosis Factor Alpha in Rabbits Infected with *Sarcoptes scabiei*. 43
SM Rizki, LT Suwanti and NDR Lastuti

Pharmacology

- Effect of Alkaloid of *Achyranthes aspera* Linn. (Caryophyllales: Amaranthaceae) on Increasing Caspase 9, Caspase 3 and Apoptosis in Mice with Breast Cancer. 51
W Wurlina, DK Meles, I Mustofa, E Safitri, S Zakaria, A Basori, DKSC Putri and N Suwasanti

Theriogenology

- Effect of Aluminum Silicate on the Spermatozoa, Plasma Membrane and Seminiferous Tubules of Mice Exposed to *Fusarium graminearum* (Sordariomycetes: Hypocreales: Nectriaceae) 59
Samik, S Mulyati, T Hernawati and E Safitri

Research Notes

Microbiology

- Isolation and Identification of Lactic Acid Bacteria from the Digestive Tract of Kampung Chicken (*Gallus gallus domesticus*). 67
B Yulianto, WP Lokapirnasari
- In Vitro* pH Tolerance, Bile Salt Resistance and Antimicrobial Activity of *Lactobacillus plantarum* Isolated from Crossbred Cattle.73
WP Lokapirnasari, AM Sahidu, L Maslachah, K Soepranianondo, AB Yulianto, D Afikasari, TB Pribadi and I Hariyati

Nutrition

- Amino Acid Sequence of Signal Transducers and Activators Transcription Proteins From Broilers. 79
A Ma'ruf, NMR Widjaja, N Hidajati and R Damayanti

Parasitology

- Antigenic Protein Profile of *Anisakis* spp. Larvae Isolated from Mackerel Tuna Fish (*Euthynnus* sp.)..... 85
ZN Wastomi, NDR Lastuti, R Ernawati, LT Suwanti, S Koesdarto, M Mufasirin and HM Raharjo
- Morphological Detection of the Intestinal Parasite *Blastocystis* sp. in Fresh and Cultured Feces of Pet Sugar Glider (*Petaurus breviceps*) in Surabaya, Indonesia. 91
F Natalia, LT Suwanti, E Suprihati, Kusnoto, S Koesdarto and P Srianto

Pathology

- Comparative Histopathologic Changes in Rabbit (*Oryctolagus cuniculus*) Skin in Relation to Degree of Infestation with *Sarcoptes scabiei*. 97
A Azhimah, NDR Lastuti, A Arimbi, D Legowo, P Hastutiek and LR Yustinasari

Pharmacology

- Effect of Sapogenin from Sambiloto (*Andrographis paniculata*) (Lamiales: Acanthaceae) on Creatinine and BUN Levels and on Gentamicin-Induced Nephrotoxicity in Rats. 103
S Zakaria, W Wurlina, DK Meles, I Mustofa, M Hariadi, S Susilowati, E Safitri, A Basori, DKSC Putri and N Suwasanti

Public Health

- Identification of Shiga Toxin-Producing *Escherichia coli* in Raw Milk Samples from Dairy Cows in Surabaya, Indonesia..... 109
MH Effendi, N Harijani, SM Yanestria and P Hastutiek
- Tetracycline Resistance Gene in *Streptococcus agalactiae* Isolated from Bovine Subclinical Mastitis in Surabaya, Indonesia. 115
MH Effendi, A Oktavianto and P Hastutiek

Theriogenology

- Bacterial Isolates from the Cervical Mucus of Dairy Cattle at Follicular and Luteal Phases. 121
K Sudrajad, SP Madyawati, W Tyasningsih, R Rimayanti, P Srianto and OS Widodo

Human Chorionic Gonadotropin (hCG) from Urine of Pregnant Women for <i>In Vitro</i> Maturation of Madura Cattle Oocytes.....	127
<i>HA Hermadi, RTS Adikara, M Hariadi and E Safitri</i>	
Effect of Bovine Seminal Protein on the Quality of Frozen Spermatozoa from Goats.	133
<i>S Susilowati, IN Triana, TW Suprayogi, A Arimbi and W Wurlina</i>	
Editorial Policies.	139
Guidelines for Authors.	141

ORIGINAL ARTICLE**VIABILITY OF RABBIT ADIPOCYTE STEM CELLS CULTURED UNDER DIFFERENT OXYGEN CONCENTRATIONS *IN VITRO*****Erma Safitri¹, Pudji Srianto¹, Thomas Valentinus Widiyatno¹, Willy Sandhika² and R. Heru Prasetyo^{*2}**¹*Faculty of Veterinary Medicine; ²Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia***ABSTRACT**

The aim of this research was to determine if rabbit adipocyte stem cells (r-ASCs) can still be viable before transplantation when these are cultured *in vitro* under hypoxic preconditioning (3% O₂ concentration). Samples were divided into two groups, one under hypoxic preconditioning and the other in hyperoxia (20% O₂ concentration). Observations were made through flow cytometry, immunofluorescence and immunocytochemical analyses. Flow cytometric analysis showed that in r-ASCs hypoxic culture, the levels of CD90+, CD44+ and CD45- were unaltered, but changed under hyperoxic culture, indicated by the down-regulation of CD90+ and CD44+, and up-regulation of CD45-. Meanwhile, immunocytochemical and immunofluorescence analysis showed that under hypoxic preconditioning, r-ASCs culture expressed quiescent cells with p63 as marker, but the latter remained unexpressed in hyperoxic culture. In conclusion, hypoxic preconditioning with 3% O₂ concentration supported r-ASCs in sustaining viability before transplantation in rabbit.

Key words: hyperoxia, hypoxic preconditioning, rabbit, r-ASCs, viability

Philipp. J. Vet. Med., 55(SI): 1-8, 2018

INTRODUCTION

Sustaining the viability of stem cells is necessary before transplantation for use in several diseases, especially in cases involving degenerative tissues. This viability is what makes stem cells become adaptive and useful in achieving successful transplantation therapies.

The results of the study by Geng (2003) showed that the viability of mesenchymal stem cells (MSCs) injected in rats was very low, 99% of the injected MSCs were already undergoing apoptosis after day 4. This remarkably low viability may be associated with the damaged microenvironment of the tissue or organ,

failing to confer resilience to the stem cells. Increasing the viability, then, of transplanted stem cells is essential in improving the success rate in stem cell therapy.

The adaptability and viability of MSCs from bone marrow can be achieved by subjecting them to hypoxic preconditioning *in vitro* culture before transplantation (Safitri *et al.*, 2013; Safitri *et al.*, 2014a). Hypoxic preconditioning has proved that MSCs from bone marrow did not only confer viability and adaptability but also allowed stem cells to express quiescence *in vitro* as observed *in vivo*. Stem cell quiescence *in vivo* is known to inhabit the system long-term (Safitri *et al.*, 2014b).

The long-term maintenance of MSCs

***FOR CORRESPONDENCE:**

(email: rheru_prasetyo@yahoo.co.id)

depends on the interaction with their optimal niches, especially specific microenvironments (Arai and Suda, 2008). The specific microenvironments with optimal niches is critical to ensure lifelong tissue maintenance and to protect the stem cell pool from senescent cells (premature exhaustion) (Tsai *et al.*, 2011), and apoptosis (Wang *et al.*, 2008) under various stresses.

This study was conducted to determine the viability of rabbit adipocyte stem cells (r-ASCs) subjected to hypoxic preconditioning *in vitro* with 3% oxygen concentration. This study serves as a new approach in sustaining viability in ASCs, based on the unaltered expression of CD90+, CD44+, and CD45- and quiescent cells expression with p63 as marker.

MATERIALS AND METHODS

Animal studies were performed in accordance with the procedures approved by the Animal Care and Ethical Clearance Committee of the Faculty of Veterinary Medicine, Universitas Airlangga. Experiments also conformed with the National Research Council's guidelines (239-KE) through an ethical seminar. The research was conducted at an experimental laboratory within the Institute of Tropical Diseases and Veterinary Medicine Faculty, Universitas Airlangga.

Isolation and culture of r-ASCs

Adipose tissue was first aspirated from the peritoneal region of one 6-month old male rabbit (New Zealand strain). Aspirate was stored and kept at 4°C and transported to the experimental laboratory at the Institute of Tropical Diseases, Universitas Airlangga for *in vitro* culture. Adipose tissue was first washed and separated from the mixture of stomach fluid, blood and fats, and further separated using a sterile separator funnel. Samples were washed with sterile saline solution and heated at 37°C repeatedly until clear. The funnel was reversed for 4-5 min then closed; its position was kept still for 3-5 min to allow for separation. For tissue digestion, the same volume as that of the sample was prepared, along with warm sterile saline

buffer containing 500 CDU/ml (equivalent to 0.5 Wunsch units/ml) of collagenase. Washed fat was poured from the separatory funnel into a sterile bottle which can hold four times the aspirate volume. Saline buffer or collagenase mixture was added, covered and placed on a warm shaker that was prewarmed at 35-38°C for 20±5 min. Frequency and amplitude of the shaking was set at a point that would allow mixing but prevent the separation of the floating adipose tissue from the collagenase liquid.

Tissue was then transferred into a sterile glass separatory funnel and allowed to settle for 5-10 min to begin phase separation. The next stage of the stopcock was opened, and the non-floating fraction, through a sterile 265 mm filter, was obtained in a sterile beaker. Aliquots of the non-floating solution were then collected in a beaker glass and placed in a 50 ml centrifuge tube and mixed at 400 g for 5 min at room temperature with low to medium speed. Supernatant aspirate (top layer) was slowly poured in a tube without pellets from the cell. Finally, the suspension was filtered in a 100 mm cell strainer and collected in a sterile tube (Rantam *et al.*, 2009).

In vitro r-ASCs culture in hypoxic and hyperoxic preconditioning

Samples were centrifuged, then supernatant and cell suspension were aspirated and transferred into a 100 µl stromal inoculation of the cell medium placed in a plate. About 500 mg of adipose tissue were added with stromal medium until the container was full. Seventy-two hours after plating, aspirate was added in the medium. Next, the cell was washed with pre-warmed PBS (1% antibiotic added in solution), doing an up and down pipetting to gently clear the cells from tissue and blood cell fragments. Fresh stromal medium was added, just until well capacity of culture plate was reached. The subsequent medium was changed every 2-3 days until the cell has formed confluence of 80-90%, then the cells were harvested. This was done by removing the medium well first and storing it in a sterile tube under sterile media conditions to be used for subsequent cell culture applications. Next, 250-500 µl of cells

were added to the sterile warm PBS, letting the PBS be above the cell for 2 min. Furthermore, PBS was mixed with 500 μ l trypsin/EDTA solution (0.5%) and incubated for 5-10 min. At this point, observation under the microscope showed that 90% of the cells were removed. Five hundred μ l stromal medium were then added to fill the serum in solution. This was used to neutralize trypsin reaction.

The medium which contains the cell suspension was transferred from the well into a sterile tube, centrifuged at 300 g for 5 min. Small amounts of aspirate of supernatant and cell suspension were mixed with stromal medium. Furthermore, a counting process was done using a cell dilution aliquot in trypan blue (for a 1:7 dilution; 12.5 μ l cell suspension was added for every 87.5 μ l trypan blue. Cells were counted using a hemocytometer. After this, cells were placed back in the appropriate culture plate cells (Rantam *et al.*, 2009). Passage was conducted thrice, then cells were subjected to two preconditioning treatments: (1) hypoxic chamber with only 3% O₂ placed in an incubator with 5% CO₂; and (2) hyperoxic environment with 20% O₂ concentration.

Flow cytometry observation of CD90+ and CD44+ expression

Flow cytometry analysis was utilized to determine the expressions of CD90+ and CD44+. First, fat-free aspirate was centrifuged at 6000 rpm for 15 min at 4°C. The precipitate was mixed with cytoperm/cytofix two times the amount of cell number obtained. This mixture was centrifuged again and formed a supernatant and a pellet. BD wash was added to the pellet at four times the amount of cell number obtained from the first centrifugation. Mixture was centrifuged and added with lysis buffer at the amount twice that of the first obtained cell number. Subsequently, labeled antibody was added to each sample, arranging five tubes at once and processing in a similar manner. Wash tube was added with single staining with CD90 FITC and double staining with CD44 PE and CD45 PerCP. Finally, all samples were stored in 4°C in a darkroom and were analyzed with flow cytometry for 1 h (Macey, 2007).

Immunocytochemical analysis

Immunocytochemical (ICC) analysis was performed to determine the expression of p63 as marker of quiescent cells. Before the ICC methods were carried out, cytological preparation and further examination through immunocytochemical technique using monoclonal antibodies p63 was conducted. Observation on p63 expression was done using a light microscope. P63 expression is indicated by the number of cells with brownish chromogen discoloration (Kumar and Rudbeck, 2009).

Immunofluorescence analysis

Immunofluorescence analysis was performed to determine the expression of quiescent stem cells using p63 protein as marker (Safitri *et al.*, 2014b). After hypoxic preconditioning of r-ASCs, indirect immunofluorescence was done to detect p63 protein expression. The r-ASCs cell culture was harvested, placed in a 15 ml tube and fixed with methanol. After 15 min, each culture was coupled with a rabbit anti-p63 protein/P51A polyclonal antibody reagent, FITC conjugated primary antibodies (bs-0723R-FITC-Biossusa), were washed with PBS, then dropped on an object glass, and analyzed under a fluorescence microscope (Rantam *et al.*, 2014).

Statistical analysis

Expressions of CD90+ and CD44+ with flow cytometry and p63 with immunocytochemical method were statistically analyzed using SPSS 15 for Windows XP with the level of significance set at 0.01 and confidence level at 99%. Comparative hypothesis testing was done by testing data normality with Kolmogorov-Smirnov test, doing homogeneity of variance test, analysis of variance (ANOVA), and post-hoc test (least significant difference test) using Tukey's HSD at 5%.

Phenotype expression results were analyzed by immunofluorescence. Phenotype expressions analyzed were done on p63 expression as the marker for quiescence stem cells.

RESULTS AND DISCUSSION

The level of expressions of CD90+, CD44+, and CD45- in cells analyzed with flow cytometry showed that under hypoxic preconditioning in 3% O₂ concentration, r-ASCs culture were undifferentiated. Meanwhile, under 20% O₂ concentration, cells differentiated, as indicated

by the decreased expression of CD90+ and CD44+ (Fig. 1A-1C).

Based on Figure 1, the phenotype showed no significant differences between hypoxic treatments and control, suggesting that stem cells cultured in this study remained undifferentiated. Attaining undifferentiated cells was one of the main objectives of this




Fig. 1. Flow cytometric analysis of rabbit adipocyte stem cells (r-ASCs). A: culture under hypoxic preconditioning (3% O₂ concentration) showing positive expression of CD90+ and CD44+ (85.50%) and negative expression of CD45-; B: control, r-ASCs before hypoxic preconditioning culture, showing positive expression of CD90+ and CD44+ (75.56%) and negative expression of CD45-; C: culture under hyperoxia (20% O₂ concentration), showing positive expression of CD90+ and CD44+ (51.91%) and negative expression of CD45-.

study, since this is a key characteristic of long-term maintenance in stem cells. The results of this study is consistent with the report of Halim *et al.* (2010), which stated that undifferentiation is one of the unique characteristics of stem cells that sets it apart from other cells of the body.

The existence of this unique undifferentiated stem cells means that the *in vitro* cell culture in this study can survive much longer than the progenitor or the mature stem cells. This is based on the studies by Takubo (2012) and Elliason *et al.* (2010), where they emphasized that long-term maintenance (LTM) must be manifested by stem cell cultures to ensure the viability of transplanted stem cells. LTM, according to both researchers, can be attained when the stem cells are in the G0 phase when transplanted. G0 stage is the stage in which cultured cells are not bound to undergo the rest of the cell division (G1/S/G2/M) (Hermitte *et al.*, 2006; Arai and Suda, 2008). According to Morrison and Spradling (2008), Mohyeldin *et al.* (2010) and Suda *et al.* (2011), this stage is characterized by the presence of dormant or quiescent cells, and it is likely to occur if there is a conducive microenvironment for the stem cells.

The unique feature of stem cells is its ability to make it possible for any undifferentiated cell to become any kind of cell upon transplantation. This is the case of stem cells *in vivo*, such that, when under undifferentiated conditions, cells can differentiate into more than one cell type (multipotent/pluripotent). Thus, if undifferentiated cells' viability *in vitro* can be sustained, then cells that need to be transplanted can differentiate into any type of desired cell.

Immunocytochemical staining and immunofluorescence analysis enabled the expression of p63, the viability factor of mesenchymal stem cells. This positive expression was evident in r-ASCs cultured in hypoxic preconditioning at 3% O₂ concentration (Fig. 2A and 2B). Meanwhile, p63 expression was absent in cultured cells subjected to 20% O₂ concentration (Fig. 3A and 3B).

Hypoxic preconditioning at 3% O₂ concentration caused r-ASCs to remain undifferentiated and viable. This feature in

stem cells is important to ensure long-term tissue maintenance (Arai and Suda, 2008) and to protect the stem cell pool from premature exhaustion or turning into senescent cells (Tsai *et al.*, 2011) and undergoing apoptosis (Wang *et al.*, 2008). The lifelong maintenance of stem cells largely depends on the interaction with their specific microenvironment called niche (Arai and Suda, 2008).

In this study, hypoxic preconditioning with 3% O₂ concentration *in vitro* niche resembles the *in vivo* microenvironment. Specifically, under hypoxic preconditioning, inducible factor-1 α (HIF-1 α), a transcription factor, is stabilized. This means that HIF-1 α levels are elevated in stem cells, which regulate their metabolism (Simsek *et al.*, 2010; Takubo *et al.*, 2010).

The one mechanism relating to the favorability of hypoxic conditions for stem cell maintenance is the oxygen-dependent hydroxylation of amino acids on specific α subunit of the hypoxia-inducible transcription factors (HIFs) (Watt *et al.*, 2009). The activity and stability of HIF-1 α protein subunits eventually allow their targeting for proteosomal degradation. In this study, as oxygen levels drop below 3%, hydroxylation becomes inhibited, stabilizing the HIF1 α subunits to bind with HIF1 β , and then bind to hypoxia response element (HRE) in target genes, and, in association with transcriptional co-activators, turn these genes on.

Genes activated by HIFs include those encoding proteins involved in cell proliferation, self-renewal and survival. Self-renewal markers like SOX2 and OCT4 sustain undifferentiated state (Mohyeldin *et al.*, 2010).

Being undifferentiated is the basic characteristic of ASCs. In this study, undifferentiated state is indicated by the level of CD90+, CD44+, and CD45- expression in r-ASCs culture. Flow cytometric analysis of r-ASCs culture under hypoxic preconditioning with 3% O₂ concentrations showed positive expression of CD90+ and CD44+ (85.50%) (Fig. 1A), which was not significantly different ($P > 0.05$) with the control (before hypoxic preconditioning). The control group showed positive expression of CD90+ and CD44+ (75.56%) (Fig. 1B). Meanwhile, under




Fig. 2. Rabbit adipocyte stem cells (r-ASCs) cultured under hypoxic preconditioning (3% O₂ concentration). A: Immunocytochemical analysis: positive expression of p63 (red arrow head); B: Immunofluorescence analysis: positive expression of p63 (green fluorescence).




Fig. 3. Rabbit adipocyte stem cells (r-ASCs) cultured under hyperoxia (20% O₂ concentration). A: Immunocytochemical analysis: negative expression of p63 (red arrow head); B: Immunofluorescence analysis: negative expression of p63 (absence of green fluorescence).

hyperoxia (20% O₂ concentration), CD90+ and CD44+ expression was less pronounced (51.91%) (Fig. 1C).

This study shows that hypoxic preconditioning with 3% oxygen concentration enabled the maintenance of viability in r-ASCs before transplantation, as evident in the unaltered expressions of CD90+, CD44+, and CD45- and expression of quiescent cells with p63 as marker.

ACKNOWLEDGMENT

This research was partially supported by the funding from the Directorate General of Higher Education (DIKTI) 2017, Ministry of National Education, Indonesia.

REFERENCES

- Arai F and Suda T. 2008. *Quiescent Stem Cells in the Niche*. Stembook, ed. The Stem Cell Research Community, StemBook.
- Eliasson P, Rehn M, Hammar P, Larsson P, Sirenko O, Flippin LA, Cammenga J and Jonsson JI. 2010. Hypoxia mediates low cell-cycle activity and increases the proportion of long-term-reconstituting hematopoietic stem cells during *in vitro* culture. *Experimental Hematology* 38(4): 301-310.
- Geng YJ. 2003. Molecular mechanisms for cardiovascular stem cell apoptosis and growth in the hearts with atherosclerotic coronary disease and ischemic heart failure. *Annals of the New York Academy of Sciences* 1010(1): 687-697.
- Halim DH, Murty F, Sandra A, Boediono T, Djuwantono B and Setiawan B. 2010. *Stem Cell Basic Theory and Clinical Application*. Jakarta: Erlangga Press.
- Hermitte F, dela Grange PB, Belloc F, Praloran V and Ivanovic Z. 2006. Very low O₂ concentration (0.1%) favors G₀ return of dividing CD34+ cells. *Stem Cells* 24(1): 65-73.
- Kumar GL and Rudbeck L. 2009. *Immunohistochemical Staining Methods* (5th ed.). Carpinteria, CA: Dako North America.
- Macey MG (ed.). 2007. *Flow Cytometry: Principles and Applications* (3rd ed.). Jakarta: Humana Press.
- Mohyeldin A, Garzon-Muvdi T and Quiñones-Hinojosa A. 2010. Oxygen in stem cell biology: a critical component of the stem cell niche. *Cell Stem Cell* 7(2): 150-161.
- Morrison SJ and Spradling AC. 2008. Stem cells and niches: mechanisms that promote stem cell maintenance throughout life. *Cell* 132(4): 598-611.
- Rantam FA, Fezdiansyah N and Purwati A. 2009. *Stem Cell Exploration: Methods of Isolation and Culture* (1st ed.). Surabaya: Airlangga University Press.
- Rantam FA, Ferdiansyah X and Purwati X. 2014. *Mesenchymal and Hematopoietic Stem Cells Model Application* (2nd ed.). Surabaya: Airlangga University Press.
- Safitri E, Utama S, Bumi C, Mulyani SWM, Susilowati H, Retnowati E, Purwati, Prasetyo RH, Hariadi M and Rantam FA. 2013. The role of adaptive MSCs an attempt regeneration of spermatogenesis process using by hypoxia precondition *in vitro*. *Journal of Animal Production Advances* 3(11): 318-322.
- Safitri E, Utama S, Bumi C, Mulyani SWM, Retnowati E, Purwati, Prasetyo RH, Hariadi M, Aulani'am, Mahyudin F and Rantam FA. 2014a. Hypoxic preconditioning for viable and self renewing mesenchymal stem cells (MSCs) as the regeneration of spermatogenesis process. *Advances in Natural and Applied Sciences* 8(8): 42-46.
- Safitri E, Utama S, Prasetyo RH, Hariadi M and Rantam FA. 2014b. Quiescence stem cells (QSCs) for the treatment of infertility in male mice. *Asian Academy Research Journal of Multidisciplinary* 1(28): 187-199.
- Simsek T, Kocabas F, Zheng J, Deberardinis RJ, Mahmoud AI and Olson EN. 2010. The distinct metabolic profile of hematopoietic stem cells reflects their location in a hypoxic niche. *Cell Stem Cell* 7(3): 380-390.
- Suda T, Takubo K and Semenza GL. 2011. Metabolic regulation of hematopoietic stem cells in hypoxic niche. *Cell Stem Cell* 9(4): 298-310.
- Takubo K, Goda N, Yamada W, Iriuchishima H, Ikeda E and Kubota Y. 2010. Regulation of the HIF-1 α level is essential for hematopoietic stem cells. *Cell Stem Cell* 7(3): 391-402.
- Takubo K. 2012. The hypoxia regulatory system in hematopoietic stem cells. *Advances in Hematopoietic Stem Cell Research*.
- Tsai CC, Chen YJ, Yew TL, Chen LL, Wang JY, Chiu CH and Hung SC. 2011. Hypoxia inhibits senescence and maintains mesenchymal

- stem cell properties through downregulation of E2A-p21 by HIF-TWIST. *Blood Journal* 117(2): 459-469.
- Wang J, Chen TL, Jiang J, Shi H, Gui C, Luo RH, Xie XJ, Xiang MX and Zhang X. 2008. Hypoxic preconditioning attenuates hypoxia/reoxygenation-induced apoptosis in mesenchymal stem cells. *Acta Pharmacologica Sinica* 29(1): 74-82.
- Watt SM, Tsaknakis G, Forde SP and Carpenter L. 2009. Stem cells, hypoxia and hypoxia-inducible factors. In: Rajasekhar VK, Vemuri MC (eds.) *Regulatory Networks in Stem Cells. Stem Cell Biology and Regenerative Medicine*. Humana Press.

Author Index

- Adikara RTS, 127
 Afikasari D, 73
 Arimbi, 97, 121, 133
 Azhimah A, 97
 Basori A, 35, 51, 103
 Damayanti R, 79
 Effendi MH, 97, 109, 115
 Ernawati R, 17, 51, 85
 Hariadi M, 35, 91, 103, 127
 Harijani N, 97, 109
 Hariyati I, 73
 Hastutiek P, 97, 97, 109, 115
 Hermadi HA, 127
 Hernawati T, 59
 Hidajati N, 79
 Kartika D, 17
 Koesdarto S, 51, 57, 85, 91
 Kusnoto, 91
 Lastuti NDR, 43, 85
 Legowo D, 97
 Lokapirnasari WP, 67, 73
 Ma'ruf A, 79
 Madyawati SP, 121
 Maslachah L, 73
 Meles DK, 35, 51, 103
 Mufasirin, 51, 85
 Mulyati S, 59
 Mustofa I, 25, 35, 51, 103
 Natalia F, 91
 Oktavianto A, 115
 Prasetyo RH, 1
 Pribadi TB, 73
 Rahardjo AP, 17
 Raharjo HM, 51, 85
 Rahmahani J, 9, 17
 Rahmawati IL, 17
 Rantam FA, 9, 17
 Rimayanti R, 121
 Rizki SM, 43
 Safitri E, 1, 35, 51, 59, 103, 127
 Sahidu AM, 73
 Samik A, 59
 Sandhika W, 1
 Soepranianondo K, 73
 Srianto P, 1, 57, 91
 Sudrajad K, 121
 Suprayogi TW, 121, 133
 Suprihati E, 57
 Susilowati S, 91, 103, 133
 Suwanti LT, 43, 85, 91
 Suwarno, 9
 Suwasanti N, 35, 51, 103
 Triana IN, 133
 Tyasningsih W, 121
 Utama S, 25
 Wastomi ZN, 51, 85
 Widiyatno TV, 1
 Widjaja NMR, 79
 Widjaja NS, 17
 Widodo OS, 121
 Wurlina, 25, 35, 51, 103, 133
 Yanestria SM, 109
 Yulianto AB, 67, 73
 Yustinasari LR, 97

ACKNOWLEDGMENTS

The editorial staff wishes to thank the following who served as evaluators on an ad-hoc capacity for their critical review of manuscripts submitted to the journal:

- Dr. Marietta C. Amatorio, College of Veterinary Medicine, Benguet State University, Philippines
- Dr. Edwin C. Atabay, Philippine Carabao Center at Central Luzon State University (CLSU), Philippines
- Dr. Ayasan, Çukurova Agricultural Research Institute, Turkey
- Dr. Vasudevan Bakthavatchalu, Division of Comparative Medicine, Massachusetts Institute of Technology, USA
- Dr. Jose Arceo N. Bautista, Animal and Dairy Sciences Cluster, College of Agriculture, UPLB
- Dr. Esmeraldo M Cabana, College of Veterinary Science and Medicine, CLSU
- Dr. Gerry A. Camer, College of Veterinary Medicine, University of Eastern Philippines
- Dr. Joseph F. Dela Cruz, College of Veterinary Medicine, UPLB
- Dr. Rio John T. Ducusin, College of Veterinary Medicine, UPLB
- Dr. Salcedo L. Eduardo, College of Veterinary Medicine, UPLB
- Dr. Marianne Leila S. Flores, College of Veterinary Medicine, UPLB
- Dr. Gemerlyn G. Garcia, College of Veterinary Science and Medicine, CLSU
- Dr. Mary Joy N. Gordoncillo, OI Sub-Regional Representation for South-East Asia
- Dr. Hiromitsu Katoh, Osaka Prefecture University, Osaka, Japan
- Dr. Balasubramanian Manickam, Seventh Wave Laboratories LLC, Chesterfield, Montana, USA
- Dr. Carmencita D. Mateo, College of Agricultural and Food Science, UPLB
- Dr. Claro T. Mingala, Philippine Carabao Center National Headquarters and Gene Pool, Philippines
- Dr. Noraine P. Medina, College of Veterinary Science and Medicine, CLSU
- Dr. Anantharaman Muthuswamy, Wisconsin National Primate Research Center, Wisconsin, USA
- Dr. Mildred A. Padilla, College of Veterinary Medicine, UPLB
- Dr. Michelle Grace V. Paraso, College of Veterinary Medicine, UPLB
- Dr. Alessandra Pelagalli, Facoltà di Medicina Veterinaria, Università degli Studi di Napoli "Federico II", Napoli, Italy
- Dr. Antonio A. Rayos, Animal and Dairy Sciences Cluster, College of Agriculture, UPLB
- Dr. Frances C. Recuenco, Hokkaido University Graduate School of Medicine, Sapporo, Hokkaido, Japan
- Dr. Cesar C. Sevilla, College of Agricultural and Food Science, UPLB
- Dr. Luzviminda T. Simborio, College of Veterinary Medicine, CMU
- Dr. Guangliang (Johnny) Wang, John Hopkins University School of Medicine, Baltimore, USA

INDEXES TO VOLUME 55, Special Issue, December 2018

Keyword Subject Index

- Achyranthes aspera*, 35, 51
 alkaloid, 51, 69
 aluminum silicate, 59
 amino acid, 9
Anisakis spp., 85
 antigenic protein, 85
 antigenic site, 9, 17
 antimicrobial activity, 73
 apoptosis, 103
 apoptotic cell, 51
 bile salt, 73
Blastocystis sp., 91
 bovine, 19, 133
 breast cancer, 51
 broiler, 79
 caspase 3, 51
 caspase 9, 51
 congestion, 69
 culture medium, 91
 cytokine, 43
 dairy cattle, 121
 degeneration, 35
 dog, 9
Escherichia coli, 109
Euthynnus sp., 85
 follicular phase, 121
 frozen spermatozoa, 19, 133
Fusarium graminearum, 59
Gallus gallus domesticus, 67
 gentamicin, 103
 G-gene, 9
 goat, 19, 133
 hcg, 127
 histopathologic changes, 97
 hyperoxia, 1
 hypoxic preconditioning, 1
 immunohistochemistry, 43
in vitro maturation, 127
 Indonesia, 9, 17
 Kampung chicken, 67
 kidney, 35
 lactic acid bacteria, 67, 73
 lentogenic strain, 17
 luteal phase, 121
 Madura beef cattle, 127
 mice, 35
 multiplex PCR, 109
 milk urea nitrogen, 25
 native chicken, 17
 NCD, 17
 necrosis, 35, 103
 non-specific bacteria, 121
 oocytes, 127
 pH tolerance, 73
 plasma membrane, 59
 protozoan, 91
 rabbit, 1, 97
 rabies virus, 9
 r-ASCs, 1
 renal tubular cells, 103
 reproductive efficiency, 25
 sambiloto, 103
Sarcoptes scabiei, 43, 97
 scabies, 43, 97
 seminal protein, 19, 133
 seminiferous tubule, 59
 shiga toxin, 109
 smallholder dairy farmers, 25
 sperm, 59
 spermatozoa quality, 121, 133
 staining, 57
 STAT-1, 79
 STAT-3, 79
Streptococcus agalactiae, 115
 stx2 gene, 109
 staining, 91
 subclinical mastitis, 115
 sugar glider, 91

synthetic protein, 79
T-cell epitopes, 17
tetO gene, 115
tetracycline resistance, 115
TGF- β , 43

third-stage larvae, 85
TNF- α , 43
toxicity, 35
urine of pregnant women, 127
viability, 1

The Philippine Journal of Veterinary Medicine

Volume 55

Special Issue

December 2018



The 2nd Veterinary Medicine International Conference Surabaya, Indonesia 4-5 July 2018



Philippine Journal of Veterinary Medicine


4

Country Philippines · [SJR Ranking of Philippines](#)
Subject Area and Category [Vet-Ins](#) / [Veterinary \(miscellaneous\)](#)
Publisher University of the Philippines
Publication type Journals
ISSN 00317765
Coverage 2006-ongoing


H Index

[Join the conversation about this journal](#)

Quantiles



SJR



Citations per document



This website uses cookies to ensure you get the best experience on our website