

HUMAN CHORIONIC GONADOTROPIN FROM URINE OF PREGNANT WOMEN FOR IN VITRO MATURATION OF MADURA CATTLE OOCYTES

by Erma Safitri

Submission date: 07-Mar-2023 02:58PM (UTC+0800)

Submission ID: 2031026302

File name: Bukti_C18_Human_Chorionic_Gonadotropin_from....pdf (279.45K)

Word count: 3398

Character count: 17771

RESEARCH NOTE**HUMAN CHORIONIC GONADOTROPIN FROM URINE OF PREGNANT WOMEN FOR *IN VITRO* MATURATION OF MADURA CATTLE OOCYTES**

Herry Agoes Hermadi¹, Raden Tatang Santanu Adikara²,
Mas'ud Hariadi¹ and Erma Safitri^{1*}

¹Department of Veterinary Reproduction; ²Department of Veterinary Anatomy, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

ABSTRACT

The purpose of this study was to test whether human chorionic gonadotropin (hCG) from urine of pregnant women can be used for *in vitro* maturation of Madura cattle oocytes. Urine samples were collected from 50 healthy pregnant women gestating for 1.5 to 3.5 months. Molecular weights of hCG were 37 kDa and 22 kDa. HCG levels measured via ELISA had 10 times the deposition, with an average of 27,333 mIU/l at 1.5 months and 105,667 mIU/l at 3.5 months, respectively. T-test showed no significant difference in oocyte maturation rate between the control group with patent hCG and hCG from pregnant women at $P > 0.05$. This study illustrates that hCG from pregnant women at 1.5 to 3.5 months can aid in *in vitro* maturation of Madura cattle oocytes.

Key words: hCG, *in vitro* maturation, Madura beef cattle, oocytes, urine of pregnant women

Philipp. J. Vet. Med., 55(SI): 127-132, 2018

INTRODUCTION

Pregnancy tests, quantitative blood tests and the most sensitive urine tests in humans usually detect hCG between 6 and 12 days after ovulation (Wilcox *et al.*, 1999). However, it should be considered that the total hCG level varies a lot within the first four weeks of pregnancy, which makes tests susceptible to error (Butler *et al.*, 2001). A 35% increase in hCG level over 48 h is the proposed minimum increment, consistent with an appropriate intrauterine pregnancy test (Kirk *et al.*, 2013).

Human chorionic gonadotropin (hCG) is not specific-species and not limited to human hormones, *i.e.*, urine from pregnant women also holds therapeutic effects for receptive animals. In fact, Santibañez *et al.* (2014) suggested that intrauterine injection effect of hCG may be directly involved in *in vitro*

fertilization (IVF) process in humans before embryo transfer. HCG is a hormone the placenta produces after implantation (Gregory and Finlay, 1999; Cole, 2009) and this can be detected in several pregnancy tests.

Human chorionic gonadotropin is a glycoprotein composed of 237 amino acids. The molecular mass of intact, α (alpha) and β (beta) hCG are 36.7, 14.5 and 22.7 kDa, respectively (Gam and Latiff, 2005). It is heterodimeric with alpha subunits identical to luteinizing hormone (LH), follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH) and beta subunit unique to hCG. The α subunit contains 92 amino acids, while β subunit contains 145 amino acids, encoded by six high homologous genes composed of tandems and pairs on chromosome 19q13.3-CGB (Steel and Torrie, 1960; Canfield *et al.*; 1987; Agrawal *et al.*, 2000).

HCG treatment, in terms of embryonic

*FOR CORRESPONDENCE:

(email: erma-s@fkh.unair.ac.id)

development and oocyte collection, has been shown to be a more promising treatment to female infertility compared with recombinant human FSH. In the study by Ng *et al.* (2001), 85% of the oocytes reached metaphase at IVF treatment 2; consequently, it was tested whether there would be an additional benefit to administering FSH on top of hCG in priming *in vitro* maturation (IVM). Third to fifth cycles were treated with 75 IU FSH for 6 days, while 33 cycles were given 10,000 IU hCG 36 h before oocyte retrieval and fertilization via IVF and ICSI. The resulting embryo was replaced on day 2 or 3. A total of 1528 immature oocytes were found. The overall maturation and fertilization rates were 74.2 and 72.8%, respectively. After embryo transfer, 23 pregnancies (33.8%).

The use of hCG in animals, specifically on follicular growth and development *in vivo* *in vitro*, has been investigated. Due to its similarity to LH, hCG may also be used clinically to induce ovulation in the ovaries. For this reason, some organizations collect urine from pregnant women to extract hCG. Using hCG, comparative follicles begin to occur simultaneously, but many cases become atretic. During the luteal phase of the follicular growth cycle, one of the follicles becomes dominant. Notably, biochemical changes in the development of subordinate follicles to dominant follicles, when analyzed for chemical changes, are different from those in FSH, LH and receptor (Zemitis *et al.*, 2015).

In the study of molecular hCG, folliculogenesis is highly dependent on the influence of gonadotropin hormone; antral folliculi responsible for FSH-LH. Inhibin, activin, insulin-like growth factor I (IGF I) and protein walls (bonds with proteins) directly affect granulosa cells and theca cells that stimulate follicular growth and oogenesis (Gardner *et al.*, 2004). Injection of 1,000 IU hCG exogenously within 24 h of sow postpartum farrowing induced ovulation in 41%-75% of the population at 7-10 days after injection (Anderson *et al.*, 1999; Kirkwood *et al.*, 1999). Postpartum sow ovaries have potentially estrogenic medium-sized follicles (4 to 5 mm), and some sows have exhibited estrous behavior (De Rensis *et al.*, 1993; Sesti and Britt, 1993; Sesti and Britt, 1994; Langendijk *et al.*, 2007). But the postpartum estrous behavior observed at 2-4 days postpartum farrowing is anovulatory,

which is likely due to the inability to generate a pre-ovulatory LH surge (Sesti and Britt, 1993).

Medical experts up to this day have been relying on LH activity-dependent alone in inducing final oocyte maturation, which overlooks the redundancy of the natural waves of mid-cycle FSH surge. Thus, this needs to be resolved. Evidence from clinical research suggests that in an undetermined subset of patients, the LH and FSH waves are doubly advantageous compared to the LH spike in the form of a human chorionic gonadotropin (hCG) trigger. Double waves can be triggered by gonadotropin-releasing hormones, agonist boluses, causing endogenous LH and FSH flare-ups and resembling natural waves of concomitant hCG to ensure adequate exposure to LH activity (Kol and Humaidan, 2010). Further research is needed to characterize hCG from urine of pregnant women, which can then be potentially used for *in vitro* maturation of Madura cattle oocytes.

The purpose of this research was to extract hCG from urine of pregnant women gestating at 1.5 to 3.5 months, characterize and determine the molecular weight of the hCG protein and perform a biological potential test to determine its effect on the maturation level of Madura cattle oocytes.

MATERIALS AND METHODS

HCG isolation, identification and purification

Urine samples were taken from fifty healthy pregnant women gestating for 1.5 to 3.5 months. Samples were obtained by a midwife from a clinic in Surabaya, East Java, Indonesia. In the morning, 100 ml of urine was collected from each pregnant woman and these were centrifuged at 3,000 rpm for 15 min at 4°C to separate the metabolite cells. Thirty mg/100 ml activated charcoal powder was added, and the mixture was centrifuged at 3,000 rpm at 4°C for 20 min (Green and Leake, 1987) until mixture was homogeneous. Charcoal removes toxic substances, dyes and steroid hormones; it also absorbs and inactivates organic chemicals. Supernatant was filtered through an Erlenmeyer tube using a filter paper, producing a 50 ml supernatant. This was poured into Sephadex G-100 filtration device and poured into a chamber with running buffer

to carry out electrophoresis (Sakakibara *et al.*, 1987). Furthermore, hCG extract obtained from urine extract was processed through SDS-PAGE and ELISA (Gam and Latiff, 2005). A running gel was inserted through the well roughly less than the upper limit. Around 1 ml of butanol was added, and the gel was left for 25 min until it has solidified. Butanol was discarded, and the gel was cleansed with PBS and dried with Whatman paper. Comb was removed and remnants of the gel with buffer were cleaned.

HCG concentration

HCG concentration was determined through ELISA based on the sandwich principle (Mahaputra and Mustafa, 2003). The microtiter wells were coated with a monoclonal antibody directed towards a unique antigenic site of the hCG molecule. An aliquot of sample containing endogenous hCG was incubated in the coated well. After washing, a second incubation followed with an enzyme conjugate, an anti-hCG antibody conjugated with horseradish peroxidase. After incubation, the unbound conjugate was washed off. The amount of bound peroxidase was proportional to the concentration of hCG in the sample. Having added the substrate solution, color intensity developed was proportional to the concentration of hCG in the sample (Kariman *et al.*, 2011).

In vitro maturation of cattle oocytes

Extracted hCG was used to investigate its potential in inducing *in vitro* maturation in Madura beef cattle oocytes. There were two steps: the control group was given TCM 199 + 0.5 IU pregnant mare serum gonadotrophin (PMSG) + 0.5 IU hCG (Chorulon, Intervet, Holland), while the experimental group received the same treatment but with 0.5 IU hCG from gestating women.

Ovaries from slaughterhouses were washed 2 to 3 times with physiological saline solution, placed in a glass above water bath at 37°C, taken one by one with sterile tweezers and dried with sterile tissue paper. Using a 10 ml sterile disposable syringe, samples were washed with 1.5 ml of oocyte washing medium. Suction was done by stabbing the adjacent part of the ovary parenchyma in the next follicle bubble and directing the needle tip to the follicle (approximately 5 mm in diameter) near the point of needling (without

removing the needle first). After each follicle fluid was siphoned off (about 3-4 ml), this was carefully transferred into test tubes to avoid mechanical damage to the oocyte and placed in a water bath. After the oocyte had dropped to the bottom of the tube, precipitate was evaluated by placing it on a large petri dish and examining under a stereo microscope (Mahaputra and Mustofa, 2003).

Once an oocyte was found, this was collected with a modified pastry pipette (of the same diameter as the oocyte), placed on a smaller petri dish with washing medium and examined again under a microscope to determine its quality (Mahaputra and Mustafa, 2003). Afterwards, all oocytes were washed 3-4 times with washing medium and was finally washed with 2.5-3 ml of tissue culture medium 199 (TCM 199). Oocytes with no cumulus debris were transferred into the maturation medium, dropping 100 µl into a petri dish. A total of 10 oocytes were used in this study, prepared 2 h earlier in incubator with 5% CO₂. Four 100 µl drops of maturation medium were placed in each 35 mm sterile petri dish. Each drop of medium can culture as much as 5 oocytes. Media were then covered with mineral oil and placed in an incubator containing 5% CO₂ set at 39°C and humidity of 95-100% for 24 h. One percent aceto-orcein stain was added to determine maturity level of egg cell (Chen *et al.*, 2013).

To collect data on egg maturity, harvested eggs were placed and covered on the glass object, then placed in a fixative solution for 48 h. These were stained with 1% aceto-orcein for 2-3 min, washed with solution and viewed under an inverted microscope. Assessment of egg maturity level was based on germinal vesicle (GV), germinal vesicle breakdown (GVBD), metaphase I and metaphase II criteria (de Oliveira *et al.*, 2016). Mature oocytes were identified by locating those which can resume meiosis (extruding a first polar body) and can undergo fertilization (Fig. 1).

Data analysis

Data on identification, isolation, purification and measurement of hCG concentration from the urine of pregnant women at 1.5-3.5 months were processed descriptively. Student's t-test (Steel and Torrie, 1960) was used to determine the difference on oocyte maturation rate between patent hCG (Chorulon, Intervet, Holland) and

13
hCG from urine of pregnant women.

RESULTS AND DISCUSSION

HCG isolation, identification and purification

13
SDS-PAGE results showed that hCG from the urine of pregnant women had molecular sizes of 37 kDa and 22 kDa (Fig. 2). Gam and Latiff (2005) detected other bands from non-reduced fractionally desialylated-hCG sample: 43.5, 38.5, 29.45 and 20.85 kDa, which, after being reduced, resulted to an uncompounded pledge at 35.2 kDa.

Calculation of hCG concentration

The human pregnant urine ammonium sulphate precipitated concentrations of hCG at age 1.5 and 3.5 month were 27,333 mIU/l and 105,66 mIU/l, respectively. The measurement of hCG provides a specific test for pregnancy. Lyophilized hCG, although stable at room temperature for 3 weeks, should be stored at 2-8°C (Suthar and Shah, 2009). It is advised to reconstitute it in sterile 18 M-cm H₂O at a concentration of 1,000 IU/ml, which can be further diluted to other aqueous solutions.

In vitro maturation of cattle oocytes

Oocytes that matured *in vitro* are capable of resuming meiosis, extruding a first polar body and can undergo fertilization (Fig. 1). In this research, the rate of maturation of cow oocytes in TCM 199 + 0.5 IU PMSG + 0.5 IU hCG (Chorulon, Intervet, Holland) was 91.15±8.57, while maturation rate in TCM 199 + 0.5IU PMSG + 0.5 IU hCG from pregnant

18
women was 90.54±12.86. T-test showed no significant difference between groups (P>0.05) (Table), suggesting that hCG from pregnant woman can be used for maturation process of cow oocytes.

The dairy cattle industry has perfected the application of the first reproductive biotechnology, *i.e.*, artificial insemination (AI), a success story which made use of embryo transfer technology (ETT). In addition, emerging researchers have taken interest in the field of transvaginal oocyte recovery (TVOR) and *in vitro* embryo production (IVEP). IVF has paved the starting point for the production of generative material for many sophisticated reproduction techniques, like sperm microinjection into mature oocytes. In several countries, commercial IVF facilities are already being serviced by cattle ET operators. Also, various research groups have reported on the modification of TVOR (Suthar and Shah, 2009).

This study tested the use of hCG from urine of pregnant women in inducing maturation of cattle oocytes. Recorded molecular weight of hCG via SDS-PAGE was 37 and 22 kDa. HCG concentrations via ELISA assay had 10 times the deposition, with average level of 27,333 mIU/l at 1.5 months and 105,667 mIU/l at 3.5 months. T-test revealed no significant difference (P>0.05) between groups. This study illustrates that hCG extracted from urine of pregnant women at 1.5 to 3.5 months can be used to induce *in vitro* maturation in Madura cattle oocytes.




Fig. 1. *In vitro* maturation culture of Madura beef cattle oocytes at different stages. A: addition of hCG to maturation culture; B: immature oocyte; C mature oocyte; D: egg cells at metaphase stage.




Fig. 2. SDS-PAGE shows molecular weight of hCG from urine of pregnant woman at 37 and 22 kDa.

Table. Rate of maturation in cattle oocytes induced by different HCG sources.

hCG used	n	Mean \pm SD
TCM 199 + 0.05 PMSG + 0.5 IU hCG 0.5 hCG (Chorulon, Intervet, Holland)	50	91.15 \pm 8.57 ^a
TCM 199 + 0.5 IU PMSG + 0.5 IU hCG from pregnant women	50	90.54 \pm 12.86 ^a

ACKNOWLEDGMENT

This research was partially supported by funding from the Directorate General of Higher Education (DIKTI), Ministry of National Education, Indonesia.

REFERENCES

- Agrawal R, Holmes J and Jacobs HS. 2000. Follicle-stimulating hormone or human menopausal gonadotropin for ovarium stimulation in *in vitro* fertilization cycles: a meta-analysis. *Fertility and Sterility* 73: 338-343.
- Armstrong TA, Flowers WL and Britt JH. 1999. Control of the weaning-to-estrus interval in sows using gonadotrophins and prostaglandins during lactation. *Journal of Animal Science* 77(9): 2533-2539.
- Butler SA, Khanlian SA and Cole LA. 2001. Detection of early pregnancy forms of human chorionic gonadotropin by home pregnancy test devices. *Clinical Chemistry* 47(12): 2131-2136.
- Canfield RE, O'Connor JF, Birken S, Krichevsky A and Wilcox AJ. 1987. Development of an assay for a biomarker of pregnancy and early fetal loss. *Environmental Health Perspectives* 74: 57-66.
- Chen X, Chen SL, He YX and Ye DS. 2013. Minimum dose of hCG to trigger final oocyte maturation and prevent OHSS in a long GnRH α protocol. *Journal Huazhong University Science Technology Medical Sciences* 33(1): 133-136.
- Cole LA. 2009. New discoveries on the

- biology and detection of human chorionic gonadotropin. *Reproductive Biology and Endocrinology* 7(8): 1-37.
- de Oliveira SA, Calsavara VF and Cortés GC. 2016. Final oocyte maturation in assisted reproduction with human chorionic gonadotropin and gonadotropin-releasing hormone agonist (dual trigger). *JBRA Assisted Reproduction* 20(4): 246-250.
- De Rensis F, Hunter MG and Foxcroft GR. 1993. Suckling induced inhibition of luteinizing hormone secretion and follicular development in the early postpartum sow. *Biology of Reproduction* 48(5): 964-969.
- Gam LH and Latiff A. 2005. SDS-PAGE electrophoretic property of human chorionic gonadotropin (hCG) and its β -subunit. *International Journal of Biological Sciences* 1(3): 103-109.
- Gardner DK, Weissman A, Howles CM and Shoham Z. 2004. *Textbook of Assisted Reproductive Techniques* (4th ed.). Florida: CRC Press.
- Green B and Leake RE. 1987. *Steroid Hormones: A Practical Approach* (2nd ed.). Oxford: IRL Press.
- Gregory JJ and Finlay JL. 1999. Alpha-fetoprotein and beta-human chorionic gonadotropin: their clinical significance as tumour markers. *Drugs* 57(4): 463-467.
- Kariman N, Hedayati M, Taheri Z, Fallahian M, Salehpour S and Alavi Majd SH. 2011. Comparison of ELISA and three rapid HCG dipsticks in diagnosis of premature rupture of membranes. *Iran Red Crescent Medicine Journal* 13(6): 415-419.
- Kirk E, Bottomley C and Bourne T. 2013. Diagnosing ectopic pregnancy and current concepts in the management of pregnancy of unknown location. *Human Reproduction Update* 20(2): 250-261.
- Kirkwood RN, Henry SC, Tokach LM and Foxcroft GR. 1999. Human chorionic gonadotropin at parturition fails to consistently induce ovulation in sows. *Swine Health Production* 7: 69-71.
- Kol S and Humaidan P. 2010. LH (as HCG) and FSH surges for final oocyte maturation: sometimes it takes two to tango? *Reproductive Biomedicine Online* 21(5): 590-592.
- Langendijk P, Dieleman SJ, van den Ham CM, Hazeleger W, Soede NM and Kemp B. 2007. LH pulsatile release patterns, follicular growth and function during repetitive periods of suckling and non-suckling in sows. *Theriogenology* 67: 1076-1086.
- Mahaputra L and Mustafa I. 2003. Performance of serum cattle and horse estrus supplement of oocyte maturation media on *in vitro* fertilization of Madura cattle. *Jurnal Biosains Pasca Sarjana* 4(3): 113-117.
- Ng EH, Lau EY, Young WS and Ho PC. 2001. hMG is as good as recombinant human FSH in term of oocyte and embryo quality: a prospective randomized trial. *Human Reproduction* 16(2): 319-325.
- Sakakibara R, Tominaga N, Sakai A and Ishiguro M. 1987. Electrophoretic separation of proteins on agarose gel—application for direct immunization with a gel piece containing an antigen. *Analytical Biochemistry* 162(1): 150-155.
- Santibañez A, García J, Pashkova O, Colín O, Castellanos G, Sánchez AP and Jara JFD. 2014. Effect of intrauterine injection of human chorionic gonadotropin before embryo transfer on clinical pregnancy rates from *in vitro* fertilisation cycles: a prospective study. *Reproductive Biology and Endocrinology* 12(1): 9.
- Sestis LA and Britt JH. 1993. Influence of stage of lactation, exogenous luteinizing hormone-releasing hormone, and suckling on estrus, positive feedback of luteinizing hormone, and ovulation in sows treated with estrogen. *Journal of Animal Science* 71(4): 989-998.
- Sesti LA and Britt JH. 1994. Secretion of gonadotropins and estimated releasable pools of gonadotropin-releasing hormone and gonadotropins during establishment of suckling-induced inhibition of gonadotropin secretion in the sow. *Biology of Reproduction* 50(5): 1078-1086.
- Steel RGD and Torrie JH. 1960. *Principles and Procedures of Statistics*. New York: McGraw Hill.
- Suthar VS and Shah SG. 2009. Bovine *in vitro* embryo production: an overview. *Veterinary World* 2(12): 478-479.
- Wang W, Zhang XH, Wang WH, Liu YL, Zhao LH, Xue SL and Yang KH. 2011. The time interval between hCG priming and oocyte retrieval in ART program: a meta-analysis. *Journal of Assisted Reproduction Genetics* 28(10): 901-910.
- Wilcox AJ, Baird DD and Weinberg CR. 1999. Time of implantation of the conceptus and loss of pregnancy. *New England Journal of Medicine* 340(23): 1796-1799.
- Zemitis J, Emmy G, Bouwman P, Langendijk WHEJ, van Wettere and Kirkwood RN. 2015. Postpartum injection of human chorionic gonadotrophin: effects on sow ovarian follicles. *Journal of Swine Health and Production* 23(3): 137-139.

HUMAN CHORIONIC GONADOTROPIN FROM URINE OF PREGNANT WOMEN FOR IN VITRO MATURATION OF MADURA CATTLE OOCYTES

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	aasv.org Internet Source	1%
2	pubmed.ncbi.nlm.nih.gov Internet Source	1%
3	www.prospecbio.com Internet Source	1%
4	porkcrc.com.au Internet Source	1%
5	www.ncbi.nlm.nih.gov Internet Source	1%
6	Peter Kopp. "Chapter 5169 Hypercalciuria", Springer Science and Business Media LLC, 2022 Publication	1%
7	patents.justia.com Internet Source	1%
8	www.frontiersin.org Internet Source	

1 %

9

univmed.org

Internet Source

1 %

10

knepublishing.com

Internet Source

1 %

11

www.degruyter.com

Internet Source

1 %

12

ira.le.ac.uk

Internet Source

1 %

13

hal.archives-ouvertes.fr

Internet Source

1 %

14

www.notulaebiologicae.ro

Internet Source

1 %

15

Shahar Kol, Peter Humaidan. "LH (as HCG) and FSH surges for final oocyte maturation: sometimes it takes two to tango?",

Reproductive BioMedicine Online, 2010

Publication

<1 %

16

epubs.icar.org.in

Internet Source

<1 %

17

Chuanming. Duan, Mark E. Meyerhoff.

"Separation-Free Sandwich Enzyme

Immunoassays Using Microporous Gold

Electrodes and Self-Assembled

<1 %

Monolayer/Immobilized Capture Antibodies", Analytical Chemistry, 2002

Publication

18

erepo.unud.ac.id

Internet Source

<1 %

19

rnajournal.cshlp.org

Internet Source

<1 %

20

"EFFECT OF EPIDERMAL GROWTH FACTOR ON IN VITRO PRODUCTION OF CAMEL (CAMELUS DROMEDARIUS) EMBRYOS BY USING FROZEN SEMEN", Egyptian Journal of Animal Production, 2012

Publication

<1 %

21

Caroline Prouillac, Farah Koraichi, Bernadette Videmann, Michelle Mazallon et al. "In vitro toxicological effects of estrogenic mycotoxins on human placental cells: Structure activity relationships", Toxicology and Applied Pharmacology, 2012

Publication

<1 %

22

bmcbgenomics.biomedcentral.com

Internet Source

<1 %

23

"Abstracts for poster presentation", Theriogenology, 2002

Publication

<1 %

24

twivf.blogspot.com

Internet Source

<1 %

25 C. Simon. "Similar endometrial development in oocyte donors treated with either high- or standard-dose GnRH antagonist compared to treatment with a GnRH agonist or in natural cycles", Human Reproduction, 07/29/2005
Publication

26 Robabeh Taheripanah, Marzieh Zamaniyan, Atefeh Moridi, Anahita Taheripanah, Narges Malih. "Comparing the effect of gonadotropin-releasing hormone agonist and human chorionic gonadotropin on final oocytes for ovulation triggering among infertile women undergoing intrauterine insemination: An RCT", International Journal of Reproductive BioMedicine, 2017
Publication

27 perdami.or.id
Internet Source

28 pure-oai.bham.ac.uk
Internet Source

29 rep.bioscientifica.com
Internet Source

30 Xuemei Liu, Ding Ma, Wenjuan Wang, Qinglan Qu, Ning Zhang, Xinrong Wang, Jianye Fang, Zhi Ma, Cuifang Hao. "Intrauterine administration of human chorionic gonadotropin improves the live birth rates of

patients with repeated implantation failure in frozen-thawed blastocyst transfer cycles by increasing the percentage of peripheral regulatory T cells", Archives of Gynecology and Obstetrics, 2019

Publication

31

Huiying Jie, Mingpeng Zhao, Odai Ali Mohammad Alqawasmeh, Carol Pui Shan Chan, Tin Lap Lee, Tinchiu Li, David Yiu Leung Chan. "rescue immature oocytes – a literature review ", Human Fertility, 2021

<1 %

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

HUMAN CHORIONIC GONADOTROPIN FROM URINE OF PREGNANT WOMEN FOR IN VITRO MATURATION OF MADURA CATTLE OOCYTES

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
