

KEH FKH Unair



**KOMISI ETIK HEWAN  
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA  
*Animal Care and Use Committee (ACUC)***

**KETERANGAN KELAIKAN ETIK  
“ ETHICAL CLEARENCE ”**

**No : 064.KEH.023.02.2014**

**KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE)  
FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA,  
TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG  
DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :**

**PENELITIAN BERJUDUL** : Alternative Pemanfaatan Madu untuk Induksi Stem Cell  
pada Tikus Putih (*Rattus norvegicus*) yang Mengalami  
Malnutrisi

**PENELITI UTAMA** : Erma Safitri

**UNIT/LEMBAGA/TEMPAT PENELITIAN** : Fakultas Kedokteran Hewan Universitas Airlangga

**DINYATAKAN** : LAIK ETIK

Surabaya, 23 Pebruari 2014

**Mengetahui  
Dekan FKH-Unair**



**Prof. Romziah Sidik, Ph.D., drh.  
NIP. 195312161978062001**


**Ketua,**

**Dr. E. Bimo Aksono, M.Kes., Drh.  
NIP. 196609201992031003**

Tech Science Press

# BIOCELL

ISSN 0327-9545 BIOCELL (Printed)  
ISSN 1667-5746 BIOCELL (Online)







## Editorial Board

BIOCELL

ISSN 0327-9545 (Printed)

ISSN 1667-5746

### Editor-in-Chief:

Prof. Alfredo Castro-Vazquez, Instituto de Histología y Embriología de Mendoza (IHEM), CONICET-Universidad Nacional de Cuyo), ARGENTINA

### Associate Editors:

- Prof. Hikmet Hakan Aydin, Department of Medical Biochemistry, Ege University, TURKEY
- Prof. Gregorio Bigatti, Laboratorio de Reproducción y Biología Integrativa de Invertebrados Marinos (LARBIM), Instituto de Biología de Organismos Marinos (IBIOMAR), CCT CONICET-CENPAT, ARGENTINA
- Prof. Juan Carlos Cavicchia, Instituto de Histología y Embriología de Mendoza (IHEM), CONICET-Universidad Nacional de Cuyo, ARGENTINA
- Prof. María Isabel Colombo, Instituto de Histología y Embriología de Mendoza (IHEM), CONICET-Universidad Nacional de Cuyo, ARGENTINA
- Prof. Miguel Walter Fornés, Instituto de Histología y Embriología de Mendoza (IHEM), CONICET-Universidad Nacional de Cuyo, ARGENTINA
- Prof. Francesco Marotta, ReGenera R&D International for Aging Intervention (Milano-Beijing) & VCC Preventive Medical Promotion Foundation, ITALY
- Prof. Luis S. Mayorga, Instituto de Histología y Embriología de Mendoza (IHEM), CONICET-Universidad Nacional de Cuyo, ARGENTINA
- Prof. Mahmoud Moustafa, Department of Biology, Faculty of Science, King Khalid University, SAUDI ARABIA
- Prof. Alicia Mabel Seltzer, Instituto de Histología y Embriología de Mendoza (IHEM), CONICET-Universidad Nacional de Cuyo, ARGENTINA
- Prof. Wan-Xi Yang, College of Life Sciences, Zhejiang University, CHINA

### Editorial Board:

- Prof. Alejandro Aballay, Department of Molecular Microbiology & Immunology, School of Medicine, Oregon Health & Science University, PORTLAND
- Dr. Vishnu Agarwal, Department of Biotechnology, Motilal Nehru National Institute of Technology, INDIA
- Prof. Osama Mohamed Ahmed, Physiology Division, Zoology Department, Faculty of Science, Beni-Suef University, EGYPT
- Dr. Handan Ak, Department of Medical Biochemistry, Ege University, TURKEY
- Dr. Muhammad Akram, Department of Eastern Medicine, Directorate of Medical Sciences, Faculty of Life Sciences, Government College University, PAKISTAN
- Prof. Subramanian Babu, VIT School of Agricultural Innovations and Advanced Learning (VAIAL), Vellore Institute of Technology, INDIA
- Prof. Sônia Nair Bão, Departamento de Biologia Celular, Instituto de Ciências Biológicas, Universidade de Brasília, BRAZIL
- Prof. Claudio Barbeito, Laboratory of Descriptive, Comparative and Experimental Histology and Embryology, School of Veterinary Sciences, National University of La Plata, ARGENTINA
- Prof. Manuel A. Barbieri, Department of Biology, Florida International University, USA
- Dr. Alfonso Blázquez-Castro, Department of Physics of Materials, Faculty of Sciences, Autonomous University of Madrid, SPAIN
- Prof. Néstor Oscar Bianchi, Instituto Multidisciplinario de Biología Celular (IMBICE, CONICET-Universidad Nacional de La Plata), ARGENTINA

- Prof. Rubén Bottini, Instituto de Biología Agrícola de Mendoza (IBAM, CONICET-Universidad Nacional de Cuyo), ARGENTINA
- Prof. Benzhi Cai, Department of Pharmacy, the Second Affiliated Hospital of Harbin Medical University, CHINA
- Prof. Hernandes F. Carvalho, Department of Structural and Functional Biology, Institute of Biology, State University of Campinas (UNICAMP), BRAZIL
- Prof. Orlando José Castejón, Instituto de Investigaciones Biológicas, Universidad del Zulia, VENEZUELA
- Dr. Marta Castiglia, Department of Surgical, Oncological and Oral Sciences, Section of Medical Oncology, University of Palermo, ITALY
- Prof. Hector Chemes, Centro de Investigaciones Endocrinológicas (CEDIE, CONICET), ARGENTINA
- Prof. Daniel Ramón Ciocca, Instituto de Biología y Medicina experimental de Cuyo (IMBECU, CONICET), ARGENTINA
- Prof. Zong Jie Cui, Institute of Cell Biology, Beijing Normal University, CHINA
- Dr. Néstor Raúl Curvetto, Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS, CONICET), ARGENTINA
- Prof. Hesam Dehghani, The Research Institute of Biotechnology, Ferdowsi University of Mashhad, IRAN
- Prof. Mohammad Hashemi, School of Medicine, Zahedan University of Medical Sciences, IRAN
- Dr. Peter John, Atta-Ur-Rahman School of Applied Biosciences (ASAB), National University of Sciences & Technology, PAKISTAN
- Prof. Doo Hun Kim, Department of Chemistry, College of Natural Science, Sookmyung Women's University, KOREA
- Prof. Kyung Min Kim, Division of Plant Biosciences, School of Applied BioSciences, College of Agriculture & Life Sciences, Kyungpook National University, KOREA
- Prof. Guodong Li, Department of General Surgery, the Fourth Affiliated Hospital of Harbin Medical University, CHINA
- Prof. Daniel M. Lombardo, Institute of Research and Technology in Animal Reproduction, University of Buenos Aires, ARGENTINA
- Dr. Ricardo Williams Masuell, Instituto de Biología Agrícola de Mendoza (IBAM, CONICET-Universidad Nacional de Cuyo), ARGENTINA
- Prof. Victor Benno Meyer-Rochow, Department of Biology, University of Oulu, FINLAND
- Prof. Carlos R. Morales, Department of Anatomy and Cell Biology, McGill University, CANADA
- Prof. Kasim Ocakoglu, Department of Energy Systems Engineering, Faculty of Technology, Tarsus University, TURKEY
- Prof. Roman R. Poznanski, Faculty of Medicine, Sultan Zainal Abidin University, Malaysia
- Prof. Susana Puntarulo, Instituto de Bioquímica y Medicina Molecular (IBIMOL), CONICET-Universidad de Buenos Aires, ARGENTINA
- Dr. Narasimha Reddy Parine, Genome Research Chair, Department of Biochemistry, College of Science, King Saud University, SAUDI ARABIA
- Dr. Carlos Bernardo Passera, Cátedra de Fisiología Vegetal, Universidad Nacional de Cuyo, ARGENTINA
- Dr. Eduardo L. Rodríguez-Echandía, Instituto de Biología y Medicina Experimental de Cuyo (IMBECU, CONICET), ARGENTINA
- Dr. Antoni Francesc Roig, Instituto Argentino de Nivología, Glaciología y Ciencias Ambientales (IANIGLA, CONICET-Universidad Nacional de Cuyo-Gobierno de Mendoza), ARGENTINA
- Prof. Jose Russo, Fox Chase Cancer Center, USA
- Dr. Prabodh Sehajpal, Department of Molecular Biology, Guru Nanak Dev University, INDIA
- Dr. Anquan Shang, Department of Laboratory Medicine, Tongji Hospital, Tongji University School of Medicine, CHINA
- Prof. Alberto J. Solari, II Cátedra de Histología, Facultad de Medicina, Universidad de Buenos Aires, ARGENTINA
- Prof. Virginie Sottile, Wolfson Centre for Stem Cells, Tissue Engineering and Modelling (STEM), The University of Nottingham, UK
- Amit Kumar Srivastava, Department of Biotechnology, Indian Institute of Technology Roorkee, INDIA
- Prof. Juan Carlos Stockert, Departamento de Biología, Facultad de Ciencias, Universidad Autónoma de Madrid, SPAIN

- Prof. Sergey Suchkov, Center for Personalized Medicine, Sechenov University; Dept for Translational Medicine, Moscow Engineering Physical Institute (MEPhI); Dept for Clinical Immunology, A.I.Evdokimov Moscow State University of Medicine & Dentistry, RUSSIA
- Dr. Wenjun Tu, Peking Union Medical College, Chinese Academy of Medical Sciences, CHINA
- Prof. Dahms Hans-Uwe, Department of Biomedical Science and Environmental Biology, Kaohsiung Medical University, TAIWAN
- Dr. Yirong Wang, School of Life Sciences, Peking University, CHINA
- Dr. Dan Weng, School of Environment and Biological Engineering, Nanjing University of Science and Technology, CHINA
- Dr. Haineng Xu, Ovarian Cancer Research Center, University of Pennsylvania Perelman School of Medicine, USA
- Dr. Xiaoxiang Yan, Departments of Cardiology, Shanghai Jiaotong University School of Medicine, CHINA
- Dr. Chuanbin Yang, School of Chinese Medicine, Hong Kong Baptist University, CHINA
- Prof. Binwu Ying, Laboratory Medicine, West China Hospital, Sichuan University, CHINA
- Prof. Haozhe Piao, Department of Neurosurgery, Liaoning Cancer Hospital & Institute, Cancer Hospital of China Medical University, CHINA
- Prof. Naufal Zagidullin, Department of internal Diseases, Bashkir State Medical University, RUSSIA
- Prof. Hao Yang, Translational medicine center, Honghui Hospital, Xi'an Jiaotong University, CHINA
- Prof. Dawei Zhang, College of Life Science, Sichuan University, CHINA
- Prof. Weili Zhang, State Key Laboratory of Cardiovascular Diseases, FuWai Hospital, Peking Union Medical College & Chinese Academy of Medical Sciences, CHINA
- Prof. Rui Zhang, Department of Colorectal Surgery, China Medical University Cancer Hospital & Institute, CHINA

(Sorting in alphabetical order by last name)



## Vol 43, No 4 (2019)

Open Access Subscription Access

### Table of Contents

#### Articles

[Isolation and pathogenicity test of Entomopathogenic fungi on Colorado Potatobeele \[\(Leptinotarsa decemlineata \[Say\] \(Coleoptera: Chrysomelidae\)\] under the laboratory conditions](#)

Engin Kiliç

PDF

[Correlation between MiRNA and osteogenic differentiation of bone marrow mesenchymal stem cells](#)

Chen Fei, Ji-Cheng Wang, Wei Wei, Yan Zhuang, Bin-Fei Zhang, Kun Zhang

PDF

[A potential of prognostic 7-genes expression signatures for predicts survival in patient with colorectal cancer](#)

Liewen Lin, Ligang Xia, Wenbiao Chen

PDF

[Alantolactone sensitizes H1650 lung adenocarcinoma cells to gefitinib via inhibition of STAT3 activation](#)

Bashir Ahmad, Tahir Mehmood, Muhammad Khan, Tonghui Ma

PDF

[Cognitive and memory enhancing activities of tannins on scopolamine-induced learning and memory impairment in rats](#)

Ju Hyun Park, Chang Doo Lee, Jung Kyu Lee, Chang Jong Kim, Ji-Yun Lee

PDF

[Progress in the treatment of secondary deformities after hypospadias surgery](#)

Qiyu Liu, Yangqun Li, Zhe Yang, Ning Ma, Weixing Wang, Lisi Xu

PDF

[The ever-challenging diagnosis of langerhans cells histiocytosis of bone in adult: A case report](#)

Han Feng, Chen Qi, Qu Wei

PDF

[Acetylated mangostin attenuates intrinsic apoptotic and anti-transforming activities of mangostin](#)

Han-Heom NA, Seung-Young KIM, Keun-Cheol Kim

PDF

[Effectivity of Honey to regenerate the production of testosterone by induction of endogenous stem cells of rat with low libido due to malnutrition](#)

Erma Safitri, Thomas V. Widiyatno, Willy Sandhika, R.Heru Prasetyo

PDF

ISSN: 1667-5746

---

# BIOCELL

---

ISSN 1667-5746  
ELECTRONIC

2019 43(4-1): 235-239



## Effectivity of honey to regenerate the production of testosterone by induction of endogenous stem cells of rat with low libido due to malnutrition

Erma Safitri<sup>1</sup>, Thomas V. Widiyatno<sup>2</sup>, Willy Sandhika<sup>3</sup>, R. Heru Prasetyo<sup>4,\*</sup>

<sup>1</sup>Department of Reproduction Veteriner, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, 60115, Indonesia, E-mail address: rma\_fispro@yahoo.com

<sup>2</sup>Department of Pathology Veteriner, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, 60115, Indonesia, E-mail address: thvwidiyatno@gmail.com

<sup>3</sup>Department of Pathology Anatomy, Faculty of Medicine, Universitas Airlangga, Surabaya, 60131, Indonesia, E-mail address: willysandh@gmail.com

<sup>4</sup>Department of Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya, 60131, Indonesia, E-mail address: rheru\_prasetyo@yahoo.co.id

### ABSTRACT

The aim of the study was to know effectivity of honey from bee forest (*Apis dorsata*) to regenerate the production of testosterone by induction of endogenous stem cells of rat with low libido due to malnutrition. 40<sup>th</sup> rats as animal model were divided 4<sup>th</sup> groups: normal male rats without honey (C-) and 3<sup>th</sup> groups rats with low libido due to malnutrition were consists of without honey (C+), with 30% (v/v) honey (T1) and 50% (v/v) honey (T2) during 10 days. Low libido condition was induced by fasting without food during 5<sup>th</sup> days. Results: The improvement of libido was based on immunoreactivities of testosterone were detected in testis tissue. Testosterone expression as an immune response for regenerate signal in group C-, C+, T1 and T2 respectively is  $2.95^c \pm 0.15$ ;  $0.19^a \pm 0.55$ ;  $0.27^a \pm 0.45$  and  $2.05^{bc} \pm 0.35$ . T1 and C+ were decreased significantly ( $p < 0.05$ ) compared with C- and T2, while T2 didn't show decrease significantly ( $p > 0.05$ ) compared with C-. The libido level were based 3 category of libido: the number of copulations, time of reaction and period of copulation. The T1 group was not increased of libido, while in the T2 group, there was an increased of libido that significant different ( $p < 0.05$ ) compared with T1 and C+. The conclusion is the 50% (v/v) honey from bee forest (*Apis dorsata*) can to be use regenerate the production of testosterone by induction of endogenous stem cells and improvement of libido of rat.

**Keywords:** Honey, *Apis dorsata*, testosterone, libido.

### 1. Introduction

One of the major cases of low libido in male rat can be caused by malnutrition with the testis tissue degeneration (Safitri<sup>a</sup> et al., 2016; Prasetyo and Safitri, 2016). Therapy for degenerative testis tissue was using stem cells in now and next few decades is very interesting and greatly increased sharply, because stem cells potential is very promising for utilization as a treatment (Caplan and Correa, 2011). Nevertheless, the complexity of the isolation method, in vitro culture process and transplant procedure with booster treatment are very expensive (Hariadi and Safitri, 2019). The innovation therapy is needed as an effort to degenerative testis tissue in male. The honey as bee product is a nutrient gotten from bees (Safitri<sup>b</sup> et al., 2016; Hasib et al., 2017). It could serve as an antioxidant and as an antibacterial (Naqvi et al., 2013). Antioxidant is an important substance that protects individuals from free radicals such as reactive oxygen species (ROS). Adequate consumption of an antioxidant can reduce the prevalence of cardiovascular failure, cancers, digestive tract disorder, cataract and other degenerative diseases (Silva et al., 2013; Prasetyo and Hestianah, 2017), It also leads to degenerative testicular (Safitri<sup>a</sup> et al., 2016). Consumption of honey is also effective in the treatment of diarrhea with low immune response (Prasetyo and Safitri, 2016) and the reproductive system disorder as well (Safitri<sup>b</sup> et al., 2016; Parwata et al., 2010).

As a conceptual solution, further study is needed to explore the benefits of the honey from the use of bee forest (*Apis dorsata*), to induce of endogenous stem cells as an alternative for stem cells therapy.

### 2. Materials and methods

\* Corresponding author: Department of Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya, 60131, Indonesia, E-mail address: rheru\_prasetyo@yahoo.co.id

### 2.1. Low libido condition

Low libido model of male rat was available through food fasting 5<sup>th</sup> days long, but drinking of water still administrated (Safitri<sup>a</sup> et al. 2016). The experiment animals was used in this study are healthy, 8-10 week old male rats (*Rattus norvegicus*), Wistar strain with a body weight of 250-300g each. They were placed in an individual plastic cage in experimental animal laboratory at Faculty of Veterinary Medicine, Universitas Airlangga. The study was approved by ethical committee vide Ethical Clearance (*Komisi Etik Penelitin, Fakultas Kedokteran Hewan Universitas Airlangga*), Animal Care and Use Committee (ACUC). They were divided into 4 groups, each group consist of 10 rats. Group C-: normal male rats without honey, Group C+: low libido male rats without honey, Groups T1 and T2: the low libido of rats were given 30% (v/v) and 50% (v/v) of honey product from bee forest (*Apis dorsata*) for 10 days.

### 2.2. Immunoreactivity of testosterone by Immunohistochemical (IHC)

The testosterone expression was performed by immunohistochemical (IHC) method. Before IHC observation were made histological preparation, by way of a incision was made transversely of testicular tissue from paraffin block. Furthermore, examination through IHC method using a monoclonal testosterone antibody (Monoclonal, P2G1, Thermo Fisher Scientific, Pittsburgh, PA, USA). That was done to determine the expression of testosterone. Observation of testosterone were made with 200x magnification of microscope (Nikkon H 600L Microscope; digital camera DS Fi2 300 megapixel), and the expression of each variable was indicated by score of cells with brownish discoloration in each incision (Hariadi and Safitri, 2019).

### 2.3. Libido observaton

The libido observation of male rats through single mating with the female rat. Estrus synchronization was performed on the female rat using the PMSSG-hCG hormone (Bioworld, 220606401-MIH9827, Thermo Fisher Scientific, Pittsburgh, PA, USA), this was done in order to provide same reproductive organs in estrus phase. The detection of the capability of the libido of male rats were done by the close monitoring of the female rats. The detection of the libido is done after 1 cycle of spermatogenesis process which takes about 35days and they were fed normally. The libido data was taken based on the observations of the male rats and the female rats in the cage. (Safitri et al., 2014; Canale and Postonia, 2000). The libido level were based 3 category of libido: the number of copulations, time of reaction and period of copulation These observations were made on the 4<sup>th</sup> day, after a one hour mating, after which the male rats were separated from the female rats.

### 2.4. Statistical analysis

Expression of score of testosterone and male rat libido that classified into 3 categories (the number of copulations, time of reaction and period of copulation) were statistically analyzed using SPSS 15 for Windows XP with the evel of significance 0.05 (=0.05) and the confidence level 99% ( $\alpha=0.01$ ). Test data normality with the Kolmogorof Smirnov test, homogeneity of variance test, Anova and post-hoc test using the Tukey HSD 5%.

## 3. Results

The improvement of libido was based on immunoreactivities of testosterone were detected in testis tissue. Testosteron expression as a immune response for regenerate signal of Leydig cells in testis in group: C-, C+, T1 and T2 respectively is  $2.95^c \pm 0.15$ ;  $0.19^a \pm 0.55$ ;  $0.27^a \pm 0.45$  and  $2.05^{bc} \pm 0.35$  (Table 1, Figure 1).

The observations of the male rat libido were classified into 3 categories (Safitri et al., 2014; Canale and Postonia, 2000): Number of mating per unit time, reaction time, and the mating period (time between one mating with the next mating). The result of the libido test after being observated was based on the statistics as shown on Table 2.

## 4. Discussion

The results of this study showed that the administration of a dose of 50% (v/v) honey product from *Apis dorsata* in drinking water for 10 days in the T2 group can be used to treat low libido of male rats. Observation for libido from the male rats, are divided into three categories (Safitri et al., 2014; Canale and Postonia, 2000): number of mating per unit time, reaction time or time at the introduction of the male and female animals until the first mating; and the mating period or time of one mating with the next mating. The control positive group (C+) experienced a decreased libido, because the adrenal cortex became ineffective in producing dehydroepiandrosterone (DHEA) due to malnutrition. Low levels of DHEA in the blood can be acause of decreased body stamina, fatigue, and also decreased libido. DHEA was produced by the renal adrenal cortex (Heckbert and Heian, 2002) and the Leydig cells (Hafez and Hafez, 2013) is the most potent precursor of steroid

hormones such as testosterone. It also occurs in the degenerative group of testes who have received 30% honeybee product (T1) for 10 days. This suggests that 30% dose of honey has not been able to restore libido as in the negative control group (C-).

**Table 1**  
Score of Testosterone expression in rat testis with IHC method in several treatments.


No	Treatments	Average score of testosterone ± SD
1.	Negative control group (C-) : no fasted (fed standard ad libitum) and without honey	2.95 <sup>c</sup> ± 0.15
2.	Positive control group (C+) : fasted for 5 <sup>th</sup> days and without honey	0.19 <sup>a</sup> ± 0.55
3.	Trial group 1 (T1) : fasted for 5 <sup>th</sup> days and 30% (v/v) honey during 10 <sup>th</sup> days	0.27 <sup>a</sup> ± 0.45
4.	Trial group 2 (T2) : fasted for 5 <sup>th</sup> days and 50% (v/v) honey during 10 <sup>th</sup> days	2.05 <sup>bc</sup> ± 0.35

<sup>a,b,c,d</sup> Different superscripts in the same column was significantly different ( $P < 0.05$ ).

**Table 2**  
Libido test based on 3 category of libido.

No	Treatments	Number of mating per unit of time (60 minutes) ±SD	3 Category of Libido	
			Reaction time or time at the introduction of the male and female animals until the first mating (in 60 minutes observation)±SD	Mating period or time of one mating with the next mating (during 60 minutes) (average in minutes) ± SD
1.	Negative control group (C-) : no fasted (fed standard ad libitum) and without honey	5.2 <sup>c</sup> ± 0.6	2.495 <sup>b</sup> ± 0.37	13.176 <sup>b</sup> ± 1.102
2.	Positive control group (C+) : fasted for 5 <sup>th</sup> days and without honey	0 <sup>a</sup> ± 0	0 <sup>a</sup> ± 0	0 <sup>a</sup> ± 0
3.	Trial group 1 (T1) : fasted for 5 <sup>th</sup> days and 30% (v/v) honey during 10 <sup>th</sup> days	0 <sup>a</sup> ± 0	0 <sup>a</sup> ± 0	0 <sup>a</sup> ± 0
4.	Trial group 2 (T2) : fasted for 5 <sup>th</sup> days and 50% (v/v) honey during 10 <sup>th</sup> days	3.1 <sup>b</sup> ± 0.497	5.56 <sup>c</sup> ± 1.75	18.64 <sup>c</sup> ± 1.245

<sup>a,b,c,d</sup> Different superscripts in the same column was significantly different ( $P < 0.05$ ).



**Fig. 1.** Immunohistochemical analysis from response immune signal of testicular tissue based on Testosterone expression (brown chromogen) on several treatments 200× magnification (Nikkon H600L Microscope; digital camera DS Fi2 300 megapixel). The different superscripts indicate significant difference at  $p < 0.05$ . A. the negative control group (C-) was on the score 2.95<sup>c</sup> ± 0.15; B. The positive control group (C+) was on the score 0.19<sup>a</sup> ± 0.55; C. The group 30% (v/v) honey (T1) was on the score 0.27<sup>a</sup> ± 0.45; D. The group 50% (v/v) honey (T2) was on the score 2.05<sup>bc</sup> ± 0.35.

DHEA was increased as a precursor of steroid hormones (testosterone), it is also responsible for fat metabolism and acts as an enzyme inhibitor of glucose 6-phosphat dehydrogenase, which has a role as a biocatalyst that changes glucose to fat. The increase in DHEA allows for an increase in the amount of free ATP in the body, thereby, increasing the body stamina (Joshi et al., 2018), libido and fertility (Hafez and Hafez, 2013).

The libido process begins from the stimulation in the hypothalamus, where dopamine is produced as neurotransmitters and neurohormones that affect the activities and sexual behavior in individuals. The stimuli received by the sensory nerve triggers the acetaminoline in the stimulation of the nitric oxide secretion from endothelial cells to activate the cGMP. Activation of the cGMP causes the muscles of the corpus cavernosus, and the penis to be relaxed. This causes the occurrence of dilatation of the penis arteriole so that the blood will flow in erectile tissue of penis. The erection of erectile tissue from the penis by the blood will cause the veins to be depressed and inhibit the release of blood and water so that there is an increase of turgor in the penis, which leads to erection (Guyton and Hall, 1997).

Lack of glucose in rats with malnutrition and degenerative testicles, were found in the positive control group (C+) and the 30% honey bee product group (T1). This caused less fuel and energy sources. Glucose is universal for all cells proliferation including spermatogenesis process. The unavailability of glucose leads to the absence of carbon source for synthesis of most other compounds such as amino acids, nucleic acids, fatty acids, cholesterol, and steroid hormones such as testosterone. In addition, glucose is needed as a precursor for a variety of other glucoses such as glycosaminoglycans, lactose and nucleotides (Marks et al., 2000).

However, the existence of malnutrition due to the control positive group (C+) means there is a deficiency of vitamin B complex and unavailability of glucose from the food ration, plus imperfect metabolic process, it can be assumed as free energy or ATP in the body. This would also decrease the stamina of the male rat so that there would be a manifestation of decreased libido.

The effectivity of honey from bee forest (*Apis dorsata*) to regenerate the production of testosterone by induction of endogenous stem cells of rat with low libido due to malnutrition and increase of libido of rat based on 3 categories could be improvement.

## 5. Conclusion

The conclusion is the 50% (v/v) honey from bee forest (*Apis dorsata*) can to be use regenerate the production of testosterone by induction of endogenous stem cells and improvement of libido of rat based on 3 category (the number of copulations, time of reaction and period of copulation)

## References

- Canale, D., Postonia, S., 2000. Libido and Hormones. 21-23pp. CNS Spectrum, East Melbourne.
- Caplan, A.I., Correa, D., 2011. The MSC: An injury drugstore. Cell Stem Cell, 6, 11-15.
- Guyton, A., Hall, J.E., 1997. Buku Ajar Fisiologi Kedokteran. Editor Iraati Setiawan, E.G., Jakarta, Indonesia. 1283-1288.
- Hafez, E.S.E., Hafez, B., 2013. Reproduction in farm animals. Wiley-Blackwel, Philadelphia, Pennsylvania, USA. 221-232.
- Hariadi, M., Safitri, E., 2019. Comparison of biotechnological culture of hypoxia-conditioned rat mesenchymal stem cells with conventional in vitro culture of normoxia-conditioned rat mesenchymal stem cells for testicular failure therapy with low libido in rats. Vet. World, 12, 916-924.
- Hasib, A., Muhamad, R., Reksa, T.Y., Artha, A.U., Safitri, E., 2017. Utilization of Sumbawa tropical forest honey *Apis dorsata* to improve fertility of Indonesai oriental magpie robin (*Copsychus saularis*). KnE Life Sci., 619-626.
- Heckbert, L., Heian, J.R., 2002. Acute the dehydroepiandrosteron (DHEA) effect on sexual arousal in post menopusal women in: Women Health Gend. Base. Med., 2, 155-162.
- Joshi, K., Hassan, S.S., Ramaraj, P., 2018. Differential biological effects of dehydroepi androsterone (DHEA) between mouse (B16F10) and human melanoma (BLM) cell lines. Dermato-Endocrin, 9, 1-10.
- Marks, D.B., Allan, D.M., Collen, M.S., 2000. Biokimia Kedokteran Dasar. Sebuah Pendekatan Klinis. 165-177pp. Alih Bahasa Braun UP, Penerbit Buku Kedokteran ECG, Jakarta, Indonesia.
- Naqvi, S.A.R., Mahmood, N., Naz, S., Hussain, Z., Sherazi, T.A., Khan, Z.A., 2013. Antioxidant and antibacterial evaluation of honey bee hive extracts using *in vitro* models. Mediterr. J. Nutr. Metab., 6, 247-253.
- Parwata, I.M.O.A., Ratnayani, K., Listya, A., 2010. Aktivitas aantiradikal bebas serta kadar beta karoten pada madu randu (*Ceiba pentandra*) dan madu kelengkeng (*Nephelium longata* L.). J. Kim., 4, 54-62.
- Prasetyo, R.H., Hestianah, E.P., 2017. Honey can repair damage of liver tissue due to protein energy malnutrition through induction of endogenous stem cells. Vet. World, 10, 711-715.

- Prasetyo, R.H., Safitri, E., 2016. Effects of honey to mobilize endogenous stem cells in efforts intestinal and ovarian tissue regeneration in rats with protein energy malnutrition. *Asian Pac. J. Reprod.*, 5, 198-203.
- Safitri, E., Utama, S., Prasetyo, R.H., Hariadi, M., Rantam, H.A., 2014. Quiescence stem cells for the treatment of infertility in male mice. *Asian Acad. Res. J. Multidisc.*, 1, 123-127.
- Safitri, E., Utama, S., Widiyatno, T.V., Sandhika, W., Prasetyo, R.H., 2016. Auto-regeneration of mice testicle seminiferous tubules due to malnutrition based on stem cells mobilization using honey. *Asian Pac. J. Reprod.*, 5, 31-35.
- Safitri, E., Widiyatno, T.V., Prasetyo, R.H., 2016. Honeybee product therapeutic as stem cells homing for ovary failure. *Vet. World*, 9, 1324-1330.
- Silva, T.M.S., Santos, F.P.D., Evangelista-Rodrigues, 2013. A phenolic profile, antioxidant activity and palynological analysis of stingless bee honey from Amazonas. *Northern Braz. J. Food Chem*, 1141(4), 3552-3558.