
## E-1SSN: 1309-100X

# Journal of

International

# **Dental and Medical**

Research



2021 - Vol. 14 – No. 2 http://www.jidmr.com

### **Editorial Board of JIDMR**

Prof. Dr. Izzet YAVUZ Editor-in-Chief and General Director

#### **Advisory Board**

Prof. Dr. Refik ULKU Editor for Medicine Prof. Dr. Zulkuf AKDAG Editor for Biomedical Research

Prof. Dr. Ozkan ADIGUZEL Associate Editor

Gajanan Kiran KULKARNI (CANADA) Betul KARGUL (TURKEY) Diah Ayu MAHARANI (INDONESIA) Francisco Cammarata- Scalisi (Venezuela) Myroslav Goncharuk- Khomyn (UKRAINE)	Ferranti WONG (UK) Zeki AKKUS (TURKEY) Michele CALLEA (ROME, ITALY) Zelal ULKU (TURKEY)	Moschos A. PAPADOPOULOS (GREECE) Lindawati S KUSDHANY (INDONESIA) Yasemin YAVUZ (TURKEY) Yuliya MAKEDONOVA (RUSSIA) Nik Noriah Nik HUSSEIN (MALAYSIA)
--	--	---

Guvenc BASARAN (TURKEY)

### Editorial Board

Abdel Fattah BADAWI (EGYPT) Abdurrahman ONEN (TURKEY) Ahmet YALINKAYA (TURKEY) Ahmet DAG (TURKEY) Ali Al-ZAAG (IRAQ) Ali BUMIN (TURKEY) Ali FADEL (EGYPT) Ali GUR (TURKEY) Ali Kemal KADIROGLU (TURKEY) Ali Riza ALPOZ (TURKEY) Ali Riza Tunçdemir (TURKEY) Allah Bakhsh HAAFIZ (USA) Alpaslan TUZCU (TURKEY) Alpen ORTUG (TURKEY) Armelia Sari WIDYARMAN (INDONESIA) Ashish AGGARWAL (INDIA) Avse GUNAY (TURKEY) Aziz YASAN (TURKEY) Awiruth KLAISIRI (THAILAND) Balasubramanian MADHAN (INDIA) Benik HARUTUNYAN (ARMENIA) Betul KARGUL (TURKEY) Betul URREHMAN (UAE) Bugra OZEN (TURKEY) Carlos Menezes AGUIAR (BRAZIL) Cemil SERT (TURKEY) Chiramana SANDEEP (INDIA) Christine Bettina STAUDT (SWITZERLAND) Cihan AKGUL (TURKEY) Claudia DELLAVIA ( ITALY ) Diah Avu MAHARANI (INDONESIA) Dinesh Rokaya (NEPAL) Edoardo BAUNER (ROMA) Emmanuel Joao N. Leal da SILVA (BRAZIL) Emin Caner TUMEN (TURKEY) Emrullah BAHSI (TURKEY) Ertunc Dayı (TURKEY) Fadel M. ALI (EGYPT)

Guven ERBIL (TURKEY) Halimah AWANG (MALAYSIA) Halit AKBAS (TURKEY) Heloisa Fonseca MARAO (BRAZIL) Hilal TURKER (TURKEY) Huseyin ASLAN (TURKEY) Igor BELYAEV (SWEDEN) Ilhan INCI (ZURICH) Ilker ETIKAN (TURKEY) Isil TEKMEN (TURKEY) Isin ULUKAPI (TURKEY) Jalen DEVECIOGLU KAMA (TURKEY) Kemal CIGDEM (TURKEY) Kemal NAS (TURKEY) Kewal KRISHAN (INDIA) King Nigel MARTYN(HONG KONG, CHINA) Kursat ER (TURKEY) Levent ERDINC (TURKEY) Luca TESTARELLI (ROME) Lucianne Cople MAIA (BRAZIL) Luciane Rezende COSTA (BRAZIL) Marri Sai ARCHANA (INDIA) Manoj KUMAR (INDIA) Marcelo Rodrigues AZENHA (BRAZIL) Marcia Cancado FIGUEIREDO (BRAZIL) Marco MONTANARI (ITALY) Margaret TZAPHLIDOU (GREECE) Maria Elisa Oliveira dos SANTOS (BRAZIL) Medi GANIBEGOVIC (BOSNIA and HERZEGOVINA) Mehmet DOGRU (TURKEY) Mehmet Emin ERDAL (TURKEY) Mehmet Sinan DOGAN (TURKEY) Mehmet Ünal (TURKEY) Mehmet Zulkuf AKDAG (TURKEY) Meral ERDINC (TURKEY) Michele CALLEA (ITALY)

Mohamed TREBAK (USA)

Nezahat AKPOLAT (TURKEY) Nihal HAMAMCI (TURKEY) Nik Noriah Nik HUSSEIN (MALAYSIA) Nicola Pranno (ROME) Nurten AKDENIZ (TURKEY) Nurten ERDAL (TURKEY) Orhan TACAR (TURKEY) **Ozant ONCAG (TURKEY)** Ozgur UZUN (TURKEY) Ozkan ADIGUZEL (TURKEY) Rafat Ali SIDDIQUI (PAKISTAN) Refik ULKU (TURKEY) Sabiha Zelal ULKU (TURKEY) Sabri BATUN (TURKEY) Sadullah KAYA (TURKEY) Saul Martins PAIVA (BRAZIL) Sedat AKDENIZ (TURKEY) Seher GUNDUZ ARSLAN (TURKEY) Selahattin ATMACA (TURKEY) Selahattin TEKES (TURKEY) Serdar ERDINE (TURKEY) Serdar ONAT (TURKEY) Sergio Adriane Bezerra DE MOURA (BRAZIL) Serhan AKMAN (TURKEY) Sertac PEKER (TURKEY) Seyed Amir Danesh Sani (USA) Seyit Burhanedtin ZİNCİRCİOĞLU (TURKEY) Shailesh LELE (INDIA) Sinerik N. AYRAPETYAN (ARMENIA) Smaraqda KAVADIA (GREECE) Sossani SIDIROPOULOU (GREECE) Stefano Di CARLO (ROME) Sunit Kr. JUREL (INDIA) Stephen D. SMITH (USA) Susumu TEREKAWA (JAPAN) Suha TURKASLAN (TURKEY) Suleyman DASDAG (TURKEY) Taskin GURBUZ (TURKEY)

#### Fahinur ERTUGRUL (TURKEY)

Feral OZTURK (TURKEY) Feridun BASAK (TURKEY) Ferranti WONG (UNITED KINGDOM) Feyzi Çelik (TURKEY) Feyzullah Uçmak (TURKEY) Figen SEYMEN (TURKEY) Filippo BATTELLI (ITALY) Filiz Acun KAYA (TURKEY) Flavio Domingues Das NEVES (BRAZIL) Folakemi OREDUGBA (NIGERIA) Francesca De Angelis (ITALY) Gajanan Kiran KULKARNI (CANADA) Gamze AREN (TURKEY) Gauri LELE (INDIA) Gonul OLMEZ (TURKEY) Gulsen YILMAZ (TURKEY) Gulten UNLU (TURKEY)

Mohammad Khursheed Alam (KSA) Mohammad SUHRON (INDONESIA) Mohammed Mustahsen URREHMAN (UAE) Moschos A. PAPADOPOULOS (GREECE) Mostaphazadeh AMROLLAH (IRAN) M.S. Rami REDDY (INDIA) Muhammad FAHIM (INDIA) Mukadder ATMACA (TURKEY) Murat AKKUS (TURKEY) Murat SOKER (TURKEY)

Mustafa KELLE (TURKEY)

Mustafa ZORTUK (TURKEY)

Muzeyyen YILDIRIM (TURKEY)

Neval Berrin ARSERIM (TURKEY)

#### Ufuk ALUCLU (TURKEY)

Ugur KEKLIKCI (TURKEY) Xiong-Li YANG (CHINA) Vatan KAVAK (TURKEY) Yasar YILDIRIM (TURKEY) Yasar YILDIRIM (TURKEY) Yavuz SANISOGLU (TURKEY) Yu LEI (USA) Yuri LIMANSKI (UKRAINE) Zafer C. CEHRELI (TURKEY) Zafer C. CEHRELI (TURKEY) Zeynep AYTEPE (TURKEY) Zuhal KIRZIOGLU (TURKEY)



#### Journal of International Dental and Medical Research / ISSN: 1309-100X

#### **TABLE OF CONTENTS / 2021; 14 (2)**

#### DENTISTRY

**EXPERIMENTAL ARTICLE** 

- Alendronate Associated with Bovine Bone Graft in Bone Defect Repair: A Histomorphometric Study Douglas Bertazo Musso, Condrado Dias do Nascimento Neto, Natália Marreco Weigert, Stela Maris Wanderley Rocha, Robson Almeida de Rezende, Elizabeth Pimentel Rosetti, Rossiene Motta Bertollo, Martha Chiabai Cupertino Castro, Daniela Nascimento Silva Pages 467-473
- 2. Effect of Silver Nanoparticles Synthesized Using Betel Leaf Extract Added into Orthodontic Adhesive on the Bracket's Tensile Bond Strength Yanuarti Retnaningrum, Ananto Ali Alhasyimi Pages 474-480

**EXPERIMENTAL ARTICLE** 

**EXPERIMENTAL ARTICLE** 

3. Reliability of Two Electronic Shade-Matching Devices Ali A. Razooki Al-Shekhli, Isra'a Al Aubi Pages 481-484

**EXPERIMENTAL ARTICLE** 

4. Polishing of zirconia reinforced lithium silicate press ceramics. An in vitro study Elena Vasileva, Angelina Vlahova, Ilian Hristov, Stoyan Yankov, Zlatina Tomova, Zhivko Georgiev Pages 485-488

**EXPERIMENTAL ARTICLE** 

5. Property Test of Phosphate and Hydroxyl Groups from Lates Carcarifer Fish Scale as a Candidate for Synthetic Hydroxyapatite using the Ftir Method Dian Agustin Wahjuningrum, Setyabudi, Anuj Bhardwaj, Syania Edinda Febriyanti, Nadia Liliani Soetjipta, Latief Mooduto Pages 489-493

**EXPERIMENTAL ARTICLE** 

- 6. Effects of Different Prophylaxis Procedures on Titanium Implant Fixture: A Scanning Electron Microscopy Study Zul Fahmi Bahari, Raja Azman Raja Awang, Akram Hassan Pages 494-499
  EXPERIMENTAL ARTICLE
- 7. Pressure absorbability between polymethyl-methacrylate and thermoplastic nylon denture base materials

Elis Crystal, Hubban Nasution, Ika Andryas, Putri Welda Utami Ritonga, Siti Wahyuni, Ricca Chairunnisa, Ariyani Dallmer

Pages 500-504

**EXPERIMENTAL ARTICLE** 

 Evaluation of the Changes Created by Endosteal Implants Installed at Different Lengths, Angles and Diameters on the Maxilla and Mandible Using Three-Dimensional Modeling and Finite Elements Stress Analysis Nedim Güneş, Rezzan Güner

Pages 505-513

**EXPERIMENTAL ARTICLE** 

TÜRKİYE ATIF DİZİNİ

 Mechanical Evaluation of Anadara Granosa Scaffold with Various Gelatin Concentrations for Bone Regeneration Meinar Nur Ashrin, Widyasri Praningrum, Fitria Rahmitasari, Teofilus Timon Lirungan, Ryski Dea Citra Anindita, Rima Parwati Sari

Pages 514-518

copus

JUEST INDEX ( COPERNICUS

IOURNALS MASTER LI

J Int Dent Med Res

Türkiye Klinikleri Academic Keys



#### Journal of International Dental and Medical Research / ISSN: 1309-100X

#### **TABLE OF CONTENTS / 2021; 14 (2)**

**57. HED (Hypohidrotic Ectodermal Dysplasia): A Review** Michele Callea, Francisco Cammarata Scalisi, Izzet Yavuz, Mehmet Sinan Dogan, Colin Eric Willoughby, Murali Dharan Bashyam Pages 785-789

REVIEW

**REVIEW** 

**REVIEW** 

- 58. Academic Stress in Oral Diseases Of Students Gayane Mkhoyan, Ellina Velichko, Olga Bait Said, Dmitriy Serebrov Pages 790-796
- 59. The Effect Vitamin D Supplementation on Severity and Mortality Risk of COVID-19: A Systematic Review Indah Bachti Setyarini, Nurul Ratna Mutu Manikam, Ninik Mudjihartini Pages 797–805

#### MEDICINE

EXPERIMENTAL ARTICLE

60. Stress-Protected Effect of Torasemide in Acute Immobilization Stress in Rats Serhiy M. Bilash, Valentyna P. Bilash, Mykhailo M. Koptev, Natalia M. Deviatkina, Nataliia I. Vynnyk, Borys M. Filenko, Sergii A. Proskurnya, Natalia V. Roiko Pages 806-811

**EXPERIMENTAL ARTICLE** 

61. Mechanism of Centella asiatica extract in Increasing Alveolar Macrophages Apoptosis in Rat Tuberculosis Models Arifa Mustika, Anny Setijo Rahaju, Sri Agus Sudjarwo, Mangestuti agil, Ni Made Mertaniasih Pages 812-819

**EXPERIMENTAL ARTICLE** 

62. The Expression of Visceral Fat Uncoupling Protein-1 is Higher in Moderate-Intensity Swimming than in Low or High-Intensity Swimming in Mice Chabib Fachry Albar, Soebagijo Adi Soelistijo, Muhammad Miftahussurur, Purwo Sri Rejeki Pages 820-824

**CLINICAL ARTICLE** 

63. Combined Method of Treatment of Dentists in the Early Stages of Osteochondrosis of the Cervical Spine

Zhandarov K.A, Ogarev E.V, Chekanov M.S, Vasil'ev Yu.L, Meylanova R.D, Kryuchko P.V, Dydykin S.S Pages 825-834

**CLINICAL ARTICLE** 

64. Biofeedback Exercise and its Relation to Pelvic Floor Muscle Strength: An Experiment at 3 Weeks and 6 Weeks Postpartum Eighty Mardivan Kurniawati, Maria Diah Zakiyah, Rwahita Satyawati, Dharmanta, Gatut Hardianto, Hari

Eighty Mardiyan Kurniawati, Maria Diah Zakiyah, Rwahita Satyawati Dharmanta, Gatut Hardianto, Hari Paraton, Nur Anisah Rahmawati Pages 835-840

**CLINICAL ARTICLE** 

TÜRKİYE ATIF DİZİNİ

Academic Keys

65. The Degree of Pineal Gland Calcification in the Aged People is Associated with Changes in the Internal Structure Ivan I. Starchenko, Ruslana M. Grinko, Anastasiia D. Shkodina, Borys M. Filenko, Nataliia I. Vynnyk, Nataliia

V. Roiko, Serhiy A. Proskurnya, Svitlana M. Sovhyria, Valentyna V. Cherniak, Victoriia I. Babenko Pages 841-844

Türkiye Klinikleri

J Int Dent Med Res

UEST INDEX COPERNICUS

TOURNA

copus

#### Mechanism of Centella asiatica extract in Increasing Alveolar Macrophages Apoptosis in Rat Tuberculosis Models

Arifa Mustika<sup>1</sup>\*, Anny Setijo Rahaju<sup>2</sup>, Sri Agus Sudjarwo<sup>3</sup>, Mangestuti agil<sup>4</sup>, Ni Made Mertaniasih<sup>5</sup>

1. Department of Anatomy, Histology, and Pharmacology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia.

- 2. Department of Anatomical Pathology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia.
- 3. Department of Pharmacology, Faculty of Veterinary Medicine Universitas Airlangga, Surabaya, Indonesia.
- 4. Department of Phytochemistry and Pharmacognosy, Faculty of Pharmacy Universitas Airlangga, Surabaya, Indonesia.
- 5. Department of Microbiology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia.

#### Abstract

Mycobacterium tuberculosis has the capacity to manipulate host defense pathway, particularly the ability to inhibit apoptosis of infected host cells. Centella asiatica is a medicinal plant which is able to reduce growth of Mycobacterium tuberculosis and enhance the cellular immune response and apoptosis in cancer cells. The research is to determine effect and mechanism of the extract of Centella asiatica in enhancing macrophages apoptosis from rat tuberculosis models. Rat tuberculosis models were divided randomly into 4 groups. Group 1, 2, and 3 were given extracts of Centella asiatica at 375mg/kgbw, 750mg/kgbw, and 1500mg/kgbw daily for 14 days. The 4 group is a control group. Apoptosis, expression of Bcl 2, Bax, Caspase 8 were performed by immunohistochemistry.

The study shows reduce expression of Bcl-2, increase expression of Bax and Caspase 8 also enhance the macrophages apoptosis. Ethanol extract of Centella asiatica have active ingredients to enhance of apoptosis alveolar macrophages from rat lung tissue that were infected by Mycobacterium tuberculosis through the increase of Caspase 8's expression, Bax'expression and decreasing of Bcl-2's expression.

Experimental article J Int Dent Med Res 2021; 14(2): 812-819) Keywords: Apoptosis, Centella asiatica, Macrophage, Rat tuberculosis models.

Received date:14 March 2021Accept date:09 May 2021

#### Introduction

Immune response plays a fundamental for outcome of tuberculosis bacterial role infection. It is believed that the failure in treating tuberculosis is because of the failure of patients' immune response <sup>1–4</sup>. In tuberculosis infection, macrophages there is а anti-apoptotic phenomenon which is characterized by an increase of various anti-apoptotic protein such as Bcl-2 protein and a decrease of pro-apoptotic <sup>5–11</sup>. *Mycobacterium* protein namely Bax tuberculosis can multiply in macrophages by various process. Therefore, inhibition process of apoptosis in macrophages by Mycobacterium *tuberculosis* is a virulence factor for bacteria <sup>12–15</sup>.

Mechanism of tuberculosis immune

\***Corresponding author:** Arifa Mustika,

Division of Pharmacology and Therapy, Department of Anatomy, Histology, and Pharmacology, Faculty of Medicine Universitas Airlangga, JI Moestopo 47 Surabaya, Indonesia. E-mail : arifa-m@fk.unair.ac.id pathogenesis as mentioned above become the foundation in conducting a research on a more appropriate method of tuberculosis treatment <sup>16</sup>. New treatment of tuberculosis should also enhance immune response, especially to increase apoptosis in infected macrophages. The advantage of this apoptosis is to increase the elimination of bacteria without any intracellular material that comes out into extracellular so that bacteria cannot spread out of the cell. In addition, this apoptotic process will minimize inflammatory reaction that damage surrounding tissue <sup>17–19</sup>.

Centella asiatica is a plant that is widely available in several countries including Indonesia and used as folk medicine as antimicrobial, antiinflammation and wound healing agent. The biggest compounds of this plant are madecasic acid, Asiatic acid, madecasoside and asiaticoside. Other Ingredients are asiatiquercetin-3-glycoside, kämpferol-3-glycoside<sup>20,21</sup>. Ethanol extract of Centella asiatica inhibits the growth of Mycobacterium tuberculosis in vitro and increases immune response <sup>22-24</sup>. Water extract and ethanol extract of Centella asiatica increase

 $Volume \cdot 14 \cdot Number \cdot 2 \cdot 2021$ 

levels of Tumor Necrosis Factor- $\alpha$  (TNF- $\alpha$ ) in healthy macrophage cell cultures 25. Those extracts also increase cell expression of Cluster of Differentiation (CD) 4, CD8, macrophages and production of cytokines TNF- $\alpha$ , interferon- $\gamma$  (IFN- $\gamma$ ) and inhibit transforming growth factor- $\beta$  (TGF- $\dot{\beta}$ ) in macrophage cells  $2^{2\overline{6},2\overline{7}}$ . Asiatic acid, one of the chemical compounds in Centella asiatica, induces macrophage apoptosis through the increase of cytochrome-c and caspase-3 protein<sup>28</sup>. Numerous studies have shown that extracts of both Centella asiatica and its derivative, Asiatic acid, induce apoptosis in various cancer cells<sup>27–31</sup>. This study aimed to determine the effect of ethanol extract of Centella asiatica on protein expression of Bax, BCI-2, apoptosis Caspase 8. and of alveolar macrophages in rat tuberculosis models.

#### Materials and methods

<u>Preaparation of extract of Centella asiatica</u> Centella asiatica plant was used for this study and it was taken from Balai Materia Medika, Batu, Malang after determination was performed in Purwodadi Botanical Garden, Pasuruan. The extraction was conducted using maceration by soaking Centella asiatica in 70% ethanol.

#### Preparation of Bacteria

Mycobacterium tuberculosis strain H<sub>37</sub> RV (ATCC27294) obtained from tuberculosis of Tropical Disease Institute laboratory Universitas Airlangga was used. The bacterial suspension was made by taking bacterial colonies which were then put into glass tubes, calibrated before being used containing 2 drops of 0.05% tween 80. Afterwards, ± 6 ml aquadest was added and then vortexed. The suspension was compared to Mc Farland standard (1 Mc Farland is equal to 10<sup>8</sup>/ ml *Mycobacterium* tuberculosis <sup>32,33</sup>).

**Chemicals** 

The chemicals used in this study were HCI injection (HameIn ketamine Pharmaceuticals), ether (EMS catalogue no. 1180), absolute alcohol, alcohol 96%, alcohol 80%, alcohol 70% (Brataco), agua destilata (Brataco), carboxy methyl cellulose sodium (Brataco), Kieselgel 60F 254 (Merck, catalog no. 0345629), xylol, n-Hexana, ethylacetate, anisaldehide sulfuric acid, Hematoxylin Eosin staining (EMS catalogue no. 14653), Phosphate Buffered Saline (PBS) (Brataco), formaldehyde

(Brataco), Ziehl-Neelsen staining (EMS catalogue no. 484k), Middlebrook 7H9 and 7H10 medium (BD Biosciens catalogue no. 295964), fungizone (Sigma, catalog no. A2942), Apoptag detection kit (Millipore corporation, catalog no. S1701), primary antibody for Bcl-2 was anti mouse monoclonal antibody Bcl-2 (Bioworld Technology Inc., catalog no. BS1511), primary antibody for Bax which was rabbit Ig G Bax antibody (species reactivity to rat and mouse) (Novus Biologicals, catalog no. NB120 -7977), primary antibody for Caspase-8 was rabbit IgG caspase 8 antibody (Novus Biologicals, catalog no. NB120-15552), and Immunohistochemistry Kit (secondary antibody, streptavidim-HRP, DAB from Daco LSAB catalog No. K0673).

#### Rats Tuberculosis Models

The Experimental animals used were male rats (Rattus norvegicus Wistar strain) at 2-3 months of age with body weight between 125-200 gram obtained from the animal unit of Universitas Gadjah Mada. Rats were kept in BSL cabinet in Faculty of veterinary Medicine Universitas Airlangga and were given ad libitum access to food and water. Ketamine were given for anesthesia to all rats before they were infected with *M.tuberculosis*. 50µl *Mycobacterium* tuberculosis suspension (1 ml contains 10<sup>8</sup> Mycobacterium tuberculosis) was injected into the trachea <sup>7,34,35</sup>. After injection, the rats were kept and given food and drink ad libitum for up to 28 days. On the 29<sup>th</sup> day, the rats were randomly sacrificed and their lungs were taken for culture that they were infected to prove with M.tuberculosis while other rats will be used for rat tuberculosis models.

**Treatment** 

Rat tuberculosis models were divided into 4 group and began to receive ethanol extract of Centella asiatica. Group 1 was given a dose of 375 mg/kgbw of ethanol extract of Centella asiatica, group 2 was given a dose of 750 mg/kgbw of ethanol extract of Centella asiatica, group 3 was given a dose of 1,500 mg/kgbw of ethanol extract of Centella asiatica. The extract was administered orally using sonde once a day for 14 days. Group 4, a control group, was given a suspension of 1% CMC Na in aquadest orally using sonde once a day for 14 days. On the 15<sup>th</sup> day after treatment, rats from both treatment groups and control group were sacrificed. The right lung tissue was put in 10% buffered formalin for histopathological examination using

 $Volume \cdot 14 \cdot Number \cdot 2 \cdot 2021$ 

Immunohistochemistry. The method used for examination of macrophage apoptosis in this study was immunohistochemical staining using *Terminal dUTP Nick-End Staining Labeling* (TUNEL) assay.

#### Data Analysis

Data were assayed for normality and homogeneity of variance. For comparison, ANOVA was used for comparative test, while *Spearman* correlation and path analysis was used for correlation test.


#### Ethical Approval

This study was ethically approved by Ethics Committee of Faculty of Veterinary Medicine, Universitas Airlangga (Ethical Clearance No. 129-KE) before conducting the study.

#### Results

The effect of Centella asiatica ethanol extract on

Bcl-2 protein expression in alveolar macrophage The results showed a significant difference between treatment groups and control group. Highest mean of Bcl-2 protein expression in alveolar macrophage was found in the control group (Table 1) (Figure 1.)




G1= 375mg/kgbw; G2 = 750mg/kgbw; G3 = 1500mg/kgbw; G4 = control group\* p < 0.05

**Table 1.** The average of Expression of Bax, Bcl-2,Cas-8 protein and apoptosis of alveolarmacrophages of rat lung tissue.

<u>The effect of Centella asiatica ethanol</u> <u>extract on Bax protein expression in alveolar</u> <u>macrophage</u>

The results showed a significant difference between treatment groups and control group. Highest mean of Bax protein expression

of alveolar macrophages was found in the second group (Table 1) (Figure 2).



**Figure 1.** Expression of Bcl-2 protein from alveolar macrophages in rat lung tissue. The image above shows the effect of ethanol extract of *Centella asiatica* of 375 mg/kgbw (a), 750 mg/kgbw (b), 1500 mg/kgbw (c), and control group (d) on expression of Bcl-2. Examination was conducted by using immunohistochemistry. Brown cytoplasm shows positive expression of Bcl-2 protein (yellow arrow) whereas colorless cytoplasm shows negative result (red arrow). The amount of Bcl-2 protein expression is the highest in the control group. 400x magnification.




Figure 2. Expression of Bax protein of alveolar macrophages of rat lung tissue. The image

above shows the effect of ethanol extract of *Centella asiatica* at a dose of 375 mg/kgbw (a), 750 mg/kgbw (b), 1500 mg/kgbw (c), and control group (d) on expression of Bax. Examination was conducted by using immunohistochemistry. Brown cytoplasm shows positive expression of Bax protein (yellow arrow) whereas colorless cytoplasm shows negative result (red arrow). The amount of Bax protein expression is the highest in the group that received ethanol extract of *Centella asiatic* of 750 mg/kgbw. 400x magnification.

## The effect of *Centella asiatica* ethanol extract on Caspase-8 expression of alveolar macrophage

The results showed a difference between treatment groups and control group. Highest mean of Caspase-8 protein expression of alveolar macrophages was found in group 2 (Table 1) (Figure 3).




Figure 3. Expression of Caspace-8 protein of alveolar macrophages of rat lung tissue. The image above shows the effect of ethanol extract of Centella asiatica at a dose of 375 mg/kgbw (a), 750 mg/kgbw (b), 1500 mg/kgbw (c), and control group (d) on expression of Caspace-8 protein. Examination conducted was by using immunohistochemistry. Brown cytoplasm shows positive expression of Caspace-8 protein (vellow arrow) whereas colorless cytoplasm shows negative result (red arrow). The amount of Caspace-8 protein expression is the highest in the group that received ethanol extract of Centella 815 siatic at a dose of 750 mg/kgbw. 400x magnification.

The effect of *Centella asiatica* ethanol extract in alveolar macrophage apoptosis

The effect of ethanol extract of *Centella* asiatica on apoptosis of alveolar macrophage in *Mycobacterium tuberculosis* - infected rat lung tissue showed a significant difference between treatment groups and control group. Highest mean of alveolar macrophage apoptosis was found in group 2 (Table 1) (figure 4).




Figure 4. Apoptosis of alveolar macrophage cell of rat lung tissue. The image above shows the effect of ethanol extract of Centella asiatica of 375 mg/kgbw (a), 750 mg/kgbw (b), 1500 mg/kgbw (c), and control group (d) on expression Caspace-8 protein. Examination of was conducted by using Tunnel assay (Apoptag detection kit). Brown nucleus shows positive result (yellow arrow) whereas colorless nucleus shows negative result (red arrow). The amount of apoptosis is the highest in the group receiving ethanol extract of Centella asiatic at a dose of 750 mg/kgbw. 400x magnification.

Pathway analysis was conducted to determine the correlation between ethanol extract of *Centella asiatica* and Bcl 2 protein expression, Bax protein expression, caspase-8 protein expression, apoptosis and the number of bacteria. The results of pathway analysis showed that ethanol extract of *Centella asiatica* significantly decreased Bcl-2 protein expression of alveolar macrophage (p = 0.044), increased Bax protein expression significantly (p = 0.001),

as well the increase of Caspase-8 protein (p = 0.000). the significant increase of Caspase 8 protein expression affected the increase of Bax protein expression (p = 0.000) and the increase of apoptosis of alveolar macrophage (p = 0.000). Ethanol extract of *Centella asiatica* increased the apoptosis of alveolar macrophage (p = 0.000).




Figure 5. Results of Path Analysis.

#### Discussion

The results of this study showed that ethanol extract of *Centella asiatica* decreased the Bcl-2 protein expression in rat tuberculosis models. The ethanol extract of *Centella asiatica* did not linearly decrease Bcl-2 protein expression of alveolar macrophages cell of rat lung tissue with increasing given doses. The ethanol extract of *Centella asiatica* at a dose of 375 mg/kgbw, 750 mg/kgbw and 1500 mg/kgbw showed a significant difference in the decrease of Bcl-2 protein expression of alveolar macrophage when compared to the control group. However, there was no significant difference between treatment groups.

There was an increase of Bcl-2 expression in alveolar macrophage by 65% when compared to treatment group. This finding was consistent with previous study by Adolfo (2006) which stated that Mycobacterium tuberculosis has the ability to increase the expression of Bcl-2 protein in the alveolar macrophages cell. Bcl-2 protein expression in alveolar macrophage increases from 12.5% in the first days after infection to 32% at day 28 after infection <sup>36</sup>. One of the active ingredients of ethanol extract of Centella asiatica is Asiatic acid. This ingredient is believed to has a significant role to decrease Bcl2 protein expression in alveolar macrophage. Bcl-2 is a member of Bcl protein family that has a very important role in apoptosis inhibition of macrophage cell by *Mycobacterium tuberculosis*<sup>6,11,37</sup>. One possible mechanism of *Mycobacterium tuberculosis* to increase Bcl-2 protein expression in alveolar macrophage is through the increase of cytokines interleukin-10 secretion <sup>6</sup>. However, this ability can be inhibited by IFN-y <sup>38,39</sup>.

Bcl-2 protein is a competitive inhibitor against Bax protein receptor, BH3 domain. Bcl-2 protein bonding makes Bax protein unable to bind BH3 domain, which makes Bax protein is not activated and this leads to the inhibition of the apoptotic process <sup>11</sup>. BH3-only protein has a function to receive stimuli originated from the outside cell such as drug or radiation. These stimuli activate BH3-only protein and work directly in releasing bond the between Bax and Bcl-2 protein and decrease Bcl-2 expression <sup>9,10</sup>. In this study, ethanol extract of Centella asiatica at a dose of 375 mg/kgbw has not reached the therapeutic dose required to increase Bax protein expression while at a dose of 1500 mg/kgbw there may have been a down-regulation or desensitization of the receptor. Hence, the effective dose of ethanol extract of Centella asiatica to increase Bax protein expression is 750 mg/kgbw.

Asiatic acid, one of the active ingredient of Centella asiatica ethanol extract, plays an important role in the increase of Bax protein expression in alveolar macrophage. Previous study showed that Asiatic acid increases apoptosis of melanoma cell through increased Bax protein expression <sup>30</sup>. The results of path analysis showed that ethanol extract of Centella asiatica had a strong correlation with a decrease Bcl-2 protein expression of alveolar of macrophage and an increase of Bax protein expression of alveolar macrophage. However, correlation between these two proteins and apoptotic proteins was weak, meaning that the two did not directly increase apoptosis of alveolar macrophages but through other pathways which was not examined in this study(Figure 5).

In normal condition without any stimulation, Bax protein presents in cytosol cell in an inert state. Appropriate stimulus can change protein structure so that Bax protein can perform insertions on the target membrane, which are at the outer membrane of mitochondria and

 $Volume \cdot 14 \cdot Number \cdot 2 \cdot 2021$ 

endoplasmic reticulum. Interactions between Bax protein and any protein mentioned afore occur in BH3 domain and trigger multilevel changes of Bax protein and this phase is an early stage of apoptotic process through intrinsic pathway. BH3-only proteins receive stimuli originated from outside cell such as drug or radiation. These stimuli make BH3-only proteins activated and work directly in releasing the bond between Bax and Bcl-2 protein and decrease Bcl-2 expression. Stimuli from outside such as medicinal materials, irradiation and stress can affect interactions between pro-apoptosis proteins and antiapoptosis proteins <sup>40,41</sup>.

The graphic between Centella asiatica ethanol extract doses and caspase-8 protein expression of alveolar macrophage showed an inverted U-shaped curve with highest peak in dose of 750mg/kgbw. Ethanol extract of Centella asiatica at a dose of 375 mg/kgbw is not reach the therapeutic dose require to increase caspase-8 protein expression whereas downregulation or desensitization of the receptor probably already happens at a dose of 1500 mg/kgbw. Caspase-8, is an active form from procaspase-8. Activation of procaspase-8 into caspase-8 occurs through extrinsic pathway involving Death receptor transduction signal. The *Death receptor* will bind to appropriate ligands to form a death domain which has a significant role in transmitting death signal from cell surface to intracellular. Death domains to have been identified included FasL/FasR, TNF-a/TNFR1, Apo3L/DR3, Apo2L/DR4 and Apo2L / DR5 <sup>40,42,43</sup>.

Activation of procaspase-8 into caspase-8 in this study happened because ethanol extract of Centella asiatica improved the interaction between TNF- $\alpha$  ligand and transmembrane TNFR1 receptors. This finding was in accordance with previous study which stated that ethanol extract of Centella asiatica increases levels of TNF- $\alpha$  <sup>25,27</sup>. The increased levels of TNF- $\alpha$ causes the bond between TNF- $\alpha$  ligand to its receptors, TNFR1, to form death domain in cytoplasm cell. Death domain along with the death effector domains forms an adapter protein that is TNF Receptor-Associated Death Domain (TRADD). Part of the complex, death effector domains, binds to procaspase-8 to form a complex called Death-Inducing Signaling (DISC). DISC complex Complex makes activation of procaspase-8 become caspase-8 autolytical. When caspase-8 is activated, it will

trigger apoptosis of alveolar macrophages<sup>40,42</sup>.

*Mycobacterium tuberculosis* has an ability to increase secretion of TNFR2 proteins that can inactivate the bond between TNF- $\alpha$  and TNFR2 so that TNF- $\alpha$  cannot bind to TNFR1 and form a *death domain* <sup>43</sup>. Increased TNFR2 secretion occurs because of the induction of cytokine IL-10. Therefore, inhibition on IL-10 will decrease TNFR2 secretion. The results of path analysis showed that caspase-8 increased not only apoptosis but also Bax protein expression in intrinsic pathway. We suggest that there is a communication between extrinsic and intrinsic pathways on apoptosis.

There was a significant increase of of alveolar apoptosis macrophage in Mycobacterium tuberculosis - infected rat lung tissue in treatment groups compared to control group. However, the graphic between *Centella* asiatica ethanol extract doses and apoptosis of alveolar macrophage protein expression of alveolar macrophage showed an inverted Ushaped curve with highest peak in dose of 750mg/kg bw. We suggest that ethanol extract of *Centella asiatica* given at a dose of 375 mg/kgbw did not achieve a therapeutic dose required to increase apoptosis of alveolar macrophage whereas receptor desensitization or downregulation probably already happens at a dose of 1500 mg/kgbw.

Each cell will respond to any stress or stimuli received through a variety of ways from activation of life signal to initiation of cell death. The purpose of this process is to maintain homeostasis so cells not needed will be eliminated. The mechanism depends on a variety of exogenous factors and ability of cells to cope with stress or stimulus<sup>44</sup>. In this study, the stimulus was provided in the form of natural ingredient and this material showed an increased in apoptosis of alveolar macrophages. The results of path analysis showed that there was a strong correlation between ethanol extract of Centella asiatica and the increase of caspase-8 protein expression and there was also a strong correlation between the increase of caspase-8 and apoptosis expression of alveolar macrophage. This proved that the mechanism of increased apoptosis by ethanol extract of Centella asiatica was through the extrinsic pathway.

*Mycobacterium tuberculosis* death mechanism in apoptotic process is through an

increase in Nitric Oxide (NO) and through the phagosome fusion process of into phagolysosome <sup>39,45</sup>. NO can kill intracellular Mycobacterium tuberculosis through a variety of mechanisms namely directly killing DNA of bacteria, affecting copper metabolism, virulence targets of microbacterial or serving as a second messenger. In addition, Nitric Oxide (NO) is known to have the ability to induce apoptosis in macrophages and NO production was induced by IFN-y through the increase of expression of Nitric Oxide Synthetase 2 (NOS2) <sup>46</sup>. Research conducted by Herbs in 2011 shows that NO induced bv IFN-γ increases macrophage apoptosis infected Mycobacterium with tuberculosis and kills the bacteria. Previous study showed that ethanol extract of Centella asiatica increased IFN-y production so that ethanol extract of Centella asiatica which, in turn, also increases NO production is alleged to cause death mechanism of intracellular Mycobacterium *tuberculosis*<sup>46</sup>. Apoptotic process is suspected to have links with architectural change of host system causing fusion between vesicular phagosome and lysosome previously inhibited by Mycobacterium tuberculosis<sup>47</sup>.

#### Conclusions

The present research clearly demonstrated that the ethanol extract of *Centella asiatica* have active ingredients to enhance of apoptosis alveolar macrophages from rat lung tissue that were infected by *Mycobacterium tuberculosis* through increasing of Caspase 8's expression, Bax'expression and decreasing of Bcl-2's expression.

#### Acknowledgements

We thank the Directorate of Research and Community Services, the Directorate General of Higher Education, and Ministry of Education and Culture in Indonesia for the financial support.

#### **Declaration of Interest**

The authors report no conflict of interest.

#### References

1. Ellner JJ. Immunoregulation in TB: Observations and implications. Clin Transl Sci. 2010;3(1):23–8.

- Ahmad S. Pathogenesis, immunology, and diagnosis of latent mycobacterium tuberculosis infection. Clin Dev Immunol. 2011;11(1);1-18.
- de Martino M, Lodi L, Galli L, Chiappini E. Immune Response to Mycobacterium tuberculosis: A Narrative Review. Front Pediatr. 2019;7:1–8.
- Sia JK, Rengarajan J. Immunology of Mycobacterium tuberculosis Infections . Gram-Positive Pathog. 2019;7(4):1056–86.
- Mustafa T, Bjune G, Jonsson R, Hernandez Pando R, Nilsen R. Increased expression of Fas ligand in human tuberculosis and leprosy lesions: A potential novel mechanism of immune evasion in mycobacterial infection. Scand J Immunol. 2001;54(6):630–9.
- Mogga SJ, Mustafa T, Sviland L, Nilsen R. Increased Bcl-2 and reduced Bax expression in infected macrophages in slowly progressive primary murine Mycobacterium tuberculosis infection. Scand J Immunol. 2002;56(4):383–91.
- Rodrigues MF, Barsante MM, Alves CCS, Souza MA, Ferreira AP, Amarante-Mendes GP, et al. Apoptosis of macrophages during pulmonary Mycobacterium bovis infection: Correlation with intracellular bacillary load and cytokine levels. Immunology. 2009;128(1):691–9.
- Chen L, Shi W, Li H, Sun X, Fan X, LeSage G, et al. Critical role of toll-like receptor 9 in morphine and Mycobacterium tuberculosis-induced apoptosis in mice. PLoS One. 2010;5(2):3–9.
- Aguiló N, Uranga S, Marinova D, Martín C, Pardo J. Bim is a crucial regulator of apoptosis induced by Mycobacterium tuberculosis. Cell Death Dis. 2014;5(7):1–10.
- Agarwal RG, Sharma P, Nyati KK. microRNAs in mycobacterial infection: Modulation of host immune response and apoptotic pathways. Immune Netw. 2019;19(5):1–22.
- Abdalla AE, Ejaz H, Mahjoob MO, Alameen AAM, Abosalif KOA, Elamir MYM, et al. Intelligent mechanisms of macrophage apoptosis subversion by Mycobacterium. Pathogens. 2020;9(3):1–11. The results of this study showed th
- Zhang J, Jiang R, Takayama H, Tanaka Y. Survival of virulent Mycobacterium tuberculosis involves preventing apoptosis induced by Bcl-2 upregulation and release resulting from necrosis in J774 macrophages. Microbiol Immunol. 2005;49(9):845–52.
- Behar SM, Divangahi M, Remold HG. Evasion of innate immunity by mycobacterium tuberculosis: Is death an exit strategy? Nat Rev Microbiol. 2010;8(9):668–74.
- Divangahi M, Desjardins D, Nunes-Alves C, Remold HG, Behar SM. Eicosanoid pathways regulate adaptive immunity to Mycobacterium tuberculosis. Nat Immunol. 2010;11(8):751–8.
- Miller JL, Velmurugan K, Cowan MJ, Briken V. The type I NADH dehydrogenase of Mycobacterium tuberculosis counters phagosomal NOX2 activity to inhibit TNF-α-mediated host cell apoptosis. PLoS Pathog. 2010;6(4):1–14.
- Lam A, Prabhu R, Gross CM, Riesenberg LA, Singh V, Aggarwal S. Role of apoptosis and autophagy in tuberculosis. Am J Physiol - Lung Cell Mol Physiol. 2017;313(2):L218–29.
- 17. Yang S, Li F, Jia Š, Zhang K, Jiang W, Shang Y, et al. Early secreted antigen ESAT-6 of mycobacterium tuberculosis promotes apoptosis of macrophages via targeting the microrna155-SOCS1 interaction. Cell Physiol Biochem. 2015;35(4):1276–88.
- Yao K, Chen Q, Wu Y, Liu F, Chen X, Zhang Y. Unphosphorylated STAT1 represses apoptosis in macrophages during Mycobacterium tuberculosis infection. J Cell Sci. 2017;130(10):1740–51.
- Schaaf K, Smith SR, Duverger A, Wagner F, Wolschendorf F, Westfall AO, et al. Mycobacterium tuberculosis exploits the PPM1A signaling pathway to block host macrophage apoptosis. Sci Rep. 2017;7:1–16.
- Gray NE, Alcazar Magana A, Lak P, Wright KM, Quinn J, Stevens JF, et al. Centella asiatica: phytochemistry and mechanisms of neuroprotection and cognitive enhancement. Phytochem Rev. 2018;17(1):161–94.

- Irham WH, Tamrin, Marpaung L, Marpongahtun. Bioactive Compounds in Pegagan Leaf (Centella asiatica L. Urban) for Wound Healing. J Phys Conf Ser. 2019;1232(1):5–10.
- Salmiwanti S, Ilyas A, Saleh A. Isolasi Senyawa Metabolit Sekunder Fraksi N- Heksana Dari Daun Pegagan (Centellaasiatica L.) dan Uji Antibakteri Terhadap Mycobacterium tuberculosis. Al-Kimia. 2016;4(2):52–63.
- Suresh M, Rath PK, Panneerselvam A, Dhanasekaran D, Thajuddin N. Anti-mycobacterial effect of leaf extract of Centella asiatica (Mackinlayaceae). Res J Pharm Technol.2010;3:872–6.
- 24. Delbo M, Calapai G. Assessment report on Centella asiatica (L.) Urban, herba. Eur Med Agency. 2010;44:1–44.
- Punturee K, Wild CP, Vinitketkumneun U. Thai medicinal plants modulate nitric oxide and tumor necrosis factor-α in J774.2 mouse macrophages. J Ethnopharmacol. 2004;95(2–3):183–9.
- Lyu SY, Park WB. Production of cytokine and NO by RAW 264.7 macrophages and PBMC in vitro incubation with flavonoids. Arch Pharm Res. 2005;28(5):573–81.
- 27. Sun B, Wu L, Wu Y, Zhang C, Qin L, Hayashi M, et al. Therapeutic Potential of Centella asiatica and Its Triterpenes: A Review. Front Pharmacol. 2020;11:1–24.
- Cho MK, Sung MA, Kim DS, Park HG, Jew SS, Kim SG. 2-Oxo-3,23-isopropylidene-asiatate (AS2006A), a wound-healing asiatate derivative, exerts anti-inflammatory effect by apoptosis of macrophages. Int Immunopharmacol. 2003;3(10–11):1429– 37.
- Lee YS, Jin DQ, Kwon EJ, Park SH, Lee ES, Jeong TC, et al. Asiatic acid, a triterpene, induces apoptosis through intracellular Ca2+ release and enhanced expression of p53 in HepG2 human hepatoma cells. Cancer Lett. 2002;186(1):83–91.
- Park BC, Bosire KO, Lee ES, Lee YS, Kim JA. Asiatic acid induces apoptosis in SK-MEL-2 human melanoma cells. Cancer Lett. 2005;218(1):81–90.
- Tang XL, Yang XY, Jung HJ, Kim SY, Jung SY, Choi DY, et al. Asiatic acid induces colon cancer cell growth inhibition and apoptosis through mitochondrial death cascade. Biol Pharm Bull. 2009;32(8):1399–405.
- Saunders BM, Cheers C. Intranasal infection of beige mice with Mycobacterium avium complex: Role of neutrophils and natural killer cells. Infect Immun. 1996;64(10):4236–41.
- 33. Sato K, Akaki T, Tomioka H. Differential potentiation of antimycobacterial activity and reactive nitrogen intermediateproducing ability of murine peritoneal macrophages activated by interferon-gamma (IFN-γ) and tumour necrosis factor-alpha (TNF- α). Clin Exp Immunol. 1998;112(1):63–8.
- Chandra G, Selvaraj P, Jawahar MS, Banurekha V V., Narayanan PR. Effect of vitamin D3 on phagocytic potential of macrophages with live Mycobacterium tuberculosis and lymphoproliferative response in pulmonary tuberculosis. J Clin Immunol. 2004;24(3):249–57.
- 35. Mustika A, Rahaju AS, Indrawati R. Penurunan Kerusakan Jaringan Paru Terinfeksi Tuberkulosis oleh Ekstrak Pegagan Melalui Peningkatan Ekspresi Tissue Inhibitor of Matrix Metalloproteinase-1. J Vet. 2015;15(4):530–40.
- Ríos-Barrera VA, Campos-Peña V, Águilar-León D, Lascurain Ledesma R, Meraz-Ríos MA, Moreno J, et al. Macrophage and T lymphocyte apoptosis during experimental pulmonary tuberculosis: Their relationship to mycobacterial virulence. Eur J Immunol. 2006;36(2):345–53.
- Chen YJ, Chao KSC, Yang YC, Hsu ML, Lin CP, Chen YY. Zoledronic acid, an aminobisphosphonate, modulates differentiation and maturation of human dendritic cells. Immunopharmacol Immunotoxicol. 2009;31(3):499–508.
- Kornfeld H, Mancino G, Colizzi V. The role of macrophage cell death in tuberculosis. Cell Death Differ. 1999;6(1):71–8.
- Herbst S, Schaible UE, Schneider BE. Interferon gamma activated macrophages kill mycobacteria by nitric oxide induced apoptosis. PLoS One. 2011;6(5):e19105
- 40. Galluzzi L, Vitale I, Aaronson SA, Abrams JM, Adam D, Agostinis P, et al. Molecular mechanisms of cell death: Recommendations of the Nomenclature Committee on Cell Death 2018. Cell Death Differ. 2018;25(3):486–541.

- Cairra<sup>~</sup>o, Fa<sup>'</sup>tima; and Domingos, Pedro M. Apoptosis: Molecular Mechanisms. Encyclopedia of Life Sciences (ELS). 2010;1;1-8.
- 42. Elmore S. Apoptosis: A Review of Programmed Cell Death. Toxicol Pathol. 2007;35(4):495–516.
- Lam A, Prabhu R, Gross CM, Riesenberg LA, Singh V, Aggarwal S. Role of apoptosis and autophagy in tuberculosis. Am J Physiol - Lung Cell Mol Physiol. 2017;313(2):L218–29.
- Samali A, Fulda S, Gorman AM, Hori O, Srinivasula SM. Editorial: Cell stress and cell death. Int J Cell Biol. 2010;2010:2–3.
- 45. Jamwal S V., Mehrotra P, Singh A, Siddiqui Z, Basu A, Rao KVS. Mycobacterial escape from macrophage phagosomes to the cytoplasm represents an alternate adaptation mechanism. Sci Rep. 2016;6:1–9.
- 46. Jamaati H, Mortaz E, Pajouhi Z, Folkerts G, Movassaghi M, Moloudizargari M, et al. Nitric oxide in the pathogenesis and treatment of tuberculosis. Front Microbiol. 2017;7:1–11.
- 47. Queval CJ, Brosch R, Simeone R. The macrophage: A disputed fortress in the battle against Mycobacterium tuberculosis. Front Microbiol. 2017;8:1–11.



### Source details

Journal of International Dental and Medical Research	CiteScore 2021 <b>1.1</b>	(j
Scopus coverage years: from 2009 to Present		
Publisher: Ektodermal Displazi Grubu		
ISSN: 1309-100X	,	(j
Subject area: (Dentistry: General Dentistry)	0.212	
Source type: Journal		
	SNIP 2021	(i)
View all documents > Set document alert Save to source list Source Homepage	0.400	
Subject area: (Dentistry: General Dentistry)	SJR 2021 0.212 SNIP 2021 0.460	

CiteScore CiteScore rank & trend Scopus content coverage

 $\sim$ 

i Improved CiteScore methodology
 CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data
 papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. Learn more >

# **1.1** = 1,171 Citations 2018 - 2021 1,039 Documents 2018 - 2021

CiteScoreTracker 2022 <sup>①</sup>

754 Citations to date 839 Documents to date 0.9 = Last updated on 05 May, 2022 • Updated monthly

Calculated on 05 May, 2022

CiteScore 2021


#### CiteScore rank 2021 ①

Category	Rank	Percentile
Dentistry		
General Dentistry	#79/109	27t

View CiteScore methodology  $\succ$  CiteScore FAQ  $\succ$  Add CiteScore to your site  ${\cal S}^{\!\!\!O}$ 

Q

#### Journal of International Dental and Medical Research 8





KOMISI ETIK PENELITIAN FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA Animal Care and Use Committee (ACUC)

#### **KETERANGAN KELAIKAN ETIK**

"ETHICAL CLEARENCE"

#### No: 129-KE

#### KOMISI ETIK PENELITIAN (ANIMAL CARE AND USE COMMITTEE) FAKULTAS KEDOKTERAN HEWAN UNIVERSITAS AIRLANGGA SURABAYA, TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA :

PENELITIAN BERJUDUL : Mekanisme Peningkatan Apoptosis Makrofag Terinfeksi Mycobacterium tuberculosis pada Jaringan Paru Tikus Dengan Infeksi Mycobacterium tuberculosis oleh Pemberian Ekstrak Etanol Centella asiatica (Pegagan) Melalui Peningkatan Protein Bcl-2, Bax, Caspase-8

PENELITI UTAMA

UNIT/LEMBAGA/TEMPAT : Program Pasca Sarjana PENELITIAN Universitas Airlangga

DINYATAKAN

: LAIK ETIK

: Arifa Mustika

Mengetahui, Dekan FKH-Unair,

Prof. Romzlah Sidik, Ph.D.,drh. NIP. 130687305 Surabaya, 13 Mei 2011

Ketua

Dr. E. Bimo Aksono, M.Kes., Drh. NIP. 132014464