

Variation of histopathological features in colonic mucosal biopsy with clinical diagnosis of suspicious inflammatory bowel disease in Dr. Soetomo General Academic Hospital, Surabaya periode 2015-2019

by Ariadna Anggi Pasang

Submission date: 21-Nov-2022 05:31PM (UTC+0800)

Submission ID: 1960196181

File name: oetomo_General_Academic_Hospital,_Surabaya_periode_2015-2019.pdf (528.8K)

Word count: 3777

Character count: 21919

How to Cite:

Pasang, A. A., Rahniayu, A., & Rahaju, A. S. (2022). Variation of histopathological features in colonic mucosal biopsy with clinical diagnosis of suspicious inflammatory bowel disease in Dr. Soetomo General Academic Hospital, Surabaya period 2015 - 2019. *International Journal of Health Sciences*, 6(S5), 808–818. <https://doi.org/10.53730/ijhs.v6nS5.8759>

Variation of histopathological features in colonic mucosal biopsy with clinical diagnosis of suspicious inflammatory bowel disease in Dr. Soetomo General Academic Hospital, Surabaya period 2015 - 2019

Ariadna Anggi Pasang

Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

Alphania Rahniayu

Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

Corresponding Author E-mail: alphania-r@fk.unair.ac.id

Anny Setijo Rahaju

Department of Anatomical Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

Abstract---Inflammatory bowel diseases (IBDs) are chronic inflammatory diseases that often relapse and divided into two types, Crohn's disease (CD) and ulcerative colitis (UC). Histopathological findings in colonic mucosal biopsy with clinical diagnosis suspicious IBD can vary and overlap. Therefore, criteria and guidelines have been created to improve the diagnostic accuracy. This descriptive observational study was performed retrospectively with cross sectional approach. 122 samples of colonic mucosal biopsies with clinical diagnosis of suspicious IBD were retrieved from histopathological archives in the Anatomical Pathology Laboratorium of Dr. Soetomo Hospital, Surabaya during period 1st January 2015 - 31st December 2019. The most common histopathological feature found in colonic mucosal biopsies with clinical diagnosis of suspicious IBD was crypt distortion (97/79.50% samples), and the least was irregularity of surface epithelium (30/24.59% samples). 10 of 122 samples was concordant with the final diagnosis of IBD. Knowledges regarding the variations of histopathological features in colonic mucosal biopsy specimens with clinical diagnosis of suspicious IBD, can improve the diagnostic accuracy.

Keywords—IBD, colonic biopsy histopathological, suspicious inflammatory.

Introduction

Colitis is an inflammation in colonic mucosa that can occur either acutely or chronically, which caused by infection, inflammation, ischemic, drugs, or other idiopathic diseases (Nielson & Seidelin, 2012; Puspitarini & Prijambodo, 2022). Inflammatory bowel diseases are the most common of chronic colitis that often relapse; can involve any segment of gastrointestinal tract and divided into two types, CD and UC (Villanacci et al., 2021; Fatimah et al., 2021). The incidence of IBD in Asia has increased significantly in the last two decades, ranged between 4.2 and 3.1 per 100,000 population (Ng et al., 2016; Simadibrata & Adiwinata, 2017).

The criteria and guidelines have been created to improve the diagnostic accuracy of IBD when the colonic biopsy was used as the initial surveillance in patients with the chief complaint of diarrhea. Histopathological diagnosis of IBD established based on combination of microscopic findings and clinical history which include patient's age, symptoms, duration of symptoms, and colonoscopy results (Lang-Schwarz et al., 2021).

Chronic colitis, in this case is IBD, determined histopatologically from presence of chronicity features, as follows: crypt distortion, crypt atrophy, chronic inflammation, basal lymphoplasmacytosis, granulomas, and paneth cell metaplasia (Feakins, 2013). The histopathological features in biopsy samples can vary and overlap. Therefore, the completes of clinical information and examination history are important to make the correct diagnosis and plan the effective therapeutic options (Villanacci et al., 2021; Kalishah et al., 2022).

Crohn's disease and ulcerative colitis differ from each other in aspects of epidemiology, clinical presentation, endoscopy, and histopathology features as well as the complication and management. The correct pathological approach and adequate biopsy samples can also contribute to classify the CD and UC (Villanacci et al., 2021).

At this time, studies about variation of histopathological features in colonic mucosal biopsy with clinical diagnosis of IBD are still limited. This study describes variation of histopathological features in colonic mucosal biopsy specimen with clinical diagnosis of IBD.

Materials and Methods

This research descriptive observational study had been performed with a cross sectional approach, using histopathological archives from colonic mucosal biopsy specimen with clinical diagnosis suspicious of IBD in the Anatomical Pathology Laboratory of Dr. Soetomo Hospital Surabaya during the period from 1st January 2015 - 31st December 2019, which had endoscopy results and had not received the previous IBD therapies. The slides were reviewed by two pathologists to

evaluate histopathological features of IBD (basal lymphoplasmacytosis, crypt distortion, transmucosal lymphoplasmacytic, crypt atrophy, mucin depletion, irregularity of surface epithelium), and presented as frequency distribution in table. This study had been approved by the Committee of Health Research Ethic at Dr. Soetomo General Academic Hospital, Surabaya, Indonesia (0915/LOE/301.4.2/V/2021).

Results and Discussions

There were 122 samples of colonic mucosal biopsy with clinical diagnosis of suspicious IBD. Most samples in this study were aged between 51 – 60 years with mean age was 43.23 years. Most samples were male (76 samples, 62.29%).

Based on the slide review, we found the variation of histopatological features in colonic mucosal biopsy, as follows: crypt distortion (97/122 samples, 79.50%), lymphoid aggregates (87/122 samples, 71.31%), transmucosal lymphoplasmacytic (45/122 samples, 36.88%), crypt atrophy (37/122 samples, 30.32%), mucin depletion (36 samples, 29.50%), basal lymphoplasmacytosis (32/122 samples, 26.22%), and irregularity of mucosal surface epithelium (30/122 samples, 24.59%) (Table 1). The variation of histopathological features in colonic mucosa biopsy with clinical diagnosis of IBD, is depicted in Figure 1. We did not find granuloma feature in this study samples. We also could not assess the Paneth cells metaplasia in this study due to lack of information about where the location of biopsy samples was taken, whether from the left or right side of colon.

There were 2/122 samples (1.63%) each for amebic colitis, eosinophilic colitis, and amyloidosis (Figure 2A, B, C). Congo-red stain was performed for definitive diagnosis of amyloidosis (Figure 2D). There was 1/122 sample (0.81%) with histopathological features of chronic colitis with mild dysplasia (Figure 2E). There were 13/122 samples (10.65%) with interpretation of non-specific colitis because from the microscopic features, only obtained the lymphoid aggregates and erosion of the surface epithelium, but there are no chronicity features. There were 7/122 samples (5.73%) with non-diagnostic because from the microscopic feature, there was no muscularis mucosae.

Based on the medical record from Internal Medicine Department, there were 10/122 (8.19%) samples that had final diagnosis with IBD, 9 samples with UC and 1 sample with CD. All the samples showed the chronicity features that supported to IBD diagnosis (Table 2).

Table 1
Variation of histopathological features in colonic mucosa biopsy with clinical diagnosis of IBD

Variation of histopathological features	Number of cases from 122 samples	Percentages
Crypt distortion	97	79.50%
Lymphoid aggregates	87	71.31%
Transmucosal lymphoplasmacytic	45	36.88%

Crypt atrophy	37	30.32%
Mucin depletion	36	29.50%
Basal lymphoplasmacytosis	32	26.22%
Irregularity of surface epithelium	30	24.59%




Figure 1. Variation of histopathological features in colonic mucosa biopsy with clinical diagnosis of IBD. (A) Crypt distortion (HE, 200x). (B) Lymphoid aggregates (HE, 100x). (C) Transmucosal lymphoplasmacytic (HE, 100x). (D) Crypt atrophy (HE, 100x). (E) Mucin depletion (HE, 100x). (F) Basal lymphoplasmacytosis (HE, 200x). (G) Irregularity of surface epithelium (arrow) (HE, 400x).




Figure 2. (A) Amebic colitis (HE, 400x). (B) Eosinophilic colitis (HE, 400x). (C) Amyloidosis (HE, 200x). (D) Congo-red is positive in amyloid deposits (Congo-red, 200x). (E) Chronic colitis with mild dysplasia (HE, 400x).

Table 2
Final diagnosis vs morphology finding parameter

Final diagnosis	Morphology finding parameter						
	Trans mucosal lympho plasmacytic	Crypt distortion	Crypt atrophy	Basal lympho plasmacytosis	Irregularity of surface epithelium	Lymphoid aggregate	Mucin depletion
5 Ulcerative colitis	✓	✓	✓	✓	✓	✓	✓
Ulcerative colitis	✓	✓	✓		✓	✓	

Ulcerative colitis	✓	✓	✓	✓	✓	✓	✓
Ulcerative colitis	✓	✓	✓	✓	✓	✓	✓
Ulcerative colitis	✓	✓	✓	✓	✓	✓	✓
Ulcerative colitis	✓	✓	✓	✓	✓	✓	✓
Ulcerative colitis	✓	✓	✓	✓	✓	✓	✓
Ulcerative colitis	✓	✓	✓	✓	✓	✓	✓
Ulcerative colitis	✓	✓	✓	✓	✓	✓	✓
Crohn's disease	✓	✓	✓	✓	✓	✓	✓

Most cases were 51 – 60 years old, with male predominance (76/62.29% samples). Ulcerative colitis tends to occur in adolescent and young adult, although there is a second incidence peak among middle-aged men. In CD, most patients are diagnosed in second to fourth decades of life, with incidence peak between 20 and 30 years (Odze & Goldblum, 2015). Ulcerative colitis in Asia is usually found between ages of 35 – 45 years, and CD is diagnosed at a younger age than UC (Ng, 2014). However, about 10 -15% of IBD cases can be found at the age of > 60 years. The incidence peak of a bimodal age for IBD, occurs between ages of 50 – 70 years (Val, 2011). The incidence of CD in Asia is generally more common in male, and gender distribution in UC is equal. The lifestyle factors, such as smoking, a high-fat and low-fiber dietary and environmental sanitation problems, are the several factors that can increase the risk of IBD (Alatab et al., 2020).

Study from Gajendran et al. in 2019 reported that IBD diagnosis shows 2 or 3 from 4 microscopic changes, such as crypt distortion, decreased crypt density, irregularity mucosal surface, and diffuse transmucosal inflammation (Gajendran et al., 2019). A strong indicator for IBD diagnosis is focal or diffuse of basal lymphoplasmacytosis (Geboes & Van Eyken, 2009). Basal lymphoplasmacytosis is one of the earliest of histopathological features of IBD and has a high predictive value compared with other non-IBD colitides (Canavese et al., 2017). Crypt distortion was the most common histopathological finding in this study (97/79.50% samples). Study by Surawicz and Belic reported that crypt distortion is also one of the characteristics of IBD besides basal lymphoplasmacytosis (Surawicz & Belic, 1984). Various studies showed distorted architectural crypt in UC could occur ranging from 57% to 100% cases (Dhakhwa et al., 2016).

Lymphoid aggregates can be one of the histology indicators for IBD, especially CD (Villanacci et al., 2020; Avisiena et al., 2019). Some of studies revealed that lymphoid aggregates also can be seen in normal colon (Assarzaghan et al., 2017). No granuloma was found in all study samples. Study by Turner et al. showed that only 9% from 10,000 patients with CD had non-necrotizing epithelioid granuloma appearance. Colonic biopsy specimens tend to involve only the mucosa, occasionally part of the muscularis mucosa, but rarely from the underlying submucosa. Therefore, biopsy specimen-based would almost never find granulomas located below the muscularis mucosa (Turner et al., 2014). The metaplasia of Paneth cells could not be assessed in this study because there was no information about where the location of biopsy samples was taken. Paneth cells are normal component located at the base of the crypt in small intestine and proximal colonic mucosa. The Paneth cells in the distal colon are a metaplastic adaptation to chronic mucosa damage (Feakins, 2013).

Diagnostic uncertainty is commonly found for IBD or to distinguish UC and CD. This is caused by many factors, such as therapy, disease severity, and experience of clinicians and pathologists in patient management, that can disrupt the correct diagnosis. Pathologists often did not get the sufficient information about patient's clinical and endoscopy result. The final diagnosis can only be made if the overall supporting clinical information is obtained (Jenkins et al., 1997; Farmer et al., 2000).

The adequate of taking biopsy samples from endoscopic sampling is also can increase the correct diagnosis of IBD. Optimal sampling for newly diagnosed IBD needs at least two biopsy samples in the terminal ileum and in each segment of the colonic sites. The specimens are separated according to the sampling location and provides information, as follows: terminal ileum (bottle 1), ascending and transverse colon (bottle 2), descending and sigmoid colon (bottle 3), and rectum (bottle 4) (Villanacci et al., 2021).

The term of non-specific colitis refers to inflammation in colon that microscopically does not have characteristic features of any specific form of colitis and usually found in pathology report from colonoscopy biopsy. This term is used for colonic inflammation that does not have specific pathological features from common cause of colitis (Emara et al., 2019). In this study, the samples with the lymphoid aggregate features and erosion on the surface epithelium were included into the non-specific colitis interpretation.

There were 2 samples each for amoebiasis, eosinophilic colitis, and amyloidosis. Chronic inflammation and crypt architectural changes were also found in amoebiasis cases in this study. Study by Singh et al. revealed that architectural changes and chronic inflammation could also be found in amebic colitis as in IBD, but the architectural changes and inflammation in amebic colitis were only mild or moderate (Singh et al., 2015). Diagnostic histopathologically of eosinophilic colitis is determined by the existence of eosinophil clusters in lamina propria (> 60 eosinophils/ 10 HPFs), eosinophilic cryptitis, crypt abscess, and crypt distortion can be found as well (Bates, 2012; Walker et al., 2019). Amyloidosis is a disease that has the similar pathogenesis with other unrelated diseases, wherein all the insoluble fibrillar proteins converge in the extracellular tissue of various organs and cause the organ dysfunction. Amyloidosis consists of primary and secondary amyloidosis. The secondary amyloidosis is caused by extracellular deposition of fibrils from serum amyloid A (SAA) protein. The secondary amyloidosis can also occur in IBD and shows the uncontrolled and persistent of inflammation activity. Amyloidosis occurs approximately 0.9% in CD and 0.7% in UC. The reason why CD is more prone to have complication of amyloidosis than UC is not known but may be related to degree of systemic inflammation that is higher in CD than UC, especially in association with abscesses and fistulae (Sattianayagam et al., 2009). The serum concentrations in acute phase proteins, C-reactive protein dan SAA protein, are also higher in CD than UC (Saverymuttu et al., 1986). In this study, 1 case of amyloidosis was found in patient with a history of SLE.

The histopathological features of chronic colitis with mild dysplasia also obtained in this study. Patients with long-standing IBD have an increased risk for

colorectal cancer. Dysplasia is the best marker for increased risk of malignancy. The identification and grading of dysplasia are the basis for the management of IBD (Villanacci et al., 2021).

Non-diagnostic interpretation was found in 7 samples because from the microscopic feature there was no muscularis mucosae. Ideal mucosa samples that are obtained from colonoscopy consist of mucosa and slight of superficial submucosa (Montgomery et al., 2012). Optimal assessment can be obtained if the biopsy specimens contain of all the full thickness of mucosa and muscularis mucosae (Jenkins et al., 1997). If there is no muscularis mucosae, we also cannot assess the basal lymphoplasmacytosis because it is located between the base of crypt and muscularis mucosae, and basal lymphoplasmacytosis is a strong indicator for IBD diagnosis (Geboes & Van Eyken, 2009).

There were 10/122 samples that had final diagnosis with IBD, 9 samples with UC and 1 sample with CD. The final diagnosis we got from medical record data of Internal Medicine Department. We have performed re-evaluations for all samples that met the inclusion criteria. Samples with final diagnosis IBD (UC or CD) showed the morphology with chronicity features that also supported to IBD diagnosis. Histopathological diagnosis is not the golden-standard modality for diagnosis IBD. The diagnosis is based on the combination of clinical symptoms, endoscopy, laboratory tests, and histopathology. After receiving the pathology report that suspicious of IBD, clinicians should correlate the report with patient clinical symptoms and all the supporting examinations, such as laboratory and endoscopy for the final diagnosis. The diagnostic procedure that involves a multidisciplinary approach, aims to rule out any differential diagnoses, such as infection or any other types of colitis (Kellermann & Riis, 2021).

Conclusion

Based on the review of colonic mucosal biopsy specimens with clinical diagnosis of suspicious IBD from 2015 until 2019, the most patients were male, and the largest age group was at the age group of 51 - 60 years. Knowledges regarding the variations of histopathological features in colonic mucosal biopsy specimens with clinical diagnosis of suspicious IBD, can improve the diagnostic accuracy and start an appropriate treatment.

Acknowledgements

The authors would like to thank to Gastroentero-Hepatology Division, Department of Internal Medicine of Dr. Soetomo General Hospital Surabaya, Indonesia, for endoscopy data contribution for this study.

References

- Alatab, S., Sepanlou, S., Ikuta, K., Homayoon, V., Bisignano, C., & Safiri, S. (2020). The global, regional, and national burden of inflammatory bowel disease in 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *The Lancet Gastroenterology and Hepatology*, 5(1), 17–30.

- Assarzagdegan, N., Montgomery, E., & Pezhouh, M. K. (2017). Colitides: diagnostic challenges and a pattern based approach to differential diagnosis. *Diagnostic Histopathology*, 23(12), 536–543.
- Avisiena A, Nusi IA, Maimunah U, Rahaju AS, Setiawan PB, Purbayu H, Widodo B, Miftahussurur M, Vidyani A, Thamrin H. (2019) Diagnostic values of helicobacter pylori stool antigen immunochromatographic method compared to histopathology in dyspepsia patient. *The New Armenian Medical Journal*, 13(1), 13-19.
- Bates, A. W. H. (2012). Diagnosing eosinophilic colitis: Histopathological pattern or nosological entity? *Scientifica*, 12, 1–9.
- Canavese, G., Villanacci, V., Antonelli, E., Cadei, M., Sapino, A., Rocca, R., Daperno, M., Suriani, R., Di Santo, M. G., Cassoni, P., Bernardini, N., & Bassotti, G. (2017). Eosinophilia – associated basal plasmacytosis: An early and sensitive histologic feature of inflammatory bowel disease. *Apmis*, 125(3), 179–183.
- Dhakhwa, R., Shrestha, H., & Acharya, I. (2016). Histopathological evaluation of ulcerative colitis in colonoscopic biopsies. *Journal of Pathology of Nepal*, 6(11), 932–936.
- Emara, M. H., Salama, R. I., Hamed, E. F., Shoriet, H. N., & Abdel-Aziz, H. R. (2019). Non-specific colitis among patients with colitis: Frequency and relation to inflammatory bowel disease, a prospective study. *Journal of Coloproctology. Sociedade Brasileira de Coloproctologia*, 39(4), 319–325.
- Farmer, M., Petras, R. E., Hunt, L. E., Janosky, J. E., & Galandiuk, S. (2000). The importance of diagnostic accuracy in colonic inflammatory bowel disease. *American Journal of Gastroenterology*, 95(11), 3184–3188.
- Fatimah S, Rahaju AS, Rahniayu A. (2021) Role of Claudin-4 and Matrix Metalloproteinase-2 in Tumor Invasion of Colorectal Adenocarcinoma. *Research Journal of Pharmacy and Technology*, 14(9), 4795.
- Feakins, R. M. (2013). Inflammatory bowel disease biopsies: Updated British Society of Gastroenterology reporting guidelines. *Journal of Clinical Pathology*, 66(12), 1005–1026.
- Gajendran, M., Loganathan, P., Jimenez, G., Catinella, A. P., Ng, N., Umopathy, C., Ziade, N., & Hashash, J. G. (2019). A comprehensive review and update on ulcerative colitis. *Elsevier Inc.*, 65(12), 100851.
- Geboes, K., & Van Eyken, P. (2009). Inflammatory bowel disease unclassified and indeterminate colitis: The role of the pathologist. *Journal of Clinical Pathology*, 62(3), 201–205.
- Jenkins, D., Balsitis, M., Gallivan, S., Dixon, M. F., Gilmour, H. M., Shepherd, N. A., Theodossi, A., & Williams, G. T. (1997). Guidelines for the initial biopsy diagnosis of suspected chronic idiopathic inflammatory bowel disease. The British Society of Gastroenterology Initiative. *Journal of Clinical Pathology*, 50(2), 93–105.
- Kalishah, J., Husada, D., Arfijanto, M. V., & Wahyu Widodo, A. D. (2022). Biofilm Formation and Antimicrobial Resistance of Escherichia coli in vitro Towards Ceftriaxone and Cefixime. *Current Internal Medicine Research and Practice Surabaya Journal*, 3(1), 5–8.
- Kellermann, L., & Riis, L. B. (2021). A close view on histopathological changes in inflammatory bowel disease, a narrative review. *Digestive Medicine Research*, 4, 3–3.

- Lang-Schwarz, C., Agaimy, A., Atreya, R., Becker, C., Danese, S., Fléjou, J. F., Gaßler, N., Grabsch, H. I., Hartmann, A., Kamarádová, K., Köhl, A. A., Lauwers, G. Y., Lugli, A., Nagtegaal, I., Neurath, M. F., Oberhuber, G., Peyrin-Biroulet, L., Rath, T., Riddell, R., Rubio, C. A., Sheahan, K., Tilg, H., Villanacci, V., Westerhoff, M., & Vieth, M. (2021). Maximizing the diagnostic information from biopsies in chronic inflammatory bowel diseases: Recommendations from the Erlangen International Consensus Conference on inflammatory bowel diseases and presentation of the IBD-DCA score as a proposal for a new i. *Virchows Archiv*, 478(3), 581–594.
- Montgomery, E., Voltaggio, L., & Canto, M. I. (2012). Much ado about very little (lamina propria)? *Gastrointestinal Endoscopy*, 75(1), 19–22.
- Ng, S. C. (2014). Epidemiology of inflammatory bowel disease: Focus on Asia. *Best Practice and Research: Clinical Gastroenterology*, 28(3), 363–372.
- Ng, W. K., Wong, S. H., & Ng, S. C. (2016). Changing epidemiological trends of inflammatory bowel disease in Asia. *Intestinal Research*, 14(2), 111–119.
- Nielson, O. H., and Seidelin, J. B. (2012). Non-IBD and noninfectious colitis. *Geriatric Gastroenterology*, 493–499.
- Odze, R. D. & Goldblum, J. R. (2015). Surgical pathology of the GI tract, liver, biliary tract and pancreas, surgical pathology of the GI tract, liver, biliary tract, and pancreas.
- Puspitarini, E. D., Prijambodo, P. (2022). Relationship between Sarcopenia in Abdomen CT Scan Results with C-Reactive Protein Level in Colorectal Cancer Patients at Dr. Soetomo General Academic Hospital Surabaya. *Biomolecular and Health Science Journal*, 5(1), 42–46.
- Sattianayagam, P. T., Hawkins, P. N., & Gillmore, J. D. (2009). Systemic amyloidosis and the gastrointestinal tract. *Nature Reviews Gastroenterology and Hepatology*. *Nature Publishing Group*, 6(10), 608–617.
- Saverymattu, S. H., Hodgson, H. J. F., Chadwick, V. S., & Pepys, M. B. (1986). Differing acute phase responses in Crohn's disease and ulcerative colitis. *Gut*, 27(7), 809–813.
- Simadibrata, M., & Adiwinata, R. (2017). Current issues of gastroenterology in Indonesia. *Acta medica Indonesiana*, 49(3), 270–278.
- Singh, R., Balekuduru, A., Simon, E. G., Alexander, M., & Pulimood, A. (2015). The differentiation of amebic colitis from inflammatory bowel disease on endoscopic mucosal biopsies. *Indian Journal of Pathology and Microbiology*, 58(4), 427–432.
- Surawicz, C. M., & Belic, L. (1984). Rectal biopsy helps to distinguish acute self-limited colitis from idiopathic inflammatory bowel disease. *Gastroenterology*, 86(1), 104–113.
- Turner, K., Genta, R. M., Lujan, G., Robiou, C., & Sonnenberg, A. (2014). Significance of the epithelioid granuloma in biopsies of Crohn's colitis. *Inflammatory Bowel Diseases*, 20(12), 2271–2275.
- Val, J. H. (2011). Old-age inflammatory bowel disease onset: A different problem? *World Journal of Gastroenterology*, 17(22), 2734–2739.
- Villanacci, V., Reggiani-Bonetti, L., Caprioli, F., Saragoni, L., Salviato, T., Mescoli, C., Canavese, G., Manenti, S., Spada, E., Baron, L., Leoncini, G., Cadei, M., Battista, S., & Armuzzi, A. (2020). Histopathology of inflammatory bowel disease — Position statement of the pathologists of the Italian Group for the Study of Inflammatory Bowel Disease (IG-IBD) and Italian Group of

- Gastrointestinal Pathologists (GIPAD-SIAPEC). *Digestive and Liver Disease. Editrice Gastroenterologica Italiana*, 52(3), 262-267.
- Villanacci, V., Reggiani-Bonetti, L., Salviato, T., Leoncini, G., Cadei, M., Albarello, L., Caputo, A., Aquilano, M. C., & Battista, S. (2021). Histopathology of IBD Colitis. A practical approach from the pathologists of the Italian Group for the study of the gastrointestinal tract (GIPAD). *Digestive and Liver Disease. Editrice Gastroenterologica Italiana*, 39-53.
- Walker, M. M., Potter, M. D., & Talley, N. J. (2019). Eosinophilic colitis and colonic eosinophilia. *Current Opinion in Gastroenterology*, 35(1), 42-50.

Variation of histopathological features in colonic mucosal biopsy with clinical diagnosis of suspicious inflammatory bowel disease in Dr. Soetomo General Academic Hospital, Surabaya periode 2015-2019

ORIGINALITY REPORT

15%	11%	11%	1%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	link.springer.com Internet Source	1%
2	www.sciencescholar.us Internet Source	1%
3	curis.ku.dk Internet Source	1%
4	indonesianjournalofcancer.or.id Internet Source	1%
5	geneticassociationdb.nih.gov Internet Source	1%
6	Joan Tosca Cuquerella, Marta Maia Bosca-Watts, Rosario Anton Ausejo, Sandra Tejedor Alonso et al. "Amyloidosis in Inflammatory Bowel Disease: A Systematic Review of Epidemiology, Clinical Features, and Treatment", Journal of Crohn's and Colitis, 2016 Publication	1%

7	Internet Source	1 %
8	academic.oup.com Internet Source	1 %
9	"Abstracts—APASL 2013", Hepatology International, 2013 Publication	1 %
10	sloap.org Internet Source	1 %
11	"The First China-Indonesia Joint International Symposium on Hepatobiliary Medicine and Surgery", Hepatology International, 2008 Publication	<1 %
12	"Practical Gastrointestinal Pathology", Springer Science and Business Media LLC, 2021 Publication	<1 %
13	"Soft Tissue and Bone Pathology", Histopathology, 2022 Publication	<1 %
14	Ondrej Fabian, Lukas Bajer. "Histopathological assessment of the microscopic activity in inflammatory bowel diseases: What are we looking for?", World Journal of Gastroenterology, 2022 Publication	<1 %

15 Gabriella Canavese, Vincenzo Villanacci, Elisabetta Antonelli, Moris Cadei et al. "Eosinophilia - associated basal plasmacytosis: an early and sensitive histologic feature of inflammatory bowel disease", APMIS, 2017
Publication <1 %

16 Martin Farmer. "The importance of diagnostic accuracy in colonic inflammatory bowel disease", The American Journal of Gastroenterology, 11/2000
Publication <1 %

17 coek.info
Internet Source <1 %

18 publikationen.ub.uni-frankfurt.de
Internet Source <1 %

19 Colitis, 2014.
Publication <1 %

20 epdf.pub
Internet Source <1 %

21 oamjms.eu
Internet Source <1 %

22 repub.eur.nl
Internet Source <1 %

23 vital.seals.ac.za:8080
Internet Source <1 %

24 "Morson and Dawson's Gastrointestinal Pathology", Wiley, 2013 <1 %
Publication

25 onlinelibrary.wiley.com <1 %
Internet Source

26 www.nature.com <1 %
Internet Source

27 "Endoscopy in Inflammatory Bowel Disease", Springer Science and Business Media LLC, 2015 <1 %
Publication

28 Cynthia Puspa Pitaloka, Absa Secka, Ernawati Ernawati, Agus Sulistyono, Hermanto Tri Juwono, Erry Gumilar Dachlan, Aditiawarman. "Characteristics Shifting of Heart Disease in Pregnancy: A Report from Low Middle-Income Country", Journal of Public Health Research, 2021 <1 %
Publication

29 repository.unair.ac.id <1 %
Internet Source

30 Lauge Kellermann, Lene Buhl Riis. "A close view on histopathological changes in inflammatory bowel disease, a narrative review", Digestive Medicine Research, 2021 <1 %
Publication

Exclude quotes On

Exclude bibliography On

Exclude matches Off

Variation of histopathological features in colonic mucosal biopsy with clinical diagnosis of suspicious inflammatory bowel disease in Dr. Soetomo General Academic Hospital, Surabaya periode 2015-2019

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11
