



SURAT KETERANGAN

Nomor : 1883/UN3.1.6/KP/2023

Yang bertanda tangan dibawah ini :

Nama : Prof. Dr. Mustofa Helmi Effendi, drh., DTAPH  
NIP : 196201151988031002  
Pangkat/Golongan : Pembina (Gol. IV/a)  
Jabatan : Wakil Dekan III

Dengan ini menerangkan bahwa :

Nama : Dr. Erma Safitri, drh., M.Si  
NIP : 196907231999032001  
Pangkat/Golongan : Penata Tk. I (Gol. III/d)  
Jabatan : Lektor

Telah melaksanakan penelitian dengan judul sebagai berikut :

No	Judul Karya Ilmiah	Tahun Pelaksanaan Penelitian
1	Immunomodulatory Activity of Black Jinten Oil ( <i>Nigella sativa</i> ) as Macrophage Activator for <i>Salmonella typhimurium</i> Infected Rat	2020
2	Screening the Reproductive Tract of Dairy Cattle for Pathogenic Micros	2019
3	Human Chorionic Gonadotropin (hCG) from Urine of Pregnant Women to Manipulate in vivo Ovulation and Pregnancy of Madura Cows	2019
4	Anti Early Embryonic Protein (EEP) for Pregnancy Test by Microtiter Strip in Dairy Cows	2019
5	The Effect of Feeding High Level of Protein on Reproductive Performance of Bali Starling.	2019
6	Antisperm Antibody in Repeat Breeder Friesian Holstein Cows at KPSP Setia Kawan Nongkojajar, Tutur District, Pasuruan, Indonesia.	2019
7	Diagnosis of Single and Twin Pregnancy, and Early Embryo Mortality Through Progesterone Level Test on Local Does.	2019
8	Improvement of Pregnancy Rate in Bali Cows with the Combination of Equine Chorionic Gonadotropine (eCG) from Local Pregnant Mare with PGF2 $\alpha$ .	2019
9	Progesterone Profile of Dairy Cows which Experienced the Failure of Pregnancy to Artificial Insemination (AI).	2019
10	Effect of Heat Shock Protein (HSP) in Post Thaw Baluran Bull Semen	2018
11	Potency of Mycotoxin Binders on MDA Level, Expressions of Caspase 9 and Caspase 3 in The Uterus of Mice Exposed to Zearalenone	2017



ASEAN  
University  
Network





# UNIVERSITAS AIRLANGGA

## FAKULTAS KEDOKTERAN HEWAN

Kampus C Mulyorejo Surabaya 60115 Telp. (031) 5992785, 5993016 Fax (031) 5993015

Laman: <http://www.fkh.unair.ac.id>, e-mail: [info@fkh.unair.ac.id](mailto:info@fkh.unair.ac.id)

12	Polymorphism of Growth Hormone Gene in The Artificial Insemination Result of Madura Cattle with Limousin Semen as a Reference for Genetic Selection	2018
13	Implementation of fotogrametry techniques as body mass estimation of indo-pacific bottle nose dolphin (Tursiops aduncus) in bali dolphin lodge	2020
14	Uji Sensitivitas Kebuntingan Sapi Perah Menggunakan Pregnancy Specific Protein B (PSPB) Microtiter Strip dan Progesteron sebagai Gold Standard	2007
15	Estimation of Equine Chorionic Gonadotropin (eCG) concentrate in the Blood Sera of Pregnant Mare	2014
16	Efek Pemberian L-Arginin Terhadap Gambaran Histologi Jumlah Spermatisit Primer pada Mencit (Mus musculus) Setelah Terpapar Suhu Panas	2019
17	Anti Prolactine Overcomes Heat Stress on Laying Hen.	2008
18	Unnatural Forced Moulting in The Laying Hen as Cause of Zoonosis from Salmonella Enteritidis	2009
19	Case Study: Dystocia on Beef Cattle in Kunir Regency of Lumajang District, East Java, Indonesia in 2015 and 2016	2017
20	Teratogenic Effect of Congenital Toxoplasmosis in Chicken Embryo	2017

Adapun penelitian tersebut layak dilakukan, meskipun belum ada ***Ethical Clearence*** karena menggunakan hewan coba yang minimal dan menghasilkan output yang sangat baik.

Demikian surat keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional **Guru Besar**

Surabaya, 3 April 2023

Wakil Dekan III,

Prof. Dr. Mustofa Helmi Effendi, drh., DTAPH

NIP 196201151988031002



ASEAN  
University  
Network



PROCEEDING



*10<sup>th</sup> Congress and International Conference  
of Indonesian Society for Microbiology*

Surabaya, November 19<sup>th</sup>-21<sup>th</sup>, 2009

**Recent Advances of Microbiology  
in Health, Bio-Industri, Agriculture and Environment**

**Editors :**

Fedik Abdul Rantam  
Koesnandar  
Soewarno  
Purwati  
Eryk Hendrianto

Wm  
✓

# PROCEEDING



## *10<sup>th</sup> Congress and International Conference of Indonesian Society for Microbiology*

Surabaya, November 19<sup>th</sup> - 21<sup>th</sup>, 2009

### **Editors :**

Fedik Abdul Rantam

Koesnandar

Soewarno

Purwati

Eryk Hendrianto

**Proceeding**

**10<sup>th</sup> Congress and International Conference of Indonesian Society for Microbiology**

**Editor :**

**Fedik Abdul Rantam, Koesnandar, Soewarno, Purwati, Eryk Hendrianto**

©2009 International Symposia Conference of Indonesian Society for Microbiology  
Surabaya-Indonesia

504 + x hal

All rights reserved this book or any part there of may not be reprinted or reproduced or utilited in any form or by an electronic, mechanical, or other means, including photocopying and recording, or in any infomation storage in retrieval system, without permission in writing from publisher

**ISBN 978-602-97750-0-6**

**Publisher:**

**Indonesian Society for Microbiology & Airlangga University  
Surabaya, November 2009**

*Surabaya, November 19<sup>th</sup>, 2009*

## OPENING REMARK CHAIRMAN OF ORGANIZING COMMITTEE

Microbiology-based industrial development to date is a fundamental need in the development of health and welfare of the whole community. Microorganisms are the foundation exploration and engineering that have a competitive and productive, so has the value of a significant advantage. Basic of this development based on the role of molecular biotechnology that utilizes microorganisms. The microorganism is use for industrial, environmental remediation, diagnostic, therapeutic materials, materials for vaccine and gene therapy vectors.

Congress of this proceeding are :

Theoretically Molecular exploration can be developed and applied in the industrial world for that health is a critical point of this approach to creating superior products based on microorganisms.

This Proceeding has discussed of about the latest developments bioindustri role in the development of microbiology, health, therapeutic, food technology, agriculture and environment. The focus of the discussion which will be discussed, lactic acid, enzymes, vaccines, diagnostics, immunotherapy, probiotics, food and environment-based bioindustri and microbiology.

PERMI in espousing the construction industry based on these microorganisms in cooperation with the International Microbiology society (IMS), the Association of lactic acid, Institute of Tropical Disease (ITD) as a center of medical biotechnology, Airlangga University, which is a tropical disease research center that focuses on education, research and service in health field. Related to those things need to be addressed so that the development of scientifically-based industrial technology on microorganisms are the main targets in the development of biotechnology that leads to its role in the bioindustri, health, food security and environment.

Surabaya, November 19<sup>th</sup> 2009

Chairman of the Committee,



Prof. Dr. Fedik A. Rantam, DVM

# CONTENTS

Pag

1. Opening Remark Chairman of Indonesian Society for Microbiology .....
2. Opening Remark Chairman of Organizing Committee .....
  
- HEALTH :**
3. Resistance of Antimicrobial pattern from Bench to Bedside:  
How to manage  
*Usman Hadi*.....
4. Nucleoprotein Gene Analysis of Rabies Virus Isolates Cross Indonesian to Design of Vaccines Candidate  
*Suwarno*.....
5. The Role of Cytokine in Dengue Haemorrhagic Fever (DHF) Pathogenesis  
*Purwati*.....
6. New Treatment Approach on HIV Infection  
*Suharto*.....
7. Cloning, Expression and Purification of Protease Human Immunodeficiency Virus Type I  
*Henny Saraswati*.....
8. The Increase of Allergic Reaction in Ovalbumin-induced Allergic Rats Stimulated with *Porphyromonas Gingivalis* Lipopolysaccharide (Verification of The Dental Plaque-Induced Asthma Hypothesis)  
*Haryono Utomo*.....
9. Butyric Acid Effect as A Result of Carbohydrate Fermentation by The Expression of P53 Colonic Microflora and Apoptotic Colonic Epithelial Cells of BALB / c Male After Inducible 9.10-dimethyl-1, 2-benz (a) anthracene (DMBA)  
*Cherry Siregar*.....
10. Nuclear Translocation of NF- $\kappa$ B induced by *Streptococcus intermedius* Intermedilysin  
*Heni Susilowati*.....
11. Antimicrobial Activity of Etanol Extract of Mahkota Dewa (*Phaleria macrocarpa*) Fruits and Leaves Against *Pseudomonas aeruginosa* by Agar Dilution and Scanning Electron Microscopy  
*Muhammad Ali Shodikin*.....
12. Power Antibacterial Turmeric Extract (*Curcuma domestica* Val.) Toward a clinical isolate of *Neisseria gonorrhoeae*  
*Maryani*.....
13. The Genetic Relationship of *C.albicans* between Diabetes Mellitus patients and Non-DM with Oral Candidiasis  
*Retno Pudji Rahayu*.....

14. Ability of Adhesin 61 kDa <i>Outer Membrane Protein Chlamydia pneumoniae</i> To Enhance Macrophage Activity <i>Sri Murwani</i> .....	99
15. Application of <i>Ribosome Inactivating Protein</i> (RIP) Isolated from <i>Annona squamosa</i> L. Seed as Alternative Mycoplasmaside <i>Widhianto, Tricahyadi</i> .....	106
16. Isolation and Identification Using 16S DNA of Mycorrhizosphere Bacteria and Their Antagonistic Potential Against Fungal Pathogen <i>Ganoderma boninense</i> in Vitro <i>Yenni Bakhtiar</i> .....	111
17. The Effect of Extract Kaffir Lime Peels ( <i>Citrus hystrix DC</i> ) in Inhibiting the Colonization of <i>Salmonella thymurium</i> in Small Intestine of Mice (in order to get drug candidate for typhoid fever) <i>Zulvikar Syam Bani Ulhaq</i> .....	123
18. TLC-Densitometry Validation Method for Determining Concentration in Tablet Gentamicin Topical <i>Achmad Burhanudin</i> .....	130
19. Effect of oxygenated water and probiotic administration on fecal microbiota of rats <sup>1</sup> <i>Ingrid S. Surono</i> .....	144
20. Study of Screening and Isolation Antibiotic from Marine Actinomycetes (Isolate A11) <i>Rofiq Sunaryanto</i> .....	151
21. Effectivity <i>Bacillus thuringiensis</i> soil isolate from Yogyakarta as larvacide against <i>Aedes aegypti</i> invitro <i>Lilis Suryani</i> .....	163
22. <i>Mycobacterium leprae</i> in Source Water Houses Patients and Non-endemic area of leprosy sufferers in East Java <i>Ratna Wahyuni</i> .....	170
23. The Effect of Red Fruit Oil ( <i>Pandanus Conoideus</i> ) to Apoptosis of Endothelial Cells Exposed with Falciparum Malaria Patient Serum and Neutrophil <i>Yunita Armiyanti</i> .....	179
24. Prevalence Distribution of Intestinal Parasite Infestation in Under Five Years Children with Severe Malnutrition in Kasihan, Bantul, Yogyakarta Based on Risk Factors <i>Tri Wulandari Kesetyaningsih</i> .....	195
<b>AGRICULTURE :</b>	
25. Cloning, Expression and Characterization of Regulator Protein for Lovastatin Biosynthesis in <i>Escherichia coli</i> <i>Dudi Hardianto</i> .....	202
26. Anticancer activity of Endophytic fungi <i>Curvularia lunata</i> BioMCC-FE00283 isolated from <i>Cibotium barometz</i> <i>Erwahyuni E. Prabandari</i> .....	213



41. Production of Xylooligosaccharides from Corn cob Using Xylanolytic Enzyme Produced by <i>Bacillus subtilis</i> AQ1 <i>Dadang Suhendar</i> .....	343
42. In Vitro Screening of <i>Dumortiera hirsuta</i> Extracts Against <i>Candida albicans</i> <i>Junairiah</i> .....	350
43. The Ability Policulture Lactic Acid Bacteria from Sauerkraut by Various Starters In Fermenting Tomato Juice <i>Dyah Fitri Kusharyati</i> .....	355
44. Fermentable Substrate from Cellulosic Material Saccharification by Using Microbial Enzymes and Pretreatment <i>Lasam Soeroso</i> .....	363

#### INDUSTRI:

45. The Role of <i>Rhizopus oligosporus</i> on Antioxidant Biosynthesis During Tempe Fermentation <i>Suyanto Pawiroharsono</i> .....	375
46. Aspergillus Biodiversity and Their Role in Industrial Enzyme Production : mini review <i>Isworo Rukmi</i> .....	387
47. Study of Probiotic Bacteria Growth ( <i>Lactobacillus casei</i> and <i>Lactobacillus plantarum</i> B2) in Soy Milk and Skim Milk Fermentation Medium <i>Elok Zubaidah</i> .....	399
48. Transformation of Autotrophic Into Heterotrophic Microalgae by Cell Biotechnology to Increase Lipid Production for Biodiesel Source from Blooming Microalgae <i>Hermin Pancasakti Kusumaningrum</i> .....	409
49. Production of Polyunsaturated Fatty Acids with <i>Rhizomucor miehei</i> by Submerged Fermentation <i>Ignatius Srianta</i> .....	418
50. Isolasi dan Identifikasi Bakteri Asam Laktat pada Tape Biji Teratai ( <i>Nymphae pubescens Willd</i> ) <i>Rita Khairina</i> .....	428
51. Molecular Characterization of Bacterial Isolates from the Coastal Region of South Jember - Indonesia: Diversity and Carbon Substrates Utilization <i>Kartika Senjarini</i> .....	437
52. The Effect of Incubation Time of Tapioca Liquid Waste Fermentation by <i>Candida utilis</i> on Production Characteristic of Single Cell Protein <i>Osfar Sjojfan</i> .....	447
53. Isolasi dan Identifikasi <i>Azotobacter vinelandii</i> dari Rhizosfer Serta Uji Potensi untuk Memproduksi Biopolimer Alginat <i>Nurosid</i> .....	458
54. Extracellular Protease Activity of Proteolytic Bacteria Isolated from Coturnix Japonica Digestive Tract in Soybean Meal Medium <i>Inge Libertina</i> .....	465

55. Amylase Production by <i>Aspergillus niger</i> Using Raw Starch <i>Suraya</i> .....	475
56. Extracellular Amylase Activity of Amylolytic Bacteria Isolated from <i>Coturnix japonica</i> Digestive Tract in Corn Flour Medium <i>Winda Kusuma Wardhani</i> .....	483
57. Unnatural Forced Moulting in The Laying Hen as Cause of Zoonosis from <i>Salmonella enteritidis</i> .....	493
<b>APPENDIX</b> .....	<b>499</b>

# COMMITTEE

## 10<sup>th</sup> CONGRESS AND INTERNATIONAL SYMPOSIA CONFERENCE OF INDONESIAN SOCIETY FOR MICROBIOLOGY

### 1. Steering Committee

Chairman : Prof. Dr. Harianto Danutirto, DEA.  
Secretary : Dr. Ir. Koesnandar M. Eng.  
Members : Prof. Amin Subandrio, MD., Ph.D.  
Prof. Dr. Ir. Antonius Suwanto, MSc  
Prof. Dr. Ir. Sutriswati Endang Rahayu, MSc.  
Prof. Sam Suharto, MD, Sp.MK. (K)  
Dr. Ir. Students Setyahadi  
Dr. Nasronudin, MD. SpPD (K)  
Prof. Dr. Kuntaman, M.S. MD (SpMK)  
Prof. Dr. Abdul Fedik Rantam, DVM.

### 2. Organizing committee

Chairman : Prof. Dr. Abdul Fedik Rantam, DVM  
Secretary : Dra. Yenni Bachtiar, M.Agr.Sc.,  
Dr. Retno Indrawati, M. Si, drg.  
Purwati, dr., SpPD  
Treasurer : Dra. Diana Nurani.,  
Dr. Diah S. Ernawati, drg., MSI., SpOM  
Funds : Drs. Nicodemus, MM.,  
Lindawati Palilingan, dr., MS  
Scientific : Dr. Suwarno, MSI. DVM.  
M. Inge Lucida dr., MS., Ph.D.  
Protocol & gathering night:  
Dr. Prihatini Widiyanti, M. Kes., Drg  
Dr. AT. Soelih Estoepangestie, drh., DVM.  
Food and Beverage:  
Tutik Kusumaningsih, drg. M.Kes.  
Dr. Anggraini Rahayu, SKM., Msi  
Transportation : Bayu, dr.  
Budi Rahardjo, drg. M. Kes.  
Equipment : Arja Mukti Diamond, dr.  
Wartono SH  
Publications and documentation:  
Agung Dwi Wahyu Widodo, dr., Si  
M. Anam Al-Arif, MP., DVM.

# UNNATURAL FORCED MOULTING IN THE LAYING HEN AS CAUSE OF ZONOSIS FROM *SALMONELLA ENTERITIDIS*

**Erma Safttri, Pudji Srianto, Trilas Sardjito**

Reproduction Veteriner – Faculty of Veteriner-Unair  
Kampus C, Jl. Mulyorejo, Unair. e-mail: [rma\\_fispro@telkom.net](mailto:rma_fispro@telkom.net)

## **Abstract**

The aims of this reseach is to give information to the farmers and people about the bad consequences of consuming the poultry products that use unnatural forced moulting. The research was carried out by examining *Salmonella enteritidis* bacteria in the laying hens feces after treatment.

The laying hens devided in three groups and treatment with unnatural forced moulting, dietary feed carried out during 2, 3 and 4 days, and the another group given natural forced moulting treatment by 200 ìg/ml each hen á Prl single dose compared with the group without dietary feed and á Prl.

The result showed that unnatural forced moulting with dietary feed during 2, 3 and 4 days decrease laying hen immune function. It was founded *Salmonella enteritidis* in laying hens feces after treatment. The *Salmonella enteritidis* can be the causal factor of zoonosis for human.

**Key words :** Unnatural forced moulting, natural forced moulting, laying hen,  $\alpha$  Prl, *Salmonella enteritidis*, zoonosis

## Introduction

Problem of stopping egg production when laying hens enter molting period cause many loss of benefits, economically and time (Safitri, 2005, Safitri, 2008). First molting period of laying hens is at 22<sup>nd</sup> – 24<sup>th</sup> months. The duration of this period is 60 -75 days (Darmana and Sitanggang, 2002, Sudarso and Siriwa, 2002). If there is no action against this period, it will take longer time for laying hens to produce again eggs, about 80-100 days (Marhiyanto, 2000, Indarto, 1989, Juli, 1982).

Bell and Kunney (2003) and Avma (2003) found that to overcome molting period in United States, it has been done natural forced molting and unnatural forced molting. The forced molting method carried out in 3 ways : dietary feed, giving low nutrient feed (protein, calcium, and natrium), use of drugs and methalibure, chlordamine, high dose iodium, aluminum, and zinc.

In Indonesia, it has been done many efforts to overcome molting period : dietary feed (unnatural forced molting). It was carried out for about 30 days (Safitri, 2008, Barton, 2003, Sainsbury, 1995). Long unnatural forced molting can cause stress to laying hens, decrease of immune function, and laying hens can be attacked easily by diseases (Poultry, 2003, Alodan and Mashalay, 1999).

One of the diseases follows unnatural forced molting is Salmonellosis caused by *Salmonella enteridis* (Sukamto and Hendarti, 2008, Fact, 2001, Webster, 1999). Salmonellosis is infectious disease and zoonosis to human. Since 2000, it has been banned the use of unnatural forced molting by dietary feed (Avma, 2003).

Butcher and Miles (2002) found that unnatural forced molting decreased T cell in blood. The decrease of T cell can cause the immune decrease and increase the sensitivity to the disease, especially Salmonellosis. Turner and Bagnara (1998), Knobil et al (1988) and Hafez (2000), found high prolactin hormone at molting hens. Safitri (2005) and Safitri et al (2006) found an alternative to overcome molting period by natural method to arabic hens and laying hens (use of  $\alpha$  Prl). It is effective to resist molting period and make laying hens produce again eggs.

The present research is carried out to give information about the method to overcome molting period by natural forced molting as substitute of unnatural forced molting method.

## Materials and methods

Fifty laying hens in molting period were grouped and placed in cage randomly by five treatments (each treatment with 10 repetitions):

- PO (control) : ten laying hens in molting period without natural forced molting and without unnatural forced molting
- P1 : ten laying hens in molting period were treated by natural forced molting, by giving single dose of  $\alpha$  Prl 200  $\mu$ g/mL each laying hens

- P2 : ten laying hens in molting period were treated by unnatural forced molting, by dietary feed for 2 days consecutive without feed and only drink given
- P3 : ten laying hens in molting period were treated by unnatural forced molting, by dietary feed for 3 days consecutive without feed and only drink given
- P4 : ten laying hens in molting period were treated by unnatural forced molting by dietary feed for 4 days consecutive without feed and only drink given

In the fifth day, it has been checked *Salmonella* in 50 laying hens feces. The method was by taking one colony round shaped and transparent from MCA media, then growth at SSA media, and put at 37°C for 24 hours. Identification test were done to the colony. To ensure that the colony growth at SSA media were *Salmonella enteritidis*, biochemical tests were done to various media (TSIA, SIM, citrate, urea, and sweets).

#### **Triple Sugar Iron Agar (TSIA) Media**

The growth of *Salmonella* in TSIA media was done by taking *Salmonella* from pure culture by sterile needle then put it in the bottom of agar, and then media was putted at 37°C for 24 hours. The growth in this media is to evaluate the ability of *Salmonella* to ferment glucose, sucrose, and the formation of H<sub>2</sub>S and gas (Cappucino and Sherman, 1983).

#### **Sulfide Indol Motility (SIM) Media**

The growth in SIM media was done by taking *Salmonella* from pure culture then put it in the half part of agar and the put the media at 37°C for 24 hours. Observation was done by adding chloroform and Kovach reagent, both at the same quantity. Positive result was shown by indol circle (red).

#### **Citrat Agar**

The growth in citrate agar was done by streaking in media. The objective of this growth is to identify whether *Salmonella* uses carbon from citrate. Positive result was shown by the change of color to blue and negative result was shown by the change of color to green.

#### **Urea Agar**

Pure culture was taken to streak in media. The objective of this test to identify whether *Salmonella* produces enzyme that can hydrolyze urea to ammonia. The positive result was shown by the change of media to red color.

- P2 : ten laying hens in molting period were treated by unnatural forced molting, by dietary feed for 2 days consecutive without feed and only drink given
- P3 : ten laying hens in molting period were treated by unnatural forced molting, by dietary feed for 3 days consecutive without feed and only drink given
- P4 : ten laying hens in molting period were treated by unnatural forced molting by dietary feed for 4 days consecutive without feed and only drink given

In the fifth day, it has been checked *Salmonella* in 50 laying hens feces. The method was by taking one colony round shaped and transparent from MCA media, then growth at SSA media, and put at 37°C for 24 hours. Identification test were done to the colony. To ensure that the colony growth at SSA media were *Salmonella enteritidis*, biochemical tests were done to various media (TSIA, SIM, citrate, urea, and sweets).

#### **Triple Sugar Iron Agar (TSIA) Media**

The growth of *Salmonella* in TSIA media was done by taking *Salmonella* from pure culture by sterile needle then put it in the bottom of agar, and then media was putted at 37°C for 24 hours. The growth in this media is to evaluate the ability of *Salmonella* to ferment glucose, sucrose, and the formation of H<sub>2</sub>S and gas (Cappucino and Sherman, 1983).

#### **Sulfide Indol Motility (SIM) Media**

The growth in SIM media was done by taking *Salmonella* from pure culture then put it in the half part of agar and the put the media at 37°C for 24 hours. Observation was done by adding chloroform and Kovach reagent, both at the same quantity. Positive result was shown by indol circle (red).

#### **Citrat Agar**

The growth in citrate agar was done by streaking in media. The objective of this growth is to identify whether *Salmonella* uses carbon from citrate. Positive result was shown by the change of color to blue and negative result was shown by the change of color to green.

#### **Urea Agar**

Pure culture was taken to streak in media. The objective of this test to identify whether *Salmonella* produces enzyme that can hydrolyze urea to ammonia. The positive result was shown by the change of media to red color.

### Sweets

The growth in sweets media was done by taking Salmonella the put it in media. The objective of this growth is to identify whether Salmonella ferments sweets (Anonymous, 1999).

### Experiment Design and Statistical Analysis

In this present research, complete random design is used. The results were analyzed by Analysis of Variant (ANOVA) and least significant difference 5% level test if there was the different result (Steer and Torrie, 1995).

### Results and Discussion

The results of experiments are shown in the table 1.

Table 1 Identification of Salmonella in different treatment samples

Repetition	Treatment				
	Control	P1	P2	P3	P4
1	-	-	+	+	+
2	-	-	-	+	+
3	-	-	+	+	+
4	-	-	+	+	+
5	-	-	+	+	+
6	-	-	-	+	+
7	-	-	+	+	+
8	-	-	+	+	+
9	-	-	+	+	+
10	-	-	-	+	+
Sum	0 <sup>a</sup>	0 <sup>a</sup>	7 <sup>b</sup>	10 <sup>c</sup>	10 <sup>c</sup>

- : negative, no Salmonella found

+: positive, Salmonella found

The positive results in the table shown that there were Salmonella enteriditis in the tested feces. It is identified by biochemical test by TSIA, SIM, Citrat agar, urea agar, and sweets methods as shown in table 2.



**Table 2. Identification of *Salmonella enteritidis* in different media**

Media	Result
TSIA	+
Base	+
Acid	+
H <sub>2</sub> S	+
Gas	+
SIM	-
Motility	-
Indol	-
H <sub>2</sub> S	+
Citrate	-
Urea	-
Glucose	+
Lactose	-
Manitol	+
Maltose	+
Sucrose	-

According to statistical analysis, one way ANOVA, it was found there was no significant difference ( $p < 0,01$ ) between results of control and natural forced molting ( $\alpha$  Pri 200 mg/L each hen). Between results of control and P1,P2,P3, there was significant different ( $p < 0,01$ ), then it is necessary to analyze with least significant difference in order to determine the most important treatment that influence the existence of salmonella.

Butcher and Miles (2002) found that dietary feed will decrease T cell in blood circulation. By this decrease, it will decrease immune reaction and increase the sensitivity to certain disease. The disease that occurs especially is *Salmonella enteritidis*. According to Fact (2001), Webster (1999), this disease is very dangerous because it is zoonosis for human.

*Salmonella enteritidis*, the cause of Salmonellosis, is 1-2,5 micron length and 0,3-0,5 micron width, negative gram, non motile, without spore formation, and no capsule. The optimum growth of *Salmonella enteritidis* is at 37°C and in *Salmonella shigella* agar forms segmented colony, clear, and transparent (Hagan and Bruner, 1981).

Salmonellosis is infectious disease that spreads all over the world. Mammalian can be infected by this disease: rabbit, human (Ressang, 1984).

Salmonellosis spreads via oral wit feed, inhalation of polluted air (feather, dust, infected feces). Another way of spreading is congenital via egg (Hosfad, 1984). Factors that enhance the spreading are bad air circulation system, bad sanitation system, bad feed supply, another disease that may give synergic effect (Anonymous, 1981).

Salmonella enteritidis reaches optimum growth at 37°C and aerobic or anaerobic facultative. It can produce H<sub>2</sub>S, acid and gas from glucose, ferments manose, but can not ferment lactose. Maltose can be fermented rarely, sucrose not fermented. Salmonella enteritidis does not have urease enzyme and can not use carbon from citrate (Hofad, 1984, Kingscote, 1989).

Isolation of Salmonella enteritidis needs selective media. These media are Salmonella Shigela Agar (SSA), Mac Concey Agar (MCA), and Endo Agar. The growth of Salmonella enteritidis in SSA is manifested by colony formation which is smooth, round, transparent, and without color. Diameter of colony is 1 to 4 mm. Salmonella enteritidis in TSIA media gives the change of red color (base), yellow (acid), H<sub>2</sub>S formation (black), gas formation (burst medium). The growth of Salmonella enteritidis in SIM manifested by white color (non motile). Black formation shows H<sub>2</sub>S and negative inol test (Jackson and Simmon, 1981, Hofstad, 1984).

The present research gives information to farmers, and people the danger of poultry products produced by unnatural forced molting to overcome molting. Information can be presented in case of Salmonella enteritidis, found in laying hens feces.

## Conclusion

Unnatural forced molting by dietary feed to overcome molting can cause case of salmonella enteritidis which is zoonosis to human.

## Daftar Pustaka

- Anonimus. 1981. Pedoman Pengendalian Penyakit Hewan Menular Jilid I Cetakan ke-2. Direktorat Kesehatan Hewan. Direktorat Jendral Peternakan. Departemen Pertanian. Jakarta. 73-80.
- Anonimus. 1997. Pedoman Praktikum Bakteriologi. Fakultas Kedokteran Hewan. Universitas Airlangga. Surabaya. 34-35.
- Allen T. 2002 . Information Resources on Induced Molting in Chicken 1902-2002. Animal Welfare Information Center. Email: [awic@nal.usda.gov](mailto:awic@nal.usda.gov)<http://www.nal.usda.gov/awic/> Down load : 25 Januari 2004.

- Alodan M.A., and Mashaly M.M., 1999. Effect of Induced Molting in Laying Hens on Production and Immune Parameters. *Education and Production. Poultry Science*. 78:171-177. Down load : 27 Januari 2004.
- Avma H.A. 2003 . The Animal Welfare and Food Safety Issues Associated with the Forced Molting of Laying Birds. *AVMAHurtsAnimals: Vets Without Hearts*. United Poultry Concerns,Inc. [UPC.online.org](http://UPC.online.org). Down load:20 Pebruari 2004.
- Barton J., DVM. 2003. Molting. *Palm Beach Country Poultry Fanciers Association*. Florida. [http://www.metimes.com/issue99-24/eg/chicken.htm.p. 3](http://www.metimes.com/issue99-24/eg/chicken.htm.p.3).
- Bell and Kuney. 2003. Forced Molting of Laying. Birds. *Poultry Organization*. p. 8-10. [Poultry.org](http://Poultry.org) is an educational effort of Farm Sanctuary. Down load : 2 Maret 2004.
- Butcher G.D. and Miles R. 2002. Salmonella Control and Molting of Egg-Laying Flocks-Are They Compatible. *University of Florida. Cooperative Extension Service. Institute of Food and Agriculture Sciences*. P. 1-3. Down load : 25 Januari 2004.
- Cappucino, J.G. and N. Sherman. 1983. *Microbiology a Laboratorymanual*. Canada. 82-84.
- Darmana W. dan M. Sitanggang. 2002. Kiat Mengatasi Permasalahan Praktis. Meningkatkan Produktivitas Ayam Arab Petelur. Cetakan I. *Agro Media Pustaka*. Jakarta. H.
- Fact 2001. Nears Major Food Safety Goal. *Chicago IL 60614 (773) 525-4952. PO BOX 14599. Info@FACT.cc.www.fact.cc/se-Main.htm*. Down load : 21 April 1004
- Hagan , W.A. and D.W. Brunner. 1981. *The Infectious Disease of Domestic Animal with Special Reference to Etiology and Biology Therapy*. 3<sup>rd</sup> Ed. *Camstock Publishing Associate, A. Division of Cornell University Press*. Thaca. New York. 90-93.
- Hosfad, M.S. 1984. *Disease of Poultry*. 8<sup>th</sup> Ed. *Angus and Robertson*. Sidney Melbourne. 231-250.
- Indarto, P. 1989. *Beternak Unggas Berhasil*. Penerbit CV. Armico. Bandung. Hal. 63-69.
- Jackson, C.A.W. and G.C. Simmons. 1981. *Standard Diagnostic Technique for Salmonellosis*. 1<sup>st</sup> Ed. *The Australian Bureum of Animal Health*. Australia. 1-10.
- Jull, M.A. 1982. *Poultry Husbandry*. TMH Edition 1972, Reprinted 1982. *Tata McGraw-Hill Publishing Company Ltd*. New Delhi. 119-149.
- Kingscote, B. 1989. *Veterinary Microbiology Introduction to Bacteria and Virology*. 7<sup>th</sup> Ed. *The Iowa State University Press*. Ames. Iowa. USA. 285-309.
- Marhiyanto B. 2000. *Sukses Beternak Ayam Arab*. Cetakan I. *Difa Publisher*. Indonesia. H. 9-11 & 88-97
- Poultry. O. 2003. Forced Molting of Laying. Birds. *Poultry Organization*. p. 1-7. [Poultry.org](http://Poultry.org) is an educational effort of Farm Sanctuary. Down load : 2 Maret 2004.
- Ressang, A.A. 1984. *Patologi Khusus Veteriner*. NV. Denpasar. 591-593.
- Safitri E. 2005. Metode pembuatan antiprolaktin pada hewan coba kasmbing local sebagai penghambat proses moulting pada ayam ras petelur. *Berkala penelitian Hayati. Journal of Biological Researches*. 11(1):26-29.

- Safitri E., Misaco W., Anwar H. 2008. Penentuan dosis isolate prolaktin hasil purifikasi serum darah itik fase moulting untuk produksi anti prolaktin. *Media Kedokteran Hewan*. 24(3): 177-182.
- Sainsbury D. 1995. *Poultry Health and Management; Chicken, Turkey, Ducks, Geese and Quail*. 3 Th. Ed. University of Cambridge. 195
- Steel, R.G.D dan J.H. Torrie. 1995. *Principles and procedures of statistics A biometrical Approach*. International Student Edition. Me Graw-Hill Koga Kusha, LTD. P. 137-167
- Sudarso Y dan Siriwa A. 2001. *Ransum Ayam dan Itik*. Penebar Swadaya. Anggota ICAPI. Jakarta. Hal. 47-48.
- Sukanto R. dan Hendarti, G.A. 2008. Efek torced moulting pada unggas sebagai penyebab kasus zoonosis *Salmonella enteritidis* (SE). *Veterinaria Medika*. 2 (1) : 156-1161
- Webster A.B. 1999. Commercial Egg Tip-the Induced Molt: A Critical Control Point for Hazard Minimization of Salmonella Enteritidis Contamination of Eggs. Institute of Agriculture and Natural. *Poultry News*. Winter 1999. P2-4. <http://ianr.www.snl.edu/ianr/asdk/newslet.htm>

--- oOo ---