

## Re: [vmic-1] Submission Acknowledgement

---

From: Veterinary Medicine International Conference Universitas Airlangga (vmic@fkh.unair.ac.id)

To: rma\_fispro@yahoo.com

Date: Wednesday, June 28, 2017 at 09:50 PM GMT+7

---

Dear VMIC participants

Hereby I would like to inform you about the preparations for The 1st Veterinary Medicine International Conference 2017.

### **For those who are doing Oral Presentation :**

The plenary session for oral presentation will be held at Tandjung Adiwinata Room 2nd floor of Veterinary Medicine Faculty Universitas Airlangga. Please prepare a power point presentation for 10 minutes presentation, please be advised that the organizing committee will be strict to the timetable and thus who unable to finish their presentation is not the committee responsibilities.

We recommend the presenter to not make presentation more than 10 slides. Please send your presentation to the [vmic@fkh.unair.ac.id](mailto:vmic@fkh.unair.ac.id) before July 3, 2017 with the subject and document's name following this specific format : presentation\_First author name. The schedule for oral presentation will be informed very soon.

### **For those who are doing Poster Presentation :**

The poster session will be held in front of Tandjung Adiwinata Room 2nd floor of Veterinary Medicine Faculty Universitas Airlangga. Please prepare the poster in A0 paper (**1200 mm X 850 mm**) and please bring the poster during registration time. Presenter should put their own poster in the space that the organizing committee provided.

I enclosed the conference schedule for your information. In case you have any question regarding the conference please do not hesitate to ask the organizing committee. Thank you very much and welcome to 1st VMIC 2017.

Best regards

2017-02-28 10:49 GMT+07:00 Veterinary Medicine International Conference Universitas Airlangga  
<[vmic@fkh.unair.ac.id](mailto:vmic@fkh.unair.ac.id)>:

Dear author, after reviewing your paper we found out that your paper contain 25% of plagiarism. Please revise your paper according to plagiarism test result. Thanks.

2017-02-27 5:20 GMT+07:00 VMIC <[vmic@fkh.unair.ac.id](mailto:vmic@fkh.unair.ac.id)>:

Assalamualaikum Erma Safitri:

Thank you for your submission, "The Effectiveness of Honey in Physiological NACL to Maintain of Viabilityand Motilityof Spermatozoa"

to THE 1st VETERINARY MEDICINE INTERNATIONAL CONFERENCE. With the online conference management system that we are using, you will be able to track its progress through the editorial process by logging in to the conference web site:

Submission URL:

[http://conference.unair.ac.id/index.php/vmic1/vmic-1/author\\_submission/1212](http://conference.unair.ac.id/index.php/vmic1/vmic-1/author_submission/1212)

Username: 1erma\_safitri2

If you have any questions, please contact me. Please following us at our website and our social media platform to keep update for this up coming event. Thank you for considering this conference as a venue for your work.

VMIC

THE 1st VETERINARY MEDICINE INTERNATIONAL CONFERENCE

---

THE 1st VETERINARY MEDICINE INTERNATIONAL CONFERENCE THE 1stVETERINARYMEDICINE  
INTERNATIONAL CONFERENCE

<http://conference.unair.ac.id/index.php/vmic1/vmic-1/index>

--

The Veterinary Medicine International Conference  
Universitas Airlangga  
Surabaya  
Indonesia  
+62 878 523 413 77 (whatsapp)

--

The Veterinary Medicine International Conference  
Universitas Airlangga  
Surabaya  
Indonesia  
+62 878 523 413 77 (whatsapp)



Scientific Schedule.xlsx  
13.7kB

**PROGRAM SCHEDULE**  
**THE FIRST VETERINARY MEDICINE INTERNATIONAL CONFERENCE 2017**  
**in Collaboration with ADPRC-OHCC**  
**“New Wave Bio-X Veterinary Medicine”**  
**“Strengthen on One Health, Biomedical, Reproduction, Nutrition and Clinical Science”**  
**FACULTY OF VETERINARY MEDICINE UNIVERSITAS AIRLANGGA**  
**SURABAYA - INDONESIA**

**Day 1: Wednesday , 12 July 2017**

12.00 – 13.00 am	<b>Registration and coffee break</b>
13.00 – 13.20 am	<b>Opening Remarks:</b>
	<ol style="list-style-type: none"> <li>1. Chairman of The VMIC</li> <li>2. Dean of Faculty of Veterinary Medicine Universitas Airlangga</li> <li>3. Coordinator of INDOHUN University of Indonesia</li> <li>4. Rector of Universitas Airlangga</li> </ol>
13.20 – 13.30	<b>Opening Ceremony</b>
13.30 – 15.30	<b>KEYNOTE LECTURE-1 at Tandjung Adiwinata 2nd floor</b>
	Chairman: Prof. Bambang Sektiari, DVM Co-chairman: Dr. Hani Plumeriastuti, DVM. M.Sc.
45'	<b>Prof. Graeme B Martin, BSc (Agric) PhD</b> ( <i>University of Western Australia</i> ) <i>Science communication</i>
30'	<b>Prof. Saul Tzipori, DVM, PhD.</b> ( <i>Tufts University, USA</i> ) <i>Evolutionary and Environmental Factors Contributing to Emergence of New Diseases</i>
30'	<b>Dr. Khaled Hussein, DVM, Ph.D, MRCVS</b> ( <i>ECO Animal Health Ltd, UK</i> ) <i>Food chain and food safety</i>
15'	<i>Discussion</i>
15.30 - 17.45	<b>KEYNOTE LECTURE-2 at Tandjung Adiwinata 2nd floor</b>
	Chairman : Dr. Soelih Estoepangestie, DVM Co-Chairman: Dr. Wiwik Misaco,
30'	<b>Prof. Datin Paduka Dr. Aini Ideris</b> ( <i>Vice Chancellor-University Putra Malaysia</i> ) <i>Vaccine development to prevent infectious disease</i>
30'	<b>Prof. Dr. Koichi Sato, DVM, PhD</b> ( <i>University of Yamaguchi, Japan</i> ) <i>Cell physiological research of intestinal stem cell</i>
30'	<b>Assoc. Prof. Takeshi Haga, DVM, PhD.</b> ( <i>University of Tokyo, Japan</i> ) <i>Diversity and Pathogenicity of Papillomaviruses</i>
30'	<b>Valérie Cai</b> ( <i>SEPPIC, France</i> ) <i>Comparison of adjuvant technologies for poultry vaccines</i>
15'	<i>Discussion</i>
17.45 - 21.00	<b>Closing Remark and Gala Dinner</b>

**Day 2: Thursday , 13 July 2017**

07.30 – 08.30	<b>Registration</b>
08.30 - 10.15	<b>KEYNOTE LECTURE-3 at Tandjung Adiwinata 2nd floor</b> Chairman: Prof. Mas'ud Hariadi Co-chairman: Dr. Tita Damayanti, M.Sc., DVM 30' <b>Assoc. Prof. Dr. Naoki Miura, DVM, Ph.D</b> ( <i>Univeristy of Kagoshima, Japan</i> ) <i>The application of small non-coding RNA in the veterinary field.</i> 30' <b>Assoc. Prof. Tamaki Okabayashi, Ph.D.</b> ( <i>University of Miyazaki, Japan</i> ) Enhancement of the livestock productivity and reinforcement of countermeasure for prevention of animal infectious diseases in Southeast Asia by a cooperation with research bases with a hub function 30' <b>Prof Hong-Kean Ooi</b> ( <i>Azaba Univeristy, Japan</i> ) <i>Abortion in ruminants and neosporosis</i> 15' <i>Discussion</i>
10.15 - 10.30	<b>Coffe Break</b>
10.30 -12.15	<b>KEYNOTE LECTURE-4 at Tandjung Adiwinata 2nd floor</b> Chairman : Dr. Budi Utomo, MD., M.Kes Co-chairman : Dikky Eka Mandala, DVM, MSc 30' <b>Prof. Dr. Kuntaman, MD., SpMK (K)</b> ( <i>University of Airlangga</i> ) <i>Strategy to combat Antimicrobial Resistance in human</i> 30' <b>Dr. A. T. Soelih Estoepangestie, DVM.</b> ( <i>University of Airlangga</i> ) AMR Foodborne bacteria of wet market in Surabaya 30' <b>Dr. Mustofa Helmi Effendi, DVM. DTAPH.</b> ( <i>University of Airlangga</i> ) Methicillin Resistant Staphylococcus aureus in Surabaya 15' <i>Discussion</i>
12.15 - 13.00	<b>LUNCH</b>
13.00. 14.45	<b>KEYNOTE LECTURE-5 at Tandjung Adiwinata 2nd floor</b> Chairman : Dr. Atik Choirul Hidajah, M.Kes Co-chairman : Havan Yusuf, DVM, MSc. 30' <b>Prof. Agus Suwandono , MD., MPH.</b> ( <i>University of Diponegoro</i> ) The One Health Systems Mapping and Analysis Resource Toolkit (OH-SMART) 30' <b>Dr. Ni Nyoman Sri Budayanti, MD., Sp.MK</b> ( <i>Univeristy of Udayana</i> ) <i>The rule of travel Medicine caused by infectious disease</i> 30' <b>Prof. Dr. Fedik A. Rantam, DVM.</b> ( <i>University of Airlangga</i> ) <i>Hybrid of IgG against all serotype to dengue virus using mesenchymal stem cell</i> 15' <i>Discussion</i>
14.45-15.00	<b>Coffe Break</b>
15.00 - 16.30	<b>PLENARY SESSION</b>
16.30	<b>Closing Ceremony</b>









  
**Knowledge E**  
Engaging minds

 **KnE Life Sciences**

## The Veterinary Medicine International Conference (VMIC)

12-14 July 2017

ISSN 2413-0877



[www.KnEpublishing.com](http://www.KnEpublishing.com)



# The Veterinary Medicine International Conference (VMIC)

	<p><b>KnE Life Sciences</b> / The Veterinary Medicine International Conference (VMIC) VMIC—The Veterinary Medicine International Conference—is one of the world's leading conference focusing on a wide array of topics including Veterinary Medicine and Biomedical Science. It offers a stimulating venue for scientists, researchers, lecturers, general practitioners, and others to broaden their social scientific network. This conference aimed at improving human and public health by improving agricultural and food systems, advancing biomedical and comparative medical research, preventing and addressing zoonotic diseases, kit diagnostic, enhancing environmental and ecosystem health, and helping manage the 21st-century public health challenges.</p> <p><b>Conference date:</b> 12–14 July 2017 <b>Location:</b> Surabaya, East Java, Indonesia <b>Editors:</b> Sri Agus Sudjarwo, Fedik A. Rantam, rer. nat. Gunawan Indrayanto, Muchammad Yunus, Rimayant, Wiwik Misaco, Tita Damayanti Lestari, Mustofa Helmi Effendi, Dikky Eka Mandala Putranto, and Shafia Khairani <b>Organizer:</b> Faculty of Veterinary Medicine, Universitas Airlangga, Indonesia <b>Sponsors:</b> Ministry of Research, Technology and Higher Education of the Republic of Indonesia, Universitas Airlangga, ROMINDO PRIMA VETCOM, Gadjah Mada University, Bogor Agriculture Institute, ILRI (International Livestock Research Institute), Kagoshima University, Miyazaki University, Faculty of Veterinary Science, Chulalongkorn University, College of Veterinary Medicine, Tarlac Agricultural University, Yamaguchi University, Erasmus MC, Kasetsart University, Wyndham Veterinary Clinic, GERBU – Germany, USAID <b>Published:</b> 29 November 2017 <b>ISSN:</b> 2413-0877</p>
<p>PUBLISHED</p> <ul style="list-style-type: none"><li>• 2017-12-03</li></ul>	

## TABLE OF CONTENTS

### **Toll-Like Receptors (TLRs) Play Role in Adaptive Immunity in Rabbits Immunized by *Sarcoptes scabiei* Proteins**

Pages 1-9 Nunuk Dyah Retno Lastuti, Fedik Abdul Rantam, Poedji Hastutiek, Dony Chrismanto

### **Preservation Effect of Grouper (*Epinephelus sp*) Fillet Against Survival of Anisakidae**

Pages 10-20 Hartanto M. Raharjo, Setiawan Koesdarto, A.T Soelih Estoepangestie, Kusuma Wardhani

### **Acanthocephalan in *Xenochrophis piscator* Snake in Sidoarjo Indonesia**

Pages 21-27 Inggarsetya Syah Audini, Lucia Tri Suwanti, Setiawan Koesdarto, E. Djoko Poetranto

### **The Identification Blood Parasites On Pig (*Susdomesticus*) In Polewali Mandar District**

Pages 28-33 Silvana Arfin, Lucia Muslimin, Adryani ris

### **Spirometra in *Ptyas mucosus* Snake in Sidoarjo, Indonesia**

Pages 34-40 Garindra Tiara Pranashinta, Lucia Tri Suwanti, Setiawan Koesdarto, E. Djoko Poetranto

**Teratogenic Effect of Congenital Toxoplasmosis in Chicken Embryo** ✓

Pages 696-602 Lucia Tri Suwanti, Mufasirin Mufasirin, Hani Plumeriastuti, Erma Safitri

**Case Study : Dystocia on Beef Cattle in Kunir Regency of Lumajang District, East Java, Indonesia in 2015 and 2016** ✓

Pages 603-608 Rosiana Febriani, Widya P Lokapirnasari, Tjuk I Restiadi, Imam Mustofa, Herry A Hermadi, Erma Safitri

**The Effectiveness of Honey in Physiological NaCl to Maintain of Viability and Motility of Spermatozoa** ✓

Pages 609-618 Elsa Agustina, Herry Agoes Hermadi, Hario Puntodewo S, Tatik Hernawati, Indah Norma Triana, Erma Safitri

**Utilization of Sumbawa Tropical Forest Honey Apis Dorsata to Improve Fertility of Indonesia Oriental Magpie Robin (Copsychus saularis) as Effort Animal Population Increase** ✓

Pages 619-626 Abdullah Hasib, Risaldi Muhamad, Talita Yuanda Reksa, Alvina Ulimaz Artha, Erma Safitri

**Utilization of Honey Apis dorsata as Antiosteoporosis on Requirements of Bone Calcium Ash Density on Ovariohysterectomized White Rat (Ratus norvegicus)**

Pages 627-632 Muhammad Huda Ramadhan Ibrahim, Abdullah Hasib Hasib, Siti Nur Rohmah, Salsabilla Abani, Samsi Yordan, Ira Sari Yudaniayanti

**Increased Integrity of Plasma Membrane and Acrosome Cap Spermatozoa Limousin Cattle at Post Thawing in Frozen Media by adding Seawater Extract**

Pages 633-641 Nur Faidah, Tatik Hernawati, Mirni Lamid, Ismudiono Ismudiono, Tri Wahyu Suprayogi, Sri Mulyati

**The Relation of Body Temperature and Vaginal Cytology Examination in Time Artificial Insemination Rate Fat-tailed Sheep (Ovis Aries) in The District Sidoarjo East Java**

Pages 642-649 Rhendyka Prasetya Anggriawan, Suzanita Utama, Hana Eliyani

**Effect of Laser Acupuncture Shoot on Ova Point of Male Mojosari Duck (Anas platyrhynchos) on The Number of Sertoli and Leydig Cells**

Pages 650-657 Yuanara A, Abdul Samik, Ira S Y/i, Tatang S Adikara, Eka P Hestianah, Suzanita Utama

**Insulin-Like Growth Factor-I (IGF-I) from Crossbred Pregnant Mare Serum to Increase Follicle Number of Mice (Mus musculus)** ✓

Pages 658-667 Abdullah Abdullah, Tjuk I Restiadi, Nunuk DR Lastuti, Tita Damayanti, Wurlina Wurlina, Erma Safitri

**Morphological Identification Nematode Tanqua tiara Found on Gastric Varanus salvatorat EastJava**

Pages 668-676 Alfiana Laili Dwi Agustin, Setiawan Koesdarto, Bambang Sektiari Lukiswanto, Lucia Tri Suwanti, Zainal Arifin, Emmanuel Djoko Putranto

## Conference Paper

# THE EFFECTIVENESS OF HONEY IN PHYSIOLOGICAL NaCl TO MAINTAIN OF VIABILITY AND MOTILITY OF SPERMATOZOA

Elsa Agustina<sup>1</sup>, Herry Agoes Hermadi<sup>2</sup>, Hario Puntodewo S<sup>3</sup>, Tatik Hernawati<sup>2</sup>, Indah Norma Triana<sup>2</sup>, and Erma Safitri<sup>2</sup>

<sup>1</sup>Student of Veterinary Medicine Faculty Surabaya, 60115, Indonesia

<sup>2</sup>Veterinary Reproduction Department of Veterinary Medicine Faculty, Surabaya, 60115, Indonesia

<sup>3</sup>Veterinary Public Health Department of Veterinary Medicine Faculty, Surabaya, 60115, Indonesia

## Abstract

This research was conducted to determine whether there is influence of various concentrations of mead physiological saline as diluent on the viability and motility fat-tailed sheep during storage 5°C. This research is compiled using the trial completely randomized design (CRD). Differences in the treatment of thinning the honey in physiological saline are PO (0%), P1 (0.2%), P2 (0.4%), P3 (0.6%), P4 (0.8%). The treatment stored in a refrigerator at a temperature of 5°C and then observed every 0, 24, and 48 hours. Data were analyzed by ANOVA followed by Duncan's Multiple Range Test. The conclusions obtained from this research there is the effect of adding various concentrations of NaCl physiological mead on the viability and motility fat tailed sheep during storage 5°C and conscentration mead NaCl physiological to maintain viability and motility best tail sheep are treated P3 (0.6% solution = 0.6 ml honey in 99.4 ml physiological NaCl) at the time T24 (24 hours).

**Keywords:** Fat-Tailed Sheep, Honey, Motility, Physiological NaCl, Spermatozoa, Viability.

## 1. Introduction

The efforts to increase local livestock production are necessary to support the demand for meat consumption. Fat-tailed sheep is one of the genetic resources of livestock that has the potential to be developed in the provision of meat. Sheep is one type of livestock that plays a large role in the life of farmers, because sheep are suitable for cultivation at the village level. Fat-tailed sheep have several advantages such as the ability to give birth to twins [1], fast adult age, good reproduction rate and high

Corresponding Author:

Erma Safitri

erma-s@fkh.unair.ac.id

Received: 03 October 2017

Accepted: 10 October 2017

Published: 29 November 2017

Publishing services provided  
by Knowledge E

© Elsa Agustina et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Selection and Peer-review under the responsibility of the VMIC Conference Committee.

 OPEN ACCESS

adaptability to the environment [2]. The population of sheep in Indonesia reaches 16,509.33 tails per year [3].

One of the many ways to increase livestock productivity is by introducing and implementing artificial insemination technology (AI). AI is a way of inserting spermatozoa into the female reproductive organs with a specific tool with the help of human, including semen collect, assessment, dilution, until the result of Artificial Insemination. AI technology in sheep and goats is still less widely applied in Indonesia. Due to various obstacles, such as timeliness, insemination techniques, and semen quality. The quality of semen is determined by several factors such as preservation techniques (storage-management) and the diluent composition used [4].

Semen dilution is an attempt to increase the volume of semen, reduce spermatozoa density and maintain the survival of spermatozoa up to a certain time under conditions of storage below or above freezing. Dilution and storage of semen is an attempt to maintain spermatozoa fertility over a longer period to extend the viability, motility, and fertility of spermatozoa [5].

Some commonly used diluents are egg yolks, milk, coconut water. Other ingredients are often used for the dilution of the semen NaCl solution. NaCl solution gives buffer properties (maintaining pH = 7) at room temperature and isotonic with liquid cell. Storage of semen with a physiologic NaCl diluent solution can only be used no more than 60 minutes after shelter because it lacks the energy source required by the spermatozoa. It is necessary to add other ingredients that provide energy or nutrients so as to prolong the viability of spermatozoa and spermatozoa motility in storage media [4].

The energy required by the spermatozoa is provided by simple sugars (monosaccharides). Honey contains simple sugars (monosaccharides) needed by spermatozoa to keep alive. Honey as an addition of energy ingredients or nutrients from physiologic NaCl diluents is expected to support the survival and movement of spermatozoa in the storage process [6].

Various studies have been conducted to determine the effect of adding honey to the diluents of various animal species such as to dilute semen etawa goat breed [7], comet fish [8], catfish [9], turkey [10]. Until now the effect of adding honey to dilute semen sheep has never been done. This study aims to determine the effect of adding various concentrations of honey in physiological NaCl to viability and motility of fat-tailed sheep spermatozoa during storage of 5 °C.

## 2. Materials and Methods

### 2.1. Design Research

The diluent used in this study was the honey diluent in physiological NaCl with various concentrations including P<sub>0</sub> (0%), P<sub>1</sub> (0.2%), P<sub>2</sub> (0.4%), P<sub>3</sub> (0.6%), and P<sub>4</sub> (0.8%) are repeated 4 times. Concentration of diluent solution in% units equivalent to ml / 100ml: P<sub>0</sub> (control) = 0 ml of honey in 100 NaCl Physiological; P<sub>1</sub> (0.2% solution) = 0.2 ml of honey in 99.8 ml NaCl Physiological; P<sub>2</sub> (0.4% solution) = 0.4 ml of honey in 99.6 ml NaCl Physiological; P<sub>3</sub> (0.6% solution) = 0.6 ml of honey in 99.4 ml NaCl Physiological; P<sub>4</sub> (0.8% solution) = 0.8 ml of honey in 99.2 ml NaCl Physiological.

Spermatozoa are collected using an artificial vagina. Spermatozoa that have been accommodated macroscopic and microscopic examination. Macroscopic observations were observed in volume, color, odor and pH of spermatozoa. Microscopic observations include determination of spermatozoa concentration, determination of live percentage (viability) of spermatozoa and motility of spermatozoa. Sperm dilution is done by using sperm ratio: diluent = 1: 9. Observations in every 0, 24, and 48 hours.

### 2.2. Design Research and Data Analysis

The design used Completely Randomized Design (RAL) with five treatments with four repetitions. The data obtained were analyzed using ANOVA (Analysis of Variant). If there is a significant effect then proceed with Duncan's test with a significant level of 5% [11].

## 3. Results

### 3.1. Fresh Sperm Quality of Fat-tailed sheep

Initial inspection of fresh semen quality is very important to know as the determination of semen feasibility to be processed further. The sheep semen to be diluted is first examined macroscopically and microscopically. Macroscopic examination results obtained an average volume of 0.8 ml of semen with a dense white color, the distinctive smell of sheep semen, semen pH 7 and the consistency of viscous semen. Microscopic examination of fresh sperm of fat-tailed sheep obtained average semen concentration of 1.380 million / MI (10<sup>6</sup>), with 95% spermatozoa viability, progressive



**Figure 1:** Test of Spermatozoa Viability (400X magnification). Live spermatozoa with white head (a) and dead spermatozoa with red head (b).

sperm motility (+++), and individual motility 80%. Based on the results of macroscopic and microscopic fresh sperm fat tailed sheep are still worthy of further research.

### 3.2. Spermatozoa Viability of Fat-tailed sheep

The percentage of viability is determined by the absorption of the eosin dye mixed in the sperm. Spermatozoa is calculated by calculating transparent (colorless) spermatozoa in the staining process with eosin-negrosin. The living spermatozoa will not be colored by the dye, while the dead spermatozoa on the head will be red-purple. Examination was observed using a microscope with 400X magnification. A total of 4 times a field of view.

The result of statistical analysis from variance analysis shows that the value of F test sig for diluent solution, storage time and also interaction between dilution solution and storage time is 0.00, because it is smaller than sig level. 0.05, it can be said that the effect of dilution dilution interaction and real time observation on viability of spermatozoa. Furthermore, Duncan Distance Test is used to find out which interactions provide the highest percentage of motility during storage.

The study showed that all parameters of semen quality observed showed a significant decrease in the quality of semen ( $p < 0,05$ ) after storage of  $T_0$ ,  $T_{24}$ , and  $T_{48}$ . This situation indicates that during processing and storage there are physical and biochemical changes of the spermatozoa used. The process of storing spermatozoa at 5°C causes a decrease in viability, because during storage at 5°C the spermatozoa retain metabolism to maintain life against cold. The metabolism of spermatozoa will produce a by-product of lactic acid which will be toxic to spermatozoa [12].

TABLE 1: Mean and Standard Deviation Percentage of Spermatozoa Viability of Fat-tailed Sheep.

Jam Ke-	Mean and Standard Deviation				
	Po	P1	P2	P3	P4
Ho	86.25 <sup>a</sup> ± 1.50	84.75 <sup>a</sup> ±3.68	85.50 <sup>a</sup> ±1.00	87.00 <sup>a</sup> ± 1.41	85.50 <sup>a</sup> ±1.00
H24	7.50 <sup>a</sup> ± 5.00	40.00 <sup>b</sup> ±8.16	48.25 <sup>c</sup> ±2.36	55.50 <sup>c</sup> ±3.31	53.75 <sup>c</sup> ±4.78
H48	0.00 <sup>a</sup> ± 0.00	5.00 <sup>a</sup> ±4.08	20.00 <sup>b</sup> ±7.07	27.50 <sup>b</sup> ± 6.45	22.50 <sup>b</sup> ±5.00

Different superscript in the same column shows a marked difference (p <0.05).




Figure 2: Mean and Standard Deviation Percentage of Spermatozoa Viability of Fat-tailed Sheep.

### 3.3. Spermatozoa Motility of Fat-tailed Sheep

The percentage of motility is a representation of progressive spermatozoa activity (forward motion). The ultimate goal of dilution is that it can be used for artificial insemination activities, the progressive spermatozoa becomes an absolute standard [13]. This examination was observed using a microscope with 400X magnification. This test is focused on observing the percentage of the number of active spermatozoa in each field of view. Observation of sperm motility is determined by the number of spermatozoa moving from a field of view and percentage.

Data Tables and figures show that on observation of To (clock o) there is no significant difference between treatment Po (83.25%), P1 (81.25%), P2 (82.50%), P3 (83.00%), and P4 (82.00%). Observation of T24 (24th hour) there was a significant difference between treatments (p <0.05). The highest motility percentage for T24 was obtained at treatment P4 (55%) which was significantly different with Po (3.75%). Observation of T48 (48th hour) there was a significant difference between treatments

TABLE 2: Mean and Standard Deviation of Spermatozoa Motility Percentage of Fat-tailed Sheep.

Jam Ke-	Mean and Standard Deviation				
	Po	P1	P2	P3	P4
Ho	83.25 <sup>a</sup> ± 3.94	81.25 <sup>a</sup> ± 2.50	82.50 <sup>a</sup> ± 2.88	83.00 <sup>a</sup> ± 3.55	82.00 <sup>a</sup> ± 4.00
H24	3.75 <sup>a</sup> ± 2.50	20.00 <sup>b</sup> ± 4.08	45.00 <sup>c</sup> ± 4.08	51.25 <sup>c</sup> ± 6.29	55.00 <sup>cd</sup> ± 7.07
H48	0.00 <sup>a</sup> ± 0.00	18.75 <sup>b</sup> ± 2.09	20.00 <sup>b</sup> ± 4.08	31.25 <sup>bc</sup> ± 2.50	36.25 <sup>c</sup> ± 7.50

*Superscript yang berbeda pada kolom yang sama menunjukkan perbedaan yang nyata (p < 0,05).*




Figure 3: Mean and Standard Deviation of Spermatozoa Motility Percentage of Fat-tailed Sheep.

(p < 0.05), with the highest percentage at P4 (36.25%) which was not significantly different with P3 (31.25%), but significantly different P2 (20.00%) and P1 (18.75%). At T48 the treatment of Po (control) had no motility.

#### 4. Discussion

The percentage of viability of spermatozoa on observation can still last up to 48 hours, but the result of viability at T48 40% observation so as not to meet the minimum standard of viability for AI. According to [14] the survival spermatozoa viability test is 50% minimum for AI. Spermatozoa are able to live after exit from the testes only in the range of 1-2 hours [15]. The average percentage of Fat-tailed sheep spermatozoa viability in this study had a live percentage of more than 50% in P3 treatment (55.5%) and P4 (53.75%) on T24 observations (Table 1). The process of spermatozoa storage is done by addition of diluent solution which can spermatozoa survive by providing



nutrients as an energy source so that spermatozoa functionality can be maintained [16] (Sunarma et al., 2007). The main ingredient used by spermatozoa as a source of energy from outside the testes is fructose that is converted into lactic acid and energy with the help of the fructolysin enzyme in the process of glycolysis [12]. Reference Labetubun and Shiva, (2011) states that reducing sugar can be metabolized by spermatozoa to produce energy in the form of ATP. Furthermore, spermatozoa utilize ATP as a source of energy in maintaining viability [17].

The good spermatozoa motility is seen by looking at the progressive motion of the spermatozoa. The ability of the spermatozoa propels itself forward because of the contractile substance in the central part of the spermatozoa which is passed on to the rest of the tail. Normal spermatozoa motility shows spermatozoa movements coming forward simultaneously caused by a tail motion that leads left and right. Rapid and powerful tail movements will be able to push the spermatozoa into the egg for fertilization or fertilization [18].

The storage of spermatozoa after dilution is carried out at 5°C, intended for spermatozoa metabolism to be minimized thus saving energy. The goal is to minimize spermatozoa metabolism so that when used spermatozoa still have enough energy to support during the trip within the female reproductive tract [20]. According to Susilawato et al., 1999 quoted from [20] the decline in spermatozoa motility rate is due to the reduced spermatozoa energy due to the ongoing metabolic processes and is influenced by the dilution of semen that can cause damage to the plasma membrane. This statement is supported by Beraden and Fuquay (1984) quoted from [20] that spermatozoa motility in semen storage has decreased due to the continuous metabolic processes running during storage resulting in reduced energy, resulting in sperm motility increasingly decreasing. Metabolism continually causes the accumulation of lactic acid which causes decreased motility [21].

Conditions when spermatozoa are outside the testes, spermatozoa need nutrients to survive. This study used honey as a nutrient in spermatozoa. Honey mainly contains glucose and fructose that are used as a source of energy for the survival and motility of spermatozoa. The continued decomposition of ATP into ADP in mitochondria will produce energy for sperm motility [22].

## 5. Conclusion

Based on the research that can be concluded: There is an effect of adding various concentrations of honey solution in physiological NaCl to viability and motility of fat-tailed sheep spermatozoa during storage at 5°C and concentration of honey solution in physiological NaCl which can maintain viability and motility of fat-tailed sheep spermatozoa. The best treatment was P<sub>3</sub> (0.6% soluble = 0.6 ml of honey in 99.4 ml of physiological NaCl) at the time of T<sub>24</sub> (24h).

## References

- [1] Noor, R.R., A. Djayanegara and L.Schuser. 2001. Selection to Improve Birth and Weaning Weight of Javanese Fat Tailed Sheep. Arch. Tier Dummerstorf 44(6):649-659.
- [2] Werdhani M. dan Lestari.L. 1996. Pengantar Peternakan di Daerah Tropis. Penerjemah: Darmadja. D.S.G.N Gajah Mada. Universitas Press. Yogyakarta.
- [3] Direktorat Jendral Peternakan dan Kesehatan Hewan. 2015. Populasi dan Produksi Peternakan Indonesia Tahun 2014-2015. <http://www.bps.go.id/>. Diakses 25 Mei 2016.
- [4] Rahardhianto A, N. Abdulgani, dan N. Trisyani. 2012. Pengaruh Konsentrasi Larutan Madu Dalam NaCl Fisiologis terhadap Viabilitas dan Motilitas Spermatozoa Ikan Patin (*Pangasius pangasius*) Selama Masa Penyimpanan. Institut Teknologi Sepuluh November. Surabaya.
- [5] Situmorang, P., E Triwulaningsih, A. Lubis., T. Sugiarti dan Caroline W. 2002. Optimalisasi Penggunaan Chilling Semen untuk Meningkatkan Persentase Kebuntingan Sapi Perah, Laporan Akhir T.A 2000/2001. Balai Penelitian Ternak.
- [6] Dea, L.B. 2015. The Effect of Haney to Motility and Viability Catfish (*Pangasius pangasius*) Spermatozoa. Faculty of Madicine, Lampung University.
- [7] Ekaninta, A. 2001. Pengaruh Lama Penyimpanan dan Media Pengencer Kuning Telur dengan Madu Terhadap Kualitas Sperma Kambing Peranakan Etawa (PE). Universitas Sumatra Utara. Medan.
- [8] Boedi,S.R.A., Mubarak, S., dan Rini, S. 2010. Penambahan Ekstender Madu dalam Proses Penyimpanan Sperma Beku Terhadap Motilitas dan Viabilitas Spermatozoa Ikan Komet (*Carassius auratus auratus*). Jurnal Ilmiah Perikanan dan Kelautan, 6(2): 63-70.
- [9] Arsetyo, R., Nurlita. A, dan N. Trisyani. 2012. Pengaruh Konsentrasi Larutan Madu dalam NaCl Fisiologis terhadap Viabilitas dan Motilitas Spermatozoa Ikan Patin

- (*Pangasius pangasius*) selama Masa Penyimpanan. Sains dan Seni ITS. Vol. 1: 58-63.
- [10] Wayan, B, Ni Made. D.P, dan Gusti.N.B.Trilaksana. 2015. Honey Improve the Quality of Semen Turkey During Storage. Universitas Udayana. Bali. Vol: 7 No: 2: 164-171.
- [11] (Kusriningrum, 2012)
- [12] Hidayaturrahmah. 2007. Waktu Motilitas dan Viabilitas Spermatozoa Ikan Mas (*Cyprinus Carpio L*) pada Beberapa Konsentrasi Larutan Fruktosa. Fakultas Matematika dan Ilmu Pengetahuan Alam. Universitas Lampung Mangkurat. Vol. 4 (2007) 9-18.
- [13] Solihati, N. dan P.Kune. 2009. Pengaruh Jenis Pengencer Terhadap Motilitas dan Daya Hidup Spermatozoa Semen Cair Sapi Simental. Universitas Nusantara Cendana. Kupang. <http://pustaka.unpad.ac.id/> [12 Mei 2016].
- [14] Toelihere
- [15] Jiabi, P. C. Chew, Z.A. Rashid, and R. Hassan. 2010. Application Of innovative Biotechnologies Regarding Aquaculture And fisheries Sector In Malaysia: Cryopreservation Programme. *Freshwater Fisheries Researchcenter*
- [16] Sunarma A. D.W.B. Hastuti, dan Y. Sistina. 2007. Penggunaan Ekstender Madu yang Dikombinaasikan dengan Kriprotektan Berbada pada Pengawetan Speraikan Nilem (Indonesian Sharkminnow, *Osteochilus Hasseltiivalenciennes*, 1842). Konferensi Aquaculture Indonesia (2007)
- [17] Solihati, N. 2008. "Studi Terhadap Kualitas dan Daya Tahan Hidup Spermatozoa Cauda Epididimidis Domba Garut Menggunakan Berbagai Jenis Pengencer," In *Proc. Seminar Nasional Teknologi Peternakan danVeteriner*, Puslitbang Peternakan, Bandung: 401-408.
- [18] Salisbury, C.V. and N.L. Van Demark. 1985. Fisiologi Reproduksi dan Inseminasi Buatan pada Sapi. Diterjemahkan oleh P.Djanuar. Gajah Mada University Press. Yogyakarta. 444-481.
- [19] Hernawati, M. 2005. Pengaruh Kuning Telur Ayam Buras dan Air Kelapa Muda dengan Konsentrasi yang Berbeda Terhadap Daya Hidup dan Motilitas Spermatozoa Domba. Skripsi. Fakultas Kedokteran Hewan, Universitas Airlangga Surabaya
- [20] Nuranti. 2005. Pengaruh Penambahan Heparin Pada Level yang Berbeda Terhadap Kualitas Semen Cair Kambing Boer Hasil Pemisahan Spermatozoa X dan Y. Fakultas Peternakan Universitas Hasanuddin. Makasar.
- [21] Wicaksono. A, dan Arifiantini R.I. 2009. "Uji Banding Empat Bahan Pengencer untuk Preservasi Semen Anjing Retriever," *Jurnal IlmuTernak dan Veteriner*, Vol.14, No 1:50-57.

- [22] Labetubun, J. dan Siwa, I.P. 2011. "Kualitas Spermatozoa Kauda Epididimis Sapi Bali dengan Penambahan Laktosa atau Maltosa yang Dipreservasi pada suhu 3-50C," *Jurnal Veteriner September*, Vol. 12 No. 3 Hal:200-207