

Barriers for multiparous women to using long-term contraceptive methods in Southeast Asia: case study in Philippines and Indonesia

by Agung Dwi Laksono

Submission date: 08-Nov-2022 06:26PM (UTC+0800)

Submission ID: 1948112829

File name: Artikel_Published.pdf (716.38K)

Word count: 6512

Character count: 33501

RESEARCH

Open Access



Barriers for multiparous women to using long-term contraceptive methods in Southeast Asia: case study in Philippines and Indonesia

Agung Dwi Laksono¹, Nikmatur Rohmah² and Hario Megatsari^{3*}

Abstract

Background: Multiparous women are supposed to be able to end their reproductive cycle to decrease population growth. This study aimed to analyze barriers for multiparous women to use long-term contraceptive methods (LTCM) in the Philippines and Indonesia.

Methods: The study population was women aged 15–49 years old who have given birth to a live baby > 1 in the Philippines and Indonesia. The weighted sample size was 12,085 Philippines women and 25,543 Indonesian women. To identify variables associated with the use of LTCM, we analyzed place of residence, age group, education level, marital status, employment status, and wealth status. The final step employed multinomial logistic regression.

Results: In both countries, the results showed that variables associated with non-user LTCM were younger women, living in rural areas with poor education. Women without partner and unemployed had higher probability to not use LTCM. Finally, low wealth status had a higher probability than the richest multiparous to not use LTCM.

Conclusion: The study concluded that there were six barriers for multiparous women to use LTCM in the Philippines and Indonesia. The six obstacles were living in rural areas, being younger, poor education, single, unemployed, and low wealth.

Keywords: Multiparous women, Parity, Long-term contraceptive methods, Contraception, Public health

Introduction

Population growth in Southeast Asia has outpaced growth in the U.S., U.K., and China in the last 5 years. Meanwhile, the population growth rate in Southeast Asia is 5.8%. Philippines population growth of 8.2% is the highest growth compared to other Southeast Asian countries. Indonesia's population grows by 5.9%, just below Malaysia's at 7.9% [1].

Southeast Asia's current population is 672,082,017, equivalent to 8.58% of the world's total population. Southeast Asia is ranked number 3 in the Asian region.

The population density in Southeast Asia is 154 per km², and 50% reside in urban areas. The mean age of the Southeast Asian population was 30.2 years. Based on population, Indonesia is ranked first in the Southeast Asia region and fourth globally, with 273,523,615 people. The Philippines is in second place in the Southeast Asia region and 13 globally, with 109,581,078 people. The rapid population growth in Indonesia and the Philippines indicates that the number of pregnancies that reach proper delivery in both countries is high [2, 3].

The number of pregnancies that achieve a viable delivery referred to as parity [4]. Women who have given birth more than two times are called multiparous and have given birth \geq four times called grand multiparous. The more often a woman gives birth, the more it has the

*Correspondence: hario.megatsari@fkm.unair.ac.id

³ Faculty of Public Health, University of Airlangga, Surabaya, Indonesia
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

potential to increase population growth. The crude birth rates in the Philippines and Indonesia are ranked 4th and 5th in the Southeast Asia region, namely 20.2 and 17.7 live births per 1000 population [5]. The family size in the Philippines also allows a woman to give birth more than two times. The Philippines' average ideal family size is 2.7 children for all women and 3.0 children for currently married women [6].

The high proportions of multiparous and grand multiparous cause population growth to occur rapidly. The distribution of currently married women in Indonesia is 25.57% multiparous and 4.27% grand multiparous. The remaining 62.58% were primiparous, and 7.57% never gave birth based on parity [7]. In some countries, the proportion of multiparous shows a higher number than primiparous. Among them are Nigeria, Uganda, Northwestern Ethiopia, and North Carolina [8–11]. This situation is different when we compare it to China. The proportion of multiparous women is 15.5%, primiparous 29.1, and 55.4% childless [12].

The utilization of LTCM is related to many factors, several studies showed that facts [13–15]. One of the factors is related to pregnancy, both planned and unplanned pregnancies. For example study conducted by Pasundani in 2020 [15] in Indonesia, age, education, parity, and information are all variables that have been linked to the usage of LTCM. Other study revealed the same messages that unplanned pregnancy was also factors for women utilize the LTCM [16].

Meanwhile, women who give birth more than two times have the risk of various complications of pregnancy and childbirth and even death. The risk of death from maternal causes is related to pregnancy risk and complications during birth [17]. The three causes of maternal death in the Philippines include complications of delivery, pregnancy-related hypertension, and postpartum hemorrhage [18]. Multiparous women have less time to recover their reproductive organs and less time to rest. Multiparous has an increasing burden as the number of children born. In general, the history of miscarriage and gestational diabetes was higher in the grand multiparous [19].

Previous studies found a significant association between grand multiparous and pregnancy outcome. Among them include cesarean section, fetal macrosomia, diabetes mellitus, hypertension in pregnancy caused [20]. Multiparity is still a threat to women's health problems. This threat is mainly related to the various complications that can occur during pregnancy and childbirth. Efforts to minimize this threat are to carry out pregnancy and birth control.

Long-term contraceptive methods (LTCM) are the most effective method for controlling pregnancy and

birth [21]. Including the LTCM are an intrauterine device (IUD), female sterilization, and implant/Norplant. LTCM can prevent unwanted pregnancy up to 20 times better than pills and vaginal rings [22]. LTCM is a form of birth control that is very effective and has the lowest maintenance costs (most economical) [9, 23]. Unfortunately, women attracted to LTCM but did not use it are more likely to be multiparous women [11]. Although LTCM is proven to reduce pregnancy effectively and is effective regardless of user compliance, its absorption is still low [24, 25].

A previous study in seven countries found that the widespread use of LTCM was still low, namely 16% [26]. IUDs and implants' use is still low, 1.8 and 10.4%, respectively [9]. Policymakers need to know more information about the various challenges in women's access to use LTCM. Moreover, we need to investigate this further. This study analyzes the obstacles for multiparous women to use LTCM in the Philippines and Indonesia based on the background description.

Materials and methods

Availability of data and materials

The author is unable to share the data publicly because a third party and the ICF, who own it, do not have permit to do so. The author requested The 2017 IDHS and 2017 PDHS data set to the ICF (data set of childbearing age women) and the ICF gave the access to confidential data through <https://dhsprogram.com>.

Data source

The report used secondary data as research materials from the 2017 Philippines Demographic and Health Survey (PDHS) and 2017 Indonesian Demographic and Health Survey (IDHS). The PDHS and IDHS are part of the Inner City Fund's (ICF) Global Demographic and Health Survey (DHS) survey collection. The sample population was women aged 15–49 years who had given birth in Indonesia to a live baby of over 1 (multiparous). The sampling procedure used stratification and multi-stage random sampling. The method yielded a weighted sample size of 12,085 Filipino women and 25,543 Indonesian women. The study period was from July 24th – September 30th 2017.

Data analysis

The dependent variable in this study was the use of long-term contraceptive methods (LTCM). In this study, LTCM is a modern contraceptive type consisting of IUD, female sterilization, and implant/Norplant. The study divides the use of LTCM into 2 categories, namely "Yes" and "No."

There were six independent variables involved in the analysis of this study. The six variables are a type of place of residence, age group, education level, marital status, employment status, and wealth status. The residence type consists of 2 categories, namely “urban” and “rural.” The urban-rural criteria refer to Statistics Indonesia. Age group is divided into 7 categories in 5 years, namely “15–19”, “20–24”, “25–29”, “30–34”, “35–39”, and “40–45”. Education level is the respondent’s recognition of the last diploma they have. Education level is divided into 4 categories, namely “no education”, “primary”, “secondary”, and “higher”. The study also divided marital status into two categories: “single” and “married/living with a partner.” The study divided into two categories for employment status, namely “unemployed” and “employed.”

The study determined wealth status based on the quintile of wealth owned by a household. Households were scored based on the number and type of items they have, from televisions to bicycles or cars, and housing characteristics, such as drinking water sources, toilet facilities, and primary building materials for the house’s floor. This score calculated using principal component analysis. National wealth quintiles were arranged based on household scores for each person in the household and then divided by the distribution into the same five categories, accounting for 20% of the population [27].

In the first step, the authors conducted a collinearity test. The action is to ensure no multicollinearity between the independent variables. Furthermore, the study carried out a bivariate test between the dependent and independent variables using the Chi-Square test. The study used the binary logistic regression to analyze barriers for multiparous women to use LTCM in Indonesia and Philippines in the final stage, and we used the significance $p < 0.05$. We used 95% confidence interval in the binary logistic regression. All stage statistical analyzes carried out using SPSS 22 software.

Ethical statement

The 2017 PDHS and the 2017 IDHS have received ethics approval from the national ethics commission at the Ministry of Health, Philippines, and Indonesia. The authors deleted the respondents’ identities from the dataset. Besides, respondents have provided written approval for their involvement in the study. The author has obtained permission to use Inner City Fund International’s data through its website: <https://dhsprogram.com/data/new-user-registration.cfm>.

Results

Table 1 showed the results of the co-linearity test. The analysis results displayed the tolerance value for all independent variables was > 0.10 , and the variance inflation

Table 1 The result of multicollinearity test

Independent Variables	Collinearity Statistics			
	Philippines (n = 12,085)		Indonesia (n = 25,543)	
	Tolerance	VIF	Tolerance	VIF
Type of place of residence	0.892	1.121	0.799	1.252
Age group	0.900	1.111	0.903	1.107
Education Level	0.710	1.408	0.745	1.343
Marital status	0.982	1.018	0.979	1.021
Employment status	0.928	1.077	0.955	1.047
Wealth status	0.629	1.590	0.659	1.517

Dependent variable: The use of LTCM

factor (VIF) value for all independent variables was < 10.00 . The results of this analysis informed that there were no symptoms of multicollinearity between the independent variables.

Descriptive analysis

Table 2 showed descriptive statistics of multiparous women in the Philippines and Indonesia. Based on the type of place of residence, Philippines multiparous women who live in rural areas dominated the two categories of the use of LTCM. Meanwhile, multiparous Indonesian women who use LTCM were dominated by those who live in urban areas. Meanwhile, multiparous Indonesian women who do not use LTCM were dominated by those who live in rural areas.

Based on the age group, women in the 40–44 age group occupied the multiparous Philippines and Indonesian women who use LTCM. On the other hand, the Philippines and multiparous Indonesian women who do not use LTCM are dominated by the younger age group, namely 35–39. According to the education level, the Philippines and multiparous Indonesian women with secondary education dominated the two categories of the use of LTCM.

Based on marital status, the Philippines and multiparous Indonesian women who are married/living with partners dominated the two categories of the use of LTCM. Meanwhile, based on employment status in both types, the use of LTCM is dominated by the Philippines and Indonesian employed women.

Finally, according to wealth status, multiparous Philippines women who use LTCM are dominated by the wealthy class with the lower category. On the other side, multiparous Philippines women who do not use LTCM are dominated by those with the most deficient wealth status category. Moreover, the wealthiest overwhelmed multiparous Indonesian women who use LTCM. On the

Table 2 Descriptive statistics of multiparous women in Philippines (n = 12,085) and Indonesia (n = 25,543)

Variables	The use of LTCM					
	Philippines			Indonesia		
	Yes	No	P	Yes	No	P
Type of place of residence			***0.000			***0.000
• Urban	46.2%	43.9%		53.2%	47.8%	
3 • Rural	53.8%	56.1%		46.8%	52.2%	
Age group			***0.000			***0.000
• 15–19	0.3%	0.5%		0.0%	0.1%	
• 20–24	4.7%	6.2%		1.3%	2.1%	
• 25–29	10.4%	15.0%		7.3%	9.2%	
• 30–34	16.1%	18.9%		16.5%	19.3%	
• 35–39	21.4%	21.4%		26.9%	24.6%	
• 40–44	25.6%	18.6%		28.0%	22.3%	
• 45–49	21.6%	19.5%		19.9%	22.4%	
Education Level			***0.000			***0.000
• No education	0.6%	1.4%		1.1%	2.6%	
• Primary	19.3%	21.3%		33.4%	40.9%	
• Secondary	52.6%	48.0%		48.8%	47.6%	
1 • Higher	27.5%	29.4%		16.8%	8.9%	
Marital status			***0.000			***0.000
• Single	5.4%	8.3%		1.5%	6.1%	
• Married/Living with partner	94.6%	91.7%		98.5%	93.9%	
Employment status			***0.000			***0.000
• Unemployed	45.0%	48.5%		39.2%	41.5%	
• Employed	55.0%	51.5%		60.8%	58.5%	
Wealth status			***0.000			***0.000
• Poorest	18.2%	23.2%		15.0%	19.0%	
• Poorer	23.0%	21.1%		17.3%	20.0%	
• Middle	19.7%	19.1%		19.5%	20.2%	
• Richer	21.5%	18.5%		20.3%	21.0%	
• Richest	17.7%	18.0%		27.8%	19.8%	

Note: ***p < 0.001

other hand, the more decadent wealthy status category dominated multiparous Indonesian women who do not use LTCM.

Multivariate analysis

Table 3 showed binary logistic regression results. In this final stage, the study involved all independent variables. The binary logistic regression uses “The use of LTCM: Yes” as references.

Table 3 showed that multiparous Philippines women living in rural areas are more likely to live in urban areas 1.070 times not to use LTCM (AOR 1.070; 95% CI 1.070–1.070). Meanwhile, multiparous Indonesian women living in rural areas were 1061 times more likely than those residing in urban areas not to use LTCM (AOR 1.061; 95% CI 1.061–1.062). This analysis indicates that living in

rural areas is a risk factor for multiparous women in the Philippines and Indonesia for not using LTCM.

3 According to the age group, Philippines multiparous women in the age group 35–39 and under have a higher likelihood of not using LTCM than the 45–49 age group. Philippines multiparous women in the 40–44 age group are less likely to use LTCM than the 45–49 age group. Meanwhile, multiparous Indonesian women with the age group 30–34 and under have a higher likelihood of not using LTCM than the 45–49 age group. Multiparous Indonesian women in the age group over 35–39 and above are less likely to use LTCM. This information shows that younger multiparous women in the Philippines and Indonesia have a higher risk of not using LTCM.

Based on the education level, multiparous Philippines women with no education have a higher chance of not using LTCM than those with higher education.

Table 3 The result of binary logistic regression of barriers for multiparous women to using LTCM in Philippines (n = 12,085) and Indonesia (n = 25,543)

Variables	Don't use LTCM					
	Philippines			Indonesia		
	AOR	95% CI		AOR	95% CI	
	LB	UB		LB	UB	
Place of residence: Urban	-	-	-	-	-	-
Place of residence: Rural	***1.070	1.070	1.070	***1.061	1.061	1.062
Age group: 15–19	***1.715	1.714	1.717	***2.360	2.356	2.365
Age group: 20–24	***1.467	1.466	1.467	***1.455	1.455	1.455
Age group: 25–29	***1.614	1.614	1.614	***1.231	1.231	1.232
Age group: 30–34	***1.317	1.317	1.317	***1.189	1.189	1.189
Age group: 35–39	***1.108	1.108	1.108	***0.900	0.900	0.901
Age group: 40–44	***0.806	0.806	0.806	***0.740	0.740	0.740
Age group: 45–49	-	-	-	-	-	-
Education: No education	***1.778	1.777	1.779	***3.966	3.964	3.967
Education: Primary	***0.951	0.950	0.951	***2.141	2.141	2.142
Education: Secondary	***0.800	0.800	0.801	***1.716	1.716	1.716
Education: Higher	-	-	-	-	-	-
Marital: Single	***1.699	1.699	1.699	***4.258	4.256	4.259
Marital: Married/Living with partner	-	-	-	-	-	-
Employment: Unemployed	***1.084	1.084	1.084	***1.044	1.044	1.044
Employment: Employed	-	-	-	-	-	-
Wealth: Poorest	***1.155	1.155	1.155	***1.134	1.134	1.134
Wealth: Poorer	***0.895	0.894	0.895	***1.148	1.148	1.148
Wealth: Middle	***0.960	0.959	0.960	***1.094	1.094	1.095
Wealth: Richer	***0.883	0.883	0.883	***1.172	1.172	1.172
Wealth: Richest	-	-	-	-	-	-

Note: *p < 0.05; **p < 0.01; ***p < 0.001

Multiparous Philippines women with primary and secondary education have a lower possibility of not using LTCM than those with higher education. On the other hand, multiparous Indonesian women with any education level have a higher probability of not using LTCM than those with higher education. This analysis indicates that inadequate education is a risk factor for multiparous women in the Philippines and Indonesia for not using LTCM.

According to marital status, the Philippines and multiparous Indonesian women with a single group (never in a union, widowed, divorced) have a higher probability than those with status married/living with partners not to use LTCM. The result shows that not having a partner is a risk factor for multiparous women in the Philippines and Indonesia for not using LTCM.

On the other side, unemployed multiparous women in the Philippines and Indonesia have a higher probability of not using LTCM than those employed. This analysis shows that the unemployed are a risk factor for

multiparous women in the Philippines and Indonesia for not using LTCM.

Finally, based on wealth status, the poorest Philippines multiparous women have a higher likelihood of not using LTCM than the richest one. Multiparous Philippines women with other wealthy groups were less likely not to use LTCM than the richest. On the other hand, multiparous Indonesian women with any wealthy status have a higher chance of not using LARC than the richest. This analysis indicated that low wealth status has a higher probability than the richest multiparous in the Philippines and Indonesia not to use LTCM.

Discussion

The study results found that living in rural areas was a risk factor for multiparous women in the Philippines and Indonesia for not using LTCM. Empirically, rural areas are often left behind compared to urban areas in development in all fields, including the health sector

[28]. This situation impacts the availability of information and services to access LTCM for multiparous women [29, 30].

The study found that younger multiparous women in the Philippines and Indonesia had a higher risk of not using LTCM. This finding is in line with studies in India [31]. This condition is likely because older multiparous women tend to want to stop their reproductive period. Meanwhile, those who are young still have the desire to have more children [32, 33].

The research found low education to be a risk factor for multiparous women in the Philippines and Indonesia for not using LTCM. Women with better education levels better understand their needs. Moreover, with better education, women also better understand the risk factors for any action or decision they take [34, 35], including better understanding the myths circulating about LTCM [36]. This reason is why multiparous women with higher education prefer to use LTCM [37–39]. Higher education levels have been found in several previous studies to positively determine health performance [38]. Conversely, several studies found poor education to be a barrier to higher-quality implementation in the health sector [40–42].

On the other hand, not having a partner was a risk factor for multiparous women in the Philippines and Indonesia for not using LTCM. In the context of countries with Eastern customs such as the Philippines and Indonesia, sexual activity for women who do not have a partner is a disgrace [43, 44]. This condition encourages women to control their sexual activity. This situation is likely to suppress the need for LTCM in unmarried multiparous women.

Meanwhile, the unemployed was a risk factor for multiparous women in the Philippines and Indonesia for not using LTCM. This situation may be due to employed multiparous women who prefer practical and effective contraceptive methods [45, 46]. Previous studies have shown the higher effectiveness of LTCM than non-LTCM methods [47]. On the other hand, LTCM is more effective for employed multiparous women than the forgetfulness factor caused by being busy. Other study revealed that women empowerment factors, one of it is employment, related to the utilization of LTCM [13].

Finally, the study found low wealth status to have a higher probability than the richest multiparous in Indonesia not to use LTCM. The previous studies saw poverty as a barrier to performance output in many health sectors, especially those related to and requiring costs [27, 48]. This situation also applies to poor multiparous women's access to LTCM [49, 50]. This cost is a barrier that should encourage the government to realize the financing mechanism through insurance for LTCM [38, 51, 52].

In summary, this research shows that six variables prevent a multiparous woman from using LTCM, namely: living in the rural area, younger age, poor education, single, unemployed and poor. The implication of this situation for a country is an uncontrolled population explosion, leading to repeated unfavorable conditions in society, including poverty, ignorance, and high unemployment [53]. Another negative situation is the risk of an unqualified future generation because their parents cannot optimally carry out child care [54].

Previous research has recommended strategies to increase the use of LTCM, including reducing costs to get LTCM services, increasing public understanding of LTCM [55], and increase the skill capacity of health workers [56]. We can develop a strategy by cooperating with relevant stakeholders to make the LTCM program a success in an area [57]. From the findings above, appropriate policy recommendations include: providing subsidies for LTCM services for the poor, initiating an insurance model for LTCM, educating the public massively, being innovative, creative, and measurable, and periodically updating the skills capacity of health workers.

Study limitation

This study has a limitation: the study's variable is limited and depends on the availability of secondary data (DHS data). However, the study has a positive impact on the maternal and child health programs in Indonesia, such as this study can be estimated to the national level with correct weight to describe the level of the problem at the national level. Furthermore this study can be used by the government to develop policy based on the scientific evidence, especially related to LCTM and how the strategy of the government to deliver it to the specific population in the country.

Conclusions

The research results concluded that there were six barriers for multiparous women to using LTCM in the Philippines and Indonesia. The six obstacles are living in rural areas, being younger, poor education, single, unemployed, and low wealth. Based on the findings of the research, the author recommend to the Philippines and Indonesia government to integrate the public health program and family planning program that can address to the rural areas, teenagers, low education, unemployed group and poor people.

Abbreviations

LTCM: Long-term contraceptive methods; IUD: Intrauterine device; PDHS: Philippines Demographic and Health Survey; IDHS: Indonesian Demographic and Health Survey; ICF: Inner City Fund; VIF: Variance inflation factor; AOR: Adjusted odds ratio.

Acknowledgments

The author would like to thank Inner City Fund International, who has agreed to allow the authors to analyze the 2017 PDHS and the 2017 IDHS data in this article.

Authors' contributions

ADL developed the proposal, analyzed and interpreted the data. NR and HM was a significant contributor in conducting the study, interpreting the data and writing the manuscript. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The authors cannot publicly share the data because a third party and authors do not have permission to share it. The 2017 IDHS data set name requested from the ICF data set of childbearing age women is available from the ICF contact <https://dhsprogram.com/data/new-user-registration.cfm> for researchers who meet the criteria for access to confidential data.

11

Declarations

Ethics approval and consent to participate

The 2017 IDHS has passed ethical clearance from the National Ethics Committee. The authors delete all respondents' identities from the dataset. Respondents have provided written approval for their involvement in the study through filling the informed consent. The author has obtained permission for the use of data for this study through the website: <https://dhsprogram.com/data/new-user-registration.cfm>. All methods were performed in accordance with the relevant guidelines and regulations.

13

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹National Research and Innovation Agency, The Republic of Indonesia, Jakarta, Indonesia. ²Faculty of Health Science, Muhammadiyah University of Jember, East Java, Indonesia. ³Faculty of Public Health, University of Airlangga, Surabaya, Indonesia.

Received: 20 March 2021 Accepted: 14 July 2022

Published online: 27 July 2022

References

- Pusparisa Y. Southeast Asia's rapid population growth (Pesatnya Pertumbuhan Populasi Asia Tenggara); 2019. p. 1–2.
- Worldometer. South-Eastern Asia Population (LIVE). 2020.
- Worldometer. Current world Population. 2020.
- Creinin M, Simhan H. Can we communicate gravidity and parity better? *Obstet Gynecol*. 2020;113(3):709–11.
- Moore M. Crude birth rate in Southeast Asia in 2019, by country (in births per 1,000 population); 2020.
- Philippine Statistics Authority and Inner City Fund. Key Findings from the Philippine National Demographic and Health Survey 2017. 2018. 1–20.
- Gafar A, Suza DE, Efendi F, Pramono AP, Susanti IA, Mishbahatul E. Determinants of contraceptive use among married women in Indonesia [version 1; peer review: 2 approved]. *F1000Research*. 2020;6(APR):1–9.
- Solanke BL, Banjo OO, Oyinloye BO, Asa SS. Maternal grand multiparity and intention to use modern contraceptives in Nigeria. *BMC Public Health*. 2018;18(1):1–15.
- Anguzu R, Sempeera H, Sekandi JN. High parity predicts use of long-acting reversible contraceptives in the extended postpartum period among women in rural Uganda. *Contracept Reprod Med*. 2018;3(1):1–7.
- Gashaye KT, Tsegaye AT, Abebe SM, Woldetsadik MA, Ayele TA, Gashaw ZM. Determinants of long acting reversible contraception utilization in Northwest Ethiopia: an institution-based case control study. *PLoS One*. 2020;15(10 October):1–19.
- Zerden ML, Tang JH, Stuart GS, Norton DR, Verbiest SB, Brody S. Barriers to receiving long-acting reversible contraception in the postpartum period. *Women's Heal Issues*. 2015;25(6):16–21.
- Luo Z, Gao L, Anguzu R, Zhao J. Long-acting reversible contraceptive use in the post-abortion period among women seeking abortion in mainland China: intentions and barriers. *Reprod Health*. 2018;15(1):1–9.
- Yaya S, Uthman OA, Ekholuenetale M, Bishwajit G. Women empowerment as an enabling factor of contraceptive use in sub-Saharan Africa: a multilevel analysis of cross-sectional surveys of 32 countries. *Reprod Health*. 2018;15(1):214. <https://doi.org/10.1186/s12978-018-0658-5>.
- Ontiri S, Ndirangu G, Kabue M, Biesma R, Stekelenburg J, Ouma C. Long-acting reversible contraception uptake and associated factors among women of reproductive age in rural Kenya. *Int J Environ Res Public Health*. 2019;16(9):1–11.
- Nita Adhani Pasundani KB. Determinant of the use of long-term contraceptive method: an analysis of 2017 Indonesian demographic health survey. *Indian J Public Heal Res*. 2020;11(5 SE):739–45 Available from: <https://medicopublication.com/index.php/jiphrd/article/view/9426>.
- Getahun DS, Wolde HF, Muchie KF, Yeshita HY. Utilization and determinants of long term and permanent contraceptive methods among married reproductive age women at Janamora district, Northwest Ethiopia. *BMC Res Notes*. 2018;11(1):836 Available from: <https://pubmed.ncbi.nlm.nih.gov/30477564>.
- World Health Organization. World health statistics overview 2019: monitoring health for the SDGs, sustainable development goals. Geneva; 2019.
- Papp L, Kleiman M. Maternal Morbidity and Mortality Why do Filipino Mothers Die? 2020. p. 1.
- Al-Shaikh GK, Ibrahim GH, Fayed AA, Al-Mandeel H. Grand multiparity and the possible risk of adverse maternal and neonatal outcomes: a dilemma to be deciphered. *BMC Pregnancy Childbirth*. 2017;17(1):1–7.
- Alsammami M, Ahmed S. Grand multiparity: risk factors and outcome in a tertiary hospital: a comparative study. *Mater Socio Medica*. 2015;27(4):244.
- Li L, Tran D, Zhu H, Balimane P, Willett G, Zhao P, et al. Use of model-informed drug development to streamline development of long-acting products: can these successes be translated to long-acting hormonal contraceptives? *Annu Rev Pharmacol Toxicol*. 2021;61:745–56.
- Ilmiyah DE, Budihastuti UR, Nurhaeni DA, Mudigdo A. Determinants of Women's choice of a long-acting reversible contraceptive in Boyolali. *Central Java J Matern Child Heal*. 2018;03(03):225–32.
- Olson EM, Kramer RD, Gibson C, Wautlet CK, Schmuhl NB, Ehrenthal DB. Contraception in Wisconsin. *WMJ*. 2019;117(4):149–55.
- Mazza D, Bateson D, Frearson M, Goldstone P, Kovacs G, Baber R. Current barriers and potential strategies to increase the use of long-acting reversible contraception (LARC) to reduce the rate of unintended pregnancies in Australia: an expert roundtable discussion. *Aust New Zeal J Obstet Gynaecol*. 2017;57(2):206–12.
- Garrett CC, Keogh LA, Kavanagh A, Tomnay J, Hocking JS. Understanding the low uptake of long-acting reversible contraception by young women in Australia: a qualitative study. *BMC Womens Health*. 2015;15(1):1–10.
- Faustmann T, Crocker J, Moeller C, Engler Y, Caetano C, Buhling KJ. How do women and health care professionals view hormonal long-acting reversible contraception? Results from an international survey. *Eur J Contracept Reprod Heal Care*. 2019;24(6):422–9.
- Wulandari RD, Qomarrudin MB, Supriyanto S, Laksono AD. Socioeconomic disparities in hospital utilization among elderly people in Indonesia. *Indian J Public Heal Res Dev*. 2019;10(11):1800–4.
- Laksono AD, Wulandari RD, Efendi F. Determinants of hospital utilisation among urban poor societies in Indonesia. *Int J Innov Creat Chang*. 2020;12(9):375–87.
- Seran AA, Antaria MD, Haksama S, Setijanangrum E, Laksono AD, Prahastuti Sujo AD. Disparities of the use of hormonal and non-hormonal contraceptive drugs in urban and rural areas in Indonesia and the world. *Syst Rev Pharm*. 2020;11(9):66–73.
- Seran AA, Laksono AD, Sujo AD, Masruroh II, Marasabessy N, et al. Does contraception used better in urban areas?: an analysis of the 2017

- IDHS (Indonesia demographic and health survey). *Syst Rev Pharm*. 2020;11(11):1892–7.
31. Yadav K, Agarwal M, Shukla M, Singh JV, Singh VK. Unmet need for family planning services among young married women (15–24 years) living in urban slums of India. *BMC Womens Health*. 2020;20(1):187.
 32. Yadav K, Agarwal M, Singh JV, Singh VK. Determinants of non-use of family planning methods by young married women (15–24 years) living in urban slums of Uttar Pradesh. *Indian J Community Heal*. 2017;29(1):55–66.
 33. Paul R, Huysman BC, Maddipati R, Madden T. Familiarity and acceptability of long-acting reversible contraception and contraceptive choice. *Am J Obstet Gynecol*. 2020;222(4):S884.e1–9.
 34. Wulandari RD, Laksono AD. The association of socioeconomic status on early marriage among women in rural Indonesia (Hubungan status ekonomi terhadap pernikahan dini pada perempuan di perdesaan Indonesia). *J Kesehat Reproduksi*. 2020;11(2):15–24.
 35. Onyowo UG, Namo IS. Effect of Early Marriage on Girl-Child's Further Education in Okpoku Local Government Area, Nigeria: Implications for Counseling. *KIU J Humanit*. 2019;4(1):65–72.
 36. Richards MJ, Coleman-Minahan K, Sheeder J. Long-acting reversible contraceptive attitudes and acceptability in adolescents and young adults: a key to patient-centered contraceptive counseling. *J Pediatr Adolesc Gynecol*. 2020;33(6):673–80.
 37. Aduloju OP, Akintayo AA, Adefisan AS, Aduloju T. Utilization of Long-Acting Reversible Contraceptive (LARC) Methods in a tertiary hospital in southwestern Nigeria: A Mixed Methods Study. *J Obstet Gynecol India*. 2021.
 38. Eeckhaut MCW, Sweeney MM, Feng L. Desire for sterilization reversal among U.S. females: increasing inequalities by educational level. *Perspect Sex Reprod Health*. 2018;50(3):139–45.
 39. Hayford SR, Kissling A, Guzzo KB. Changing educational differentials in female sterilization. *Perspect Sex Reprod Health*. 2020;52(2):17–27.
 40. Laksono AD, Wulandari RD. The Barrier to Maternity Care in Rural Indonesia. *J Public Heal From Theory to Pract*. 2020.
 41. Wulandari RD, Laksono AD, Rachmawaty T, Rukmini. Barriers to antenatal care utilization in Indonesia. *Medico-Legal Updat*. 2021;21(1):828–34.
 42. Rohmah N, Yusuf A, Hargono R, Laksono AD, Masruroh II, et al. Determinants of teenage pregnancy in Indonesia. *Indian J forensic. Med Toxicol*. 2020;14(3):2080–5.
 43. Himawan KK, Bambling M, Edirippulige S. What does it mean to be single in Indonesia? Religiosity, social stigma, and marital status among never-married Indonesian adults. *SAGE Open*. 2018;8(3).
 44. Himawan KK. Singleness, sex, and spirituality: how religion affects the experience of being single in Indonesia. *Ment Heal Relig Cult*. 2020;23(2):1–12.
 45. Bahamondes L, Fernandes A, Monteiro I, Bahamondes MW. Long-acting reversible contraceptive (LARCs) methods. *Best Pract Res Clin Obstet Gynaecol*. 2019;66:28–40.
 46. Raccach-Tebeka B, Plu-Bureau G. Long-acting reversible contraception. *Rev Prat*. 2018;68(4):387–91.
 47. Allen S, Barlow E. Long-acting reversible contraception. *Pediatr Clin N Am*. 2017;64(2):359–69.
 48. Wulandari RD, Putri NK, Laksono AD. Socioeconomic Disparities in Antenatal Care Utilisation in Urban Indonesia. *Int J Innov Creat Chang*. 2020;14(2):498–514.
 49. Beshar I, So J, Chelvakumar M, Cahill EP, Shaw KA, Shaw JG. Socioeconomic differences persist in use of permanent vs long-acting reversible contraception: An analysis of the National Survey of Family Growth, 2006 to 2010 vs 2015 to 2017. *Contraception*. 2021.
 50. Laksono AD, Matahari R, Wulandari RD. Factors related to the choice of contraceptive methods among the poor in Indonesia. *Syst Rev Pharm*. 2020;11(9):195–200.
 51. Krashin JW, Stuart GS, Garrett J, Spector H, Bryant AG, Charm S, et al. Contraception insurance coverage and receipt of long-acting reversible contraception or depot medroxyprogesterone acetate on the day of abortion. *Obstet Gynecol*. 2017;130(1):109–17.
 52. Fuerst MF, George KE, Moore JE. Long-Acting Reversible Contraception in Medicaid: Where Do We Go From Here? *Women's Heal Issues*. 2020.
 53. Gerland P, Raftery AE, Ševčíková H, Li N, Gu D, Spoorenberg T, et al. World population stabilization unlikely this century. *Science*. 2014;346(6206):234–237 Available from: <http://science.sciencemag.org/content/346/6206/234.abstract>.
 54. Laksono AD, Wulandari RD. "Children are assets": Meta synthesis of child values in the Lani and Acehese ("Anak adalah Aset": Meta Sintesis Nilai Anak pada Suku Lani dan Suku Aceh). *J Kesehat Reproduksi*. 2019;10(1):11–20.
 55. Shoupe D. LARC methods: entering a new age of contraception and reproductive health. *Contracept Reprod Med*. 2016;1(1):4. <https://doi.org/10.1186/s40834-016-0011-8>.
 56. Cleland J, Ali M, Benova L, Daniele M. The promotion of intrauterine contraception in low- and middle-income countries: a narrative review. *Contraception*. 2017;95(6):519–28 Available from: <http://www.sciencedirect.com/science/article/pii/S0010782417300975>.
 57. Kroelinger CD, Morgan IA, DeSisto CL, Estrich C, Waddell LF, Mackie C, et al. State-Identified Implementation Strategies to Increase Uptake of Immediate Postpartum Long-Acting Reversible Contraception Policies. *J Women's Heal*. 2018;28(3):346–56. <https://doi.org/10.1089/jwh.2018.7083>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions



Barriers for multiparous women to using long-term contraceptive methods in Southeast Asia: case study in Philippines and Indonesia

ORIGINALITY REPORT

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

10%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	ejournal2.litbang.kemkes.go.id Internet Source	1%
2	refubium.fu-berlin.de Internet Source	1%
3	wiredspace.wits.ac.za Internet Source	1%
4	Anu Rammohan, Niyi Awofeso, Renae C Fernandez. "Paternal education status significantly influences infants' measles vaccination uptake, independent of maternal education status", BMC Public Health, 2012 Publication	1%
5	Jia-li Ma, Li Jiang, Ping Li, Ling-ling He, Hongshan Wei. "Long-term outcomes of synchronous splenectomy and hepatectomy or radiofrequency ablation for hepatocellular carcinoma and esophagogastric variceal bleeding", Research Square Platform LLC, 2021	1%

6	oamjms.eu Internet Source	1 %
7	Li Li, Doanh Tran, Hao Zhu, Praveen Balimane et al. "Use of Model-Informed Drug Development to Streamline Development of Long-Acting Products: Can These Successes Be Translated to Long-Acting Hormonal Contraceptives?", Annual Review of Pharmacology and Toxicology, 2021 Publication	1 %
8	www.library.mcgill.ca Internet Source	1 %
9	mafiadoc.com Internet Source	1 %
10	www.ghspjournal.org Internet Source	<1 %
11	Sri Sumarmi, Agung Dwi Laksono. "Pregnancy Readiness in Married Female Adolescents from Different Ethnic Groups in Probolinggo District, Indonesia", Research Square, 2020 Publication	<1 %
12	nrl.northumbria.ac.uk Internet Source	<1 %
13	protocolexchange.researchsquare.com Internet Source	<1 %

- | | | |
|----|---|------|
| 14 | journals.lww.com
Internet Source | <1 % |
| 15 | Alfian Gafar, Dewi Elizadiani Suza, Ferry Efendi, Eka Mishbahatul Mar'ah Has, Ahmad Putro Pramono, Ika Adelia Susanti.
"Determinants of contraceptive use among married women in Indonesia", F1000Research, 2020
Publication | <1 % |
| 16 | dhsprogram.com
Internet Source | <1 % |
| 17 | repository.unair.ac.id
Internet Source | <1 % |
| 18 | Desy Nuri Fajarningtiyas, Dedik Sulistiawan, Margareth Maya Parulianta Naibaho, Riza Fatma Arifa. "Pattern and Determinant of Contraceptive Use among Women in Indonesia from 2007 to 2017: Evidence from Demographic and Health Survey", Open Access Macedonian Journal of Medical Sciences, 2021
Publication | <1 % |
| 19 | Kusse Urmale Mare, Ezana Abrha, Ebrahim Mohammed Yesuf, Setognal Birara Aychiluhm et al. "Factors affecting utilization of long-acting reversible contraceptives among sexually active reproductive-age women in | <1 % |

the pastoral community of Northeast
Ethiopia: A community-based cross-sectional
study", Women's Health, 2022

Publication

20

heanoti.com

Internet Source

<1 %

21

journals.rudn.ru

Internet Source

<1 %

22

Joana Raquel Raposo dos Santos. "Ciência de dados e políticas públicas de saúde: exemplos práticos", Universidade de Sao Paulo, Agencia USP de Gestao da Informacao Academica (AGUIA), 2020

Publication

<1 %

23

lje.org.pk

Internet Source

<1 %

24

www.scilit.net

Internet Source

<1 %

25

Amelo Bolka Gujo, Assefa Philipos Kare. "Utilization of Long-Acting Reversible Contraceptives and Associated Factors Among Reproductive Age Women Attending Governmental Health Institutions for Family Planning Services in Wondo Genet District, Sidama, National Regional State, Southern Ethiopia", Health Services Research and Managerial Epidemiology, 2021

<1 %

26

Chien-Ming Luo, Mu-Yang Hsieh, Chi-Hong Cheng, Chiu-Hui Chen, Min-Tsun Liao, Shao-Yuan Chuang, Chih-Cheng Wu. "Association of Frailty With Thrombosis of Hemodialysis Vascular Access: A Prospective Taiwanese Cohort Study", *American Journal of Kidney Diseases*, 2022

Publication

<1 %

27

Deni W. Kurniawan, Ilana R. A. Chertok, Zelalem T. Haile. "The Relationship Between Place of Birth and Early Breastfeeding Initiation in Indonesia", *Journal of Human Lactation*, 2020

Publication

<1 %

28

cmsproject.com
Internet Source

<1 %

29

pdffox.com
Internet Source

<1 %

30

www.rchiips.org
Internet Source

<1 %

31

www.science.gov
Internet Source

<1 %

32

Otobo I. Ujah, Russell S. Kirby. "Long-Acting Reversible Contraceptive Use by Rural–Urban Residence among Women in Nigeria, 2016–

<1 %

2018", International Journal of Environmental Research and Public Health, 2022

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Barriers for multiparous women to using long-term contraceptive methods in Southeast Asia: case study in Philippines and Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
