Register Login



Home Archives Announcements Publication Ethics Editorial Team

Author Guidelines About the Journal •

Search

Home / Archives / Vol 30 No 2 (2021): June



Published: 2021-06-30

Editorial

The importance of registry for systematic review and clinical trial

Vivian Soetikno 87-8





DOI: https://doi.org/10.13181/mji.ed.215620

Abstract views: 731 PDF downloads: 509 HTML downloads: 107 EPUB downloads: 200

Basic Medical Research

Analysis of SARS-CoV-2 nucleocapsid protein sequence variations in ASEAN countries

Mochammad Rajasa Mukti Negara, Ita Krissanti, Gita Widya Pradini 89–95

△ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.oa.215304

Abstract views: 1293 PDF downloads: 691 HTML downloads: 243 EPUB downloads: 224

Clinical Research

A systematic review of intracavernosal injection of mesenchymal stem cells for diabetic erectile dysfunction

Gampo Alam Irdam, Febriyani, Nur Rasyid, Akmal Taher 96–105

☐ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.oa.204475

Abstract views: 1063 PDF downloads: 592 HTML downloads: 150 EPUB downloads: 187

Efficacy and safety of platinum chain and gold weight implants for paralytic lagophthalmos: a systematic review

Yunia Irawati, Tjahjono Darminto Gondhowiardjo, Hardyanto Soebono 106-15

☐ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.oa.214683

Abstract views: 823 PDF downloads: 475 HTML downloads: 204 EPUB downloads: 200

Delivery routes in pregnancy with COVID-19 and the risk of intrapartum vertical transmission: a meta-analysis

Razmaeda Sarastry, Crismanto Layarta, Ussisti Aladini, Besari Adi Pramono 116-22

△ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.oa.214779

Abstract views: 1164 PDF downloads: 627 HTML downloads: 156 EPUB downloads: 170

Accuracy of pediatric advanced life support method for predicting the depth of endotracheal tube in Indonesian children

Andi Ade Wijaya Ramlan, Adhrie Sugiharto, Agus Mutakim 123–8

△ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.oa.203835

Abstract views: 783 PDF downloads: 468 HTML downloads: 152 EPUB downloads: 143

Treatment delay of cancer patients in Indonesia: a reflection from a national referral hospital

Soehartati Gondhowiardjo, Sugandi Hartanto, Sigit Wirawan, Vito Filbert Jayalie, Ida Ayu Putri Astiti, Sonar Soni Panigoro, Sri Mutya Sekarutami, Andhika Rachman, Adang Bachtiar 129-37

☐ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.oa.204296

Abstract views: 1230 PDF downloads: 633 HTML downloads: 224 EPUB downloads: 162

Impact of metabolic syndrome on lower urinary tract symptoms in patients with benign prostate hyperplasia

Dyandra Parikesit, Fiastuti Witjaksono, Chaidir Arif Mochtar, Nur Rasyid, Agus Rizal Ardy Hariandy Hamid 138-42

△ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.oa.214256

Abstract views: 1094 PDF downloads: 481 HTML downloads: 107 EPUB downloads: 147

Community Research

Reliability and validity of the Indonesian version of the World Health Organization quality of life-old (WHOQOL-OLD): a Rasch modeling

Sharon Gondodiputro, Guswan Wiwaha, Melly Lionthina, Deni Kurniadi Sunjaya 143-51

☐ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.oa.215065

Abstract views: 1389 PDF downloads: 652 HTML downloads: 192 EPUB downloads: 157

Case Report/Series

A case report of acute Marchiafava-Bignami disease: a rare clinical entity in chronic alcoholism

Mohd Arif Abdul Malik Khiew, Abdul Hanif Khan Yusof Khan, Wei Chao Loh, Liyana Najwa Inche Mat, Peck Kee Chia, Wan Aliaa Wan Sulaiman, Fan Kee Hoo, Hamidon Basri 152–6

☐ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.cr.204464

Abstract views: 1171 PDF downloads: 683 HTML downloads: 121 EPUB downloads: 142

Review Article

Neurological manifestations of COVID-19: a clinical approach

Ni Made Susilawathi, Kumara Tini, Ida Ayu Sri Wijayanti, Putu Lohita Rahmawati, Dewa Putu Wisnu Wardhana, Dewa Gde Purwa Samatra, Anak Agung Raka Sudewi 157–65

△ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.rev.204821

Abstract views: 1503 PDF downloads: 911 HTML downloads: 182 EPUB downloads: 157

Brief Communication

Gastrointestinal disorders in COVID-19 patients: a great imitator

Ari Fahrial Syam

166-9

☐ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.bc.204960

Abstract views: 2252 PDF downloads: 1376 HTML downloads: 131 EPUB downloads: 150

High visibility of outdoor tobacco advertisements around health facilities in East Java, Indonesia: a geospatial analysis

Hario Megatsari, Ilham Akhsanu Ridlo, Dian Kusuma

170-4

☐ PDF ☐ HTML ☐ EPUB

DOI: https://doi.org/10.13181/mji.bc.204177

Abstract views: 997 PDF downloads: 461 HTML downloads: 110 EPUB downloads: 172

Front & Back Matter

△ PDF

Abstract views: 449 PDF downloads: 219

Make a Submission

Current Issue



Top Reviewer 2021

Congratulations to all reviewers who are listed as the top 10 best reviewers of the year 2021. You are **rewarded to have a 50% discount** on your publication fee. Please email mji@ui.ac.id for further information.

FK UNSRI/ RS dr. Mohammad Hoesin Palembang, Indonesia

Irandi Putra Pratomo

Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Indonesia - Universitas Department of Physical Medicine and Rehabilitation, Faculty Indonesia Hospital, Depok, West Java, Indonesia of Medicine, Universitas Padjadjaran, Indonesia

Vitriana Riana Tamba

University of Indonesia, Indonesia

Thariqah Salamah

Department of Radiology, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia

Praseno Hadi

Department of Pulmonology and Respiratory Medicine,

Thank you for your contributions.

Information

For Readers

For Authors

For Librarians

Advertisement


















Tersedia di seluruh **UI Publishing Official Store**



the right source

Internet addiction: a new addiction? EC MOST

Cybersex addiction: an overview of the development and treatment of a newly emerging disorder

Stress among healthcare workers during the COVID-19 pandemic and the determinant factors: a

45

A case report of tuberculous constrictive pericarditis as a sole manifestation of tuberculosis in a male adolescent

42 Agus Rizal A. H. Hamid

Current condition of social security administrator for health (BPJS Kesehatan) in Indonesia: contextual factors that affected the national health insurance

42

ABSTRACTING & INDEXING

IGKAPNYA































































Link to Download

- ICMJE Conflicts of Interest Statement Form
- Final Checklist
- Title Page
- Original Articles
- Case Reports
- Reviewer Recommendation
- Cover Letter

• MJI reference style (Need to download Mendeley Desktop)

Current Issue



Our Social Media





Section Policies | Correction and Retraction Policies | Reviewer Acknowledgment | Contact Us |

Education Tower 6th Floor, IMERI FKUI, Salemba Raya Street No. 6, Kenari, Senen, Central Jakarta, 10430, Indonesia

Platform & workflow by OJS / PKP

Register Login



Home	Archives	Announcements	Publication Ethics	Editorial Team
		Author Guidelines	About the Journal 🕶	
			Search	

Home / Editorial Team

Editor-in-chief

<u>Agus Rizal A. H. Hamid</u>, Department of Urology, Faculty of Medicine Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus <u>h-index: 9</u>

Editor-in-chief Emeritus

<u>Isnani A.S. Suryono</u>, Dept. of Histology, Faculty of Medicine Universitas Indonesia, Indonesia; Scopus <u>h-index: 2</u>

Managing Editor

Felix F. Widjaja, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus h-index: 2

Editorial Board Members

<u>Agnes Kurniawan</u>, Department of Parasitology, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus <u>h-index: 20</u>

<u>Bambang Budi Siswanto</u>, Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Indonesia/National Cardiovascular Center Harapan Kita, Indonesia; Scopus <u>h-index: 12</u>

<u>Farrokh Habibzadeh</u>, Shiraz University of Medical Sciences, Shiraz, Iran, Islamic Republic of; Scopus <u>h-index: 14</u>

<u>Grace Wangge</u>, Southeast Asian Ministers of Education Organization Regional Center for Food and Nutrition (SEAMEO RECFON/PKGR), Universitas Indonesia, Indonesia; Scopus <u>h-index: 7</u>

Haku Hotta, Konan Women's University, Kobe, Japan; Scopus h-index: 40

<u>Hans-Joachim Freisleben</u>, German-Indonesian Medical Association (DIGM), Germany; Scopus <u>hindex: 24</u>

Hans-Jürgen Mägert, Anhalt University of Applied Sciences, Germany; Scopus h-index: 24

<u>Harrina E. Rahardjo</u>, Department of Urology, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus <u>h-index: 4</u>

<u>Inge Sutanto</u>, Department of Parasitology, Faculty of Medicine, Universitas Indonesia, Indonesia, Indonesia; Scopus <u>h-index: 21</u>

<u>Jeanne A. Pawitan</u>, Department of Histology, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus <u>h-index: 8</u>

Joerg Haier, Nordakademie University of Applied Sciences, Germany; Scopus h-index: 39

<u>Jose R.L. Batubara</u>, Department of Pediatric, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus <u>h-index: 3</u>

Knut Adermann, Niedersaichsisches Institutfuir Peptid-Forschung (IPF), Germany; Scopus <u>h-index:</u> 36

Laurentius A. Pramono, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus h-index: 5

Markus Meyer, Cardiorentis AG, Germany; Scopus h-index: 36

<u>Melva Louisa</u>, Department of Pharmacology and Therapeutics, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus <u>h-index: 10</u>

<u>Nia Kurniati</u>, Department of Pediatric, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus <u>h-index</u>: 12

<u>Pradana Soewondo</u>, Department of Internal Medicine, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus <u>h-index: 17</u>

<u>Rianto Setiabudy</u>, Department of Pharmacology and Therapeutic, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus <u>h-index: 11</u>

<u>Saleha Sungkar</u>, Department of Parasitology, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus <u>h-index: 4</u>

Sentot Santoso, Institute for Clinical Immunology and Transfusion Medicine, Germany; Scopus <u>h-index: 49</u>

<u>Sri W.A. Jusman</u>, Department of Biochemistry and Molecular Biology, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus <u>h-index: 5</u>

<u>Theddeus O.H. Prasetyono</u>, Department of Plastic Surgery, Faculty of Medicine Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus <u>h-index: 8</u>

<u>Vivian Soetikno</u>, Department of Pharmacology and Therapeutics, Faculty of Medicine, Universitas Indonesia, Indonesia; Scopus <u>h-index: 19</u>

<u>Wilfred C.G. Peh</u>, Department of Diagnostic Radiology, Khoo Teck Puat Hospital, Singapore, Singapore; Scopus <u>h-index: 36</u>

Associate Editors

Apriani Oendari, Center for Community Health Education Research and Service, Inc. (CCHERS) Boston, Massachusetts, United States;

Ary I. Savitri, Medical Journal of Indonesia, Indonesia; Scopus h-index: 4

<u>Aulia Rizka</u>, Department of Internal Medicine, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus <u>h-index: 2</u>

<u>Hariyono Winarto</u>, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus <u>h-index: 4</u>

<u>Dicky L. Tahapary</u>, Department of Internal Medicine, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Indonesia; Scopus h-index: <u>8</u>

Anna M. Singal, Department of Clinical Nutrition, Universitas Indonesia Hospital, Indonesia

Assistant Editors

Dania Clarisa, Medical Journal of Indonesia, Indonesia

Devita A. Prabowo, Medical Journal of Indonesia, Indonesia

Frisky N. Salvianny, Medical Journal of Indonesia, Indonesia

Novi A. Anggraeni, Medical Journal of Indonesia, Indonesia

Windu C. H. N. Suryaningrat, Medical Journal of Indonesia, Indonesia

Production Editor

Maytias Tri Pratiwi, Medical Journal of Indonesia, Indonesia

Layout Editors

Ilham Afriansyah, Medical Journal of Indonesia, Indonesia

Tsania Faza, Medical Journal of Indonesia, Indonesia

Make a Submission

Current Issue



Top Reviewer 2021

Congratulations to all reviewers who are listed as the top 10 best reviewers of the year 2021. You are **rewarded to have a 50% discount** on your publication fee. Please email mji@ui.ac.id for further information.

Nuswil Bernolian

FK UNSRI/ RS dr. Mohammad Hoesin Palembang, Indonesia

Irandi Putra Pratomo

Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Indonesia - Universitas Indonesia Hospital, Depok, West Java, Indonesia

Vitriana

Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Universitas Padjadjaran, Indonesia

Riana Tamba

University of Indonesia, Indonesia

Thank you for your contributions.

Information

For Readers

For Authors

For Librarians

Advertisement


















Tersedia di seluruh **UI Publishing Official Store**



the right source

Internet addiction: a new addiction? EC MOST

Cybersex addiction: an overview of the development and treatment of a newly emerging disorder

Stress among healthcare workers during the COVID-19 pandemic and the determinant factors: a

45

A case report of tuberculous constrictive pericarditis as a sole manifestation of tuberculosis in a male adolescent

Agus Rizal A. H. Hamid **42**

Current condition of social security administrator for health (BPJS Kesehatan) in Indonesia: contextual factors that affected the national health insurance

42

ABSTRACTING & INDEXING

IGKAPNYA































































Link to Download

- ICMJE Conflicts of Interest Statement Form
- Final Checklist
- Title Page
- Original Articles
- Case Reports
- Reviewer Recommendation
- Cover Letter

• MJI reference style (Need to download Mendeley Desktop)

Current Issue



Our Social Media





Section Policies | Correction and Retraction Policies | Reviewer Acknowledgment | Contact Us |

Education Tower 6th Floor, IMERI FKUI, Salemba Raya Street No. 6, Kenari, Senen, Central Jakarta, 10430, Indonesia

Platform & workflow by OJS / PKP

Brief Communication

High visibility of outdoor tobacco advertisements around health facilities in East Java, Indonesia: a geospatial analysis

Hario Megatsari, 1 Ilham Akhsanu Ridlo, 2 Dian Kusuma 3



pISSN: 0853-1773 • eISSN: 2252-8083 https://doi.org/10.13181/mji.bc.204177 Med J Indones. 2021;30:170-4

Received: September 03, 2019 Accepted: November 09, 2020 Published online: April 28, 2021

Authors' affiliations:

¹Department of Health Promotion and Behavior Sciences, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia, ²Department of Health Policy and Administration, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia, ³Centre for Health Economics & Policy Innovation, Imperial College Business School, London, United Kingdom

Corresponding author:

Dian Kusuma
Centre for Health Economics & Policy
Innovation, Imperial College London,
South Kensington Campus, London SW7
2AZ

Tel/Fax: +44-20-75895111 **E-mail:** d.kusuma@imperial.ac.uk

ABSTRACT

BACKGROUND Indonesian tobacco control initiatives are minimal despite having the second-highest adult male smoking prevalence in the world, with less than 10% of districts/cities banning outdoor tobacco advertisements. This research aimed to provide evidence on the presence of outdoor tobacco advertisements near health facilities in Surabaya where there is no outdoor advertising ban.

METHODS Data collection was carried out in Surabaya from October to November 2018. Data of government (public) and private health facilities were obtained from the city health office. Two spatial data analyses were carried out: a buffer analysis near the healthcare facilities and an advertisement hotspot analysis using ArcMap 10.6.

RESULTS From 308 tobacco advertisements that were identified, there were billboards (63%), banners (31%), and videoboards (7%). Of 142 public and 1,242 private health facilities in Surabaya, 26% and 31% had advertisements within 300 m and 63% and 70% were within advertisement hotspots, respectively. Furthermore, 5% of advertisements were within 300 m from public health facilities and 21% of them were within 300 m from private health facilities.

CONCLUSIONS Outdoor tobacco advertisements were widespread throughout the city, prominently around public and private health facilities.

KEYWORDS geospatial analysis, health facilities, Indonesia, tobacco advertisement, visibility

The Indonesian government has not signed and ratified the Framework Convention on Tobacco Control despite having the second-highest prevalence of adult male smoking in the world. Andorra, the Dominican Republic, Eritrea, Liechtenstein, Malawi, Monaco, Somalia, and South Sudan are the other eight countries that have not signed and ratified the treaty. Although the other eight countries have smaller populations (with Malawi having the largest population at 19 million), Indonesia has more than 260 million inhabitants and has contributed to about 61.4 million current tobacco users worldwide. There is no

improvement in this situation according to the latest national health survey. A survey showed that the total smoking prevalence among people aged 10 years old and above remained at 29% and increased by 26% (7.2% to 9.1%) among people aged 10-18 years old from 2013-2018 ²

National and local tobacco control efforts are limited compared with the World Health Organization's comprehensive six MPOWER initiatives.³ Less than 10% of the 514 districts/cities have issued outdoor tobacco advertisement bans with varying degrees of enforcement.⁴ Although advertisements are found in

Copyright @ 2021 Authors. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are properly cited. For commercial use of this work, please see our terms at https://mji.ui.ac.id/journal/index.php/mji/copyright.

10% of the districts that ban tobacco advertisements (e.g., Banyuwangi),4 the remaining 90% of the districts do not present the implementation of tobacco advertisement bans. Surabaya is the capital of the province of East Java and the second-largest city in Indonesia with a population of over 3 million in 2017. It was among the first in the country to implement smoke-free areas in selected facilities, including health facilities that ban indoor smoking since 2008.5 Unfortunately, effective tobacco control initiatives are currently falling behind. This research aimed to provide evidence of outdoor tobacco advertisements near government (public) and private health facilities in Surabaya.

METHODS

A spatial analysis was performed on the presence and clustering of outdoor tobacco advertisements near public and private health facilities in Surabaya. Advertisement and facility are the two main parameters. Over 250 registered roads and streets (according to mayor's Regulation Number 70/2010)6 were visited by motorcycles and cars from October to November to collect advertisement data. The following variables were used for the advertisement parameter: advertisement geographical coordinates (latitude and longitude), advertisement type (videoboard, billboard, and banner), product information (brand and product name), and picture. The geographical coordinates were obtained using Samsung Galaxy Note A6's (Samsung, South Korea) My Location feature.

Data for the facility parameter were obtained from the city health office (per January 2019) for both public and private facilities. The public health facilities were as follows: provincial health office (East Java), district health office (Surabaya), hospitals, public health centers (puskesmas), and puskesmas subclinics (pustu). On the other hand, private health facilities included the following: hospitals, primary clinics, specialty clinics, beauty clinics, pharmacies, drugstores, and laboratories. A pharmacy has a pharmacist whereas drugstores do not have one. The variables used were facility name, sector (public/private), and address. Google Sheets (Google, United States) with geocoding add-ons and ArcGIS (ESRI, United States) online were used to convert address details to geographic coordinates.

The analysis was carried using ArcMap 10.6 software (ESRI) with Open Street Map as the baseline map. The following ArcMap tools had been used: (a) a geoprocessing/buffering tool to create buffers around the health facilities (100 m, 200 m, and 300 m); (b) a spatial join tool to determine the number of facilities with at least one tobacco advertisement within the facility buffers; (c) spatial join and dissolve tools to determine the number of advertisements near the health facilities; and (d) optimized hotspot analysis tool to determine the hotspots with significant levels of 99%, 95%, and 90%. Hotspot analysis using Getis-Ord Gi* statistics to identify clusters⁷ is more common in infectious disease epidemiology than in tobacco control studies.8,9 The fishnet approach (dividing areas into squares) was used in the hotspot analysis




Figure 1. The presence of outdoor tobacco advertisement around health facilities in Surabaya (2018). (a) Buffers around the health facilities; (b) health facilities within outdoor tobacco advertisements hotspots. Yellow polygons are public hospitals; the green squares are public health centers (puskesmas); the circles around them are facility buffers at 100 m, 200 m, and 300 m; the red squares are outdoor tobacco advertisements. Cold spots/hotspots are areas with a significantly higher density of outdoor tobacco advertisements. Buffers were drawn, and hotspot analysis was conducted using ArcMap

because of the absence of a smaller boundary, such as the census area.

Most of the health facilities in our research were represented as a point on the map. The provincial health offices, city health offices, and hospitals were depicted as building polygons and had larger areas. We manually drew the polygon in ArcMap using Google Maps (Google) satellite view (see Figure 1 for public hospitals). The buffers were 100 m, 200 m, and 300 m from each polygon boundary.

RESULTS

Of 142 public health and 1,242 private health facilities, around 308 advertisements were identified: billboards (63%), banners (31%), and video boards (7%). Most of the public health facilities were composed of hospitals (16 or 11%) and public health centers (124 or 87%) whereas the private facilities were mainly composed of pharmacies (761 or 61%) and primary clinics (156 or 13%).

The presence and clustering of outdoor tobacco advertisements around health facilities are shown in Figure 1. The yellow polygons represent the public hospitals, and the green squares are the public health centers (puskesmas). The buffers at 100 m, 200 m, and 300 m are represented by the circles around the facilities. The results show that medium and large outdoor tobacco advertisements (represented as red squares on the map) were common across the city as shown in Figure 1a, with certain areas having a higher number of advertisements (indicative of clustering). The hotspot analysis results indicate that the middle section of the city has a significant number of advertisements as shown in Figure 1b, represented as red hotspot areas.

The number of health facilities with at least one tobacco advertisement within 100 m, 200 m, and 300 m around each facility is displayed in Table 1. A total of 37 public health facilities had advertisements within 300 m, ranging from one provincial/city health office to 16 health centers. In comparison, 388 private health facilities had advertisements within 300 m, ranging from 17 hospitals to 229 pharmacies. In terms of proportion of total facilities, 26% of public health facilities are near advertisements within 300 m, ranging

Table 1. Number of health facilities with at least one advertisement within the buffers

Health facility	Total facility	Number of facility with at least one advertisement, n (% of total)			Number of facility in hotspot* n (% of total)	
		100-m buffer	200-m buffer	300-m buffer	99% confidence	95% confidence
Government/Public						
Provincial health office	1	0 (0)	1 (100)	1 (100)	1 (100)	1 (100)
City health office	1	1 (100)	1 (100)	1 (100)	1 (100)	1 (100)
Hospitals	16	2 (13)	6 (38)	8 (50)	10 (63)	13 (81)
Puskesmas	63	1 (2)	7 (11)	16 (25)	32 (51)	37 (59)
Pustu	61	1 (2)	4 (7)	11 (18)	34 (56)	37 (61)
Total	142	5 (4)	19 (13)	37 (26)	78 (55)	89 (63)
Private						
Hospitals	43	2 (5)	7 (16)	17 (40)	24 (56)	27 (63)
Primary clinics	156	7 (5)	22 (14)	38 (24)	97 (62)	106 (68)
Specialist clinics	86	4 (5)	17 (20)	30 (35)	62 (72)	64 (74)
Beauty clinics	92	12 (13)	17 (19)	26 (28)	61 (66)	67 (73)
Pharmacy [†]	761	69 (9)	151 (20)	229 (30)	466 (61)	524 (69)
Drugstore [†]	59	7 (12)	19 (32)	30 (51)	44 (75)	47 (80)
Lab	45	5 (11)	14 (31)	18 (40)	29 (64)	33 (73)
Total	1,242	106 (9)	247 (20)	388 (31)	783 (63)	868 (70)

Puskesmas=public health centers, pustu=puskesmas subclinics

^{*}Hotspot analysis uses Getis-Ord Gi* statistics in ArcMap. Hotspots/red spots show a significant cluster of a higher number of tobacco advertisements; †a pharmacy has pharmacists and drugstores do not have one. Confidence levels of 99% and 95% show spatial statistical significance. Buffer and calculation were conducted using ArcMap

from 18% of auxiliary health centers (pustu) to 100% of provincial/city health offices. Similarly, almost 31% of private health facilities had advertisements within 300 m, ranging from 24% of primary clinics to 51% of drugstores.

The number of health facilities within the advertisement hotspots shown in Figure 1 is displayed in Table 1. These hotspots or red spots show a significant number of tobacco advertisements clustering at 95% and 99% confidence levels. Eighty-nine public health facilities within hotspot areas ranging from one provincial/city health office to 37 puskesmas/ pustu were identified using a 95% confidence level, whereas 868 private health facilities were identified within hotspot areas ranging from 27 hospitals to 524 pharmacies. In terms of proportion of total facilities, 63% of public health facilities were within the hotspot areas, ranging from 59% of puskesmas to 100% of provincial/city health offices, whereas 70% of private health facilities were within hotspot areas, ranging from 63% of hospitals to 80% of drugstores.

We also calculated the number of advertisements around each health facility (the results are not shown but are available upon request). There were 16 advertisements within 300 m of public health facilities, ranging from two advertisements around city health offices to 34 around all puskesmas, whereas there were 65 advertisements within 300 m of private health facilities, ranging from 24 advertisements near hospitals to 208 near pharmacies. In terms of proportion of total advertisements, 5% of advertisements were within 300 m of all public health facilities, ranging from 1% of advertisements around city health offices to 11% around all puskesmas, whereas 21% of advertisements were within 300 m of all private health facilities, ranging from 8% around all hospitals to 68% around all pharmacies.

DISCUSSION

research provides empirical evidence on the significant presence of outdoor tobacco advertisements around health facilities in Surabaya that are lack of comprehensive tobacco controls, such as advertisement bans. Our findings show that although the presence of advertisements was similarly high for both public and private health facilities in terms of proportion (e.g., 26% versus 31% of facilities had at least one advertisement within 300 m), the presence of advertisements was considerably higher around private health facilities in terms of number (e.g., 37 versus 388 facilities had at least one advertisement within 300 m). The number of young people being exposed to advertisements increases as the number of health facilities joining the national health insurance (Jaminan Kesehatan Nasional) also increases.

There is a significant presence of advertisements for both public and private hospitals and clinics based on the results of the study. Residents and visitors are exposed to tobacco advertisements across the city because the majority of public and large private hospitals also provide services to the neighboring districts. Patients and customers in almost 500 health centers and clinics (including health centers, private clinics, specialty clinics, and beauty clinics) are also exposed to tobacco advertisements (with many facilities within the hotspot areas). Tobacco advertisements have a significant presence in over 800 pharmacies and drugstores across the city, with about 40% of facilities having at least one advertisement within 300 m and 80% of both facilities within the hotspot areas. These results indicate a possible effect on health behaviors and outcomes. Smoking prevalence was relatively high in Surabaya according to the latest Basic Health Research (RISKESDAS) 2018. Around 9.6% of boys and 1.8% of girls are smoking among youths (13 and 14 years old) compared with the 10.2% and 0.2% national averages; among adults (15+ years), 53.7% of men and 0.4% of women are smoking compared with the 61.4% and 2.3% of national averages.¹⁰ Data from RISKESDAS also indicate that the burden of clinical risk factors and smoking-related illnesses was also comparatively high in Surabaya. The prevalence of adults (15+ years) with hypertension (systolic blood pressure of at least 140 mmHg or diastolic blood pressure of 90 mmHg) was 30.2% in Surabaya compared with the national average of 29.8%. Also, the prevalence of adults with diabetes mellitus (reported diagnosis by a doctor) was 4.4% in Surabaya compared with the national average of 1.8%.11,12

These data prove that Surabaya and other districts/ cities without advertisement ban policies should have laws to limit the presence of tobacco advertisements. Otherwise, adult and children who will access these health facilities will be exposed to outdoor tobacco advertisements. This would weaken public health programs, such as tobacco control measures. Further research should be conducted on small- and medium-

sized outdoor and point-of-sales advertisements because our study is limited to medium to large outdoor tobacco advertisements.4

In conclusion, the results of this research show a significant presence of outdoor tobacco advertisements in the city of Surabaya, with high visibility of advertisements around public and private health facilities including provincial and city health offices, hospitals, and clinics. This can influence the government's policy to reduce smoking in public areas.

Conflict of Interest

The authors affirm no conflict of interest in this study.

Acknowledgment

None.

Funding Sources

This study was funded by Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia.

REFERENCES

- World Health Organization. Factsheet 2018 Indonesia [Internet]. World Health Orgnization; 2018 [cited 2019 Jan 29]. Available from: https://apps.who.int/iris/bitstream/handle/10665/272673/ wntd 2018 indonesia fs.pdf.Ministry of Health of The Republic of Indonesia. Basic Health Research (RISKESDAS) 2018.
- Ministry of Health of the Republic of Indonesia: National Institute of Health Research and Development; 2018. Indonesian.

- World Health Organization. WHO report on the global tobacco epidemic 2008: the MPOWER package. Geneva: World Health Organization: 2008.
- Sebayang SK, Dewi DMSK, Lailiyah S, Ahsan A. Mixed-methods evaluation of a ban on tobacco advertising and promotion in Banyuwangi District, Indonesia. Tob Control. 2019;28(6):651-6.
- Surabaya City Government. City regulation number 5/2008 on smoke free area [Internet]. Surabaya; 2008 [cited 2019 Jan 29]. Available from: https://jdih.surabaya.go.id/pdfdoc/perda 99. pdf. Indonesian.
- Mayor of Surabaya. Mayor's regulation number 70/2010 on calculation of advertisement rent [Internet]. Surabaya; 2010 [cited 2019 Jan 29]. Available from: https://jdih.surabaya.go.id/ pdfdoc/perwali 530.pdf.
- Stopka TJ, Goulart MA, Meyers DJ, Hutcheson M, Barton K, Onofrey S, et al. Identifying and characterizing hepatitis C virus hotspots in Massachusetts: a spatial epidemiological approach. BMC Infect Dis. 2017;17(1):294.
- Lessler J, Azman AS, McKay HS, Moore SM. What is a hotspot anyway? Am J Trop Med Hyg. 2017;96(6):1270-3.
- Chowell G, Rothenberg R. Spatial infectious disease epidemiology: on the cusp. BMC Med. 2018;16:192.
- Hapsari D, Nainggolan O, Kusuma D. Hotspots and regional variation in smoking prevalence among 514 districts in Indonesia: analysis of Basic Health Research 2018. Glob J Health Sci. 2020;12(10):1-32.
- Adisasmito W, Amir V, Atin A, Megraini A, Kusuma D. Geographic and socioeconomic disparity in cardiovascular risk factors in Indonesia: analysis of the Basic Health Research 2018. BMC Public Health. 2020;20:1004.
- 12. U.S. Department of Health and Human Services. The health consequences of smoking—50 years of progress: a report of the surgeon general. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014.

KEMENTERIAN PENDIDIKAN, KEBUDAYAAN, RISET, DAN TEKNOLOGI

UNIVERSITAS AIRLANGGA

FAKULTAS KESEHATAN MASYARAKAT

Kampus C Jl. Dr. Ir. H. Soekarno, Mulyorejo, Surabaya 60115 Telp. (031) 5920948 Fax (031) 5924618 Laman: https://fkm.unair.ac.id, e-mail: info@fkm.unair.ac.id

SURAT KETERANGAN Nomor: 2679/UN3.1.10/KP/2023

Yang bertanda tangan di bawah ini:

Nama

: Dr. Santi Martini, dr., M.Kes

NIP

: 196609271997022001

Pangkat / Golongan

: Pembina/Gol. IV/a

Jabatan

· Dekan

Dengan ini menerangkan bahwa:

Nama

: Hario Megatsari, SKM., M.Kes.

NIP

: 198209122008011006

Pangkat / Golongan

: Penata (III/c) / TMT 01-04-2012

Jabatan

: Lektor

Telah melaksanakan penelitian dengan judul sebagai berikut :

No.	Judul Karya Ilmiah	Tahun pelasanaan Penelitian
1	High visibility of outdoor tobacco advertisements around health facilities in East Java, Indonesia: a geospatial analysis	2021

Adapun penelitian tersebut layak dilakukan, meskipun belum ada *Uji Etical Clearence* karena menggunakan titik GPS sebagai subyek penelitian dan menghasilkan output yang sangat baik.

Demikian surat keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional Lektor Kepala.

abaya, 31 Maret 20223

Dekan

Dr_{AT}Santi Martini, dr., M.Kes NIP 196609271997022001