



Volume 23 No. 2 July 2016

p-ISSN 0853-442X
e-ISSN 2355-1462

Indonesian Journal of Urology Jurnal Urologi Indonesia

URINARY NEUTROPHIL GELATINASE-ASSOCIATED LIPOCALIN AND CREATININE SERUM BPH PATIENTS WITH ACUTE URINE RETENTION TO DETECT KIDNEY FUNCTION DISORDERS

BODY MASS INDEX RELATIONS WITH THE DIFFICULTIES AND COMPLICATIONS OF PERCUTANEOUS NEPHROLITHOTOMY

CORRELATION BETWEEN BLOOD SERUM PSA LEVEL AND MMP-2 IN PROSTATE ADENOCARCINOMA

INITIAL EXPERIENCES WITH THREE RENAL ARTERIES OF KIDNEY ALLOGRAFT IN KIDNEY TRANSPLANTATION: A CASE SERIES

ETIOLOGY OF MALE INFERTILITY IN JAKARTA

DELAYED GRAFT FUNCTION FOLLOWING LAPAROSCOPIC LIVE DONOR NEPHRECTOMY: A MULTIVARIATE ANALYSIS

RELATION BETWEEN COMPLICATING FACTORS OF HYPOSPADIA AND COMPLICATIONS AFTER TIP: A RETROSPECTIVE STUDY

PERCUTANEOUS NEPHROSTOMY IN PATIENTS WITH OBSTRUCTIVE UROPATHY DUE TO MALIGNANCY: A SURVIVAL ANALYSIS

FACTORS ASSOCIATED WITH TUMOR RECURRENCE IN STAGE 1 BLADDER TRANSITIONAL CELL CARCINOMA IN SARDJITO HOSPITAL YOGYAKARTA

THE EFFICACY OF NASOGASTRIC TUBE IN A LARGE BLOOD CLOT EVACUATION DURING CYSTOSCOPY

MATURE RETROVESICAL TERATOMA IN AN ADULT MALE: A CASE REPORT AND LITERATURE REVIEW

SENSITIVITY AND SPECIFICITY OF URINALYSIS TO DIAGNOSE UTI IN PATIENTS WITH UROLITHIASIS AT SARDJITO GENERAL HOSPITAL

THE CORRELATIONS AMONG COMPONENTS OF METABOLIC SYNDROME, PSA, AND PROSTATE VOLUME IN BPH PATIENTS

Indonesian Urological Association

JURI

Vol. 23

No. 2

Page 67 - 139

July 2016

p-ISSN 0853-442X
e-ISSN 2355-1462

Accredited No. 81/DIKTI/Kep/2011

Editorial Team

EDITOR IN-CHIEF:

Dr. dr. Tarmono, SpU (K), (Department of Urology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital, Surabaya)

FOUNDER:

Prof. Dr. dr. Sunaryo Hardjowijoto, SpU(K), (Department of Urology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital, Surabaya)

SECTION EDITOR

dr. Fikri Rizaldi, SpU(K), (Department of Urology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital, Surabaya)

JOURNAL MANAGER

dr. M. Ayodhia Soebadi, SpU(K), (Department of Urology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital, Surabaya)

ETHICS EDITOR

Dr. dr. Besut Daryanto, SpU(K), (Division of Urology/Department of Surgery, Faculty of Medicine/Brawijaya University, Saiful Anwar General Hospital, Malang)

ASSOCIATE EDITOR

BASIC SCIENCE RESEARCH

dr. Agus Rizal A.H. Hamid, PhD, SpU(K), (Department of Urology, Faculty of Medicine/Universitas Indonesia, Cipto Mangunkusumo General Hospital, Jakarta)

CLINICAL RESEARCH

Dr. dr. Wahjoe Djatisoesanto, SpU(K), (Department of Urology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital, Surabaya)

TRANSLATIONAL RESEARCH

dr. Aaron Tigor Sihombing, SpU (K), (Department of Urology, Faculty of Medicine/Padjadjaran University, Hasan Sadikin General Hospital, Bandung)

EDITORIAL BOARDS

Prof. Dr. dr. Akmal Taher, SpU(K), (Department of Urology, Faculty of Medicine/Universitas Indonesia, Cipto Mangunkusumo General Hospital, Jakarta)

Prof. Dr. dr. Soetojo, SpU (K), (Department of Urology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital, Surabaya)

dr. I Wayan Yudiana, SpU, (Division of Urology/Department of Surgery, Faculty of Medicine/Udayana University, Sanglah General Hospital, Denpasar)

dr. Indrawarman, SpU, (Division of Urology/Department of Surgery, Faculty of Medicine/Gadjah Mada University, Sardjito General Hospital, Yogyakarta)

dr. Anugrah Dianfitriani, SpU, (KU Leuven-University of Leuven, Belgium)

INTERNATIONAL EDITORIAL BOARDS

Prof. Seung-June Oh, MD, (Department of Urology, Seoul National University College of Medicine, Seoul National University Hospital, Seoul, Korea)

Dr. Sanjay Balwant Kulkarni, MBBS, MS, FRCS (Glasgow), Dip. Urology (London), (Center for Reconstructive Urethral Surgery, 3 Rajpath Society, Paud Road, Pune, India)

Prof. Eric Chung, MBBS, FRCS (Urol), (Department of Urology, Princess Alexandra Hospital, University of Queensland, Australia)

Prof. Shigeo Horie, M.D, Ph.D, (Department of Urology, Jutendo Hospital, Tokyo, Japan)



ARTICLES

URINARY NEUTROPHIL GELATINASE-ASSOCIATED LIPOCALIN AND CREATININE SERUM BPH PATIENTS WITH ACUTE URINE RETENTION TO DETECT KIDNEY FUNCTION DISORDERS

Septina Rahayu, Sunaryo Hardjowijoto, Tarmono Djojodimedjo, Mohammad Ayodhia Soebadi

[PDF](#)
BODY MASS INDEX RELATIONS WITH THE DIFFICULTIES AND COMPLICATIONS OF PERCUTANEOUS NEPHROLITHOTOMY

Fauriski Febrian Prapiska, Sunaryo Hardjowijoto, Doddy M Soebadi, Tarmono Djojodimedjo

[PDF](#)
CORRELATION BETWEEN BLOOD SERUM PSA LEVEL AND MMP-2 IN PROSTATE ADENOCARCINOMA

Anny Setijo Rahaju, Aniek Meidi, Gondo Mastutik, Sjahjenny Mustokoweni, Arifa Mustika

[PDF](#)
INITIAL EXPERIENCES WITH THREE RENAL ARTERIES OF KIDNEY ALLOGRAFT IN KIDNEY TRANSPLANTATION: A CASE SERIES

Tommie Prasetyo, Nur Rasyid, Arry Rodjani

[PDF](#)
ETIOLOGY OF MALE INFERTILITY IN JAKARTA

Ari Basukarno, Ponco Birowo, Nur Rasyid

[PDF](#)
DELAYED GRAFT FUNCTION FOLLOWING LAPAROSCOPIC LIVE DONOR NEPHRECTOMY: A MULTIVARIATE ANALYSIS

Pande Made Wisnu Tirtayasa, Gerhard Reinaldi Situmorang, Arry Rodjani, Nur Rasyid

[PDF](#)
RELATION BETWEEN COMPLICATING FACTORS OF HYPOSPADIA AND COMPLICATIONS AFTER TIP: A RETROSPECTIVE STUDY

Johannes Aritonang, Arry Rodjani, Irfan Wahyudi

[PDF](#)
PERCUTANEOUS NEPHROSTOMY IN PATIENTS WITH OBSTRUCTIVE UROPATHY DUE TO MALIGNANCY: A SURVIVAL ANALYSIS

Prahara Yuri, Sungsang Rochadi

[PDF](#)
FACTORS ASSOCIATED WITH TUMOR RECURRENCE IN STAGE 1 BLADDER TRANSITIONAL CELL CARCINOMA IN SARDJITO HOSPITAL YOGYAKARTA

donny eka putra, Trisula Utomo

[PDF](#)
THE EFFICACY OF NASOGASTRIC TUBE IN A LARGE BLOOD CLOT EVACUATION DURING CYSTOSCOPY

Caesar Khairul Wallad, Jumadi Santoso, Kuncoro Adi

[PDF](#)
MATURE RETROVESICAL TERATOMA IN AN ADULT MALE: A CASE REPORT AND LITERATURE REVIEW

Vera Vera, Ferry Safriadi

[PDF](#)
Sensitivity and Specificity of Urinalysis to Diagnose Urinary Tract Infection (UTI) in Patients with Urolithiasis at RSUP Dr. Sardjito Hospital

Awang Dyan Purnomo, Untung Tranggono

[PDF](#)
THE CORRELATIONS AMONG COMPONENTS OF METABOLIC SYNDROME PSA AND PROSTATE VOLUME IN BPH PATIENTS

Yanuar Wicaksana Sumasta, Danarto Danarto

[PDF](#)

CORRELATION BETWEEN BLOOD SERUM PSA LEVEL AND MMP-2 IN PROSTATE ADENOCARCINOMA

¹Anny Setijo Rahaju, ¹Aniek Meidi, ¹Gondo Mastutik, ¹Sjahjenny Mustokoweni, ²Arifa Mustika.

¹Department of Anatomic Pathology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital Surabaya.

²Department of Pharmacology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital Surabaya.

ABSTRACT

Objective: This study aims to prove the correlation between Prostate Specific Antigen (PSA) blood level and Matrix Metalloproteinase-2 (MMP-2) expression in patients with prostate adenocarcinoma. **Material & method:** Prostate cancer patients' data from January 2009 to May 2012 were collected at the Department of Pathology, Soetomo General Hospital Surabaya. Data collected included patient medical documents, PSA blood examination, and histopathological examination. Histopathology slides and paraffin blocks of needle biopsies, Transurethral Resection of Prostate (TURP) and radical prostatectomy of prostate cancer patients were re-read, then the samples that met the inclusion criteria were stained by immunohistochemistry using antibodies MMP-2. **Results:** Data collection was done to obtain data samples of prostate cancer patients in 2009 to 2012 comprising as many as 22 patients between the ages of 52-91 years. Prostate adenocarcinoma in age of 70-79 was found in 8 patients, with a mean age of 68 years. PSA values obtained from medical documents were between 8.6-594.41 ng/ml. Spearman's test performed in this study showed a positive correlation (one-tailed) (correlation coefficient (r) 0.431, $p < 0.05$) between blood PSA level and MMP-2 expression in patients with prostate adenocarcinoma. **Conclusion:** Blood PSA level correlates positively with MMP-2 expression in prostate adenocarcinoma.

Keywords: Prostate specific antigen, matrix metalloproteinase-2, prostate adenocarcinoma.

ABSTRAK

Tujuan: Penelitian ini bertujuan untuk membuktikan hubungan antara kadar Prostate Specific Antigen (PSA) darah dengan Ekspresi Matrix Metalloproteinase-2 (MMP-2) pada penderita Adenokarsinoma Prostat. **Bahan & cara:** Data arsip penderita kanker Prostat dikumpulkan di Departemen/SMF/Instalasi Patologi Anatomi RSUD Dr. Soetomo Surabaya, mulai Januari 2009 sampai Mei 2012. Pengumpulan data meliputi dokumen medik penderita, data pemeriksaan PSA darah, dan data pemeriksaan Histopatologi. Slide Histopatologi dan blok parafin hasil biopsi jarum, Transurethral Resection of Prostate (TURP) dan Radikal Prostatectomy penderita kanker prostat dibaca ulang, kemudian sampel memenuhi kriteria inklusi dipulas secara imunohistokimia dengan menggunakan antibodi MMP-2. **Hasil:** Pengumpulan data yang dilakukan memperoleh data sampel penderita kanker prostat pada tahun 2009 sampai 2012 sebanyak 22 orang, yang terjadi antara usia 52-91 tahun. Adenokarsinoma prostat terbanyak terjadi pada usia 70-79 yaitu 8 orang, dengan rerata pada usia 68 tahun. Nilai PSA yang didapatkan pada dokumen medik antara 8.6-594.41 ng/ml. Uji Spearman yang dilakukan pada penelitian ini diperoleh hasil korelasi yang positif (1-tailed) (koefisien korelasi (r) 0.431, $p < 0.05$) antara kadar PSA darah dan ekspresi MMP-2 pada penderita adenokarsinoma prostat. **Simpulan:** Terdapat korelasi positif antara kadar PSA darah dengan ekspresi MMP-2 pada adenokarsinoma prostat.

Kata Kunci: Prostate specific antigen, matrix metalloproteinase-2, adenokarsinoma prostat.

Correspondence: Anny Setijo Rahaju; c/o: Department of Anatomic Pathology, Faculty of Medicine/Universitas Airlangga, Soetomo General Hospital Surabaya. Jl. Mayjen. Prof. Dr. Moestopo 6-8 Surabaya 60286. Phone: +62 31 5020251, 5030252, 5030253; Fax: +62 31 5026333. Mobile phone: 081803089442. Email: anny_setijorahaju@yahoo.com.

INTRODUCTION

Prostate cancer is a malignancy of the sixth most-common of all types of malignancies in the world, and becomes the second leading cause of

death from cancer in men caused by the incidence of metastasis.^{1,2} The incidence of prostate cancer in the United States amounted to about 29% of the incidence of all cancers and 10% of the causes of death.^{2,3}

Factors contributing to the clinical application of prostate cancer, among others, are the level of Prostate Specific Antigen (PSA) in serum, Gleason grade, tumor stage, and operation edge.⁴ These factors have been studied for the development of molecular prognostic markers in prostate cancer. Some literature suggests that malignant tumor cells produce some matrix that can degrade extracellular matrix (ECM), such as Matrix Metalloproteinase (MMP) family that plays a role in the invasion and metastasis of malignancy.⁵

PSA is produced by prostate gland luminal epithelial cells in the form of ductal and acinar, then secreted into the lumen of the gland and excreted along with semen within the seminal vesicles. In prostate cancer (adenocarcinoma) changes in luminal epithelial cells can penetrate and damage basal cells. Changes in luminal epithelial cells and damage in basal cells cause PSA out of the lumen and come into the blood circulation, resulting in increased concentration of PSA in the blood. The increase can be detected in the blood and used for post-therapy screening and evaluation of prostate cancer.^{1,2,6}

Adenocarcinoma is a type of prostate cancer that is most commonly found, and in such malignancy there is an imbalance between MMP and Tissue Inhibitor Metalloproteinase (TIMP), which led the invasion into the surrounding tissue and the incidence of metastasis, both lymphogenic and hematogenous.⁷⁻⁹ MMP-2 is a member of MMP family that has a special composition and function. MMP-2 (gelatinase A), in its function of invasion and metastasis, degrades collagen IV.¹⁰ The invasion and metastasis that occurs in prostate adenocarcinoma can be assessed from the expression of MMP-2 in tumor cells.¹¹

OBJECTIVE

The aim of this study was to prove the correlation between blood serum PSA level and MMP-2 in prostate adenocarcinoma, since studies on the correlation between blood serum PSA levels and MMP-2 expression in prostate adenocarcinoma is rare.

MATERIAL & METHODS

This study was an observational analytic study with cross-sectional approach conducted by collecting data from the archive on prostate cancer

patients whose diagnosis was established histopathologically as adenocarcinoma of the prostate in the Department of Pathology, Soetomo General Hospital Surabaya, from January 2009 to May 2012. Data collected were in the form of medical documents of the patients, PSA blood examination and histopathological examination data. Histopathology slides and paraffin blocks of needle biopsies, TURP and Radical Prostatectomy from prostate cancer patients were collected, and then re-reading was done to those cases. Samples taken were patients who had data on blood serum PSA examination by ELISA and met the inclusion criteria. Immunohistochemical staining was done by using MMP-2 antibodies to eligible blocks. Results were obtained by conducting analytical testing with Spearman's correlation by comparing the patients' blood PSA with MMP-2 expression in tumor cells. The study was conducted at the Laboratory of Pathology, Faculty of Medicine, Universitas Airlangga, from May to August 2015.

Immunohistochemical staining on paraffin blocks of patients with a diagnosis of prostate adenocarcinoma was performed using antibodies MMP-2. Assessment of MMP-2 expression was done by assessing the expression in the cytoplasm or tumor cell membrane with staining intensity assessment and the percentage of stained tumor cells. Intensity scores are 0 = tumor cells not stained; 1 = light yellow; 2 = brownish yellow and 3 = brown, while the percentage of cell number is multiplied. The determination of the degree of expression was follows: 0 (negative) when the score is 0; 1+ = 1-3; 2+ = 4-6; and 3+ = 7-9.¹² Data on PSA and MMP-2 expression was tested using Spearman rho correlation test, with significant results if $p < 0.05$.

RESULTS

Data collection was done to obtain data on samples of prostate cancer patients from 2009 to 2012 as many as 22 people in ages of 52-91 years. Prostate adenocarcinoma occurred mostly in the age of 70-79, comprising 8 patients, with a mean age 68 years. PSA values obtained from medical documents were between 8.6 - 594.41 ng/ml.

All samples showed positive MMP-2 expression in the cytoplasm and tumor cells membrane, with the percentage as much as 30-70% and > 70%. The intensity varied from light yellow to brown. Then Spearman tests conducted in this study showed a positive correlation (one-tailed) (correla-

tion coefficient (r) 0.431, $p < 0.05$) between blood PSA level and MMP-2 expression in patients with prostate adenocarcinoma.




Figure 1. Preparation T. 3045/12, percentage $> 70\%$ (3+) and intensity 1 MMP-2 intensity I score +




Figure 2. Preparation T. 1662/10 percentage $> 70\%$ (3+) and intensity 2.

Table 1. MMP-2 expression in prostate adenocarcinoma.

MMP-2 expression score	%	I	score
0	0	0	0
1	0	13	13
2	4	5	5
3	18	4	4
Total	22	22	22





Figure 3. Preparation T. 3606/11 and 6324/10 percentage $> 70\%$ (3+) and intensity 3.

Blood PSA value in the sample was between 8.6–594.41 ng/ml, showing an increase in serum PSA level. Some literature mentions that the normal level is < 4 ng/ml, showing an increase in blood PSA levels compared to normal.^{13,14}

DISCUSSION

The prostate is a retroperitoneal organ that surrounds the bladder neck and urethra, a gland shaped like a pear. Normal prostate gland is composed of two layers of cells: luminal epithelial cells and basal cells. Secretory/luminal cells have a

function to produce seminal fluid in the form of serine protease (prostate specific antigen/PSA).^{1,2} Prostate cancer is most commonly in the form of adenocarcinoma, which is an invasive malignant tumor derived from luminal epithelial cells. In adenocarcinoma luminal epithelial cells change and able to penetrate basal cells, resulting in basal cell damage. Changes in luminal epithelial cells and damage in basal cells cause the produced PSA out of the lumen and enters blood circulation, resulting in increased PSA concentration in blood. The increase can be detected in blood and is used for screening and evaluation of prostate cancer post-treatment as well as indications for biopsy in patients. PSA is also a marker that is often used to diagnose prostate cancer.^{1,2,6}

Murray in 2010 also stated that blood PSA value was positively related to the number of malignant epithelial cells stained with immunohistochemical staining with PSA antibody. Increasing number of malignant epithelial cells results in increased blood PSA level, indicating that more malignant cells that damage the barrier will lead to higher levels of PSA detected in the circulation.¹⁴

The process of cell integrity disruption causes the release of PSA in circulation. This integration disorder also occurs in hyperplasia, inflammation and tumors. PSA value in benign prostate tissue: 0.5 ± 0.4 ng/ml, in BPH: $0:31 \pm 0:25$ ng/ml, so that up to now there is no definite value in cancer.⁶

Zivkovic in 2004 obtain levels of 4.0-10.0 ng/ml in benign hyperplasia, prostatitis and prostate cancer. Murray conducted a study in 2010 by measuring the weight of prostate tissue, giving results that each gram of prostate cancer tissue can increase PSA of approximately 2.3 ng/ml. Hyperplastic prostate tissue is 10 times lower than cancerous tissue. In addition, they also found that the highest PSA level was in the lumen of the prostate gland, and between gland lumen and capillary blood vessel, basal cells in the gland, prostate stromal and capillary endothelial cells they found barrier that prevents PSA to circulate in the circulation. Therefore, PSA can increase drastically in the state of prostatitis, but can return to normal after the infection resolved.¹⁴

All samples of this study showed positive expression of MMP-2, according to a study by Trudel (2003), who found that 70% of malignant epithelial cells had positive expression of MMP-2

due to the degradation of ECM by MMP-2.¹⁵ There was a positive correlation between blood PSA levels with MMP-2 expression in prostate cancer, showing that increased blood PSA level is associated with increased MMP-2 expression in prostate cancer (correlation coefficient (r) 0.431, $p < 0.05$).

Malignant cells produce matrix that actively degrades with different specific substrates that act together in the degradation of Extracellular Matrix (ECM), that is the Matrix Metalloproteinase (MMP) family, which is associated with the invasion and metastasis of tumor cells. This tumor metastases is the main cause of death in patients with cancer, particularly prostate cancer.^{8,16}

Matrix Metalloproteinase (MMP), or matrix in, is a group of free endopeptidase structure that contain zinc, in cooperation with other proteolysis enzymes, such as cysteine proteinases, aspartic proteinase and serine proteinase, which plays a role in extracellular matrix degradation.^{8,9} MMP is secreted in the form of proenzyme, then be activated by the proteolytic gap, and this process can be inhibited by Tissue Inhibitor Metalloproteinase (TIMP) in order to avoid degradation. Imbalance between MMP and TIMP is one of the causes of invasion and metastasis that occurs in malignant tumors.⁷⁻¹⁰ Such imbalance is also an important factor that participates in tumor progression.¹⁷ The expression of MMPs (MMP-2 and MMP-9) is often associated with tumor aggressiveness and overall survival, and it is also often used as a marker of malignancy.¹⁸

Degradation by MMP in malignancy process is taking place through several stages, including: 1) assisting the formation of a micro-environment through the release of extracellular matrix growth factor; 2) assisting the process of tumor angiogenesis, and enhancing the ability of the migration and invasion of tumor cells; 3) playing a role in angiogenesis at the site of metastasis; 4) playing a role in the damage to basal membrane of blood vessel walls, allowing intravasation and extravasation of tumor cells; and 5) playing a role in shaping new microenvironment at metastasis site.¹⁹

Positive correlation between blood PSA levels with MMP-2 expression in prostate cancer showed that increased blood PSA level is associated with increased MMP-2 expression in prostate cancer (correlation coefficient (r) 0.431, $p < 0.05$). Increased PSA levels in blood indicates barrier damage caused by invasion through the degradation of collagen type IV, as well as damage caused by MMP-2.²⁰

According to Amalinei, Caruntu and Balan (2007), based on the substrate specificity, sequence similarity, and domain organization, MMP is divided into 6 groups: gelatinase (MMP-2, MMP-9); matrilysins (MMP-7, MMP-26); collagenases (MMP-1, MMP-8, MMP-13); stromelysin (MMP-3, MMP-10, MMP-11); membrane-type MMP (MT-MMP → MMP-14, MMP-15, MMP-17, etc.); and other MMPs (MMP-12, MMP-19, etc.).²⁰

Gelatinase-type MMP (MMP-2 and MMP-9) plays an important role because it can degrade collagen types IV, V, VII, X, XI, XIV, gelatin, elastin, proteoglycan core proteins, myelin basic protein, fibronectin, fibrillin-1, a precursor TNF-alpha and IL-1b. MMP-2 is able to break down collagen type I, which is the main component that forms the stroma and break down the molecular structure of collagen type IV which is a most protein constituent of basal membrane and the extra cellular matrix, which is instrumental in the invasion process.⁷⁻¹⁰ Similarly, Murray in 2009 found that the expression of MMP-2 plays a role in the spread of malignant cells through invasion and metastasis.¹⁴ MMP-2 expression (gelatinase A) in some malignant tumors has been widely studied. This relates to the prognostic value. Studies on the MMP-2 have also been started in prostate cancer (adenocarcinoma).

CONCLUSION

Blood PSA level correlates positively with MMP-2 expression in prostate adenocarcinoma and increased blood PSA can be used to predict ECM damage in MMP-2-induced prostate adenocarcinoma.

REFERENCES

- 1 Epstein, JI, Algaba, F, Allbrook, WC, Bastacky, S. Acinar adenocarcinoma, in WHO classification of tumours, pathology and genetics tumours of the urinary system and male genital organs, Ed. Ebk JN, IARC Press, Lyon; 2004. p. 161-92.
- 2 Rosai, J. Male reproductive system. In Rosai and Ackerman's surgical pathology. 9th Ed. Elsevier, China. 2011; 1: 1361-87.
- 3 Epstein, JI. The lower urinary tract and male genital system, in Robbins pathologic basic of disease, 8th Ed. Saunders: Philadelphia; 2010. p. 993-1003.
- 4 Young, RH. Atlas of tumors pathology, tumors of the prostate gland, seminal vesicles, male uretra, and penis, 3rd Series. Armed Forces Institute of Pathology: Washington DC; 2000. p. 111-200.
- 5 Brehmer B, Biesterfeld S, Jakse G. Expression of matrix metalloproteinases (MMP-2 and -9) and their inhibitors (TIMP-1 and -2) in prostate cancer tissue, Prostate Cancer and Prostatic Diseases 6; 2003. p. 217-22. doi:10.1038/sj.pcan.4500657.
- 6 <http://emedicine.medscape.com/article/457394-overview>
- 7 Nelson RA, Fingleton B, Rothenberg ML, Matrisian LM, Matrix metalloproteinases: Biologic activity and clinical implications, Journal of Clinical Oncology. 2000; 218(5): 1135-49.
- 8 Duffy MJ, Maguire TM, Hill A, McDermott E, O'Higgins N. Metalloproteinases: role in breast carcinogenesis, invasion and metastasis. Breast Cancer Res. 2000; 2(4): 252-7.
- 9 Verma RP, Hansch C. Matrix metalloproteinase (MMPs): chemical-biological function and (Q) SARs. Bioorg Med Chem. 2007; 15(6): 2223-68.
- 10 Tero Leinonen, Risto Pirinen, Jan Bohm. Increased expression of matrix metalloproteinase-2 (MMP-2) predicts tumour recurrence and unfavourable outcome in non-small cell lung cancer. In Histol Histopathol. 2008; 23: 693-700.
- 11 <http://atlasgeneticsoncology.org/Genes/MMP2ID4136ch16q13.html>
- 12 Daniel Kaemmerer, Luisa Peter, Emelie Lupp. Original Article: Comparing of IRS as immunohistochemical scoring schemes in gastroenteropancreatic neuroendocrine tumors. Int J Clin Exp Pathol. 2012; 5(3): 187-94
- 13 Stadiana Zivkovic. Original article: Correlation between prostate-specific antigen and histopathological difference of prostate carcinoma. In Arch Oncol. 2004; 12(3): 148-51.
- 14 Nigel P. Murray, Gloria M. Calaf, Leonardo Badinez. P504S expressing circulating prostate cells as a marker for prostate cancer. In Oncology Reports. 2010; 24; 687-92.
- 15 Dominique trudel, Yves fradet, Franc. Significance of MMP-2 expression in prostate cancer. An immunohistochemical study. In Cancer research. 2003; 63; 8511-15.
- 16 Yixuan Gong, Uma D. Chippada-Venkata, William K. Oh. Reviews: Roles of matrix metalloproteinase and their natural inhibitors in prostate cancer progression. Cancers. 2014; 6: 1298-327.
- 17 B Brehmer, S Biesterfeld, G Jakse. Prostate cancer and prostatic disease: Expression of matrix metalloproteinase (MMP-2 and -9) and their inhibitors (TIMP-1 and -2) in prostate cancer tissue. 2003; 6; 217-22.
- 18 Romano Oguic, Vladimir Mozetic, Eleonora Cini Tesar. Research article: Matrix metalloproteinase 2 and 9 immunoexpression in prostate carcinoma at the positive margin of radical prostatectomy specimen. In Hindawi Publishing Corporation Pathology Research International; 2014. Article ID 262195; 8.
- 19 <http://www.ncbi.nlm.nih.gov/gene/4313>.

20 Amalinei C, Caruntu ID, Balan RA. Biology of metalloproteinases. Romanian Journal of Morpho-

logy and Embryology. 2007; 48(4): 323-34.



INDONESIAN JOURNAL OF UROLOGY

UNIVERSITAS ABLANINGKA

P-ISSN: 10853442X E-ISSN: 23557402 Subject Area: Health

0.294118
Impact Factor

112
Google Citations

Sinta 2
Current Accreditation

[Google Scholar](#)
[Garuda](#)
[Website](#)
[Edit or URL](#)

History Accreditation

2010 2021 2022 2023 2024 2025


Garuda [Google Scholar](#)

CORRELATION BETWEEN MATERNAL AGE, LOW BIRTH WEIGHT, PREMATURITY AND HORMONAL DRUGS USING WITH THE OCCURRENCE OF HYPOSPADIAS (CASE STUDY AT BINA SEHAT JEMBER HOSPITAL, PARU JEMBER HOSPITAL, AND BHAWANGKARA BONDOWOSO HOSPITAL)

Indonesian Urological Association [Indonesian Journal of Urology](#) Vol. 29 No. 2 (2022)

[2022](#)
[DOI: 10.30421/ijur.v29i2.1432](#)
[Accred: Sinta 2](#)

Citations Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2017
Citation	112	82
h-index	5	4
i10-index	8	0



**KOMITE ETIK PENELITIAN KESEHATAN
RSUD Dr. SOETOMO SURABAYA**



**KETERANGAN KELAIKAN ETIK
(" ETHICAL CLEARANCE ")**

424 / Panke.KKE / VII / 2017

KOMITE ETIK RSUD Dr. SOETOMO SURABAYA TELAH MEMPELAJARI SECARA SEKSAMA RANCANGAN PENELITIAN YANG DIUSULKAN, MAKA DENGAN INI MENYATAKAN BAHWA PENELITIAN DENGAN JUDUL :

" Perbandingan Ekspresi MMP-2 dan MMP-9 dengan Kejadian Metastasis Tulang pada Adenokarsinoma Prostat "

PENELITI UTAMA : Anny Setijo Rahaju, dr., Sp.PA (K)

PENELITI LAIN : 1. Sjahjenny Mustokoweni, dr., Sp.PA (K), MIAC

2. Heriyawati, dr., Sp.PA

3. Aniek Meidy, dr., Sp.PA

4. Dr. Gondo Mastutik, drh., M.Kes

5. Dr. Arifa Mustika, dr., M.Kes

UNIT / LEMBAGA / TEMPAT PENELITIAN : RSUD Dr. Soetomo Surabaya

DINYATAKAN LAIK ETIK

SURABAYA, 14 JUL 2017
KETUA


(Dr. Elizeus Hanindito, dr., Sp.An, KIC,KAP)
NIP. 19511007 197903 1 002