

Volume 57 Issue 9 (September 2023)

ISSN: 0367-6722

Indian Journal *of* Animal Research



Celebrating



Years of Success

www.arccjournals.com



**AGRICULTURAL
RESEARCH
COMMUNICATION
CENTRE**



Download Cover

Editorial Board

Indian Journal of Animal Research

Chief Editor: K.M.L. Pathak

Print ISSN: 0367-6722

Online ISSN: 0976-0555

NAAS Rating: 6.43

SJR: 0.263

Impact Factor: 0.5 (2023)

Indexing Services :

Science Citation Index Expanded, BIOSIS Preview, ISI Citation Index, Biological Abstracts, Scopus, AGRICOLA, Google Scholar, CrossRef, CAB Abstracting Journals, Chemical Abstracts, Indian Science Abstracts, EBSCO Indexing Services, Index Copernicus

Volume 57 (2023)

Current Issue | Archive | Online First

Issue 9 (September 2023)

Issue 8 (August 2023)

Issue 7 (July 2023)

View All

Submit

In this Issue (24 Articles)

Article Type

Area of Research

Download Issue



Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1113-1119

Genetic Diversity of Six Duck Populations in India

P. Veeramani, R. Prabakaran, S.N. Sivaselvam, T. Sivakumar, S.T. Selvan, S.M.K. Karthickeyan

Article ID : B-4379 | doi : 10.18805/IJAR.B-4379 | Online First : 26-03-2021

Download Pdf

Download Citation

Search on Google

Connect with



Publish With US

Submit your manuscript through user friendly platform and acquire the maximum impact for your research by publishing with ARCC Journals.

Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1120-1125

Detection of Homologous Loci in Sheep and Cattle

Haifa El-Hentat, R. Aloulou, W. Derouich, O. Abidi, R. Omrane, N. Thamri, M. Hadhli

Article ID : BF-1556 | doi : 10.18805/IJAR.BF-1556 |

Online First : 09-12-2022





Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1126-1132

Partial Deoxygenation of Semen Extender Minimizes Post-thaw Damages and Improves Freezability of Crossbred Bull Spermatozoa

Lhendup Bhutia, Abhishek Kumar, Rahul Katiyar, Vinod Gupta, M. Ramamoorthy, S.K. Bhure, N. Srivastava, J.K. Prasad, S.K. Ghosh

Article ID : B-4377 | **doi :** 10.18805/IJAR.B-4377 | **Online First :** 12-04-2021

[Download Pdf](#)
[Download Citation](#)
[Search on Google](#)

Connect with

Become a Reviewer/Member

Join our esteemed reviewers panel and become an editorial board member with international experts in the domain of numerous specializations.



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1133-1138

Cryoprotective Activity of Glycerol and Ethylene Glycol: Dynamics with Canine Sperm Integrity

Kaushal Kusum, Ramakrishna Roy, Vijay B. Sharma, Raghavendra Prasad Mishra, Neha Singh

Article ID : B-4712 | **doi :** 10.18805/IJAR.B-4712 | **Online First :** 09-11-2021

[Download Pdf](#)
[Download Citation](#)
[Search on Google](#)

Connect with

Open Access


Filling the gap between research and communication ARCC provide Open Access of all journals which empower research community in all the ways which is accessible to all.

Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1139-1142

Seasonal Influence on Oocyte Recovery Rate, Quality and *in vitro* Maturation in Cows

Aiman A. Ammari, Ahmad R. Alhimaidi, Ramzi A. Amran, Muath G. Al Ghadi, Ahmed G. Rady



Products and Services

We provide prime quality of services to assist you select



Download Pdf

Download Citation

Search on Google

Connect with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1143-1145

Assessment of Frozen Semen Quality through Foldscope Microscopy- A Novel Application of Frugal Science to Reduce the Infertility Rate

P. Mooventhan, Mamta Choudhary

Article ID : B-4699 | doi : 10.18805/IJAR.B-4699 |

Online First : 22-02-2022

Download Pdf

Download Citation

Search on Google

Connect with



Support and Policies

Finest policies are designed to ensure world class support to our authors, members and readers. Our efficient team provides best possible support for you.

Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1146-1151

DNA Polymorphisms in 3'UTR and Intron-9 Region of PPARGC1A Gene and its Association with Milk Production Traits in Gaolao Cattle

Y.B. Sathe, D.S. Kale, Jaya Singh, D.V. Patil, P.G. Koringa, J.P. Korde, M.R. Jawale

Article ID : B-4791 | doi : 10.18805/IJAR.B-4791 | Online First : 04-03-2022

Download Pdf

Download Citation

Search on Google

Connect with



Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1152-1156

Gerontological Studies on the Micrometry of Liver of Bakerwali Goat



Download Pdf

Download Citation ▾

Search on Google

Connect
with



Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1157-1167

Recombinant Glycoprotein B of *Equine herpesvirus* Type 1 Elicits Protective Immune Response against Challenge in BALB/c Mouse Model

Alok Joshi, R.P. Gupta, Selvaraj Pavulraj, Bidhan Chandra Bera, Taruna Anand, Chinmoy Maji, Jyotsana Bhatt, Raj Kumar Singh, Bhupendra Nath Tripathi, Nitin Virmani

Article ID : B-4373 | **doi :** 10.18805/IJAR.B-4373 | **Online First :** 07-06-2021

Download Pdf

Download Citation ▾

Search on Google

Connect
with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1168-1176

A Clinical Study on the Use of Supracondylar Plate in the Treatment of Distal Femoral Fractures in Dogs

E. Pravalika, K. Jagan Mohan Reddy, C. Latha, T. Madhava Rao, G. Purshotham

Article ID : B-4742 | **doi :** 10.18805/IJAR.B-4742 | **Online First :** 25-04-2022

Download Pdf

Download Citation ▾

Search on Google

Connect
with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1117-1185

Morphological and Molecular Identification of Novel Green Peach Aphids (*Myzus persicae*) (Hemiptera: Aphididae) and



Sonia L. Shrivastava

Article ID : BF-1677 | doi : 10.18805/IJAR.BF-1677 |

Online First : 12-09-2023

Download Pdf

Download Citation ▾

Search on Google

Connect with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1186-1193

Comparison of Titanium Elastic Nailing and End-threaded Intramedullary Pinning for Distal Femoral Fractures in Young Dogs

Harmanpreet Singh Sodhi, Ashwani Kumar, Arun Anand, Vandana Sangwan, Opinder Singh

Article ID : B-4454 | doi : 10.18805/IJAR.B-4454 | Online First : 24-09-2021

Download Pdf

Download Citation ▾

Search on Google

Connect with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1194-1201

Metagenomic Study on the Influence of *Enterocytozoon hepatopenaei* (EHP) Infection on the Gut Microbiota in *Penaeus vannamei*

S. Ganesh Babu, A. Uma, K. Anbu Kumar, S.A. Shanmugam, A. Kathirvel Pandian

Article ID : B-5158 | doi : 10.18805/IJAR.B-5158 | Online First : 01-08-2023

Download Pdf

Download Citation ▾

Search on Google

Connect with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1202-1208



Eimeria ahsata in Northern Cyprus Sheep for the First Time

J.T. Olaifa, E. Ozgor

Article ID : BF-1560 | doi : 10.18805/IJAR.BF-1560 |

Online First : 18-11-2022

Download Pdf

Download Citation ▾

Search on Google

Connect
with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1209-1214

Therapeutic Efficacy of Ethanolic Extract of *Curcuma longa* and its Component, *Curcumin* against Experimental Cryptosporidiosis in Mice

Alveena Ganai, Anish Yadav, Rajesh Katoch, Dibyendu Chakraborty, Pawan Kumar Verma, Meenu Katoch, Amit Kumar

Article ID : B-4708 | doi : 10.18805/IJAR.B-4708 | Online First : 22-12-2021

Download Pdf

Download Citation ▾

Search on Google

Connect
with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1215-1220

Protective Effect of Resveratrol and Vitamin B17 on 8-iso-Prostaglandin F_{2α} and Raftlin-1 Levels in an Experimental Acute Urinary Retention in Rat

I. Demırhan, E.B. Kurutas

Article ID : BF-1548 | doi : 10.18805/IJAR.BF-1548 |

Online First : 03-10-2022

Download Pdf

Download Citation ▾

Search on Google

Connect
with





Article 2023): 1221-1229

Estimating Heterogeneity and Pooled Prevalence of Classical Swine Fever in Pigs in India: A Meta-analysis

P. Prema, Y.C. Bangar, Vijay Bahadur Sharma, Sanjay Kumar, Dinesh Kumar, Med Ram Verma

Article ID : B-4374 | doi : 10.18805/IJAR.B-4374 | Online First : 12-04-2021

Download Pdf

Download Citation ▾

Search on Google

Connect
with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1227-1231

Effect of Protected Calcium Salts of Fatty Acid-Amino Acid on Growth Rate of Young Ongole Crossbred Bulls

R. Antari, Mariyono, Y.N. Anggraeny, N.H. Krishna, A.S. Putri, M.F. Ulum, E. Wina

Article ID : BF-1429 | doi : 10.18805/IJAR.BF-1429 |

Online First : 14-02-2022

Download Pdf

Download Citation ▾

Search on Google

Connect
with



Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1232-1235

Objective radiographic evaluation of cardiac size in clinically normal Zaraibi goats

Ahmed I. Abdelgalil, Elham A. Hassan, Faisal A. Torad

Article ID : B-1097 | doi : 10.18805/ijar.B-1097 | Online First : 04-07-2019

Download Pdf

Download Citation ▾

Search on Google

Connect
with





Article 2023): 1230-1233

The Percentage of Male and Female *Argulus* Infesting Cyprinidae Fish in Magelang Regency, Central Java, Indonesia

Kismiyati, Alif Rizky Andika, Kusnoto

Article ID : BF-1510 | doi : 10.18805/IJAR.BF-1510 |

Online First : 23-08-2022

Download Pdf

Download Citation ▾

Search on Google

Connect
with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1240-1243

Efficacy of Paromomycin Sulphate for the Treatment of Cryptosporidiosis in Goat Kids

Mustafa Kabu, Ahmet Cihat Tunç, Mustafa Eser

Article ID : BF-1629 | doi : 10.18805/IJAR.BF-1629 |

Online First : 29-05-2023

Download Pdf

Download Citation ▾

Search on Google

Connect
with



Full Research Indian Journal Of Animal Research, Volume 57 Issue 9 (September

Article 2023): 1244-1250

The Effects of Hawthorn (*Crataegus oxyacantha*) Fruit Extract on Performance, Carcass and Some Serum Parameters of Broilers Reared Under Heat-Stress Conditions^[2]

Hacer KAYA

Article ID : BF-1657 | doi : 10.18805/IJAR.BF-1657 |

Online First : 18-08-2023

Download Pdf

Download Citation ▾

Search on Google

Connect
with





The Percentage of Male and Female *Argulus* Infesting Cyprinidae Fish in Magelang Regency, Central Java, Indonesia

Kismiyati¹, Alif Rizky Andika¹, Kusnoto²

10.18805/IJAR.BF-1510

ABSTRACT

Background: One cause of fish disease is the *Argulus* parasite. Both male and female *Argulus* were found to infect fish. This study aims to determine the percentage difference of male and female *Argulus* that infect Cyprinidae in Magelang Regency.

Methods: The present study used a survey method. The independent variables in this study are carp (*Cyprinus carpio*), goldfish (*Carassius auratus*), comet goldfish (*Carassius auratus*), koi fish (*C. carpio koi*) and the sex of the *Argulus* parasite. The dependent variable of this study was the sex percentage of the *Argulus* parasite. The data analysis was done using analysis of variance (ANOVA) then followed by Duncan's multiple range test to find out the differences between treatments.

Result: The first ANOVA result showed no significant difference ($p>0.05$), the second ANOVA showed significantly different results ($p<0.05$) and the third ANOVA reveals no significantly different percentage ($p>0.05$). The highest infestation rate of male *Argulus japonicus* is found in koi fish (*C. carpio koi*) is 60% and the lowest is in comet goldfish, which is 38.46%. Whereas infestation of female *A. japonicus* in carp, goldfish, comet goldfish and koi fish also obtained a similar result.

Key words: *Argulus*, Cyprinidae, Fisheries, Parasite.

INTRODUCTION

Magelang Regency is one of the areas that has undergone aquaculture-based development, prioritizing the principle of efficiency, quality and sustainability (Wibowo *et al.*, 2015). The area consists of Ngluwar Sub-District, Mungkid Sub-District and Muntilan Sub-District. Cyprinidae family fish species that are cultivated in Magelang Regency are common carp (*C. carpio*), goldfish (*C. auratus*), comet fish (*C. auratus auratus*) and koi fish (*C. carpio koi*) (Badan Pusat Statistik Kab. Magelang, 2014).

The main problem in cultivating fish in Indonesia to date revolves around parasites and infectious diseases. The disease causes economic losses because it can result in less optimal fish harvest (Carella and Sirri, 2017; Das and Chandra, 2018). One cause of the disease is the infestation of parasites (Picard Sánchez *et al.*, 2020). The quality of freshwater ornamental fishes decreased due to attacks from parasites such as *Argulus* sp. (Alifuddin *et al.*, 2002).

Argulus is a crustacean branchiuran parasite that causes severe problems in aquaculture throughout the world. Around 129 species of *Argulus* (family: Argulidae) are distributed worldwide and 12 species have been described in various freshwater, brackish water, marine and ornamental fish in India (Kumar *et al.*, 2017). Among them, *A. japonicus* are considered as emerging pathogens of freshwater, brackish water and coldwater fish worldwide (Tandel *et al.*, 2021).

Morphological identification of *Argulus* sp. is mostly based on distinguishing features of an adult male such as carapace and abdominal length or width, dorsal ridges of the carapace, respiratory areas, leg pigments, abdominal lobes and incision and the presence of a small coxal at the swimming appendages (Sahoo *et al.*, 2013; Soes *et al.*, 2010), requiring experienced taxonomists.

¹Department of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia.

²Department of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga, Indonesia.

Corresponding Author: Kismiyati, Department of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia. Email: kismiyati@fpm.unair.ac.id

How to cite this article: Kismiyati, Andika, A.R. and Kusnoto (2023). The Percentage of Male and Female *Argulus* Infesting Cyprinidae Fish in Magelang Regency, Central Java, Indonesia. Indian Journal of Animal Research. 57(9): 1236-1239. doi: 10.18805/IJAR.BF-1510.

Submitted: 04-03-2022 **Accepted:** 26-07-2022 **Online:** 23-08-2022

Argulus is one of the ectoparasites that attacks the Cyprinidae (Wardany and Kurniawan, 2014). *Argulus* attacks the fins, skin, gill and the entire surface of the host body (Pramujirini, 2016). Fish that has been infested by *Argulus* looks thin, with red spots appearing on its body, causing it often to rub its body on the edge of the pool. This parasitic attack is more often deadly in young fish because the body's defense system has not yet developed (Bandilla, 2007). Male and female *Argulus* usually attack carp (*C. carpio*) (Ebrahimi *et al.*, 2018). Male and female *Argulus* is also found to attack the goldfish (*C. auratus*). Male and female *Argulus* have the same properties as goldfish (*C. auratus*) (Yıldız and Kumantas, 2002). Based on these descriptions, this study aims to determine the percentage difference of *Argulus* male and *Argulus* females infesting carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*), as well as to find the amount of male and female *Argulus* infestation on the Cyprinidae family.

MATERIALS AND METHODS

Procedures

The research used the survey method. The survey method used in this study was a survey of research locations and *Argulus* parasites in fish samples. The data were collected using the descriptive method. The description of events in this study is the male and female *Argulus* parasite infestations in Cyprinidae of Magelang Regency. This study used a completely randomized factorial design. The completely randomized factorial design was applied because the study had two different factors; (1) the *Argulus* sex and (2) the Cyprinidae fish. The independent variables in this study consist of carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*) and the sex of *Argulus*. The dependent variable of this study was the sex percentage of the *Argulus* parasite. The control variables of this study were fish size, location and environmental conditions in Magelang. The collected sample amounted to 200 fish.

Data analysis

Analysis of the data used in this study was ANOVA (analysis of variance) using SPSS v16.0. If there are significant differences, further tests would be conducted using Duncan's multiple range test (Santoso, 2008).

RESULTS AND DISCUSSION

Identification results of *Argulus* sp. which infested in carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*) in Magelang Regency is *A. japonicus*. *A. japonicus* can be distinguished from other *Argulus* sp. by looking at the morphology. *A. japonicus* is identified as having a length of 3-5 mm and a width of 2-4 mm. In the Maxilla I, there is a supporting rod totaling five to nine pieces and the Maxilla II is equipped with three hooks.

The male and female *A. japonicus* can also be distinguished based on their morphology. The males are identified as having an abdominal testis. In comparison, the females are identified by their cephalothorax ovaries and seminal receptacle in the abdomen. *A. japonicus* are found in carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*). *A. japonicus* is one of the ectoparasites that attacks the Cyprinidae family (Wardany and Kurniawan, 2014). *A. japonicus* that infest in the Cyprinidae family in Magelang are observed in Fig 1.

The percentage of male *A. japonicus* infests carp (*C. carpio*) is 51.51%, while female *A. japonicus* is 48.49%. Male *A. japonicus* infest goldfish (*C. auratus*) is 50%, with the female having the same percentage. 38.46% of male *A. japonicus* infest comet goldfish (*C. auratus auratus*), with the female infestation percentage of 61.54%. 60% of male *A. japonicus* infests koi fish (*C. carpio koi*), while the female is 40%.

Data differences between the infestations of male and female *A. japonicus* in four different species of Cyprinidae were analyzed by analysis of variance (ANOVA). The results of the first ANOVA showed results that were not significantly

different ($p > 0.05$) between the average numbers of male and female *A. japonicus* that infest carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*). The second ANOVA showed significantly different results ($p < 0.05$) between carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*) which have been infected by *A. japonicus*. The infestations analyzed using the third ANOVA were not significantly different ($p > 0.05$) between the male and female *A. japonicus* females in carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*).

Based on the data from the analysis, the two treatments did not show any interaction. Thus, the data was included in the simple treatment. The simple treatment in question involves the male and female *A. japonicus* that infest carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*).

The test results show that male *A. japonicus* dominantly infests koi fish (*C. carpio koi*) and goldfish (*C. auratus*) but in a number that is not significantly different from carp (*C. carpio*). Male *A. japonicus* were least found in comet goldfish (*C. auratus auratus*), but in a number that is not significantly different from carp (*C. carpio*). In contrast, in the female infestation of *A. japonicus*, no differences were found between the four types of fish.

It has been identified that the *Argulus* sp. infest in carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*) in Magelang is *A. japonicus*. *A. japonicus* can be distinguished from other *Argulus* sp. by looking at the range of length (3-9 mm) and width of 2-6 mm (Møller, 2009). The respiratory area in the anterior is small, with the posterior being larger, five to nine supporting rods can be found in the Maxilla I and the Maxilla II is equipped with a total of three hooks. Male *A. japonicus* is equipped with testicles in the abdomen, whereas females have ovaries. The physical difference between the male and female *A. japonicus* can be seen in the abdomen located in the posterior part of the body (Kismiyati *et al.*, 2011). Female



Fig 1: *Argulus japonicus* infested common carp (*C. Carpio*), goldfish (*C. auratus*), comet fish (*C. auratus auratus*) and koi fish (*C. carpio koi*). a. Female *Argulus japonicus*; b. Male *Argulus japonicus*.

A. japonicus has spermatheca and ovaries, while males have seminal testicles and vascular (Wardany and Kurniawan, 2014).

The percentage of male *A. japonicus* infests carp (*C. carpio*) is 51.51%, while female *A. japonicus* is 48.49%. Both male and female *A. japonicus* are known to infect fish (Walker *et al.*, 2011). The percentage of male and female *A. japonicus* found to infest carp (*C. carpio*) is almost the same. That is because carp (*C. carpio*) is one of the preferred hosts of both male and female *Argulus japonicas* (Poly, 2008).

The male and female *A. japonicus* that infest goldfish (*C. auratus*) have the same percentage of 50%. Both male and female *A. japonicus* were found to infest goldfish (*C. auratus*) (Wafer *et al.*, 2015). That is because of their same parasitic properties (Mikheev *et al.*, 2015). 38.46% of male *A. japonicus* infects comet goldfish (*C. auratus auratus*), with the female infestation percentage of 61.54%. Female *A. japonicus* is found in comet goldfish fins (*C. auratus auratus*). This is due to the wide surface and slow movement of the fins (Pramujirini, 2016). The slow-motion of fish fins makes it easy for female *A. japonicus* to break away when oviposition (Kismiyati *et al.*, 2011).

There are 60% of male *A. japonicus* infest koi fish (*C. carpio koi*) while the female is 40%. Male *A. japonicus* can be found on the surface of koi fish (*C. carpio koi*). Koi fish (*C. carpio koi*) has a broad body surface that becomes the preferred predilection for male *A. japonicus*. Male *A. japonicus* prefers large areas (Taylor *et al.*, 2006).

The average number of male and female *A. japonicus* that infest carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*) is not significantly different ($p > 0.05$). That is because of their same parasitic properties (Mikheev *et al.*, 2015). Male and female *A. japonicus* are found to attack the Cyprinidae family (Wardany and Kurniawan, 2014).

The test results show that male *A. japonicus* dominantly infests koi fish (*C. carpio koi*) and goldfish (*C. auratus*) but in a number that is not significantly different from carp (*C. carpio*). Male *A. japonicus* least infests comet goldfish (*C. auratus auratus*), but in a number that is not substantially different from carp (*C. carpio*). That is because all four types of fish have other body surface areas. Male *A. japonicus* favors large areas (Taylor *et al.*, 2006). Duncan's Multiple Range Test results also showed that no differences were found between the four types of fish infested by female *A. japonicus*. Female *A. japonicus* is often found in fish fins (Pramujirini, 2016). The female chooses fins as a place of predilection because the fin movements of carp (*C. carpio*), goldfish (*C. auratus*), comet goldfish (*C. auratus auratus*) and koi fish (*C. carpio koi*) are languid. The slow movement of fish fins makes it easy for female *A. japonicus* to break away. Female *A. japonicus* will escape from the host when oviposition (Kismiyati *et al.*, 2010).

CONCLUSION

Argulus is capable of infecting cyprinid fish with different percentages for each host genus. In carp (*C. carpio*), the

percentage of male *A. japonicus* infects more than female *A. japonicus*. In goldfish (*C. auratus*), male and female *A. japonicus* have the same percentage, while in comet goldfish (*C. auratus auratus*), the percentage of female *A. japonicus* is higher than female *A. japonicus*. Although there is a difference in the percentage of male and female *A. japonicus* in a host, both have the same detrimental effect on the host fish.

ACKNOWLEDGEMENT

Acknowledgments are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged and any potential conflicts of interest are noted.

Conflict of interest: None.

REFERENCES

- Alifuddin, M., Priyono, A., Nurfatimah, A. (2002). Parasites Inventory on Ornamental Fish Transported in Soekarno-Hatta Airport, Cengkareng, Jakarta. *Jurnal Akuakultur Indonesia*. 1: 123-128.
- Badan Pusat Statistik Kabupaten Magelang. (2014). Produksi Ikan Air Tawar Menurut Kecamatan. <http://magelangkab.bps.go.id>.
- Bandilla, M. (2007). Transmission and Host and Mate Location in the Fish Louse *Argulus coregoni* and its Link with Bacterial Disease in Fish, *Jyvaskyla Studies in Biological and Environmental Science*. *Jyvaskyla: University of Jyvaskyla Press*.
- Carella, F. and Sirri, R. (2017). Fish and shellfish pathology. *Frontiers in Marine Science*. 4: 375. <https://doi.org/10.3389/fmars.2017.00375>.
- Das, D.R. and Chandra, K.J. (2018). Seasonal variation of gill, skin, muscle, liver and kidney pathology of mrigal (*Cirrhinus cirrhosus*) in cultural pond fisheries, mymensingh, Bangladesh. *Bangladesh Journal of Veterinary Medicine*. 16(1): 121-129.
- Ebrahimi, M., Nematollahi, A., Samiei, A., Golabi, M. (2018). Ectoparasitism on freshwater fish in West Azerbaijan, Northwest of Iran. *Comparative Clinical Pathology*. 27(2): 353-356.
- Kismiyati, Fatiza, R.N., Kusdarwati, R. (2011). Pengaruh pemberian garam (NaCl) terhadap kerusakan telur *Argulus japonicus* [Effect of salt (NaCl) against damage of *Argulus japonicus* egg]. *Jurnal Ilmiah Perikanan dan Kelautan*. 3(1): 113-116.
- Kismiyati, Iskhaq, N.M., Triastuti, J. (2010). Preference of oviposition object of ectoparasite *A. japonicus*. *Jurnal Ilmiah Perikanan dan Kelautan*. 2: 165-169.
- Kumar, S., Sathish, K.T., Vidya, R., Pandey, P.K. (2017). A prospective of epidemiological intervention in investigation and management of argulosis in aquaculture. *Aquaculture International*. 25(1): 303-325.
- Mikheev, V.N., Pasternak, A.F., Valtonen, E.T. (2015). Behavioural adaptations of argulid parasites (Crustacea: Branchiura) to major challenges in their life cycle. *Parasites and Vectors*. 8(1): p. 394.
- Møller, O.S. (2009). Branchiura (Crustacea)-survey of historical literature and taxonomy. *Arthropod Systematics and Phylogeny*. 67(1): 41-55.

- Picard Sánchez, A., Estensoro, I., Del Pozo, R., Palenzuela, O.R., Piazzon, M.C., Sitjà Bobadilla, A. (2020). Water temperature, time of exposure and population density are key parameters in *Enteromyxum leei* fish to fish experimental transmission. *Journal of Fish Diseases*. 43(4): 491-502.
- Poly, W.J. (2008). Global diversity of fishlice (Crustacea: Branchiura: Argulidae) in freshwater. *Hydrobiologia*. 595(1): 209-212.
- Pramujirini, D. (2016). Predileksi *A. japonicus* Jantan Dan Betina Serta Perubahan Patologi Anatomi Tiga Jenis Ikan Hias Yang Terinfestasi Di Sentra Budidaya Ikan Hias Jawa Timur. Bachelor Thesis. Surabaya: Universitas Airlangga.
- Sahoo, P.K., Mohanty, J., Garnayak, S., Mohanty, B., Kar, B., Prasanth, H., Jena, J. (2013). Estimation of loss due to argulosis in carp culture ponds in India. *Indian Journal of Fisheries*. 60(2): 99-102.
- Santoso, S. (2008). Panduan Lengkap Menguasai SPSS 16. Jakarta: Elex Media Komputindo.
- Soes, D., Walker, P., Kruijt, D. (2010). The Japanese Fish Louse *A. japonicus* new for The Netherlands. *Lauterbornia*. 70: 11-17.
- Tandel, R.S., Chanu, K.V., Bhat, R.A.H., Dash, P., Shah, T.K., Thakuria, D. (2021). Morphometric and molecular identification of *A. japonicus* (Thiele 1900) in vulnerable Himalayan snow trout, *Schizothorax richardsonii* (Gray 1832). *Aquaculture Research*. 52(12): 6770-6778.
- Taylor, N.G.H., Sommerville, C., Wootten, R. (2006). The epidemiology of *Argulus* spp. (Crustacea: Branchiura) infections in Stillwater trout fisheries. *Journal of Fish Diseases*. 29(4): 193-200.
- Wafer, L.N., Whitney, J.C., Jensen, V.B. (2015). Fish lice (*A. japonicus*) in goldfish (*C. auratus*). *Comparative medicine*. American Association for Laboratory Animal Science. 65(2): 93-95.
- Walker, P.D., Russon, I.J., Haond, C., Velde, G.V.D., Wendelaar-Bonga, S.E. (2011). Feeding in adult *Argulus japonicus* Thiele, 1900 (Maxillopoda, Branchiura), an ectoparasite on fish. *Crustaceana*. 84(3): 307-318.
- Wardany, K.H. and Kurniawan, N. (2014). Eksplorasi ektoparasit pada ikan famili cyprinidae di kolam rumah makan wilayah malang raya. *Biotropika: Journal of Tropical Biology*. 2(2): 87-91.
- Wibowo, A.B., Anggoro, S., Yulianto, B. (2015). Status keberlanjutan dimensi ekologi dalam pengembangan kawasan minapolitan berkelanjutan berbasis perikanan budidaya air tawar di kabupaten magelang. *Saintek Perikanan: Indonesian Journal of Fisheries Science and Technology*. 10(2): 107-113.
- Yildiz, K. and Kumantas, A. (2002). *Argulus foliaceus* infection in a goldfish (*C. auratus*). *Israel Journal of Veterinary Medicine*. 57(3): 118-120.