

VOL 96 No. 09 SEPTEMBER, 2019

Print ISSN 0019 - 6479
E - ISSN 0974 - 9365

₹. 80

THE INDIAN VETERINARY JOURNAL

SINCE - 1924

Journal of the
INDIAN VETERINARY ASSOCIATION
ESTD - 1922 Regd. No. Sl. No. 96/1967 (CHENNAI)



No. 11, Pasumpon Muthuramalinga Thevar Salai (Chamiers Road),
Nandanam, Chennai - 600 035, Tamil Nadu, India
Tel. : +91 44 2435 1006
Email : ivj83@yahoo.com
ONLINE : www.ivj.org.in

THE INDIAN VETERINARY JOURNAL
(Official Organ of the Indian Veterinary Association)

EDITORIAL COMMITTEE

Dr A.V. KRISHNAN, Chief Editor

B.V.Sc., M.V.Sc (Path.)

Retd. Additional Director of A.H., TN

Dr S. SUKUMAR, Managing Editor

B.V.Sc., M.V.Sc (Vet. Micro), Ph.D. (Biotech)

Retd. Professor of Biotechnology, TANUVAS, TN

Dr V. Titus George, Editor

B.V.Sc., M.V.Sc., Ph.D. (Patho)

Retd. Prof. of Pathology, TANUVAS

Dr I. Ponnu Pandian, Editor

B.V.Sc., PGDAEM

Dr K. Venukopalan, Editor

B.V.Sc., M.V.Sc., Ph.D. (Poul.)

Retd. Prof. of Poultry Science, TANUVAS

EDITORIAL BOARD

CHAIRMAN

Dr. Chirantan Kadian

B.V.Sc & A.H.

President, Indian Veterinary Association

MEMBERS

Prof. Dr C. Balachandran

M.V.Sc, Ph.D., PGDAJ, PGDEVP, DICVP

FAO Fellow, FIAVP, FNAVS, FASAW

Vice-Chancellor

Tamilnadu Veterinary and Animal Sciences University

Madhavaram Milk Colony, Chennai - 600 051.

Dr. P.K. Panwar

BVSc & AH, MVSc,

Department of Animal Husbandry

Uttar Pradesh

Dr. Subhash Chander Ahlawat

BVSc & AH.

Department of Animal Husbandry

Haryana

**MANAGING COMMITTEE OF
INDIAN VETERINARY ASSOCIATION**

President :

Dr. Chirantan Kadian

Secretary General :

Dr. R.S. Patel

Treasurer :

Dr. I. Ponnu Pandian

Joint Secretary :

Dr. M.K. Srikanth

Vice Presidents :

Dr. L. Chandrasekaran, HQ, Tamilnadu

Dr. Arun Sirkeck, Himachal Pradesh

Dr. S.G. Yalagod, Karnataka

Dr. Ilang, Nagaland

Dr. Ramji Lal Meena, Rajasthan

Dr. Sudhir Kumar, Uttar Pradesh

Zonal Secretaries :

Dr. Amit Nain, Punjab

Dr. R. Vasantha Rayalu, Andhra Pradesh

Dr. Bhuban Ch. Sarmah, Assam

Dr. Umesh Balaji Sontakke, Maharashtra

Dr. Pankaj Kumar, Jharkhand

Managing Editor, IVJ :

Dr. S. Sukumar, Tamil Nadu

THE INDIAN VETERINARY JOURNAL

(Official organ of the Indian Veterinary Association)

Vol. 96

September 2019

No. 09

CONTENTS

GENERAL ARTICLES :

Study on Morphological Traits in Indigenous <i>Jallikattu</i> Bulls of Tamil Nadu R. Priyadharsini, A. Gopinathan, S.M.K. Karthickeyan and P.N. Richard Jagatheesan	... 09
Epitopes Prediction According to Glycoprotein Encoding Gene of Rabies Virus Local Isolates as Vaccine Candidate against Circulating Rabies Virus in Indonesia JolaRahmahani, Suwarno and Fedik Abdul Rantam	... 14
Estimates of Genetic Parameters of Semen Traits in Crossbred Bulls Gopinathan, S.N. Sivaselvam, S.M.K. Karthickeyan and R. Venkataramanan	... 17
Commercial Cuts in Etawa Goat Fed with Water Spinach Pufa and Mineral Mixture Licks Kadek Rachmawati, Romziah Sidik, Vivi Virgianty, Koesnoto Supranianondo, Tri Bhawono Dadi and Herinda Pertiwi	... 20
Occurance of Ectoparasites in Mud Crab (<i>Scylla serrata</i>) and White Shrimp (<i>Litopenaeus vannamei</i>) Widyo Witulur, Kismiyati and Putri Desi Wulansari	... 23
Antiviral Activity of Ethanolic Extract of Srikaya Seeds (<i>Annona squamosa</i> L.) Against Avian Influenza Virus Maya Nurwartanti Yunita, Adi Prijo Raharjo, Prima Ayu Wibawati and Bodhi Agustono	... 26
Identification of <i>Aeromonas hydrophila</i> Outer Membrane Protein as a Potential Vaccine Candidate for Ulcer Disease M. Gandul Atik Yuliani, Anwar Ma'ruf, Ratna Damayanti and Diyantoro	... 29
Diatom Naviculoid and Nitzschioid Composition in the Sulfur Mountains Watershed East Java, Indonesia Suciyono, A.A. Sabil, E. D. Masithah, D.D. Nindarwi, M.H. Azhar and M.F.Ulkhaq	... 31
Anti Early Embryonic Protein (EEP) for Pregnancy Test by Microtiter Strip in Dairy Cows Abdul Samik and Erma Safitri	... 37
Enhancement of Broiler Chicken Growth by Laserpuncture Treatment Lilian Soekwanto, Chairul Anwar and Rimayanti	... 40
Protective Effect of Mycotoxin Binders on Ovarian Gestation Mice Exposed by Zearalenone Ragil Angga Prastiya, Abdul Samik, M. Thohawi Elziyad Purnama and Amung Logam Saputro	... 44
Detection of Antibiotic Residue in Broiler Chicken Meat at Traditional Market in Surabaya City, Indonesia Shelly Wulandari, Diyantoro and OkySetyo Widodo	... 47
The Potency of <i>Hylocereus polyrhizus</i> Peel Extract as Protector on Lead Acetate-Induced Testicular Toxicity in Mice E. Wulandari, R. I'tishom R and S.A. Sudjarwo	... 49
Probiotic Utilization in Megacolon Dog : A Case Report Tri BhawonoDadi, Ira Sari Yudaniayanti, WiwikMisacoYuniarti, NusdiantoTriakoso, BoediSetiawan, E. Djoko Poetranto, MiyayuSonetaSofyan and Herinda Pertiwi	... 51
Honey as an Alternative to Stem Cells Therapy for Degenerated Rat Testis Due to Malnutrition Erma Safitri, Thomas V. Widiyatno, Willy Sandhika and R. Heru Prasetyo	... 53

Occurance of Ectoparasites in Mud Crab (*Scylla serrata*) and White Shrimp (*Litopenaeus vannamei*)

Widyo Witular, Kismiyati and Putri Desi Wulansari¹

Department of Fish Health Management and Aquaculture, Fisheries and Marine Faculty, Universitas Airlangga, Indonesia 60115

(Received : March, 2019 88/19 Accepted : May, 2019)

Abstract

The aim of this study was to compare the prevalence and intensity of ectoparasites which infest mud crab (*Scylla serrata*) and white shrimp (*Litopenaeus vannamei*). The prevalence of ectoparasite infestation in mud crab was 92.5% with an intensity of 801.18 per sample while the infestation in white shrimp showed 62% with the intensity of 19.98 per sample. It could be concluded that the ectoparasites were found in mud crab more frequently than in white shrimp.

Key words: ectoparasites, mud crab, white shrimp

Mud crab (*Scylla serrata*) and white shrimp (*Litopenaeus vannamei*) are becoming a commodity that featured in the export field (Aji and Supriono, 2018; Hardiyanti *et al.*, 2018). One agent of disease-causing pathogens is ectoparasite. It can attack both mud crab (*S. serrata*) and white shrimp (*L. vannamei*) (Irvansyah *et al.*, 2012; Novita *et al.*, 2016). The aim of this study was to compare the prevalence and intensity of ectoparasites which infest these two hosts.




Fig 1. *Octolasmis* infested the mud crab's gill (40x)
1. Capitulum 2. Scutum 3. Tergum 4. Penducle

Materials and Methods

Materials used in this study were samples of mud crab and white shrimp. A total of 80 mud crabs having 250 grams weight and 15cm width were taken from mud crab's enlargement pond in the District of Ujung Pangkah, Gresik, Indonesia. 150 white shrimps as samples were obtained from Tanggapan village, District of Ujung Pangkah, Gresik, Indonesia with 10-12 cm in size.

Methods used to diagnosis the ectoparasite were native methods (Lom and Dikova, 1992). Ectoparasites were observed on the surface of the body, swimming legs, and gills of mud crab and white shrimp.

The calculation for prevalence and intensity was done as per the fomula mentioned below :

$$\text{Prevalence} : \frac{\text{Number of samples infected}}{\text{Number of samples examined}} \times 100\%$$

$$\text{Intensity} : \frac{\text{Number of parasites found}}{\text{Number of samples infected}}$$




Fig 2. *Zoothamnium* infested white shrimp (*Litopenaeus vannamei*) (400x)

1. The food vacuole; 2. Stalk; 3. Epistomial; 4. Peristomial

¹Corresponding author : Email : putri.dw@fpk.unair.ac.id




Fig 3. *Epistylis* infested the mud crab and white shrimp (400x)
1. The food vacuole; 2. contractile vacuole; 3. Stalk

Results and Discussion

From the observation, ectoparasites such as *Octolasmis* (Fig 1), *Zoothamnium* (Fig 2) and *Epistylis* (Fig 3) were found infested in mud crabs; while *Zoothamnium* and *Epistylis* were found infested in white shrimps.

Based on observations conducted on 80 mud crabs, 74 (92.5%) mud crabs were found infested with ectoparasite, while from 150 white shrimps, 93 (62%) were infested by ectoparasites. The prevalence of ectoparasites is presented in Table I.

Based on the observations conducted, the intensity of ectoparasites in mud crab was 801.18 in number per sample. The intensity of ectoparasites in white shrimp was 19.98 per sample. The intensity of ectoparasites in mud crab and white shrimp is presented in Table II.

The high intensity of ectoparasites infestation in mud crab and white shrimps was caused by the source of water used that comes

from river which flow directly in to the pond without any treatment. The absence of some treatment such as a water filtration system made the entry of the ectoparasites and other pathogenic agents easy in ponds (Nurlaila *et al.*, 2016). Traditional aquaculture systems also did not apply water changes, hence decrease in water quality, would cause rapid development of ectoparasites.

Several studies had been conducted on the occurrence of ectoparasite infection in white shrimp and mud crabs. Farras *et al.*, (2017) found the prevalence of ectoparasites in the intensive-shrimp-culture system was 57.5% and 56.% in the extensive-shrimp-culture system, while Maldini *et al.*, (2017) found Chilodonellosis in the mud crab (*Scylla serrata*). Mushtaq and Mustaquim (2009) found several species of *Octolasmis* on the gills of mud crab and the infestation of *Epistylis* sp, *Zoothamnium* sp, *Vorticella* sp, and *Trichodina* sp. in white shrimp (*Litopenaeus vannamei*) (Putra *et al.*, 2018).

Sessile protozoans, like *Epistylis*, interfered with the gas exchange by blocking respiratory surfaces of the eggs and larvae, especially if found in high numbers (Lavilla and de la Peña. 2004). Epibiosis organism such as *Epistylis* sp. flourishes in low dissolved oxygen conditions. Once infested, cotton wool like growth attaches to the body as well as appendages and disrupt mobility and feeding (Jithendran *et al.*, 2010). However, little is known about the ecological relationship between farmed fish and *Epistylis* sp. This sessile peritrich is often described as an epibiont on aquatic organisms, although studies have shown deleterious effects by the ciliate on their hosts.

Table I. The prevalence of ectoparasites at mud crab and white shrimp

Sample	Number of Samples	Ectoparasites (Phylum)						Total number of samples infested	Prevalence (%)
		Octolasmis		Zoothamnium		Epistylis			
		A	B	A	B	A	B		
Mud Crab	80	71	88.75	50	62.5	69	86.25	74	92.5
White Shrimp	150	-	-	47	31,33	76	50.66	93	62

A: The number of samples infested with ectoparasites B: Prevalence (%)

Table II. The intensity of ectoparasites in mud crab and white shrimp

Sample	Number of Samples	Ectoparasites (Phylum)						Total sample infested	Average Intensity (number per sample)
		Octolasmis		Zoothamnium		Epistylis			
		A	B	A	B	A	B		
Mud Crab	80	71	12.62	50	34.5	69	821.25	74	801.18
White Shrimp	150	-	-	47	15.26	76	15.01	93	19.98

A: Samples of infested ectoparasites B: Intensity of ectoparasites (number per sample)

Epibiosis has a spatially close association between two or more organisms belonging to the same or different species. Through direct and indirect interactions, this association has major effects on the species involved and on community dynamics (Wahl and Mark, 1999). Epibiotic associations between crustaceans and ciliates are very common, since the calcified surface of the crustaceans functions as a semi-permanent substrate, providing an optimal habitat for epibionts ciliates, especially in those areas where other substrates are not suitable for long-term colonization (Bozkurt and Genc, 2009). This epibiotic associations also occurs between decapod crustaceans and stalked barnacle *Octolasmis*, where this interaction does not seem to be species-specific (Machado *et al.*, 2013). Many cases of *Octolasmis* infestation have been reported in parasites infested the mud crabs (*S. serrata*), sand lobster, bamboo lobster, *Portunus sanguinolentus* and several species of crabs (Herlinawati *et al.*, *loc. cit*; Yusgita *et al.*, 2019; Li *et al.*, 2015; Machado *et al.*, *loc. cit*).

Infestation of *Zoothamnium* also forms the epibiotic association. *Zoothamnium* sp. might have contributed towards respiratory difficulties, relative weakness and greater susceptibility of prawns to cannibalism, leading to early death also inhibits egg hatching of *Penaeus monodon* (Mustafa *et al.*, 1997; Tompo and Idris, 2016)

References

Aji, J. M. M., and Supriono, A. (2018) Analisis Komparatif Usaha Budidaya Udang Vaname Tambak Tradisional dengan Tambak Intensif di Kabupaten Situbondo. *JEPA*, **2**(4): 255-266.

Bozkurt, A. and E. Genc. (2009) First Record of the Epibiont

Protozoan *Epistylis* sp. on Zooplankton (Copepoda, Cladocera, and Rotifera) from the Asi River, Turkey. *Crustaceana*, **82**(5): 515-530.

Farras, A., Mahasri, G. and Hari Suprpto. (2017) Prevalensi Dan Derajat Infestasi Ektoparasit Pada Udang Vaname (*Litopenaeus Vannamei*) Di Tambak Intensif Dan Tradisional Di Kabupaten Gresik. *JIPK*, **9**(2) : 118-126.

Hardiyanti, A. S., Sunaryo, S., Riniatsih, I., and Santoso, A. (2018) Biomorfometrik Kepiting Bakau (*Scylla* sp.) Hasil Tangkapan di Perairan Semarang Guna Menunjang Konservasi Sumberdaya Hayati. *BULOMA*, **7**(2): 81-90.

Herlinawati, A., Sarjito, A.H.C and Haditomo. (2017) Infestasi *Octolasmis* pada Kepiting Bakau (*Scylla serrata*) Hasil Budi daya dari Desa Surodadi, Kabupaten Demak, Jawa Tengah. *JamTech*, **6**(4) : 11-19.

Irvansyah, M. Y, Abdulgani, N., and Mahasri, G. (2012) Identification and Intensity of Ectoparasites on Mud Crab (*Scylla serrata*) Juvenile Crab in Crab Pond Sedati Subdistrict, Sidoarjo District. *Sains_Seni*, **1**(1): 5-9.

Jithendran, K.P., M. Poornima, C. P. Balasubramanian and S. Kulasekarapandian. (2010) Disease of Mud Crabs (*Scylla* Spp.): An Overview. *Indian J. Fish.*, **57**(3) : 55-63]

Lavilla-Pitogo CR and de la Peña LD. (2004) Diseases in Farmed Mud Crabs *Scylla* spp.: Diagnosis, Prevention, and Control. SEAFDEC Aquaculture Department, Iloilo, Philippines, p 89.

Li, H., L.Ma, X. Yu, L.Li, C. Yang, and Y.Yan. (2015) Colonization of *Octolasmis* (Cirripedia) on The Crab *Portunus sanguinolentus* (Brachyura : Portunidae) : Impacts of The Parasitism of *Diplothyacus sinensis* (Cirripedia : Rhizocephala). *J Crustacean Biol.*, **35**(2) : 159-165.

Lom, J., and Dyková, I. (1992) *Protozoan parasites of fishes*. Elsevier Science Publishers, pp 315.

Machado, G.B. de O., F.H.C. Sanches, M.D. Fortuna and T.M. Costa. (2013) Epibiosis In Decapod Crustaceans By Stalked Barnacle *Octolasmis lowei* (Cirripedia: Poecilasmatidae). *Zoologia*, **30**(3) : 307-311.

Maldini, A., Winaruddin, M. Bakri, M. Hambal, Farida, and R. Daud. (2017) Identification of Ectoparasites in the Mud Crab

Occurance of Ectoparasites in Mud Crab ...

(*Scylla serrata*) from Alue Naga Coastal Waters Banda Aceh. *J.Med.Vet.*, **11**(2) : 114-117.

Mushtaq, S. and J. Mustaqim. (2009) The Occurance and Distribution of Stalked Barnacles of The Genus *Octomasmis* on The Gills of Mud or Mangrove Crab, Genus *Scylla*. *Crustaceana*, **82**(1) : 53-61.

Mustafa, M., M. Zafar and Y.S.A Khan. (1997) Occurance of *Zoothamnium* sp. Infection in *Penaeus Monodon* (Fabricius, 1798) of The Coastal Shrimp Culture Pond of Cox's Bazar Bangladesh. *PJMS*, **6**(1) : 109-114.

Novita, D., Ferasyi, T. R., and Muchlisin, Z. A. (2016) Intensitas dan Prevalensi Ektoparasit pada Udang Pisang (*Penaeus SP.*) dari Tambak Budidaya di Pantai Barat Aceh. *JIM FKP Unsyiah*, **1**(3): 268-279.

Nurlaila, N., Dewiyanti, I., and Wijaya, S. (2016) Identifikasi dan Prevalensi Ektoparasit pada Udang Vannamei (*Litopenaeus Vannamei*) di Kabupaten Aceh Besar. *JIM FKP Unsyiah*, **1**(3): 388-396.

Putra, M.K.P., T. A. Pribadi, and N. Setiati. (2018) Prevalensi

Ektoparasit Udang Vannamei Pada Tambak di Desa Langgenharjo Kabupaten Pati. *Life Science*, **7**(2) : 31-38.

Tompo, A. and A.P.S Idris. (2016) Prevalensi dan Identifikasi Penyebab Penyakit yang Menghambat Penetasan Telur Udang Windu (*Penaeus monodon* Fabr) di Hatcheri Kabupaten Takalar. *Jurnal PTP*, **2**: 129-134.

Wahl, M. and O. Mark. (1999) The Predominantly Facultative Nature of Epibiosis: Experimental And Observational Evidence. *Mar.Ecol.Prog.Ser.*, **187**: 59-66.

Yusgita, L., Kismiyati, S.Subekti, P.D. Wulansari, and M.K. Amin. (2019) Identification and Prevalence of The Ectoparasite Octolasmis in sand lobster (*Panulirus homarus*) and Bamboo Lobster (*Panulirus versicolor*) in Floating Net Cages in Sape, Bima Regency, West Nusa Tenggara Province, Indonesia. *IOP Conference Series : EES*, **236**. doi:10.1088/1755-1315/236/1/012099.

Indian Vet. J., September 2019, 96 (09) : 26 - 29

Antiviral Activity of Ethanolic Extract of Srikaya Seeds (*Annona squamosa* L.) Against Avian Influenza Virus

Maya Nurwartanti Yunita¹, Adi Prijo Raharjo, Prima Ayu Wibawati and Bodhi Agustono

Department of Veterinary Pathology, Faculty of Veterinary Medicine, Airlangga University, Mulyorejo st. Campus C Surabaya, East Java 60286 Indonesia.

(Received : April, 2019 108/19 Accepted : May, 2019)

Abstract

Drug resistance against influenza viruses have brought the attention of public concern. The development of new antiviral was needed. This study aim to determined the antiviral activity of ethanolic extract of srikaya seeds (EESS) on Avian influenza (AI) when inoculated in embryonated chicken eggs by hemagglutination test. Determination of AI virus titers was carried out by chorioallantoic test. The results showed that the percentage of antiviral activity of the extract at 0.35µg/ml was 59.72%, while the extract at 0.15µg/ml and 0.25µg/ml was 33.33% and 43.06%, respectively. In conclusion, 0.35µg/ml EESS is the most effective concentration that

can inhibit the AI virus.

Key words: Antiviral activity, Avian influenza, ethanolic extract, srikaya seeds

Avian Influenza is acute contagious disease and highly infectious in poultry (Sharif *et al.*, 2014). H5N1 strains outbreaks in poultry is reported in 60 countries and also caused 622 human infections (Huang *et al.*, 2013) as in chicken and zebra finch, and this repertoire has been shaped through lineage-specific duplications. Genes that are responsive to influenza A viruses were identified using the lung transcriptomes of control ducks and ones that were infected with either a highly pathogenic (A/duck/Hubei/49/05.

Avian influenza virus can mutate accord-

¹Corresponding author : Email: mayanurwantanti@fkh.unair.ac.id